

Address for correspondence:

Vineet Chopra MD, MSc

2800 Plymouth Road, Building 16, Room 432W

Ann Arbor, MI 48109

vineetc@umich.edu

Conflicts of Interest: None disclosed for all coauthors.

Funding: Dr. Chopra is supported by a career development award (1-K08-HS022835-01) from the Agency of Healthcare Research and Quality (AHRQ).

Accepted Article

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version record](#). Please cite this article as [doi:10.1002/jhm.2586](https://doi.org/10.1002/jhm.2586).

ABSTRACT

BACKGROUND: Although peripherally inserted central catheters (PICCs) are often used in adults hospitalized with pneumonia, patterns related to PICC use in this context are unknown.

METHODS: Premier's Inpatient database was used to identify patients hospitalized with pneumonia between 1 July 2007 and 30 November 2011. PICC placement was identified via billing codes. Generalized estimating equations (GEE) were used to identify factors associated with PICC placement. Hospital risk-standardized rates of PICC insertion were estimated using hierarchical generalized linear models (HGLM).

RESULTS: 545,250 patients (median age 71, range 57-82) were included. A total of 41,849 (7.7%) patients received a PICC during hospitalization (median receipt: hospital day 4). PICC recipients were younger (median age 69), had higher levels of comorbidity (Gagne score median: 4 vs. 2) and were more often diagnosed with healthcare-associated pneumonia (43.1% vs. 29.9%) than those that did not receive PICCs. The three patient variables most associated with PICC receipt included weight loss (OR 2.03, 95% confidence interval [CI]: 1.97-2.10), sepsis on admission (OR 1.80, 95%CI: 1.75-1.85), and ICU-status on hospital day one or two (OR 1.70, 95%CI: 1.64-1.75). Compared to internal medicine, admission by geriatricians and critical care physicians was associated with PICC placement (OR 1.81, 95%CI: 1.62-2.03 and OR 1.14, 95%CI: 1.05-

1.24, respectively). Risk-standardized rates of PICC utilization varied from 0.3% to 41.7%. Nearly 70% of the variability in PICC use could not be explained by available data.

CONCLUSIONS: In adults hospitalized with pneumonia, PICC use appears directed towards those with severe illness and varies substantially between hospitals.

Accepted Article

INTRODUCTION

Pneumonia is the most common cause of unplanned hospitalization in the United States.¹ Despite its clinical toll, the management of this disease has evolved markedly. Expanding vaccination programs, efforts to improve timeliness of antibiotic therapy, and improved processes of care are but a few developments that have improved outcomes for patients afflicted with this illness.^{2, 3}

Use of peripherally inserted central catheters (PICCs) is an example of a modern development in the management of patients with pneumonia.⁴⁻⁷ PICCs provide many of the benefits associated with central venous catheters (CVCs) including reliable venous access for delivery of antibiotics, phlebotomy and invasive hemodynamic monitoring. However, as they are placed in veins of the upper extremity, PICCs bypass insertion risks (e.g., injury to the carotid vessels or pneumothorax) associated with placement of traditional CVCs.⁸ Because they offer durable venous access, PICCs also facilitate care transitions while continuing intravenous antimicrobial therapy in patients with pneumonia.

However, accumulating evidence also suggests that PICCs are associated with important complications, including central line-associated bloodstream infection (CLABSI) and venous thromboembolism (VTE).^{9, 10} Further, knowledge gaps in clinicians regarding indications for appropriate use and management of complications associated with PICCs have been recognized.^{10, 11} These elements are problematic because reports of unjustified and inappropriate PICC use are growing in the literature.^{12, 13} Such concerns have prompted a number of policy

calls to improve PICC use, including Choosing Wisely® recommendations by various professional societies.^{14, 15}

As little is known about the prevalence or patterns of PICC use in adults hospitalized with pneumonia, we conducted a retrospective cohort study using data from a large network of US hospitals.

METHODS

Setting and Participants

We included patients from hospitals that participated in Premier's Inpatient dataset, a large, fee-supported, multi-payer administrative database that has been used extensively in health services research to measure quality of care and comparative effectiveness of interventions.¹⁶ Participating hospitals represent all regions of the United States and include teaching and non-teaching facilities in rural and urban locations. In addition to variables found in the uniform billing form (UB-04), Premier Inpatient also includes a date-stamped list of charges for procedures conducted during hospitalization, such as PICC placement. As PICC-specific data is not available in most nationally representative datasets, Premier offers unique insights into utilization, timing and factors associated with use of PICCs in hospitalized settings.

We included adult patients aged ≥ 18 years who were (a) admitted with a principal diagnosis of pneumonia present on admission, or secondary diagnosis of pneumonia if paired with a principal diagnosis of sepsis, respiratory failure or influenza; (b) received at least one day of antibiotics between 1 July 2007 and 30

November 2011 and (c) underwent chest X-ray or CT at the time of admission. International Classification of Disease-9-CM codes were used for patient selection. Patients that were not admitted (e.g. observation cases), had cystic fibrosis, or marked as pneumonia not present on admission were excluded. For patients who had more than one hospitalization during the study period, a single admission was randomly selected for inclusion.

Patient, Physician and Hospital Data

For all patients, age, gender, marital status, insurance, race and ethnicity were captured. Using software provided by the Healthcare Costs and Utilization Project, we categorized information on 29 comorbid conditions and computed a combined comorbidity score as described by Gagne.¹⁷ Patients were considered to have healthcare-associated pneumonia [HCAP] if they were: (a) admitted from a skilled nursing or a long-term care facility, (b) hospitalized in the previous 90 days, (c) on dialysis, or (d) receiving immunosuppressing medications, (e.g. chemotherapy or steroids equivalent to at least 20 mg of prednisone per day) at the time of admission. Information on specialty of the admitting physician and hospital-characteristics (e.g. size, location, teaching status) were sourced through Premier data.

Receipt of PICCs and Related Therapies

Among eligible adult patients hospitalized with pneumonia, we identified patients who received a PICC at any time during hospitalization via PICC-specific

billing codes. Non-PICC devices (e.g. midlines, Hickman catheters) were not included. For all insertions, we assessed day of PICC placement relative to admission date. Data on type of PICC (e.g., power-injection capable, antibiotic coating) or PICC characteristics (size, number of lumens) were not available. We used billing codes to assess use of invasive or non-invasive ventilation, vasopressors, and administration of pneumonia-specific antibiotics (e.g., beta-lactams, macrolides, fluoroquinolones, etc.). Early exposure was defined when a billing code appeared within two days of hospital admission.

Outcomes of Interest

The primary outcome of interest was receipt of a PICC. Additionally, we assessed factors associated with PICC placement and variation in risk-standardized rates of PICC use between hospitals.

Statistical Analyses

Patient and hospital characteristics were summarized using frequencies for categorical variables and medians with interquartile ranges for continuous variables. We examined association of individual patient and hospital characteristics with use of PICCs using generalized estimating equations (GEE) models with a logit link for categorical variables and identity link for continuous variables, accounting for patient clustering within hospitals (Table 1).

We then developed a multivariable hierarchical generalized linear model (HGLM) for PICC placement with a random effect for hospital. In this model, we

included patient demographics, comorbidities, sepsis on admission, type of pneumonia (e.g., HCAP vs. CAP), admitting physician specialty and indicators for early receipt of specific treatments such as guideline-recommended antibiotics, vasopressors, ventilation (invasive or non-invasive), and pneumatic compression devices for prophylaxis of deep vein thrombosis.

To understand and estimate between-hospital variation in PICC use, we calculated risk-standardized rates of PICC use (RSPICC) across hospitals using HGLM methods. These methods are also employed by the Centers for Medicare and Medicaid to calculate risk-standardized measures for public reporting.¹⁸

Because hospital rates of PICC use were highly skewed (21.2% [n=105] hospitals had no patients with PICCs), we restricted this model to the 343 hospitals that had at least 5 patients with a PICC in order to obtain stable estimates. For each hospital, we estimated a predicted rate of PICC use (pPICC) as the sum of predicted probabilities of PICC receipt from patient factors and the random intercept for hospital in which they were admitted. We then calculated an expected rate of PICC use (ePICC) per hospital as the sum of expected probabilities of PICC receipt from patient factors only. RSPICC for each hospital was then computed as the product of the overall unadjusted mean PICC rate (uPICC) from all patients and the ratio of the predicted to expected PICC rate [uPICC*(pPICC/ePICC)].¹⁹ Kruskal-Wallis tests were used to evaluate the association between hospital characteristics with RSPICC rates. To evaluate the impact of the hospital in variation in PICC use, we assessed the change in likelihood ratio of a hierarchical model with hospital random effects compared to

a logistic regression model with patient factors only. In addition, we estimated the intra-class correlation (ICC) to assess the proportion of variation in PICC use associated with the hospital, and the median odds ratio (MOR) from the hierarchical model. The MOR is the median of a set of odds ratios comparing 2 patients with the same set of characteristics treated at 2 randomly selected hospitals.²⁰⁻²² All analyses were performed using the Statistical Analysis System (version 9.3, SAS Institute, Inc., Cary, NC) and STATA 13 (StataCorp. Inc., College Station, TX).

Ethical and Regulatory Oversight

Permission to conduct this study was obtained from the institutional review board at Baystate Medical Center, Springfield, MA. The study did not qualify as human subjects research and made use of fully de-identified data.

RESULTS

Between July 2007 and November 2011, 634,285 admissions representing 545,250 unique patients from 495 hospitals met eligibility criteria and were included in the study (**Figure 1**). Included patients had a median age of 71 years (inter-quartile range [IQR] 57-82) and 53.0% were female. Most patients were Caucasian (69.2%), unmarried (51.6%) and insured by Medicare (67.9%). Patients were admitted to the hospital by internal medicine providers (43.4%) hospitalists (21.4%), and family practice providers (14.7%); notably, critical care and pulmonary medicine providers admitted 6.5% of patients. The median Gagne

comorbidity score was 2 (IQR: 1-5). Hypertension, chronic obstructive pulmonary disease, diabetes and congestive heart failure were among the most common comorbidities observed (**Table 1**).

Among eligible patients, 41,849 (7.7%) received a PICC during hospitalization. Approximately a quarter of all patients that received PICCs did so by hospital day 2; 90% underwent insertion by hospital day 11 (mean=5.4 days, median=4 days). Patients who received PICCs were younger (median, IQR: 69 years; 57-80 years) but otherwise demographically similar to those that did not receive PICCs (median, IQR: 72 years; 57–82 years). Compared to other specialties, patients admitted by critical care/pulmonary providers were twice as likely to receive PICCs (12.3% vs. 6.1%, $p<0.001$). Patients that received PICCs had higher comorbidity scores than those that did not (median Gagne comorbidity score 4 vs. 2, $p<0.001$) and were more likely to be diagnosed with HCAP (43.1% vs. 29.9%, $p<0.001$) than CAP (56.9% vs. 70.1%, $p<0.001$).

PICC recipients were also more likely to receive intensive care unit (ICU) level of care (41.4% vs. 16%, $p<0.001$) and both non-invasive (17.5% vs. 8.1%, $p<0.001$) and invasive ventilation (28.6% vs. 8.8%, $p<0.001$) upon admission. Vasopressor use was also significantly more frequent in patients who received PICCs (24.7% vs. 7.6%, $p<0.001$) compared to those that did not receive these devices. Patients with PICCs were more often discharged to skilled nursing facilities (34.5% vs. 19.3%) than those without PICCs.

Characteristics Associated With PICC Use Following Multivariable Modeling

Using HGLM with a random hospital effect, multiple patient characteristics were associated with PICC use (**Table 2**). Patients ≥ 65 years of age were less likely to receive a PICC compared to younger patients (OR 0.81, 95%CI 0.79 – 0.84). Weight loss (OR 2.03, 95% confidence interval [CI]: 1.97-2.10), sepsis on admission (OR 1.80, 95%CI: 1.75-1.85), and ICU-status on hospital day one or two (OR 1.70, 95%CI: 1.64-1.75) represented three factors most strongly associated with PICC use.

Therapy with potent parenteral antimicrobials including anti-MRSA agents (OR 1.72, 95%CI 1.67-1.76), antipseudomonal beta-lactamases (OR 1.27, 95%CI 1.23-1.31) and carbapenems (OR=1.37, 95%CI 1.31-1.44) were significantly associated with PICC use. Conversely, use of macrolides (OR=0.85, 95%CI=0.82-0.88) or respiratory fluoroquinolones (OR=0.90, 95%CI=0.87-0.92) were associated with lower likelihood of PICC use. After adjusting for antimicrobial therapy, HCAP was only slightly more likely to result in PICC use than CAP (OR 1.03, 95%CI 1.01-1.06). Compared to internal medicine providers, admission by geriatricians and critical care/pulmonary specialists was associated with greater likelihood of PICC use (OR 1.85, 95%CI=1.66-2.05 and OR 1.18, 95%CI=1.13-1.24, respectively). Admission by hospitalists was associated with a modestly lower likelihood of PICC placement (OR=0.94, 95%CI=0.91-0.98).

Hospital Level Variation in PICC Use

To ensure stable estimates of hospital PICC use, we excluded 152 facilities (31%): 10% had no patients with PICCs and 21% had < 5 patients that

received a PICC. Therefore, risk-standardized rates of PICC (RS-PICC) use were estimated for 343 of 495 facilities (69%) (**Figure 2**). In these facilities, risk-standardized rates of PICC use varied from 0.3% to 41.7%. Hospital risk-standardized rate of PICC use was significantly associated with hospital location (median 11.9% vs. 7.8% for urban vs. rural hospitals respectively, $p=0.05$). RS-PICC rates were also greater among hospitals in Southern (11.3%), Western (12.7%), and Midwest (12.0%) regions of the nation compared to those in the Northeast (8.4%) ($p=0.02$) (**Table 3**).

A likelihood ratio test comparing the hierarchical model to a logistic model with patient factors only, was highly significant ($p<0.001$) indicating that the hospital where the patient was treated had a major impact on receipt of PICC after accounting for patient factors. The MOR=2.71 which is a larger effect than we found for any of the individual patient characteristics. The proportion of variance explained by hospitals was 25% (95% CI: 22% - 28%), as measured by the ICC.

DISCUSSION

In this study of 545,250 adults hospitalized with pneumonia, we found that approximately 8% of patients received a PICC. Patients who received PICCs had more comorbidities, were more frequently diagnosed with HCAP and were more often admitted to the ICU -- where they experienced greater rates of mechanical ventilation, non-invasive ventilation, and vasopressor use compared to those that did not receive PICCs. Additionally, risk-adjusted rates of PICC use varied as

much as 10-fold across institutions. In fact, almost 70% of the total variation in rates of PICC use remained unexplained by hospital or patient characteristics. While use of PICCs is often clinically nuanced in ways that are difficult to capture in large datasets (e.g. difficult venous access or inability to tolerate oral medications) the substantial variation of PICC use observed suggests that physician and institutional practice styles are the major determinants of PICC placement during a hospitalization for pneumonia. Since PICCs are associated with serious complications and evidence regarding discretionary use is accumulating, a research agenda examining reasons for such use and related outcomes appears necessary.

The placement of PICCs has grown substantially in hospitalized patients all over the world.^{23, 24} Although originally developed for total parenteral nutrition in surgical patients,²⁵ contemporary reports of PICC use in critical illness,²⁶ diseases such as cystic fibrosis,²⁷ and even pregnancy²⁸ are now common. While PICCs are clinically invaluable in many of these conditions, growing use of these devices has led to the realization that benefits may be offset by complications.^{9, 10, 29, 30} Additionally, recent data suggests that not all PICCs may be used for appropriate reasons. For instance, in a decade-long study at a tertiary care center, changes in patterns of PICC use including shortened dwell times, multiple insertions in a single patient, and unclear indications for use were reported.¹¹ In another study at an academic medical center, a substantial proportion of PICCs were found to be "idle" or unjustified.¹² It comes as little surprise, then, that a recent multi-center study found that one out of every five clinicians did not even

know that their patient had a PICC.²⁹ While calls to improve PICC use in the hospital setting have emerged, strategies to do so are limited by data that emanates from single-center reports or retrospective designs. Indeed, no other studies reporting use of PICCs across US hospitals for any clinical condition currently exist.³¹

We found that patients with weight loss, those with greater combined comorbidity scores and those that were critically ill or diagnosed with sepsis were more likely to receive PICCs than others. These observations suggest that PICC use may reflect underlying severity of illness, as advanced care such as ventilator support was often associated with PICC use. Additionally, discharge to a skilled nursing facility was frequently associated with PICC placement, a finding consistent with a recent study evaluating the use of PICCs in these settings.³² However, a substantial proportion of PICC use remained unexplained by available patient- or hospital factors. Although our study was not specifically designed to examine this question, one possible reason may relate to unmeasured institutional factors that influence the propensity to use a PICC, recently termed as "PICC culture."³³ For example, it is plausible that hospitals with nursing-led PICC teams or interventional radiology (such as teaching hospitals) are more likely to use PICCs than those without such operators. This hypothesis may explain why urban-, larger-, and teaching hospitals exhibited higher rates of PICC use. Conversely, providers may have an affinity towards PICC use that is predicated not just by operator availability, but also local hospital norms. Understanding why some facilities use PICCs at higher rates

than others and implications of such variation with respect to patient safety, cost and outcomes is important. Study designs that use mixed-methods approaches or seek to qualitatively understand reasons behind PICC use are likely to be valuable in this enquiry.

Our study has limitations. First, we used an administrative dataset and ICD-9 codes rather than clinical data from medical records to identify cases of pneumonia or comorbidities. Our estimates of PICC use across hospitals thus may not fully account for differences in severity of illness and it is possible that patients needed a PICC for reasons that we could not observe. However, the substantial variation observed in rates of PICC use across hospitals is unlikely to be explained by differences in patient severity of illness, documentation, or coding practices. Second, as PICC removal codes were not available, we are unable to comment on how often hospitalized pneumonia patients were discharged with PICCs or received antimicrobial therapy beyond their inpatient stay. Third, although we observed a number of patient- and hospital factors were associated with PICC receipt, our study was not designed to determine the reasons underlying these patterns.

These limitations aside, our study has important strengths. To our knowledge, this is the first study to report utilization and outcomes associated with PICC use among those hospitalized with pneumonia across the US. The inclusion of a large number of patients receiving care in diverse facilities lends a high degree of external validity to our findings. Second, we used advanced modeling to identify factors associated with PICC use in hospitalized patients

with pneumonia, producing innovative and novel findings. Third, our study is the first to show the existence of substantial variation in rates of PICC use across US hospitals within the single disease state of pneumonia. Understanding the drivers of this variability is important as it may inform future studies, policies and practices to improve PICC use in hospitalized patients.

In conclusion, we found that PICC use in patients hospitalized with pneumonia is common and highly variable. Future studies examining the contextual factors behind PICC use and their association with outcomes are needed to facilitate efforts to standardize PICC use across hospitals.

Accepted Article

REFERENCES

1. Elixhauser A OP. Reasons for Being Admitted to the Hospital through the Emergency Department, 2003. Healthcare Cost and Utilization Project Statistical Brief 2. Rockville, MD: Agency for Healthcare Research and Quality. . 2006;Accessed at www.hcup-us.ahrq.gov/reports/statbriefs/sb2.pdf on 27 June 2014.
2. Suter LG, Li SX, Grady JN, et al. National Patterns of Risk-Standardized Mortality and Readmission After Hospitalization for Acute Myocardial Infarction, Heart Failure, and Pneumonia: Update on Publicly Reported Outcomes Measures Based on the 2013 Release. *Journal of general internal medicine*. May 14 2014.
3. Lee JS, Nsa W, Hausmann LR, et al. Quality of care for elderly patients hospitalized for pneumonia in the United States, 2006 to 2010. *JAMA internal medicine*. Nov 2014;174(11):1806-1814.
4. Masoorli S, Angeles T. PICC lines: the latest home care challenge. *Rn*. Jan 1990;53(1):44-51.
5. Lam S, Scannell R, Roessler D, Smith MA. Peripherally inserted central catheters in an acute-care hospital. *Archives of internal medicine*. Aug 22 1994;154(16):1833-1837.
6. Goodwin ML, Carlson I. The peripherally inserted central catheter: a retrospective look at three years of insertions. *Journal of intravenous*

nursing : the official publication of the Intravenous Nurses Society. Mar-Apr 1993;16(2):92-103.

7. Ng PK, Ault MJ, Ellrodt AG, Maldonado L. Peripherally inserted central catheters in general medicine. *Mayo Clinic proceedings*. Mar 1997;72(3):225-233.
8. Funk D, Gray J, Plourde PJ. Two-year trends of peripherally inserted central catheter-line complications at a tertiary-care hospital: role of nursing expertise. *Infection control and hospital epidemiology : the official journal of the Society of Hospital Epidemiologists of America*. Jun 2001;22(6):377-379.
9. Chopra V, Ratz D, Kuhn L, Lopus T, Chenoweth C, Krein S. PICC-associated bloodstream infections: prevalence, patterns, and predictors. *The American journal of medicine*. Apr 2014;127(4):319-328.
10. Chopra V, O'Horo JC, Rogers MA, Maki DG, Safdar N. The risk of bloodstream infection associated with peripherally inserted central catheters compared with central venous catheters in adults: a systematic review and meta-analysis. *Infection control and hospital epidemiology : the official journal of the Society of Hospital Epidemiologists of America*. Sep 2013;34(9):908-918.
11. Gibson C, Connolly BL, Moineddin R, Mahant S, Filipescu D, Amaral JG. Peripherally inserted central catheters: use at a tertiary care pediatric center. *Journal of vascular and interventional radiology : JVIR*. Sep 2013;24(9):1323-1331.

12. Tejedor SC, Tong D, Stein J, et al. Temporary central venous catheter utilization patterns in a large tertiary care center: tracking the "idle central venous catheter". *Infection control and hospital epidemiology : the official journal of the Society of Hospital Epidemiologists of America*. Jan 2012;33(1):50-57.
13. Tiwari MM, Hermsen ED, Charlton ME, Anderson JR, Rupp ME. Inappropriate intravascular device use: a prospective study. *The Journal of hospital infection*. Jun 2011;78(2):128-132.
14. McMahon LF, Jr., Beyth RJ, Burger A, et al. Enhancing patient-centered care: SGIM and choosing wisely. *Journal of general internal medicine*. Mar 2014;29(3):432-433.
15. Williams AW, Dwyer AC, Eddy AA, et al. Critical and honest conversations: the evidence behind the "Choosing Wisely" campaign recommendations by the American Society of Nephrology. *Clinical journal of the American Society of Nephrology : CJASN*. Oct 2012;7(10):1664-1672.
16. Rothberg MB, Pekow PS, Priya A, et al. Using highly detailed administrative data to predict pneumonia mortality. *PloS one*. 2014;9(1):e87382.
17. Gagne JJ, Glynn RJ, Avorn J, Levin R, Schneeweiss S. A combined comorbidity score predicted mortality in elderly patients better than existing scores. *Journal of clinical epidemiology*. Jul 2011;64(7):749-759.

18. Sjoding MW, Prescott HC, Wunsch H, Iwashyna TJ, Cooke CR. Hospitals with the highest intensive care utilization provide lower quality pneumonia care to the elderly. *Critical care medicine*. Jun 2015;43(6):1178-1186.
19. Normand SL, Shahian DM. Statistical and Clinical Aspects of Hospital Outcomes Profiling. *Statistical Science*. 2007;22(2):206-226.
20. Larsen K, Merlo J. Appropriate assessment of neighborhood effects on individual health: integrating random and fixed effects in multilevel logistic regression. *American journal of epidemiology*. Jan 1 2005;161(1):81-88.
21. Larsen K, Petersen JH, Budtz-Jorgensen E, Endahl L. Interpreting parameters in the logistic regression model with random effects. *Biometrics*. Sep 2000;56(3):909-914.
22. Sanagou M, Wolfe R, Forbes A, Reid CM. Hospital-level associations with 30-day patient mortality after cardiac surgery: a tutorial on the application and interpretation of marginal and multilevel logistic regression. *BMC medical research methodology*. 2012;12:28.
23. Lisova K, Paulinova V, Zemanova K, Hromadkova J. Experiences of the first PICC team in the Czech Republic. *British journal of nursing*. Jan 2015;24 Suppl 2:S4-S10.
24. Konstantinou EA, Stafylarakis E, Kapritsou M, et al. Greece reports prototype intervention with first peripherally inserted central catheter: case report and literature review. *Journal of vascular nursing : official publication of the Society for Peripheral Vascular Nursing*. Sep 2012;30(3):88-93.

25. Hoshal VL, Jr. Total intravenous nutrition with peripherally inserted silicone elastomer central venous catheters. *Archives of surgery*. May 1975;110(5):644-646.
26. Cotogni P, Pittiruti M. Focus on peripherally inserted central catheters in critically ill patients. *World journal of critical care medicine*. Nov 4 2014;3(4):80-94.
27. Mermis JD, Strom JC, Greenwood JP, et al. Quality improvement initiative to reduce deep vein thrombosis associated with peripherally inserted central catheters in adults with cystic fibrosis. *Annals of the American Thoracic Society*. Nov 2014;11(9):1404-1410.
28. Cape AV, Mogensen KM, Robinson MK, Carusi DA. Peripherally Inserted Central Catheter (PICC) Complications During Pregnancy. *JPEN. Journal of parenteral and enteral nutrition*. May 28 2013;38(5):595-601.
29. Chopra V, Govindan S, Kuhn L, et al. Do clinicians know which of their patients have central venous catheters?: a multicenter observational study. *Annals of internal medicine*. Oct 21 2014;161(8):562-567.
30. Chopra V, Anand S, Hickner A, et al. Risk of venous thromboembolism associated with peripherally inserted central catheters: a systematic review and meta-analysis. *Lancet*. Jul 27 2013;382(9889):311-325.
31. Chopra V, Flanders SA, Saint S. The problem with peripherally inserted central catheters. *Jama*. Oct 17 2012;308(15):1527-1528.

32. Chopra V, Montoya A, Joshi D, et al. Peripherally Inserted Central Catheter Use in Skilled Nursing Facilities: A Pilot Study. *Journal of the American Geriatrics Society*. Sep 2015;63(9):1894-1899.
33. McGill RL, Tsukahara T, Bhardwaj R, Kapetanios AT, Marcus RJ. Inpatient venous access practices: PICC culture and the kidney patient. *The journal of vascular access*. Jan 27 2015;0(0):0.

Accepted Article

TABLE 1: CHARACTERISTICS OF THE STUDY POPULATION

Characteristic	Total N (%)	No PICC N (%)	PICC N (%)	p value*
	545,250 (100)	503,401 (92.3)	41,849 (7.7)	
Demographics				
Age, Median (Q1-Q3) (years)	71 (57 - 82)	72 (57 - 82)	69 (57 - 80)	<0.001
Gender				<0.001
Male	256448 (47.0)	237232 (47.1)	19216 (45.9)	
Female	288802 (53.0)	266169 (52.9)	22633 (54.1)	
Race/Ethnicity				<0.001
White	377255 (69.2)	346689 (68.9)	30566 (73.0)	
Black	63345 (11.6)	58407 (11.6)	4938 (11.8)	
Hispanic	22855 (4.2)	21716 (4.3)	1139 (2.7)	
Other	81795 (15.0)	76589 (15.2)	5206 (12.4)	
Admitting Specialty				<0.001
Internal Medicine	236859 (43.4)	218689 (43.4)	18170 (43.4)	
Hospital Medicine	116499 (21.4)	107671 (21.4)	8828 (21.1)	
Family Practice	80388 (14.7)	75482 (15.0)	4906 (11.7)	
Critical care and Pulmonary	35670 (6.5)	30529 (6.1)	41849 (12.3)	
Geriatrics	4812 (0.9)	4098 (0.8)	714 (1.7)	
Other	71022 (13.0)	66932 (13.3)	4090 (9.8)	
Insurance				<0.001
Medicare	370303 (67.9)	341379 (67.8)	28924 (69.1)	
Medicaid	45505 (8.3)	41100 (8.2)	4405 (10.5)	
Managed care	69984 (12.8)	65280 (13.0)	4704 (11.2)	
Commercial-Indemnity	20672 (3.8)	19251 (3.8)	1421 (3.4)	
Other	38786 (7.1)	36391 (7.2)	2395 (5.7)	
Comorbidities				
Gagne combined comorbidity score, Median(Q1-Q3)	2 (1 - 5)	2 (1 - 4)	4 (2 - 6)	<0.001
Hypertension	332347 (60.9)	306964 (61.0)	25383 (60.7)	0.13
Chronic pulmonary disease	255403 (46.8)	234619 (46.6)	20784 (49.7)	<0.001
Diabetes	171247 (31.4)	155540 (30.9)	15707 (37.5)	<0.001
Congestive heart failure	146492 (26.9)	131041 (26.0)	15451 (36.9)	<0.001
Atrial Fibrillation	108405 (19.9)	97124 (19.3)	11281 (27.0)	<0.001
Renal failure	104404 (19.1)	94277 (18.7)	10127 (24.2)	<0.001
Nicotine replacement therapy/Tobacco use	89938 (16.5)	83247 (16.5)	6691 (16.0)	<0.001

Obesity	60242 (11.0)	53268 (10.6)	6974 (16.7)	<0.001
Coagulopathy	41717 (7.6)	35371 (7.0)	6346 (15.2)	<0.001
Prior Stroke (1 year)	26787 (4.9)	24046 (4.78)	2741 (6.55)	<0.001
Metastatic cancer	21868 (4.0)	20244 (4.0)	1624 (3.9)	0.16
Solid tumor w/out metastasis	21083 (3.9)	19380 (3.8)	1703 (4.1)	0.12
Prior VTE (1 year)	19090 (3.5)	16906 (3.4)	2184 (5.2)	<0.001
Chronic Liver disease	16273 (3.0)	14207 (2.8)	2066 (4.9)	<0.001
Prior Bacteremia (1 year)	4106 (0.7)	3584 (0.7)	522 (1.2)	<0.001
Nephrotic syndrome	671 (0.1)	607 (0.1)	64 (0.2)	0.03
Morbidity Markers				
Type of Pneumonia				<0.001
CAP	376370 (69.1)	352900 (70.1)	23830 (56.9)	
HCAP	168520 (30.9)	150501 (29.9)	18019 (43.1)	
Sepsis Present on Admission	114578 (21.0)	96467 (19.2)	18111 (43.3)	<0.001
Non-invasive ventilation	47913(8.8)	40599 (8.1)	7314 (17.5)	<0.001
Invasive Mechanical Ventilation	56179 (10.3)	44228 (8.8)	11951 (28.6)	<0.001
ICU Status	97703 (17.9)	80380 (16.0)	17323 (41.4)	<0.001
Vasopressor Use	48353 (8.9)	38030 (7.6)	10323 (24.7)	<0.001
Antibiotic/Medication Use				
Anti-MRSA Agent (Vancomycin)	146068 (26.8)	123327 (24.5)	22741 (54.3)	<0.001
3 rd Generation Cephalosporin	250782 (46.0)	235556 (46.8)	15226 (36.4)	<0.001
Anti-Pseudomonal Cephalosporin	41798 (7.7)	36982 (7.3)	4816 (11.5)	<0.001
Anti-Pseudomonal β -Lactam	122215 (22.4)	105741 (21.0)	16474 (39.4)	<0.001
Fluroquinolone	288051 (52.8)	267131 (53.1)	20920 (50.0)	<0.001
Macrolide	223737 (41.0)	210954 (41.9)	12783 (30.5)	<0.001
Aminoglycoside	15415 (2.8)	12661 (2.5)	2754 (6.6)	<0.001
Oral steroids	44486 (8.2)	41586 (8.3)	2900 (6.9)	<0.001
Intravenous steroids	146308 (26.8)	133920 (26.6)	12388 (29.6)	<0.001
VTE Prophylaxis with LMWH	190735 (35.0)	174612 (34.7)	16123 (38.5)	0.01
Discharge Disposition				
Home	282146 (51.7)	272604(54.1)	9542 (22.8)	<0.001
Home with Home health	71977 (13.2)	65289 (13.0)	6688 (16.0)	<0.001
Skilled Nursing Facility	111541 (20.5)	97113 (19.3)	14428 (34.5)	<0.001
Hospice	20428 (3.7)	17902 (3.6)	2526 (6.0)	<0.001
Expired	47733 (8.7)	40768 (8.1)	6965 (16.6)	<0.001
Other [^]	11425 (2.1)	9725 (1.9)	1700 (4.1)	<0.001

* p-value from GEE models that account for clustering within hospital; ^ includes: discharged/transferred to cancer center/childrens hospital; discharged/transferred to federal hospital; discharged/transferred to

swing bed; discharged/transferred to long-term care facility; discharged/transferred to psych hospital; discharged/transferred to assisted living; discharged/transferred to other health institution not in list.

Accepted Article

TABLE 2: PATIENT FACTORS ASSOCIATED WITH PICC USE

Patient characteristic	Odds Ratio (OR)	95% Confidence Intervals	
Age group (> 66 vs. ≤ 65 years)	0.82	0.79	0.84
Race/Ethnicity			
Other	1.02	0.97	1.06
Black	0.99	0.95	1.03
Hispanic	0.82	0.76	0.88
White	<i>Referent</i>		
Marital status			
Other/Missing	1.07	1.01	1.14
Single	1.02	1.00	1.05
Married	<i>Referent</i>		
Insurance payor			
Other	0.85	0.80	0.89
Medicaid	1.13	1.08	1.18
Managed care	0.95	0.91	0.99
Commercial-Indemnity	0.93	0.87	1.00
Medicare	<i>Referent</i>		
Admitting physician specialty			
Pulmonary /Critical care medicine	1.18	1.13	1.24
Family practice (FP)	1.01	0.97	1.05
Geriatric medicine (FP&IM)	1.85	1.66	2.05
Hospitalist	0.94	0.91	0.98
Other specialties	1.02	0.97	1.06
Internal medicine (IM)	<i>Referent</i>		
Comorbidities			
Congestive heart failure	1.27	1.24	1.31
Valvular disease	1.11	1.07	1.15
Pulmonary circulation disorders	1.37	1.32	1.42
Peripheral vascular disease	1.09	1.05	1.13
Hypertension	0.94	0.92	0.97
Paralysis	1.59	1.51	1.67
Other neurological disorders	1.20	1.16	1.23
Chronic lung disease	1.10	1.07	1.12
Diabetes	1.13	1.10	1.16
Hypothyroidism	1.03	1.00	1.06
Liver disease	1.16	1.10	1.23
Ulcer	1.86	1.15	3.02
Lymphoma	0.88	0.81	0.96

Metastatic cancer	0.75	0.71	0.80
Solid tumor without metastasis	0.93	0.88	0.98
Arthritis	1.22	1.16	1.28
Obesity	1.47	1.42	1.52
Weight loss	2.03	1.97	2.10
Blood loss	1.69	1.55	1.85
Deficiency anemias	1.40	1.37	1.44
Alcohol abuse	1.19	1.13	1.26
Drug abuse	1.31	1.23	1.39
Psychoses	1.16	1.11	1.21
Depression	1.10	1.06	1.13
Renal failure	0.96	0.93	0.98
Type of Pneumonia			
HCAP	1.03	1.01	1.06
CAP		<i>Referent</i>	
Sepsis (POA)	1.80	1.75	1.85
Antibiotic use			
Anti-MRSA agent	1.72	1.67	1.76
Anti-Pseudomonal Carbapenem	1.37	1.31	1.44
Non-Pseudomonal Carbapenem	1.48	1.33	1.66
Third generation Cephalosporin	1.04	1.01	1.07
Anti-Pseudomonal Cephalosporin	1.25	1.20	1.30
Anti-Pseudomonal Betalactam	1.27	1.23	1.31
Aztreonam	1.31	1.23	1.40
Non-Pseudomonal Betalactam	1.36	1.23	1.50
Beta-lactam	1.55	1.26	1.90
Respiratory quinolone	0.90	0.87	0.92
Macrolide	0.85	0.82	0.88
Doxycycline	0.94	0.87	1.01
Aminoglycoside	1.21	1.14	1.27
Vasopressors	1.06	1.03	1.10
Non-invasive ventilation	1.29	1.25	1.34
Invasive ventilation	1.66	1.61	1.72
Intensive Care Unit on Admission	1.70	1.64	1.75
Atrial Fibrillation	1.26	1.22	1.29
Upper Extremity Chronic DVT	1.61	1.13	2.28
Nicotine replacement therapy/ Tobacco abuse	0.91	0.88	0.94
Aspirin	0.94	0.92	0.97
Warfarin	0.90	0.86	0.94

LMWH - prophylactic dose	1.10	1.08	1.13
LMWH - treatment dose	1.22	1.16	1.29
Intravenous steroids	1.05	1.02	1.08
Bacteremia (prior year)	1.14	1.02	1.27
VTE (prior year)	1.11	1.06	1.18
Pneumatic compression device	1.25	1.08	1.45
Invasive ventilation (prior year)	1.17	1.11	1.24
Irritable bowel disease	1.19	1.05	1.36

KEY: HCAP=healthcare-associated pneumonia; CAP=community-associated pneumonia; POA=present on admission; MRSA=methicillin-resistant Staphylococcus Aureus; LMWH=low molecular weight heparin; VTE=venous thromboembolism

Accepted Article

TABLE 3: ASSOCIATION BETWEEN HOSPITAL CHARACTERISTICS AND RISK-STANDARDIZED RATE OF PICC USE*


Hospital Characteristic (n)	Median (IQR), %	p-value**
Bed Size		0.12
≤ 200 beds (106)	9.1 (4.8 - 16.3)	
≥ 201 beds (237)	11.6 (5.8 - 17.6)	
Rural/Urban		0.05
Urban (275)	11.9 (5.5 - 17.4)	
Rural (68)	7.8 (5.0 - 14.0)	
Region		0.02
Northeast (50)	8.4 (3.9 - 13.0)	
Midwest (69)	12.0 (5.8 - 17.4)	
West (57)	12.7 (7.6 - 17.0)	
South (167)	11.3 (4.8 - 17.8)	
Teaching Status		0.77
Non-teaching (246)	10.9 (5.0 - 17.4)	
Teaching (97)	12.0 (5.8 - 16.9)	

* Numbers indicate the percentage of patients with a PICC in each category, accounting for risk associated with PICC receipt. To ensure stable estimates, 152 facilities (31%) were excluded as 10% had no patients with PICCs and 21% had < 5 patients that received a PICC.

**Kruskal-Wallis test

IQR=inter-quartile range.


Accepted

FIGURE 1: STUDY FLOW DIAGRAM

LEGEND: PN=pneumonia, POA=present on admission; MS=missing; DRG=diagnosis-related group; PICC=peripherally inserted central catheter

Accep

FIGURE 2: OBSERVED VS. RISK-STANDARDIZED RATE OF PICC USE ACROSS 343 US HOSPITALS (RESTRICTED TO SITES WHERE > 5 PATIENTS RECEIVED PICCS)



LEGEND: Horizontal axis represents rate of PICC use whereas vertical axis represents number of hospitals. The dark shaded bars represents the observed rate of PICC use whereas the non-shaded bars reflect risk-standardized rate of PICC use.

Accepted