

Received Date : 02-Dec-2015

Revised Date : 06-Jan-2016

Accepted Date : 12-Jan-2016

Article type : Ordinary Papers

## **Allogeneic Transplantation Provides Durable Remission in a Subset of DLBCL Patients Relapsing after Autologous Transplantation.**

Timothy S. Fenske<sup>1</sup>, Kwang Woo Ahn<sup>2,3</sup>, Tara M. Graff<sup>4</sup>, Alyssa DiGilio<sup>2</sup> Qaiser Bashir<sup>5</sup>, Rammurti T. Kamble<sup>6</sup>, Ernesto Ayala<sup>7</sup>, Ulrike Bacher<sup>8,9</sup>, Jonathan E. Brammer<sup>5</sup>, Mitchell Cairo<sup>10</sup>, Andy Chen<sup>11</sup>, Yi-Bin Chen<sup>12</sup>, Saurabh Chhabra<sup>13</sup>, Anita D'Souza<sup>2</sup>, Umar Farooq<sup>14</sup>, Cesar Freytes<sup>15</sup>, Siddhartha Ganguly<sup>16</sup>, Mark Hertzberg<sup>17</sup>, David Inwards<sup>18</sup>, Samantha Jaglowski<sup>19</sup>, Mohamed A. Kharfan-Dabaja<sup>7</sup>, Hillard M. Lazarus<sup>20</sup>, Sunita Nathan<sup>21</sup>, Attaphol Pawarode<sup>22</sup>, Miguel-Angel Perales<sup>23</sup>, Nishitha Reddy<sup>24</sup>, Sachiko Seo<sup>25</sup>, Anna Sureda<sup>26,27</sup>, Sonali M. Smith<sup>28</sup>, Mehdi Hamadani<sup>1,2</sup>

<sup>1</sup>Froedtert Memorial Lutheran Hospital, Milwaukee, WI; <sup>2</sup>CIBMTR (Center for International Blood and Marrow Transplant Research), Department of Medicine, Medical College of Wisconsin, Milwaukee, WI; <sup>3</sup>Division of Biostatistics, Institute for Health and Society, Medical College of Wisconsin, Milwaukee, WI; <sup>4</sup>Medical Oncology Hematology Associates, Des Moines, IA; <sup>5</sup>Department of Stem Cell Transplantation, Division of Cancer Medicine, The University of Texas MD Anderson Cancer Center, Houston, TX; <sup>6</sup>Division of Hematology and Oncology, Center for Cell and Gene Therapy, Baylor College of Medicine, Houston, TX; <sup>7</sup>Department of Blood and Marrow Transplantation, H. Lee Moffitt Cancer Center and Research Institute, Tampa, FL <sup>8</sup>Department of Haematology/Oncology, University Medicine Goettingen, Germany; <sup>9</sup>Interdisciplinary Clinic for Stem Cell Transplantation, University Cancer

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/bjh.14046](https://doi.org/10.1111/bjh.14046)

This article is protected by copyright. All rights reserved

Centre Hamburg, Hamburg, Germany; <sup>10</sup>Division of Pediatric Hematology, Oncology and Stem Cell Transplantation, Department of Pediatrics, New York Medical College, Valhalla, NY; <sup>11</sup>Center for Hematologic Malignancies, Oregon Health and Science University, Portland, OR; <sup>12</sup>Division of Hematology/Oncology, Massachusetts General Hospital, Boston, MA; <sup>13</sup>Division of Hematology/Oncology, Medical University of South Carolina, Charleston, SC; <sup>14</sup> Department of Oncology and Blood Marrow Transplantation, University of Iowa Hospitals and Clinics, Iowa City, IA; <sup>15</sup>South Texas Veterans Health Care System and University of Texas Health Science Center San Antonio, San Antonio, TX; <sup>16</sup>Blood and Marrow Transplantation, Division of Hematology and Oncology, University of Kansas Medical Center, Kansas City, KS; <sup>17</sup>Department of Haematology, Prince of Wales Hospital, Randwick NSW, Australia; <sup>18</sup>Division of Hematology, Mayo Clinic, Rochester; <sup>19</sup>Division of Hematology, The Ohio State University Medical Center, Columbus, OH; <sup>20</sup>Seidman Cancer Center, University Hospitals Case Medical Center, Cleveland, OH; <sup>21</sup>Department of Hematology, Rush University Medical Center, Chicago, IL; <sup>22</sup>Blood and Marrow Transplantation Program, Division of Hematology/Oncology, Department of Internal Medicine, The University of Michigan Medical School, Ann Arbor, MI; <sup>23</sup>Adult Bone Marrow Transplantation Service, Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY; <sup>24</sup>Division of Hematology/Oncology, Vanderbilt University Medical Center, Nashville, TN; <sup>25</sup>Vaccine and Infectious Diseases Division, Fred Hutchinson Cancer Research Center, Seattle, WA; <sup>26</sup>Servei d'Hematologia, Institut Catala d'Oncologia, Hospital Duran I Reynals, Barcelona, Spain; <sup>27</sup>Secretary, European Group for Blood and Marrow Transplantation; <sup>28</sup>Section of Hematology/Oncology, The University of Chicago, Chicago, IL

**Corresponding author:** Mehdi Hamadani, MD, Center for International Blood and Marrow Transplant Research, Medical College of Wisconsin, 9200 W. Wisconsin Avenue, Suite C5500, Milwaukee, WI 53226, USA; Phone: 414-805-0643; Fax: 414-805-0714; E-mail: [mhamadani@mcw.edu](mailto:mhamadani@mcw.edu)

**Running head:** AlloHCT for DLBCL after AutoHCT failure

**Keywords:** DLBCL, prognostic score, allogeneic transplantation, prior autologous transplan.

**Previous Presentations:** *Results presented in part as an oral presentation at the 2015 Annual Meeting of the American Society of Hematology (Orlando, FL).*

Received 2 December 2015, accepted 12 January 2016

## **ABSTRACT**

For diffuse large B-cell lymphoma (DLBCL) patients progressing after autologous haematopoietic cell transplantation (autoHCT), allogeneic HCT (alloHCT) is often considered, although limited information is available to guide patient selection. Using the Center for International Blood and Marrow Transplant Research (CIBMTR) database, we identified 503 patients who underwent alloHCT after disease progression/relapse following a prior autoHCT. The 3-year probabilities of non-relapse mortality, progression/relapse, progression-free survival (PFS) and overall survival (OS) were 30%, 38%, 31% and 37% respectively. Factors associated with inferior PFS on multivariate analysis included Karnofsky performance status (KPS) <80, chemoresistance, autoHCT to alloHCT interval <1-year and myeloablative conditioning. Factors associated with worse OS on multivariate analysis included KPS<80, chemoresistance and myeloablative conditioning. Three adverse prognostic factors were used to construct a prognostic model for PFS, including KPS<80 (4 points), autoHCT to alloHCT interval <1-year (2 points) and chemoresistant disease at alloHCT (5 points). This CIBMTR prognostic model classified patients into four groups: low-risk (0 points), intermediate-risk (2-5 points), high-risk (6-9 points) or very high-risk (11 points), predicting 3-year PFS of 40%, 32%, 11% and 6%, respectively, with 3-year OS probabilities of 43%, 39%, 19% and 11% respectively. In conclusion, the CIBMTR prognostic model identifies a subgroup of DLBCL patients experiencing long-term survival with alloHCT after a failed prior autoHCT.

## **INTRODUCTION**

Diffuse large B-cell lymphoma (DLBCL) accounts for 30% of non-Hodgkin lymphoma (NHL) cases diagnosed in the United States annually. Following the incorporation of rituximab into treatment regimens, approximately 60% of DLBCL cases are now cured with frontline therapy. Despite overall improvements in the outcomes of DLBCL, about 30-40% of patients

develop relapsed or refractory disease. Autologous haematopoietic cell transplantation (autoHCT) became the standard-of-care for chemosensitive relapsed or refractory DLBCL after the PARMA trial showed a benefit for autoHCT over conventional second-line therapy.(Philip *et al*, 1995) More recently, the CORAL (Collaborative Trial in Relapsed Aggressive Lymphoma) study provided important information regarding outcomes of relapsed or refractory DLBCL in the rituximab-era. In this study, 53% of patients who underwent an autoHCT were event-free at 3-years.(Gisselbrecht *et al*, 2010) Contemporary registry data confirm these observations, reporting 3-year progression-free survival (PFS) rates of 45-50% following autoHCT.(Fenske *et al*, 2009, Mounier *et al*, 2012, Hamadani *et al*, 2014) These data underscore the fact that a significant subset of DLBCL patients who undergo autoHCT will eventually relapse.

The prognosis for patients with recurrent disease following autoHCT is poor, with no consensus on the optimal therapy. There is evidence to support a graft-versus-lymphoma (GVL) effect in DLBCL (Bishop *et al*, 2008, Rezvani *et al*, 2008, Hamadani *et al*, 2013), and an allogeneic (allo-) HCT is generally considered to be the only potentially curative option for DLBCL patients who relapse after an autoHCT.(Hamadani *et al*, 2013, Thomson *et al*, 2009, Sirvent *et al*, 2010, Bacher *et al*, 2012, Klyuchnikov *et al*, 2014) However, the literature is limited regarding the outcomes of alloHCT, specifically in DLBCL patients who have relapsed after an autoHCT.

With many patients undergoing autoHCT for DLBCL each year, and with approximately 40-50% of those transplants ultimately failing, the decision of whether to pursue an alloHCT for a DLBCL patient who has progressed after an autoHCT is, unfortunately, a common clinical dilemma. No prognostic models are currently available to counsel such patients regarding their expected survival outcomes following alloHCT. We therefore sought to develop a prognostic model for DLBCL patients undergoing allografting after a failed prior autoHCT, utilizing clinical factors readily available immediately before alloHCT.

## **MATERIALS AND METHODS**

### ***Data sources***

The Center for International Blood and Marrow Transplant Research (CIBMTR) is a working group of more than 500 transplantation centres worldwide that contribute detailed data on HCT to a statistical centre at the Medical College of Wisconsin. Participating centres are

required to report all transplantations consecutively; patients are followed longitudinally and compliance is monitored by on-site audits. Computerized checks for discrepancies, physicians' review of submitted data and on-site audits of participating centres ensure data quality.

Observational studies conducted by the CIBMTR are performed in compliance with all applicable federal regulations pertaining to the protection of human research participants.

The CIBMTR collects data at two levels: Transplant Essential Data (TED) and Comprehensive Report Form (CRF) data. TED data include disease type, age, gender, pre-HCT disease stage and chemotherapy-responsiveness, date of diagnosis, graft type (bone marrow- and/or blood-derived stem cells), conditioning regimen, post-transplant disease progression and survival, development of a new malignancy and cause of death. All CIBMTR centres contribute TED data. More detailed disease and pre- and post-transplant clinical information are collected on a subset of registered patients selected for CRF data by a weighted randomization scheme. TED and CRF level data are collected pre-transplant, 100 days and six months post-HCT and annually thereafter or until death. Data for the current analysis were retrieved from CIBMTR (TED and CRF) report forms.

### ***Patients***

Adult ( $\geq 18$  years) patients with relapsed/refractory DLBCL, undergoing alloHCT between 2000 and 2012 after experiencing a relapse or progression following a prior autoHCT were included in this study. Eligible donors included human leucocyte antigen (HLA)-identical siblings or adult unrelated donors (URD). Patients undergoing syngeneic or alternative donor HCT (e.g. umbilical cord blood or haploidentical) and those receiving *ex vivo* graft manipulation (T-cell depleted or CD34 selected grafts) were not included in the analysis. Patients undergoing a planned tandem auto-alloHCT (n=98) were not eligible. Patients receiving the prior autoHCT for indications other than DLBCL (n=275) were not included. Similarly, patients undergoing a post-autoHCT, allograft for indications other than relapsed or refractory DLBCL (e.g. graft failure, indolent NHL, therapy-related haematological malignancies etc.) were excluded.

### ***Definitions***

The intensity of alloHCT conditioning regimens was categorized as myeloablative or reduced intensity conditioning/non-myeloablative conditioning (RIC/NMA) using consensus criteria.(Bacigalupo *et al*, 2009) Previously established criteria for categorizing the degree of HLA matching were used for URDs.(Weisdorf *et al*, 2008) Complete remission (CR) to last

therapy line before HCT on CIBMTR forms is defined as complete resolution of all known areas of disease on radiographic [computerized axial tomography (CAT) scan] assessments, while partial remission (PR) is defined as  $\geq 50\%$  reduction in the greatest diameter of all sites of known disease and no new sites of disease. Resistant disease is defined as  $< 50\%$  reduction in the diameter of all disease sites, or development of new disease sites.

### ***Study Endpoints***

Primary outcomes were non-relapse mortality (NRM), progression/relapse, PFS and overall survival (OS). NRM was defined as death without evidence of lymphoma progression/relapse; relapse was considered a competing risk. Progression/relapse was defined as progressive lymphoma after HCT or lymphoma recurrence after a CR; NRM was considered a competing risk. For PFS, a patient was considered a treatment failure at the time of progression/relapse or death from any cause. Patients alive without evidence of disease relapse or progression were censored at last follow-up. The OS was defined as the interval from the date of transplantation to the date of death or last follow-up. Acute (Przepiorka *et al*, 1995) and chronic (Shulman *et al*, 1980) graft-versus-host disease (GVHD) was defined and graded using established criteria. Neutrophil recovery was defined as the first of 3 successive days with absolute neutrophil count (ANC)  $\geq 0.5 \times 10^9/l$  after post-transplantation nadir. Platelet recovery was considered to have occurred on the first of three consecutive days with platelet count  $20 \times 10^9/l$  or higher, in the absence of platelet transfusion for 7 consecutive days. For neutrophil and platelet recovery, death without the event was considered a competing risk.

### ***Statistical analysis***

Probabilities of PFS and OS were calculated as described previously. (Zhang *et al*, 2007) Cumulative incidence of NRM, lymphoma progression/relapse and haematopoietic recovery were calculated to accommodate for competing risks. (Zhang & Zhang, 2011) Associations among patient-, disease- and transplantation-related variables and outcomes of interest were evaluated using Cox proportional hazards regression. Backward elimination was used to identify covariates that influenced outcomes. Covariates with a  $p < 0.05$  were considered significant. The proportional hazards assumption for Cox regression was tested by adding a time-dependent covariate for each risk factor and each outcome. Covariates violating the proportional hazards assumption were added as time-dependent covariates in the Cox regression model. Interactions between the main effect and significant covariates were examined. Results are expressed as

hazard ratio (HR). The variables considered in multivariate analysis are shown in Supplemental Table 1. To evaluate the impact of GVHD on transplantation outcomes, multivariate analyses were performed using Cox proportional hazards models, where the main-effect variable was defined as the time-dependent occurrence of acute grade II-IV GVHD or chronic GVHD versus neither. Each step of model building included the main-effect. Factors with a  $p < 0.05$  were kept in the final model. The potential interactions between the main effect and all significant risk factors were tested. All statistical analyses were performed using SAS version 9.3 (SAS Institute, Cary, NC).

### ***Prognostic model for PFS***

To develop a prognostic model able to predict PFS of DLBCL patients undergoing an alloHCT after a failed prior autoHCT, a Cox regression method was used to identify potential patient- and disease-related risk factors associated with treatment failure (failure event of PFS), using backward elimination with  $p < 0.05$  to enter and remove factors from the model. The results were then confirmed using a stepwise selection procedure and a forward selection. The risk factors considered in the model-building procedure are shown in Supplemental Table 1. Risk scores between 0 and 5 were assigned based on the ratios of log HRs. The risk scores were then plotted using the Kaplan-Meier (KM) curves and fitted in the Cox proportional hazards model to classify risk scores into different risk groups based on their distribution of the KM curves and HRs of the Cox model. PFS probabilities of the developed risk groups were calculated using the KM estimates.

## **RESULTS**

### ***Patient Characteristics***

Between 2000 and 2012, 503 DLBCL patients undergoing an alloHCT after experiencing disease relapse or progression following a prior autoHCT were reported to the CIBMTR. Patient characteristics are described in Table I. Briefly, median age at alloHCT was 52 years, with the majority of patients being Caucasian/white (88%). Fifty-four per cent had advanced stage disease at diagnosis and at the time of alloHCT, 10% had bulky disease and 32% had extranodal involvement. The median number of prior therapies before alloHCT was 4. Prior to alloHCT, 74% had chemosensitive disease. RIC/NMA conditioning regimens were used in 376 subjects (75%) and peripheral blood was the most common graft source (91%). Donors were balanced

between related (50%) and unrelated (50%). Median **time interval** between **autoHCT** and **alloHCT** (TIBAA) was 15 months.

### ***Univariate Outcomes***

The probabilities of neutrophil recovery at day 28 and at day 100 were 94% (95% confidence interval; [CI]: 92-96) and 96% (95%CI: 94-98), respectively. The probabilities of platelet recovery at day 28 and day 100 were 83% (95%CI: 78-86) and 89% (95%CI: 86-92), respectively (Table II). The cumulative incidence of grade II-IV acute GVHD at day +100 was 36% (95%CI = 28-44) and chronic GVHD at 1 year was 40% (95%CI = 35-44).

Median follow-up of survivors was 55 months (range 1-149). The probabilities of NRM at 1, 3 and 5 years were 23% (95%CI: 19-27), 30% (95%CI: 26-34) and 31% (95%CI: 27-36), respectively (Figure 1A). The probabilities of disease progression/relapse at 1, 3 and 5 years were 33% (95%CI: 29-37), 38% (95%CI: 34-43) and 40% (95%CI: 36-45) (Figure 1B). The probabilities of PFS at 1, 3 and 5 years were 44% (95%CI: 40-48), 31% (95%CI: 27-36) and 29% (95%CI: 24-33), respectively (Figure 1C), and those for OS were 54% (95%CI: 49-58), 37% (95%CI: 32-41) and 34% (95%CI: 30-39), respectively (Figure 1D).

### ***Multivariate Outcomes***

On multivariate analysis, chemoresistant disease before HCT (HR=1.86, 95%CI:1.23-2.81; p=0.003) and URD transplantation (HR=1.44, 95%CI:1.04-2.00; p=0.03) were associated with a higher risk of NRM (Table III). Use of myeloablative conditioning displayed a time-varying effect on the risk of NRM. During the first 10 months post-transplant it was associated with a higher NRM (HR=1.99, 95%CI:1.34-2.95; p=0.001), but not beyond 10 months post-alloHCT (HR=0.59; p=0.23). Multivariate analysis for disease progression/relapse demonstrated that KPS <80 (HR=1.81, 95%CI:1.18-2.77; p=0.006) and chemoresistant disease (HR=2.25, 95%CI:1.51-3.36; p<0.0001) were associated with a higher risk of progression/relapse post-alloHCT (Table III). TIBAA displayed a time-varying effect on the risk of disease progression/relapse. During the first year post-alloHCT, a short (<12 months) TIBAA was associated with a higher progression/relapse risk (HR=2.28, 95%CI:1.66-3.14; p<0.0001), but not beyond first year post-alloHCT (HR=0.51; p=0.14).

Patients with KPS <80 (HR=1.79, 95%CI:1.29-2.48; p=0.0005), chemoresistant disease (HR=2.04, 95%CI:1.53-2.73; p<0.0001), short TIBAA (<12 months) (HR=1.32, 95%CI:1.06-



1.64;  $p=0.01$ ) and use of use of myeloablative conditioning (HR=1.29, 95%CI:1.09-1.63;  $p=0.03$ ) had a higher risk of therapy failure (i.e. inferior PFS) (Table III).

On multivariate analysis a higher risk of mortality (i.e. inferior OS) was associated with with KPS <80 (HR=1.86, 95%CI:1.33-2.60;  $p=0.0003$ ), chemoresistant disease (HR=1.94, 95%CI:1.44-2.61;  $p<0.0001$ ) and myeloablative conditioning (HR=1.39, 95%CI:1.09-1.78;  $p=0.008$ ). Graft type displayed a time-varying effect on the risk of mortality. During the first 3 months post-transplant, peripheral blood grafts were associated with a lower risk of mortality (HR=0.37, 95%CI:0.22-0.61;  $p<0.0001$ ), but not beyond 3 months post-alloHCT (HR=1.43;  $p=0.25$ ). (Table III).

Development of acute GVHD (HR=2.24, 95%CI:1.24-4.04;  $p=0.007$ ) and chronic GVHD (HR=1.72, 95%CI:1.06-2.82;  $p=0.03$ ) was associated with higher risk of NRM. Neither acute, nor chronic GVHD were associated with risk of disease relapse/progression (data not shown). Acute GVHD was associated with a higher risk of mortality (HR=1.85, 95%CI:1.27-2.69;  $p=0.001$ ). Chronic GVHD was not associated with mortality risk ( $p=0.54$ ).

#### ***CIBMTR Prognostic Model for PFS***

Three significant prognostic factors were included in the final model predicting post-alloHCT PFS: KPS, chemosensitivity status and TIBAA. The final model only included those patients who had no missing data regarding KPS, chemosensitivity and TIBAA ( $n=417$ ). Based on the ratios of log HRs in the final model, chemoresistant disease was assigned 5 points, KPS of <80 4 points and TIBAA <12 months was assigned 2 points (Table IV). Therefore, the total risk score for any individual patient using the 3 significant prognostic factors ranged from 0 to 11. Table IV summarizes the performance of the prognostic model. Distribution of patients by total risk score was as follows: 194 patients had a total risk score of 0 (reference category), 103 patients had a total risk score of 2 (HR=1.30 range, 0.97 to 1.76), 14 patients had a total risk score of 4 (HR=1.41 range, 0.76 to 2.62), 38 patients had a total risk score of 5 (HR=1.66 range, 1.13 to 2.46), 12 patients had a total risk score of 6 (HR=2.21 range, 1.19 to 4.11), 35 patients had a total risk score of 7 (HR=2.34 range, 1.58 to 3.47), 3 patients had a total risk score of 9 (HR=1.79 range, 0.44 to 7.24) and 18 patients had a total risk score of 11 (HR=5.47 range, 3.26 to 9.19).

**Based on the HRs and the distribution of the KM curves across the total risk score categories (Supplemental Figure 1S), we classified each patient into four prognostic risk**

groups: low-risk group (score = 0), intermediate-risk group (score = 2 to 5), high-risk group (score = 6 to 9) or very high-risk group (score = 11). Statistical significance was reached when we compared the PFS between low and intermediate group ( $p=0.01$ ), low and high-risk group ( $p<0.0001$ ) and low and very high-risk group ( $p<0.0001$ ) (Table IV). The 1-year PFS probabilities for the low, intermediate, high and very high-risk groups were 54% (95% CI=47-61), 40% (95% CI=33-48), 26% (95% CI=14-38) and 6% (95% CI=0-16), respectively. The probability for 3-year PFS was 40% (95% CI:32-47), 32% (95% CI=25-40), 11% (95% CI:2-20) and 6% (95% CI:0-16) respectively, for the three prognostic groups (Figure 2A). The prognostic model also predicted OS following alloHCT (Table IV). The 1-year OS probabilities for the low, intermediate, high and very high-risk groups were 63% (95% CI=57-70), 52% (95% CI=44-60), 38% (95% CI=25-51) and 17% (95% CI=0-34), respectively. The probability for 3-year OS was 43% (95% CI:36-51), 39% (95% CI=31-46), 19% (95% CI:8-31) and 11% (95% CI:0-26) respectively, for the three prognostic groups (Figure 2A).

#### *Impact of conditioning intensity*

Compared to RIC/NMA conditioning, the patients receiving myeloablative alloHCT were younger (median age 53 years vs. 48 years;  $p=0.0001$ ), more likely to have chemoresistant disease (19% [n=71] vs. 28% [n=35];  $p=0.04$ ) and similar KPS ( $p=0.54$ ). Table V summarizes survival outcomes of the study population stratified according conditioning intensity. In patients receiving myeloablative conditioning compared to RIC/NMA, the 5-year adjusted probabilities of PFS (27% vs. 30%;  $p=0.47$ , Figure 3A) and OS (28% vs. 37%;  $p=0.055$ , Figure 3B) were not significantly different. Restricting analysis to chemoresistant patients, the 5-year adjusted probabilities of PFS (13% vs. 18%;  $p=0.47$ , Figure 3C) and OS (15% vs. 25%;  $p=0.22$ , Figure 3D) in similar order, were not significantly different.

#### *Causes of Death*

At a median follow-up of 55 months, 325 patients were no longer alive. The most common cause of death post-alloHCT was relapsed DLBCL (N=142, 44% of all deaths). GVHD accounted for 9% (n=28) of deaths, while infections were responsible for 19% of mortality (n=61). For details please see Table 2S.

## **DISCUSSION**

Prognostic models predicting outcomes of alloHCT in DLBCL failing a prior autoHCT are currently not available. Here, we have performed a registry analysis of DLBCL patients undergoing alloHCT after a failed prior autograft. This analysis provides several important observations: (i) NRM (23% at 1-year, 30% at 3-years) remains significant following alloHCT, (ii) a prognostic model based on factors readily available prior to alloHCT (TIBAA, chemosensitivity status, KPS) was developed for pre-transplant patient counselling, (iii) GVHD increased risk of non-relapse and overall mortality without reducing risk of relapse/progression and (iv) myeloablative conditioning provides no benefit in this setting, including in the subset of patients with chemoresistant disease.

There is evidence to support a possible GVL effect in DLBCL, including long-term responses in chemoresistant patients undergoing RIC alloHCT,(Hamadani *et al*, 2013) and responses to donor lymphocyte infusion and/or withdrawal of immune suppression.(Bishop *et al*, 2008, Thomson *et al*, 2009) Because of the potential for a GVL effect in DLBCL, combined with the poor prognosis associated with relapse after autoHCT, such patients are often considered for alloHCT. Notably, in the current analysis no benefit of acute or chronic GVHD was seen, in terms of reducing the risk of disease progression/relapse. These observations are in line with another recent large CIBMTR analysis.(Urbano-Ispizua *et al*, 2015)

The decision to proceed with alloHCT in DLBCL after a failed autograft is complex because many of these patients have advanced age, impaired performance status or comorbid conditions that may limit their candidacy for alloHCT. For example, in one study, only 19% of patients who relapsed or progressed after autoHCT ultimately underwent an alloHCT.(Rigacci *et al*, 2012) Among DLBCL patients undergoing alloHCT after a failed autograft, no tools are available to estimate HCT survival outcomes for patient counselling. The CIBMTR prognostic score reported in this study is not only easy to use, but utilizes information readily available prior to alloHCT (response to last therapy before alloHCT, KPS at HCT and TIBAA). This prognostic model is not designed to be applied to DLBCL patients at the time of their initial relapse after autoHCT (e.g. to determine their candidacy for salvage therapies or for a future alloHCT), but rather as a tool to be used immediately prior to alloHCT for estimating transplantation outcomes for patient counselling.

To date, there have only been three previous studies that have focused specifically on alloHCT outcomes in DLBCL patients who progressed after a prior autoHCT (Table

VI).(Rigacci *et al*, 2012, van Kampen *et al*, 2011, Kim *et al*, 2014) These studies (which largely focused on patients who underwent alloHCT from 1995-2008) showed approximately 30-40% PFS; however each study was limited by relatively short follow-up (median 2-3 years), and limited patient numbers (30-165 patients). Potentially partly due to these limitations, these three studies had conflicting results regarding factors predicting improved PFS and OS after alloHCT. In contrast, the current study is strengthened by a large number of patients (n=503), treated in a more contemporary era (2000-2012), with a median follow up of 4.6 years.

Our study found a NRM rate of 23% at 1 year and 30% at 3 years. This is in line with other studies looking at alloHCT following a failed autoHCT in DLBCL patients, in which the rate of NRM was 17-28% at 3-5 years. (Rigacci *et al*, 2012, van Kampen *et al*, 2011, Kim *et al*, 2014) In the current study KPS <80, chemoresistant disease, a TIBAA < 1-year and myeloablative conditioning were all predictive of worse survival outcomes on multivariate analysis, generally in line with predictive factors reported in prior studies (Table IV). It is worth noting that in the European Group of Blood and Marrow Transplantation (EBMT) study,(van Kampen *et al*, 2011) a time from autoHCT to post-autograft relapse of < 1-year was predictive of PFS. In contrast we used TIBAA in this study, since the interval between autoHCT and post-autograft relapse is not captured for all patients in the CIBMTR registry. The TIBAA is not only easily imputable immediately prior to alloHCT, but (for the patients in the CIBMTR registry for whom interval between autoHCT and post-autoHCT relapse was captured) it also correlates closely with the interval between autoHCT and post-autoHCT relapse (data not shown).

We found no benefit of myeloablative conditioning in this study, even in the subset of chemoresistant patients. In fact, myeloablative conditioning was associated with increased NRM, inferior PFS as well as OS on multivariate analysis. These observations are consistent with prior CIBMTR data showing no benefit of myeloablative conditioning in chemoresistant DLBCL.(Hamadani *et al*, 2013) Our results indicate that the same holds true in the setting of DLBCL patients who have undergone a prior autoHCT.

Our study has limitations. The nature of data captured in the CIBMTR registry precludes comparison against DLBCL patients failing an autoHCT but never undergoing a subsequent alloHCT. In a recent CIBMTR study,(Hamadani *et al*, 2014) among DLBCL patients undergoing autoHCT who experienced disease relapse, the 3-year *post-relapse* OS was 19% (unpublished data). These unpublished observations should however, be used with caution to ascertain the

relative benefit of alloHCT in this setting. Other limitations of the current analysis include the lack of information regarding pre-alloHCT PET status, as well as biomarkers known to affect prognosis in DLBCL, such as cytogenetic abnormalities (*MYC*, *BCL2*, and *BCL6* rearrangements) or “cell-of-origin” profile (germinal centre versus activated B-cell). However it was recently reported that pre-alloHCT PET status in NHL does not predict PFS or OS.(Bachanova *et al*, 2015) In addition, while the presence of *MYC* rearrangement is associated with inferior PFS and OS following HCT (Thieblemont *et al*, 2011), the available literature would indicate that “cell-of-origin” profile fails to predict outcomes following HCT.(Moskowitz *et al*, 2005, Gu *et al*, 2012)

In conclusion, we were able to construct a CIBMTR prognostic model to predict PFS after alloHCT, using KPS, TIBAA and chemoresistance at alloHCT. This tool was able to discriminate 3-year PFS, ranging from 38% down to 10%. This same prognostic tool was able to discriminate 3-year OS, ranging from 43% down to 14%. This prognostic index should help provide a more accurate estimate of risks and benefits with alloHCT, when counselling DLBCL patients before a planned alloHCT. This prognostic model requires independent validation, possibly by analysing data reported to other transplantation registries (e.g. EBMT registry). The CIBMTR prognostic model is not designed to assess suitability of DLBCL patient for a future allograft, at the time of their initial post-autograft relapse. On the other hand, these data also illustrate the shortcomings of alloHCT for this patient population. Further gains will need to be achieved in reducing NRM as well as augmenting GVL effects in order for alloHCT to achieve more widespread applicability for DLBCL patients relapsing after autoHCT. Rationally designed clinical trials that integrate novel agents (such as immune checkpoint inhibitors, antibody-drug conjugates and B-cell receptor signalling inhibitors) and/or novel cellular therapies (such as chimeric antigen receptor technology) with alloHCT may help to achieve this goal.

### **Acknowledgements**

Morgan Geronime for administrative Support.

The CIBMTR is supported by Public Health Service Grant/Cooperative Agreement U24-CA076518 from the National Cancer Institute (NCI), the National Heart, Lung and Blood Institute (NHLBI) and the

National Institute of Allergy and Infectious Diseases (NIAID); a Grant/Cooperative Agreement 5U10HL069294 from NHLBI and NCI; a contract HSH250201200016C with Health Resources and Services Administration (HRSA/DHHS); two Grants N00014-13-1-0039 and N00014-14-1-0028 from the Office of Naval Research; and grants from \*Actinium Pharmaceuticals; Allos Therapeutics, Inc.; \*Amgen, Inc.; Anonymous donation to the Medical College of Wisconsin; Ariad; Be the Match Foundation; \*Blue Cross and Blue Shield Association; \*Celgene Corporation; Chimerix, Inc.; Fred Hutchinson Cancer Research Center; Fresenius-Biotech North America, Inc.; \*Gamida Cell Teva Joint Venture Ltd.; Genentech, Inc.; \*Gentium SpA; Genzyme Corporation; GlaxoSmithKline; Health Research, Inc. Roswell Park Cancer Institute; HistoGenetics, Inc.; Incyte Corporation; Jeff Gordon Children's Foundation; Kiadis Pharma; The Leukemia & Lymphoma Society; Medac GmbH; The Medical College of Wisconsin; Merck & Co, Inc.; Millennium: The Takeda Oncology Co.; \*Milliman USA, Inc.; \*Miltenyi Biotec, Inc.; National Marrow Donor Program; Onyx Pharmaceuticals; Optum Healthcare Solutions, Inc.; Osiris Therapeutics, Inc.; Otsuka America Pharmaceutical, Inc.; Perkin Elmer, Inc.; \*Remedy Informatics; \*Sanofi US; Seattle Genetics; Sigma-Tau Pharmaceuticals; Soligenix, Inc.; St. Baldrick's Foundation; StemCyte, A Global Cord Blood Therapeutics Co.; Stemsoft Software, Inc.; Swedish Orphan Biovitrum; \*Tarix Pharmaceuticals; \*TerumoBCT; \*Teva Neuroscience, Inc.; \*THERAKOS, Inc.; University of Minnesota; University of Utah; and \*Wellpoint, Inc. The views expressed in this article do not reflect the official policy or position of the National Institute of Health, the Department of the Navy, the Department of Defense, Health Resources and Services Administration (HRSA) or any other agency of the U.S. Government.

\*Corporate Members

**Author contributions:**

**Conception and design:** Timothy S. Fenske, Tara Graff and Mehdi Hamadani.

**Financial support:** CIBMTR

**Collection and assembly of data:** Alyssa DiGilio and Mehdi Hamadani.

**Data analysis:** Kwang W. Ahn, Alyssa DiGilio and Mehdi Hamadani.

**Interpretation:** All authors.

This article is protected by copyright. All rights reserved

**Manuscript writing:** First draft prepared by Timothy S. Fenske, Tara Graff and Mehdi Hamadani. All authors helped revised the manuscript.

**Final approval of manuscript:** All authors

**Disclosure of conflict of interest:** No disclosures to report.

## References:

- Bachanova, V., Burns, L.J., Ahn, K.W., Laport, G.G., Akpek, G., Kharfan-Dabaja, M.A., Nishihori, T., Agura, E., Armand, P., Jaglowski, S.M., Cairo, M.S., Cashen, A.F., Cohen, J.B., D'Souza, A., Freytes, C.O., Gale, R.P., Ganguly, S., Ghosh, N., Holmberg, L.A., Inwards, D.J., Kanate, A.S., Lazarus, H.M., Malone, A.K., Munker, R., Mussetti, A., Norkin, M., Prestidge, T.D., Rowe, J.M., Satwani, P., Siddiqi, T., Stiff, P.J., William, B.M., Wirk, B., Maloney, D.G., Smith, S.M., Sureda, A.M., Carreras, J., Hamadani, M. & Center for International Blood and Marrow Transplant Research Lymphoma Working Committee. (2015) Impact of pretransplantation (18)F-fluorodeoxy glucose-positron emission tomography status on outcomes after allogeneic hematopoietic cell transplantation for non-hodgkin lymphoma. *Biology of Blood and Marrow Transplantation : Journal of the American Society for Blood and Marrow Transplantation*, **21**, 1605-1611.
- Bacher, U., Klyuchnikov, E., Le-Rademacher, J., Carreras, J., Armand, P., Bishop, M.R., Bredeson, C.N., Cairo, M.S., Fenske, T.S., Freytes, C.O., Gale, R.P., Gibson, J., Isola, L.M., Inwards, D.J., Laport, G.G., Lazarus, H.M., Maziarz, R.T., Wiernik, P.H., Schouten, H.C., Slavin, S., Smith, S.M., Vose, J.M., Waller, E.K., Hari, P.N. & Lymphoma Working Committee of the CIBMTR. (2012) Conditioning regimens for allotransplants for diffuse large B-cell lymphoma: Myeloablative or reduced intensity? *Blood*, **120**, 4256-4262.
- Bacigalupo, A., Ballen, K., Rizzo, D., Giral, S., Lazarus, H., Ho, V., Apperley, J., Slavin, S., Pasquini, M., Sandmaier, B.M., Barrett, J., Blaise, D., Lowski, R. & Horowitz, M. (2009) Defining the intensity of conditioning regimens: Working definitions. *Biology of Blood and Marrow Transplantation : Journal of the American Society for Blood and Marrow Transplantation*, **15**, 1628-1633.

- Bishop, M.R., Dean, R.M., Steinberg, S.M., Odom, J., Pavletic, S.Z., Chow, C., Pittaluga, S., Sportes, C., Hardy, N.M., Gea-Banacloche, J., Kolstad, A., Gress, R.E. & Fowler, D.H. (2008) Clinical evidence of a graft-versus-lymphoma effect against relapsed diffuse large B-cell lymphoma after allogeneic hematopoietic stem-cell transplantation. *Annals of Oncology : Official Journal of the European Society for Medical Oncology / ESMO*, **19**, 1935-1940.
- Fenske, T.S., Hari, P.N., Carreras, J., Zhang, M.J., Kamble, R.T., Bolwell, B.J., Cairo, M.S., Champlin, R.E., Chen, Y.B., Freytes, C.O., Gale, R.P., Hale, G.A., Ilhan, O., Khoury, H.J., Lister, J., Maharaj, D., Marks, D.I., Munker, R., Pecora, A.L., Rowlings, P.A., Shea, T.C., Stiff, P., Wiernik, P.H., Winter, J.N., Rizzo, J.D., van Besien, K., Lazarus, H.M. & Vose, J.M. (2009) Impact of pre-transplant rituximab on survival after autologous hematopoietic stem cell transplantation for diffuse large B cell lymphoma. *Biology of Blood and Marrow Transplantation : Journal of the American Society for Blood and Marrow Transplantation*, **15**, 1455-1464.
- Gisselbrecht, C., Glass, B., Mounier, N., Singh Gill, D., Linch, D.C., Trneny, M., Bosly, A., Ketterer, N., Shpilberg, O., Hagberg, H., Ma, D., Briere, J., Moskowitz, C.H. & Schmitz, N. (2010) Salvage regimens with autologous transplantation for relapsed large B-cell lymphoma in the rituximab era. *Journal of Clinical Oncology : Official Journal of the American Society of Clinical Oncology*, **28**, 4184-4190.
- Gu, K., Weisenburger, D.D., Fu, K., Chan, W.C., Greiner, T.C., Aoun, P., Smith, L.M., Bast, M., Liu, Z., Bociek, R.G., Bierman, P.J., Armitage, J.O. & Vose, J.M. (2012) Cell of origin fails to predict survival in patients with diffuse large B-cell lymphoma treated with autologous hematopoietic stem cell transplantation. *Hematological Oncology*, **30**, 143-149.
- Hamadani, M., Saber, W., Ahn, K.W., Carreras, J., Cairo, M.S., Fenske, T.S., Gale, R.P., Gibson, J., Hale, G.A., Hari, P.N., Hsu, J.W., Inwards, D.J., Kamble, R.T., Klein, A., Maharaj, D., Marks, D.I., Rizzieri, D.A., Savani, B.N., Schouten, H.C., Waller, E.K., Wirk, B., Laport, G.G., Montoto, S., Maloney, D.G. & Lazarus, H.M. (2013) Impact of pretransplantation conditioning regimens on outcomes of allogeneic transplantation for chemotherapy-unresponsive diffuse large B cell lymphoma and grade III follicular lymphoma. *Biology of Blood and Marrow Transplantation : Journal of the American Society for Blood and Marrow Transplantation*, **19**, 746-753.
- Hamadani, M., Hari, P.N., Zhang, Y., Carreras, J., Akpek, G., Aljurf, M.D., Ayala, E., Bachanova, V., Chen, A.I., Chen, Y.B., Costa, L.J., Fenske, T.S., Freytes, C.O., Ganguly, S., Hertzberg, M.S.,



- Holmberg, L.A., Inwards, D.J., Kamble, R.T., Kanfer, E.J., Lazarus, H.M., Marks, D.I., Nishihori, T., Olsson, R., Reddy, N.M., Rizzieri, D.A., Savani, B.N., Solh, M., Vose, J.M., Wirk, B., Maloney, D.G., Smith, S.M., Montoto, S., Saber, W., Alpdogan, O., Cashen, A., Dandoy, C., Finke, R., Gale, R., Gibson, J., Hsu, J.W., Janakiraman, N., Laughlin, M.J., Lill, M., Cairo, M.S., Munker, R., Rowlings, P.A., Schouten, H.C., Shea, T.C., Stiff, P.J. & Waller, E.K. (2014) Early failure of frontline rituximab-containing chemo-immunotherapy in diffuse large B cell lymphoma does not predict futility of autologous hematopoietic cell transplantation. *Biology of Blood and Marrow Transplantation : Journal of the American Society for Blood and Marrow Transplantation*, **20**, 1729-1736.
- Kim, J.W., Kim, S.W., Tada, K., Fukuda, T., Lee, J.H., Lee, J.J., Kwon, J.H., Bang, S.M., Kim, I., Yoon, S.S., Lee, J.S. & Park, S. (2014) Allogeneic stem cell transplantation in patients with de novo diffuse large B-cell lymphoma who experienced relapse or progression after autologous stem cell transplantation: A Korea-Japan collaborative study. *Annals of Hematology*, **93**, 1345-1351.
- Klyuchnikov, E., Bacher, U., Kroll, T., Shea, T.C., Lazarus, H.M., Bredeson, C. & Fenske, T.S. (2014) Allogeneic hematopoietic cell transplantation for diffuse large B cell lymphoma: Who, when and how? *Bone Marrow Transplantation*, **49**, 1-7.
- Moskowitz, C.H., Zelenetz, A.D., Kewalramani, T., Hamlin, P., Lessac-Chenen, S., Houldsworth, J., Olshen, A., Chaganti, R., Nimer, S. & Teruya-Feldstein, J. (2005) Cell of origin, germinal center versus nongerminal center, determined by immunohistochemistry on tissue microarray, does not correlate with outcome in patients with relapsed and refractory DLBCL. *Blood*, **106**, 3383-3385.
- Mounier, N., Canals, C., Gisselbrecht, C., Cornelissen, J., Foa, R., Conde, E., Maertens, J., Attal, M., Rambaldi, A., Crawley, C., Luan, J.J., Brune, M., Wittnebel, S., Cook, G., van Imhoff, G.W., Pfreundschuh, M., Sureda, A. & Lymphoma Working Party of European Blood and Marrow Transplantation Registry (EBMT). (2012) High-dose therapy and autologous stem cell transplantation in first relapse for diffuse large B cell lymphoma in the rituximab era: An analysis based on data from the European Blood and Marrow Transplantation Registry. *Biology of Blood and Marrow Transplantation : Journal of the American Society for Blood and Marrow Transplantation*, **18**, 788-793.
- Philip, T., Guglielmi, C., Hagenbeek, A., Somers, R., Van der Lelie, H., Bron, D., Sonneveld, P., Gisselbrecht, C., Cahn, J.Y. & Harousseau, J.L. (1995) Autologous bone marrow transplantation as

compared with salvage chemotherapy in relapses of chemotherapy-sensitive non-hodgkin's lymphoma. *The New England Journal of Medicine*, **333**, 1540-1545.

Przepiorka, D., Weisdorf, D., Martin, P., Klingemann, H.G., Beatty, P., Hovs, J. & Thomas, E.D. (1995) 1994 consensus conference on acute GVHD grading. *Bone Marrow Transplantation*, **15**, 825-828.

Rezvani, A.R., Norasetthada, L., Gooley, T., Sorror, M., Bouvier, M.E., Sahebi, F., Agura, E., Chauncey, T., Maziarz, R.T., Maris, M., Shizuru, J., Bruno, B., Bredeson, C., Lange, T., Yeager, A., Sandmaier, B.M., Storb, R.F. & Maloney, D.G. (2008) Non-myeloablative allogeneic haematopoietic cell transplantation for relapsed diffuse large B-cell lymphoma: A multicentre experience. *British Journal of Haematology*, **143**, 395-403.

Rigacci, L., Puccini, B., Doderio, A., Iacopino, P., Castagna, L., Bramanti, S., Ciceri, F., Fanin, R., Rambaldi, A., Falda, M., Milone, G., Guidi, S., Martelli, M.F., Mazza, P., Oneto, R., Bosi, A. & Gruppo Italiano Trapianto di Midollo Osseo (GITMO). (2012) Allogeneic hematopoietic stem cell transplantation in patients with diffuse large B cell lymphoma relapsed after autologous stem cell transplantation: A GITMO study. *Annals of Hematology*, **91**, 931-939.

Shulman, H.M., Sullivan, K.M., Weiden, P.L., McDonald, G.B., Striker, G.E., Sale, G.E., Hackman, R., Tsoi, M.S., Storb, R. & Thomas, E.D. (1980) Chronic graft-versus-host syndrome in man. A long-term clinicopathologic study of 20 Seattle patients. *The American Journal of Medicine*, **69**, 204-217.

Sirvent, A., Dhedin, N., Michallet, M., Mounier, N., Faucher, C., Yakoub-Agha, I., Mohty, M., Robin, M., Tabrizi, R., Clement, L., Bilger, K., Larosa, F., Contentin, N., Huyn, A., Francois, S., Bulabois, C.E., Ceballos, P., Bourhis, J.H., Buzyn, A., Cornillon, J., Guillemin, G., de Revel, T., Bay, J.O., Guilhot, F. & Milpied, N. (2010) Low nonrelapse mortality and prolonged long-term survival after reduced-intensity allogeneic stem cell transplantation for relapsed or refractory diffuse large B cell lymphoma: Report of the Societe Francaise de Greffe de Moelle et de Therapie Cellulaire. *Biology of Blood and Marrow Transplantation : Journal of the American Society for Blood and Marrow Transplantation*, **16**, 78-85.

Thieblemont, C., Briere, J., Mounier, N., Voelker, H.U., Cuccini, W., Hirsch, E., Rosenwald, A., Jack, A., Sundstrom, C., Cogliatti, S., Trougouboff, P., Boudova, L., Ysebaert, L., Soulier, J., Chevalier, C., Bron, D., Schmitz, N., Gaulard, P., Houlgatte, R. & Gisselbrecht, C. (2011) The germinal center/activated B-cell subclassification has a prognostic impact for response to salvage

therapy in relapsed/refractory diffuse large B-cell lymphoma: A bio-CORAL study. *Journal of Clinical Oncology : Official Journal of the American Society of Clinical Oncology*, **29**, 4079-4087.

Thomson, K.J., Morris, E.C., Bloor, A., Cook, G., Milligan, D., Parker, A., Clark, F., Yung, L., Linch, D.C., Chakraverty, R., Peggs, K.S. & Mackinnon, S. (2009) Favorable long-term survival after reduced-intensity allogeneic transplantation for multiple-relapse aggressive non-hodgkin's lymphoma. *Journal of Clinical Oncology : Official Journal of the American Society of Clinical Oncology*, **27**, 426-432.

Urbano-Ispizua, A., Pavletic, S.Z., Flowers, M.E., Klein, J.P., Zhang, M.J., Carreras, J., Montoto, S., Perales, M.A., Aljurf, M.D., Akpek, G., Bredeson, C.N., Costa, L.J., Dandoy, C., Freytes, C.O., Fung, H.C., Gale, R.P., Gibson, J., Hamadani, M., Hayashi, R.J., Inamoto, Y., Inwards, D.J., Lazarus, H.M., Maloney, D.G., Martino, R., Munker, R., Nishihori, T., Olsson, R.F., Rizzieri, D.A., Reshef, R., Saad, A., Savani, B.N., Schouten, H.C., Smith, S.M., Socie, G., Wirk, B., Yu, L.C. & Saber, W. (2015) The impact of graft-versus-host disease on the relapse rate in patients with lymphoma depends on the histological subtype and the intensity of the conditioning regimen. *Biology of Blood and Marrow Transplantation : Journal of the American Society for Blood and Marrow Transplantation*, **21**, 1746-53.

van Kampen, R.J., Canals, C., Schouten, H.C., Nagler, A., Thomson, K.J., Vernant, J.P., Buzyn, A., Boogaerts, M.A., Luan, J.J., Maury, S., Milpied, N.J., Jouet, J.P., Ossenkoppele, G.J. & Sureda, A. (2011) Allogeneic stem-cell transplantation as salvage therapy for patients with diffuse large B-cell non-hodgkin's lymphoma relapsing after an autologous stem-cell transplantation: An analysis of the european group for blood and marrow transplantation registry. *Journal of Clinical Oncology : Official Journal of the American Society of Clinical Oncology*, **29**, 1342-1348.

Weisdorf, D., Spellman, S., Haagensohn, M., Horowitz, M., Lee, S., Anasetti, C., Setterholm, M., Drexler, R., Maiers, M., King, R., Confer, D. & Klein, J. (2008) Classification of HLA-matching for retrospective analysis of unrelated donor transplantation: Revised definitions to predict survival. *Biology of Blood and Marrow Transplantation : Journal of the American Society for Blood and Marrow Transplantation*, **14**, 748-758.

Zhang, X. & Zhang, M.J. (2011) SAS macros for estimation of direct adjusted cumulative incidence curves under proportional subdistribution hazards models. *Computer Methods and Programs in Biomedicine*, **101**, 87-93.

Zhang, X., Loberiza, F.R., Klein, J.P. & Zhang, M.J. (2007) A SAS macro for estimation of direct adjusted survival curves based on a stratified cox regression model. *Computer Methods and Programs in Biomedicine*, **88**, 95-101.

**Table I.** Characteristics of patients who underwent an allogeneic transplant after a failed autologous transplant for DLBCL from 2000-2012 reported to the CIBMTR. (Italicized text indicates variables available in CRF-level data patients).

<b>Number of patients</b>	<b>503</b>
<b>Number of CRF-level data patients</b>	155
<b>Number of centres</b>	133
<b>Median age at transplant, years (range)</b>	52 (19-72)
<b>Male gender</b>	305 (61)
<b>Race</b>	
Caucasian/White	444 (88)
Black	17 (3)
Others <sup>1</sup>	33 (7)
Missing	9 (2)
<b>Karnofsky Performance Score</b>	
80-100%	393 (78)
<80%	52 (10)
Missing	58 (12)
<b>Stage III/IV at Diagnosis</b>	83 (54)
<b>Remission status at HCT</b>	
Complete remission	175 (35)
Partial remission	197 (39)
Chemorefractory	106 (21)
Untreated	12 (2)
Unknown	13 (3)

<b>Rituximab prior to HCT</b>	112 (72)
<b>Radiation therapy prior to HCT</b>	98 (63)
<b>Lines of therapy prior to alloHCT</b>	
Median (range)	4 (1-7)
<b>History of transformation from indolent histology</b>	25 (16)
<b>Elevated lactate dehydrogenase at HCT</b>	52 (34)
<b>Active extranodal disease at HCT</b>	49 (32)
<b>Bone marrow involvement at HCT</b>	
No bone marrow involvement	141 (91)
Bone marrow involvement	7 (5)
Missing	7 (5)
<b>Bulky Disease (&gt;5 cm) at HCT</b>	15 (10)
<b>Conditioning regimen intensity</b>	
Myeloablative	127 (25)
Reduced intensity conditioning	376 (75)
<b>TBI in conditioning regimens</b>	
Myeloablative doses of TBI	41 (8)
<b>Graft type</b>	
Bone marrow	47 (9)
Peripheral Blood	456 (91)
<b>Type of donor</b>	
HLA-identical sibling	253 (50)
Unrelated well-matched	118 (23)
Unrelated partially matched	132 (26)
<b>Donor-Recipient CMV Status</b>	
-/+	102 (20)
Other	226 (45)
Missing	175 (35)

---

**GVHD Prophylaxis**

CNI + MMF +- others	180 (36)
CNI + MTX +-others (except MMF)	219 (43)
CNI + others (except MTX, MMF)	64 (13)
Other GVHD prophylaxis <sup>2</sup>	7 (1)
Missing GVHD prophylaxis	33 (7)
<b>Antithymocyte globulin in conditioning</b>	110 (22)
<b>Alemtuzumab in conditioning</b>	7 (1)
<b>Year of Transplant</b>	
2000-2003	111 (22)
2004-2007	154 (31)
2008-2012	238 (47)
<b>Time from autoHCT to alloHCT</b>	
Median (range)	15 (1-198)
≤12 months	201 (40)
>12 months	302 (60)
<b>Median follow-up of survivors (range), months</b>	55 (1-149)

HCT=haematopoietic cell transplantation; alloHCT=allogeneic haematopoietic cell transplantation; autoHCT= autologous haematopoietic cell transplantation; TBI=total body irradiation; CMV=Cytomegalovirus; GVHD=graft-versus-host disease;

CNI=calcineurin inhibitor; MMF=mycophenolate mofetil; MTX=methotrexate

<sup>1</sup> Asian (n=25), Native American (n=1), Pacific Islander (n=1), other (n=6)

<sup>2</sup>MMF/Campath (n=1), MMF/Sirolimus (n=1), MTX(n=3), MMF/MTX(n=1), MMF/MTX/Sirolimus (n=1)

**Table II. Haematopoietic recovery, graft-versus-host disease and survival outcomes.**

Outcomes	Evaluated (n)	Probability (95% CI)
Neutrophil recovery $>0.5 \times 10^9/l$	478	
28-day		94 (92-96)%
100-day		96 (94-98)%
Platelet recovery $\geq 20 \times 10^9/l$	374	

Outcomes	Evaluated (n)	Probability (95% CI)
28-day		83 (78-86)%
100-day		89 (86-92)%
Acute GVHD (II-IV)*	151	
100-day		36 (28-44)%
Acute GVHD (III-IV)*	151	
100-day		15 (10-21)%
Chronic GVHD	454	
6 month		26 (22-30)%
1-year		40 (35-44)%
3-year		47 (42-51)%
Extensive chronic GVHD	454	
1-year		33 (28-37)%
NRM	494	
1-year		23 (19-27)%
3-year		30 (26-34)%
5-year		31 (27-36)%
Relapse/Progression	494	
1-year		33 (29-37)%
3-year		38 (34-43)%
5-year		40 (36-45)%
Progression-free Survival	494	
1-year		44 (40-48)%
3-year		31 (27-36)%
5-year		29 (24-33)%
Overall survival	503	
1-year		54 (49-58)%

Outcomes	Evaluated (n)	Probability (95% CI)
3-year		37 (32-41)%
5-year		34 (30-39)%

\*Applies to patients with CRF-level data

GVHD = graft-versus-host disease; NRM = non-relapse mortality; CI = confidence interval.

**Table III. Multivariate Analysis Results**

<b>Non-relapse mortality</b>					
	<b>N</b>	<b>HR</b>	<b>95% CI</b>		<b>P-value</b>
			<b>Lower Limit</b>	<b>Upper Limit</b>	
<b>Chemosensitivity</b>					
CR	173	1			
PR	192	1.01	0.69	1.48	0.97
Chemoresistant	104	1.86	1.23	2.81	0.003
<b>Conditioning Regimen (≤10 months)</b>					
RIC/NMA	368	1			
MA	126	1.99	1.34	2.95	0.001
<b>Conditioning Regimen (&gt;10 months)</b>					
RIC/NMA	177	1			
MA	47	0.59	0.25	1.39	0.23
<b>Type of Donor</b>					
HLA-identical sibling	245	1			
Well-matched/partially matched	249	1.44	1.04	2.00	0.03
<b>Progression/Relapse</b>					



<b>KPS</b>					
80-100	388	1			
<80	51	1.81	1.18	2.77	0.006
<b>Chemosensitivity</b>					
CR	173	1			
PR	192	1.36	0.95	1.96	0.09
Chemoresistant	104	2.25	1.51	3.36	<0.0001
<b>Time from autoHCT to alloHCT (≤1 year from HCT)</b>					
≥12 months between auto & allo	294	1			
<12 months between auto & allo	200	2.28	1.66	3.14	<0.0001
<b>Time from autoHCT to alloHCT (&gt;1year from HCT)</b>					
≥12 months between auto & allo	146	1			
<12 months between auto & allo	66	0.51	0.20	1.25	0.14
<b>Progression free survival</b>					
<b>KPS</b>					
80-100	388	1			
<80	51	1.79	1.29	2.48	0.0005
<b>Chemosensitivity</b>					
CR	173	1			
PR	192	1.14	0.88	1.49	0.31
Chemoresistant	104	2.04	1.53	2.73	<0.0001
<b>Time from autoHCT to alloHCT</b>					
≥12 months between auto & allo	294	1			

<12 months between auto & allo	200	1.32	1.06	1.64	0.01
<b>Conditioning regimen</b>					
RIC/NMA	368	1			
MA	126	1.29	1.02	1.63	0.03
<b>Overall survival</b>					
<b>KPS</b>					
80-100	393	1			
<80	52	1.86	1.33	2.60	0.0003
<b>Chemosensitivity</b>					
CR	175	1			
PR	197	1.16	0.88	1.52	0.30
Chemoresistant	106	1.94	1.44	2.62	<0.0001
<b>Conditioning Regimens</b>					
RIC/NMA	376	1			
MA	127	1.39	1.09	1.78	0.008
<b>Graft Type (≤3 months)</b>					
Bone marrow	47	1			
Peripheral blood	456	0.37	0.22	0.61	<0.0001
<b>Graft Type (&gt;3 months)</b>					
Bone marrow	27	1			
Peripheral blood	377	1.43	0.78	2.64	0.25

N = number; HR = hazard ratio; CI = confidence interval; CR = complete response; PR = partial response; RIC = reduced intensity conditioning; NMA = ; non-myeloablative conditioning MA = myeloablative conditioning; KPS = Karnofsky performance status; HLA = human leucocyte antigen; HCT = haematopoietic cell transplantation; alloHCT = allogeneic; auto = autologous.

**Table IV. Prognostic model for progression free survival and overall survival**

Prognostic Score <sup>1</sup>	N	HR	95% CI		p-value
			Lower Limit	Upper Limit	
0	194	1			
2	103	1.30	0.97	1.76	0.08
4	14	1.41	0.76	2.62	0.28
5	38	1.66	1.13	2.46	0.01
6	12	2.21	1.19	4.11	0.01
7	35	2.34	1.58	3.47	<0.0001
9	3	1.79	0.44	7.24	0.41
11	18	5.47	3.26	9.19	<0.0001
<b>Progression-free Survival Risk Groups</b>					
Low (Score 0)	194	1			
Intermediate (Score 2,4,5)	155	1.40	1.08	1.82	0.01
High (Score 6,7,9)	50	2.28	1.61	3.22	<0.0001
Very high (Score 11)	18	5.47	3.26	9.19	<0.0001
<b>Contrast</b>					
Intermediate vs. High		0.61	0.43	0.87	0.006
Intermediate vs. Very high		0.26	0.15	0.43	<0.0001
High vs. Very high		0.42	0.24	0.73	0.002
<b>Overall Survival Risk Groups</b>					
Low (Score 0)	199	1			
Intermediate (Score 2,4,5)	156	1.34	1.02	1.76	0.03
High (Score 6,7,9)	50	2.11	1.47	3.03	<0.0001
Very high (Score 11)	18	3.94	2.32	6.68	<0.0001
<b>Contrast</b>					

Intermediate vs. High		0.63	0.44	0.91	0.02
Intermediate vs. Very high		0.34	0.20	0.58	<0.0001
High vs. Very high		0.54	0.30	0.96	0.03

N = number; HR = hazard ratio; CI = confidence interval; CR = complete response; PR = partial response; KPS = Karnofsky performance status; alloHCT=allogeneic haematopoietic cell transplantation; autoHCT= autologous haematopoietic cell transplantation;

<sup>1</sup>Prognostic score determined by following:

**KPS $\geq$ 80** = 0 point, **KPS <80** = 4 points.

**Disease status CR or PR** = 0 point, **Chemoresistant** = 5 points

**Time from autoHCT to alloHCT  $\geq$ 12 months** = 0 point, **<12 months** = 2 points.

**Table V. Allogeneic transplantation outcomes stratified according to transplantation conditioning intensity.**

	<b>Myeloablative conditioning</b> Adjusted probability (95% CI)	<b>Reduced-intensity or non-myeloablative conditioning</b> Adjusted probability (95% CI)	p-value
<b>Progression-free survival</b>	N= 126	N=368	
<b>1-year</b>	36 (26-44)%	46 (42-51)%	0.03
<b>3-year</b>	29 (21-36)%	33 (28-37)%	0.39
<b>5-year</b>	27 (19-34)%	30 (25-35)%	0.47
<b>Overall survival</b>	N=127	N=376	
<b>1-year</b>	44 (36-52)%	56 (52-61)%	0.01
<b>3-year</b>	31 (23-39)%	39 (34-44)%	0.12
<b>5-year</b>	28 (20-36)%	37 (32-42)%	0.055
<b>Chemoresistant patients only</b>			
<b>Progression-free survival</b>	N=35	N=69	
<b>1-year</b>	16 (4-27)%	33 (23-44)%	0.03
<b>3-year</b>	13 (2-24)%	20 (11-29)%	0.31
<b>5-year</b>	13 (2-24)%	18 (9-27)%	0.47

Overall survival	N=35	N=71	
1-year	23 (10-37)%	45 (33-56)%	0.02
3-year	15 (3-27)%	29 (19-39)%	0.09
5-year	15 (3-27)%	25 (15-35)%	0.22

**Table VI. Studies reporting outcomes of allogeneic transplantation in diffuse large B-cell lymphoma patients who received a prior autograft.**

Study	N	MA vs. RIC (N)	NRM	PFS	OS	Factors predicting better PFS or OS
Van Kampen <i>et al</i> (2011)	101	37 vs. 64	28% (3-year)	42% (3-year)	54% (3-year)	TIBAR>1-year, normal LDH, peripheral blood graft
Rigacci <i>et al</i> (2012)	165	49 vs. 116	28% (not specified)	31% (5-year)	39% (5-year)	Chemosensitive disease, matched sibling donors
Kim <i>et al</i> (2014)	30	7 vs. 23	17% (not specified)	38% (5-year)	43% (5-year)	Chemosensitive disease, good performance status
Current analysis	503	127 vs. 376	31% (5-year)	29% (5-year)	34% (5-year)	KPS>80, chemosensitive disease, RIC, TIBAA >1-year

MA=myeloablative conditioning; RIC=reduced intensity conditioning, NRM=non-relapse mortality, PFS=progression-free survival; OS=overall survival; LDH=lactate dehydrogenase; TIBAA=time-interval between autologous and allogeneic transplantation; TIBAR=time interval between autologous transplant and post-autograft relapse; KPS=Karnofsky performance score.

## FIGURE LEGENDS

**Figure 1. Outcomes for DLBCL patients undergoing allogeneic HCT after a prior failed autologous HCT.** Cumulative incidence of (1A) non-relapse mortality, (1B) disease progression/relapse, (1C) progression-free survival and (1D) overall survival.

**Figure 2. Prognostic index for DLBCL patients undergoing allogeneic HCT (alloHCT) after a prior failed autologous HCT (autoHCT).** Three adverse prognostic factors were used to construct a prognostic model for PFS, **including KPS <80 (4 points), interval between autoHCT and alloHCT of <1 year (2 points) and chemoresistant disease at alloHCT (5 points).** This classified patients into four groups: **low-risk (0 points), intermediate-risk (2-5 points), high-risk (6-9 points) or very high-risk (11 points).** (2A) Progression-free survival and (2B) overall survival based on CIBMTR prognostic index.

**Figure 3. Overall survival (OS) and progression-free survival (PFS) of DLBCL patients undergoing allogeneic HCT after a prior failed autologous HCT, stratified by conditioning intensity.** PFS of all patients (3A), OS of all patients (3B), PFS of chemoresistant patients (3C) and OS of chemoresistant patients (3D).

Figure 1a: Non-relapse Mortality




Figure 1b: Progression / Relapse




Figure 1c: Progression-free Survival




Figure 1d: Overall Survival




Figure 2a: Progression-free Survival




Figure 2b: Overall Survival






Figure 3a: Adjusted Progression-free Survival




Figure 3b: Adjusted Overall Survival




Figure 3c: Adjusted Progression-free Survival, Chemorefractory only subset




Figure 3d: Adjusted Overall Survival, Chemorefractory only subset

