Received Date : 29-Sep-2015

Revised Date : 16-Mar-2016

Accepted Date : 16-Mar-2016

Article type : Original Manuscript

Is a Mid-level Dental Provider Model Acceptable to Potential Patients?

Elizabeth Phillips, PhD Research Associate, University of Michigan School of Social Work 1080 S. University Ave Ann Arbor, MI 48109

H. Luke Shaefer, PhD Associate Professor, University of Michigan School of Social Work 1080 S. University Ave Ann Arbor, MI 48109

Mert N. Aksu, DDS, JD, MHSA Dean, University of Detroit Mercy School of Dentistry 2700 Martin Luther King Jr. Boulevard Detroit, MI 48208-2576

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the <u>Version of Record</u>. Please cite this article as <u>doi:</u> 10.1111/CDOE.12230

This article is protected by copyright. All rights reserved

\*Corresponding author

Abstract

Adrienne Lapidos, PhD\* Assistant Research Scientist, University of Michigan School of Social Work 1080 S. University Ave Ann Arbor, MI 48109 P 734-764-6686 F 734-763-3372 alapidos@umich.edu

Objective: This study aims to assess patient attitudes toward mid-level dental providers, known as dental therapists (DTs), by surveying those likely to be their patients. The recent adoption of accreditation standards by the Commission on Dental Accreditation has reignited a debate surrounding the state-by state legalization of DTs in the United States; while the dental profession is divided on DTs, it is important to understand how potential patients may view the DT model. Methods: A questionnaire that asks about oral health experience, and comfort with the model of a dually-trained dental therapist-hygienist, based on a provided definition, was administered to 600 patients and their waiting room companions at a large urban universitybased dental clinic. Results: 40% of respondents indicated they would be comfortable being treated by a DT for all 7 of the procedures referenced, and over 75% were comfortable with each of 5 procedures. Having caps or crowns placed was the only treatment about which respondents were evenly divided. Factors associated with greater odds of comfort with various procedures include being uninsured and being under the age of 65. Uninsured patients were 1.5 to 2 times more likely than privately insured patients to accept a DT. Conclusions: The introduction of midlevel dental providers is a strategy that the public, especially those lacking regular care, appear on the whole to be comfortable with.

**Key words**: Mid-level provider, dental therapist, expanded scope of practice, dental workforce, patient satisfaction

## Is a Mid-level Dental Provider Model Acceptable to Potential Patients?

## Introduction

Mid-level dental providers, commonly referred to as dental therapists (DTs), are members of dental teams in countries such as Australia, Canada, the United Kingdom, New Zealand, and the Netherlands. In the United States, DTs practice only in Alaska and Minnesota, and were recently approved to practice in Maine. DTs provide preventive oral health services, and also perform a limited set of irreversible procedures which in the United States have historically been performed only by dentists, such as preparing and placing fillings and routine extractions. They typically work in safety net or underserved settings under the general supervision of, or in collaborative agreements with, dentists. The recent adoption of accreditation standards by the Commission on Dental Accreditation has reignited a debate surrounding the state-by state legalization of DTs in the United States. Of the 12 states that are now exploring legislation for a new oral health workforce, 11 are considering a dental hygiene-based models<sup>1</sup>. Research indicates that DTs provide safe care, and within their scope of practice, their clinical competence is comparable to that of dentists<sup>2</sup>. Moreover, there is evidence to suggest that they improve access to dental care<sup>3</sup>, an issue of particular importance given the significant psychosocial<sup>4,5,6,7</sup> and whole health<sup>8,9,10,11</sup> consequences of poor oral health, and given that the Affordable Care Act and accountable care organizations may increase demand for dental care  $^{12, 13}$ . As oral health disparities continue to plague the American healthcare system  $^{14}$ , many advocate for the introduction of DTs with the goal of increasing access, lowering costs, and improving efficiency.

Dental therapists are often described as analogous to physician assistants (PAs) or nurse practitioners (NPs), chiefly because they are licensed medical professionals with a circumscribed scope of practice, and also because their roles were established in response to healthcare access problems during times of perceived shortages or maldistributions of doctors <sup>15</sup> and dentists <sup>16</sup>, respectively. Another similarity between these professions is the manner in which the general public's knowledge and acceptance of their role and competence has gradually evolved.

Currently, DT training and scope of practice is not well understood by the US public, or even by US dentists <sup>17</sup>. Research on the US public's view toward NPs and PAs conducted when these provider types were relatively uncommon suggests an analogous lack of understanding of their training and potential benefit, as well as misgivings about their full scope of practice <sup>18, 19, 20, 21, 22</sup>. Now that midlevel medical providers are commonplace, research indicates that patients are at least as satisfied with NPs and PAs as they are with physicians <sup>23, 24, 25</sup>. Research from the United Kingdom suggests a similar evolution with respect to DTs at the dawn of an effort to diversify the skill-mix in dentistry. While limited numbers of DTs have worked in salaried public settings in the UK since the 1960s, a program to modernize dentistry began in 2002, which included expanded training of DTs, allowing them to work in private settings. A series of studies addressed the issues of public awareness and social acceptability of DTs<sup>26, 27, 28</sup>, which found that only 10%-15% of respondents were aware of DTs, and virtually none knew their permitted duties. Once DT practice was described, however, roughly 60% of respondents were comfortable with the idea of DTs doing restorations, though they were more apprehensive about DTs treating their children. In addition, it appears that certain qualifiers, specifically improved access or lowered cost, seemed sometimes to increase support.

In light of recent efforts in a number of American states to advocate for legislative changes authorizing mid-level dental practice, it is important to gain a better understanding of the US public's feelings toward DTs; in particular, to assess the feelings of those most likely be treated by them. In the US, the debate over the introduction of DTs is highly charged, with opponents sometimes arguing that it will lead to a lower standard of care for already vulnerable populations <sup>29</sup>. Even if the evidence suggests otherwise, to the extent the public believes such practitioners represent a second tier of care, rather than a new provider trained to a single standard of care, it will be less likely to embrace them. The degree to which DTs have the potential to improve access is thus limited by the willingness of patients to be treated by them. In this context, the purpose of the current study is to present data on present-day perceptions of dental therapists within a population most likely to be served by them.

### Methods

<u>Survey</u>. The respondents in this study were adult patients and their waiting-room companions at the University of Detroit Mercy School of Dentistry general dental clinic. This

clinic is located in Detroit, and serves a primarily uninsured and publicly insured population. Though not representative of all dental patients, this sample's demographic composition (Table 1) mirrors those who are likely to be served by DTs in the jurisdictions where it is permitted. Prior to data collection, the study was submitted for Institutional Review Board (IRB) approval at the University of Michigan and the University of Detroit Mercy, and both boards deemed the study to be exempt from IRB oversight. Potential respondents were informed about the purpose of the study, the voluntary nature of participation, the fact that participation or lack thereof would have no impact on the care they received, and the confidentiality and anonymity of their questionnaire data. Interviewers wore University of Michigan badges, and stated that they were from the University of Michigan, to make it clear to potential respondents that an outside organization, not the dental clinic, was requesting participation in the survey.

The questionnaire (Appendix 1) was developed based on a review of studies on the early social acceptability of both dental and medical mid-level providers. To the extent possible, questions were drawn from commonly used dental survey tools <sup>30</sup> or adapted from previous studies <sup>18, 19, 20, 21, 22, 26, 27, 28, 31</sup>. The questionnaire asked about the oral health experiences of respondents and any dependent children, any inability to receive necessary services, and the purpose of the current clinic visit. After providing a brief description of one of the prominent DT models <sup>1</sup>, a dually-trained hygienist (Appendix 2), respondents were asked about their willingness to receive various services from such a provider. If respondents indicated discomfort with any of the procedures, they were asked follow-up questions about whether certain factors might change their minds. The questionnaire was available in English, Spanish, and Arabic.

The questionnaire was administered by trained research associates over nine visits to the clinic in the summer of 2013. The method of administration rotated every other visit, between questionnaires that were self-administered and questionnaires that were read aloud by a research associate. While each method has advantages and drawbacks <sup>32</sup>, it was thought that this rotating style would provide both breadth and depth in responses. Specifically, about twice as many self-administered questionnaires could be completed in the time it took to read one aloud, although self-administered questionnaires were more likely to suffer from missing responses and potential respondent misunderstandings. Self-administered questionnaires were completed on paper and were then collected by the researcher. To protect privacy, those waiting in line outside the clinic were only offered self-administered questionnaires. All Spanish and Arabic questionnaires were

self-administered. When questionnaires were read aloud, the DT description was read verbatim, so the information received would always be the same. An attempt was made to approach every adult in the waiting room, and, on several mornings, some of those waiting in line. Respondents were free to stop at any time.

Questionnaires were offered to 778 adults and were obtained from 628, for an 81% response rate. Due to missing data on key questions of willingness to see a DT, 28 questionnaires had to be dropped. Just over 1/3 of the questionnaires (218) were read aloud to respondents; the rest were self-administered.

<u>Data Analysis</u>. All statistical analyses were conducted using the statistical package STATA V.13, and the level of significance was set at 0.05. Descriptive statistics reporting respondents' demographics and the percentage stating various opinions were calculated. Logistic regression models were estimated to parse the factors associated with respondents' comfort with the idea of treatment by a DT. We present five models, with the same predictor variables but different dependent variables corresponding to specific treatments.

### Results

Table 1 describes the demographic characteristics of our sample. Respondents were roughly evenly divided between being uninsured, publicly insured, or privately insured; the majority had at least some college education; and about 40% had dependent children. The dental experience of respondents is reported in Table 2. Though the majority visits the dentist at least once a year, half described the condition of their mouths as fair or poor. Nearly half said there had been a time in the last year when they or somebody in their household had skipped needed dental care, with cost being by far the most important reason. About 1/3 of respondents (35%) were there accompanying somebody and did not have an appointment themselves; of these, however, over 40% said they were in need of care.

Respondents were asked if they would see the proposed dually-trained DT for care, and if they had children, whether or not they would take their children to one (Table 3). In calculating these percentages, non-responses were treated as "no" in the interest of presenting findings that are maximally conservative; that is, least supportive of dental therapy. For 5 of the 7 procedures, over 75% of respondents were comfortable with the idea. They were most comfortable receiving advice and information about their teeth, or explanations of treatment options, though nearly as many were comfortable with the ideas of seeing the proposed DT in an emergency, and having routine cavities filled. Indeed, 77% of respondents indicated they would be willing to have a DT fill their teeth. Caps or crowns was the only procedure over which respondents were evenly divided. Overall, 40% were comfortable with DTs performing all 7 procedures, and fewer than 10% were uncomfortable with all. Respondents were less comfortable with DTs providing injections for their children than for themselves, and were more comfortable with DTs providing extractions for their children than for themselves.

Those respondents who were uncomfortable receiving even one of the listed procedures from a DT were asked whether they thought any of several potential factors might change their minds (Table 4). For each query, roughly half indicated they might reconsider. Nearly 2/3 stated they would reconsider if their dentist made the treatment plan and assured them the DT could do the work. Costing less and accepting insurance were the next two highest reasons for reconsideration. Those with children appear slightly less likely to change their minds about their children's care.

Positive response to treatment from DTs differed by administration method, with selfadministered questionnaires indicating lower support than for those that were read aloud Multivariate analyses control for administration style. Findings on comfort with the various procedures, both treating non-responses as "no," and excluding missing responses, are reported separately by questionnaire administration style in Appendix 3.

Odds-ratio estimates are presented in Table 5, which describes the comfort level of various demographic groups with procedures within dental therapy scope of practice. Those comfortable with all 7 procedures are more likely to have skipped dental care due to cost barriers, and less likely to be seniors (relative to working age) and have a bachelor's degree or higher (relative to those with some college education). Specific procedures were associated with varying response patterns. For example, uninsured respondents were more comfortable with DTs providing fillings, extractions, and emergency care than were respondents with private insurance; in particular, they were nearly twice as likely as privately insured patients to be comfortable with a DT providing an extraction.

### Discussion

Dental therapists are part of the dental team in a number of countries and a limited number of American states. Wherever they have been introduced, a primary goal has been improving access to care. Lack of dental care has serious social and public health implications, with those living in or near poverty disproportionately impacted. Thirty-seven percent of US households report skipping care in a given year, with the uninsured twice as likely to do so <sup>33</sup>. Poor oral health causes children to miss school and adults to miss work, resulting in lower academic performance and lost productivity <sup>33, 5</sup>. Psychosocial consequences of dental problems include embarrassment, shyness, and feeling worthless, as well as reduced job prospects <sup>6, 7</sup>. Dental problems account for a rising number of visits to hospital emergency rooms <sup>34, 35</sup>, care that is not only expensive, but often limited to treatment for pain and infection, leaving underlying problems unaddressed <sup>36</sup>, and diverting valuable time and resources away from problems that ERs are better able to treat <sup>37</sup>.

Not only would adding a new lower-cost member to the dental team increase the number of providers, but, it is argued, it would also promote cost-effective treatment, freeing dentists to concentrate on more complex cases that take advantage of their extensive skills and training <sup>3, 38, 39</sup>. Little is known about the actual reduction in cost, and there is no evidence to show that the addition of DTs in Minnesota has resulted in a reduction in the cost of providing care. A survey conducted for the WK Kellogg Foundation in 2011 found that 78% of respondents in a nationally representative sample supported the idea of training a new "licensed dental practitioner" to provide preventive, routine dental care to those going without <sup>40</sup>. While this provides support for the idea of mid-level dental providers, it does not assess respondents' level of comfort with actually patronizing them for specific procedures. In fact, while existing literature examines the social acceptability of treatment from PAs and NPs when these mid-level medical providers were relatively unknown, and a series of studies from the UK sheds light on public opinion toward treatment by DTs when they were still uncommon, to our knowledge the current study is the first to ask similar questions about the acceptability of treatment by DTs in the US, and specifically, to ask those most likely to be their patients.

Qualitatively, our findings are similar to those from past studies that found no more than 20% of respondents refusing all proposed treatments from a new mid-level provider, and between 1/3 and nearly 2/3 being comfortable with all <sup>18, 20, 22, 28</sup>. The observed variation in comfort across procedure is also similar to prior findings, in that respondents were generally least comfortable with unknown mid-levels performing non-traditional procedures, or those considered more invasive <sup>18, 19</sup>. The fact that the respondents in our study who expressed some discomfort with DTs sometimes modified their opinions, particularly if a dentist would reassure

them or the provider were less costly, is also consistent with several prior studies <sup>18, 22, 26, 41</sup>, as are our estimates of correlations between comfort with specific procedures and demographic variables such as lack of insurance and being under 65. Dyer, Humphris and Robinson, for example, found that in the UK, males, younger participants, and those with perceived treatment need were more likely to find having their teeth restored by a DT acceptable, while those receiving some private treatment were more likely to find it unacceptable <sup>28</sup>. Shamansky et al. found that respondents who said they would use NP services were 6.4 times more likely to be dissatisfied with their current health care (based in part on questions about availability and cost) than those who said they would not use an NP <sup>22</sup>. Age and measures related to health or perceived need, as well as visit frequency or regularity, was found to be related to comfort with mid-levels in other studies as well <sup>19, 21</sup>.

It is interesting to consider why comfort with treatment by hypothetical DTs appears to be somewhat higher in our study than it was for an already existing, albeit little known, practitioner in the UK, as well as for NPs and PAs when they were first being introduced. It may be related to the fact that most people have now had experience with hygienists and medical midlevels, so the idea of being treated by a non-doctor is perhaps not as foreign as it once was. In addition, our sample was drawn from an underserved population at a dental clinic, which may mean it is a less healthy or more vulnerable sample than those of many other studies. An interesting extension of this work would be to explore the feelings of more affluent private practice patients.

It must be noted that positive response to treatment from DTs differed by administration method, with self-administered questionnaires indicating lower support than for those that were read aloud (71% vs 87% for fillings, for example). Though we cannot definitively explain this, there are several possible reasons. One possible reason is that the non-response rate to specific questions varied considerably by administration style. There were only six non-responses to the questions about own comfort with various treatments among the 218 questionnaires that were read aloud. By contrast, among the 382 self-administered questionnaires, the number of non-responses to specific questions about own comfort ranged from 40 (caps and crowns) to 25 (fillings). Reasons for higher rates of non-responses in self-administered questionnaires are unknown, and could perhaps be explained by low motivation to complete all questionnaire items. The decision was made to treat non-responses as "no" because in doing so, the findings are less

supportive of dental therapy and therefore more conservative; one result of this decision is that the mean for self-administered responses is brought down, while the mean for questionnaires that were read aloud is virtually unaffected. A second possible reason is that although the questions were designed to be neutral, it is possible those answering face-to-face might have felt approval was the "right" answer, whereas those filling it in themselves may have been more willing to say "no." Yet those hearing the description read aloud might also have achieved a better understanding of dental therapy than those reading for themselves – who may then have been more inclined to say "no." While we have reason to suspect the questionnaires that were read aloud may be more accurate, by combining the data we weight the estimates toward the selfadministered questionnaires, which are less supportive of dental therapy.

A limitation of the current study is its use of a single dental school's patients, which circumscribes its external validity. It may be the case that our convenience sample has unique features that would distinguish it from other communities likely to be served by DTs. Responses to the demographic questions do suggest that the sample mirrors the patients who are commonly served in safety-net dental settings, and future surveys in other settings may clarify the extent to which the current study is generalizable. Until the current study, little was known about the feelings of potential patients toward mid-level dental providers in the US. Understanding these feelings is important, since any improvement in access - and in related public health issues depends not just on additional or better placed providers, but on the willingness of patients to be treated by them. If the segment of the population that is most likely to benefit from the introduction of a new mid-level dental provider is unwilling to be seen by them, the primary goal of introducing DTs to the dental workforce will not be met. Despite the fact that DTs do not currently practice in Michigan, and most respondents had never heard of them, the majority of respondents in our study were on the whole quite comfortable with the idea of receiving treatment from them. In fact, when asked whether "in general, do you think it would be a good idea to let dental therapists work in Michigan?" the overwhelming response among the patients and their waiting room companions in our study (roughly 90%), was "yes."

Overall, we find no evidence that potential patients would perceive DTs to be providing second-tier care. Indeed, consistent with past findings, those with perceived need or lack of access are actually more likely to be comfortable with the idea. Additionally, to the extent that comfort with certain treatments is qualified, it should be noted that when NPs and PAs were first

introduced in the UK and the US, the public was also unsure about receiving certain treatments from those unknown practitioners. Comparing the opinions of potential DT patients in the US to early opinions toward DTs in the UK, or NPs and PAs here, provides insight into how US feelings about DTs might be expected to evolve. In as much as our findings tend to mirror those of earlier studies, if such practitioners were to become commonplace, current reservations will likely be reduced. This will particularly be the case if US dentists gain comfort with the idea, and evidence suggests that with familiarity, doctors' opinions do evolve <sup>3, 42</sup>. Trust in a primary care provider has been shown to impact views on the acceptability of care from other members of the medical team <sup>18, 41</sup>. Similarly, 2/3 of those respondents in our study expressing reservations with DTs felt that being told by their dentist that a DT could do the work could lead them to reconsider. If, on the other hand, opponents of this workforce model – including, at the moment, most state dental organizations and the American Dental Association, though not the American Association of Public Health Dentistry – continue to argue, despite evidence to the contrary <sup>3, 2</sup>, that the addition of mid-level providers to the US dental workforce will create a "two-tiered" system of care <sup>29, 43</sup>, public doubt may develop where it otherwise did not exist.

The fact that so many in the US lack access to oral health care creates serious social and public health problems. In addition to the suffering of those in need, society pays a heavy price in the form of children's learning, adults' work, misuse of hospital emergency services, worsened health outcomes, and increased medical expenses. Increasing the total number of dental providers, and more efficiently utilizing the time of highly trained physicians and dentists, are key components of most strategies to reduce barriers to care. To this end, 11 American states are currently exploring legislation for a dental hygiene-based mid-level oral health workforce <sup>1</sup>. The current study begins to suggest that the public, especially those who are lacking regular care, has a significant potential to embrace this new type of provider.

# Acknowledgments

This project was supported by a grant from the W.K. Kellogg Foundation and a grant from the Nokomis Foundation. The authors wish to thank Sophie Buchmueller, Renee Tetrick, Teague Simoncic, and the staff of the University of Detroit Mercy Dental Clinic.

### References

1. The American Dental Hygiene Association. The benefits of dental hygiene-based oral health

This article is protected by copyright. All rights reserved

provider models. [Internet]. adha.org 2015 [cited 17 December 2015]. Available from: http://www.adha.org/resources-docs/75112\_Hygiene\_Based\_Workforce\_Models.pdf

- Phillips E, Shaefer H. Dental Therapists: Evidence of technical competence. Journal of Dental Research. 2013;92:S11-S15.
- 3. Wright J, Graham F, Hayes C, Ismail A, Noraian K, Weyant R et al. A systematic review of oral health outcomes produced by dental teams incorporating midlevel providers. The Journal of the American Dental Association. 2013;144:75-91.
- 4. Jackson S, Vann W, Kotch J, Pahel B, Lee J. Impact of poor oral health on children's school attendance and performance. Am J Public Health. 2011;101:1900-1906.
- 5. Gift H, Reisine S, Larach D. The social impact of dental problems and visits. Am J Public Health. 1992;82:1663-1668.
- 6. Guarnizo-Herreño C, Wehby G. Children's dental health, school performance, and psychosocial well-being. The Journal of Pediatrics. 2012;161:1153-1159.e2.
- 7. Hyde S, Satariano W, Weintraub J. Welfare dental intervention improves employment and quality of life. Journal of Dental Research. 2006;85:79-84.
- 8. Bui T, Markham C, Ross M, Mullen P. Examining the association between oral health and oral HPV infection. Cancer Prevention Research. 2013;6:917-924.
- 9. Desvarieux M, Demmer R, Jacobs D, Papapanou P, Sacco R, Rundek T. Changes in clinical and microbiological periodontal profiles relate to progression of carotid intima-media thickness: the oral infections and vascular disease epidemiology study. Journal of the American Heart Association. 2013;2:e000254-e000254.
- 10. Scannapieco F. Pneumonia in nonambulatory patients: the role of oral bacteria and oral hygiene. Journal of the American Dental Association. 2006;137:S21-S25.
- Bensley L, VanEenwyk J, Ossiander E. Associations of self-reported periodontal disease with metabolic syndrome and number of self-reported chronic conditions. Preventing chronic disease. 2011;8.

- Yarbrough C, Vujicic M, Nasseh K. More than 8 million adults could gain dental benefits through Medicaid expansion. Health Policy Institute Research Brief, American Dental Association. 2014;February.
- Sullivan L. Dental insurance but no dentists. The New York Times [Internet]. 2012 [cited 15 December 2014];. Available from: http://www.nytimes.com
- 14. Dye B, Li X, Thornton-Evans G. Oral health disparities as determined by selected Healthy People 2020 oral health objectives for the United States. Hyattsville, MD: National Center for Health Statistics.; 2012.
- 15. US Congress, Office of Technology Assessment. Nurse practitioners, physician assistants, and certified nurse-midwives: a policy analysis. Washington, DC: US Government Printing Office; 1986.
- 16. Nash D, Friedman J, Mathu-Muju K. A review of the global literature on dental therapists: In the context of the movement to add dental therapists to the oral health workforce in the United States. Battle Creek, Michigan: W.K. Kellogg Foundation; 2012.
- 17. To'olo G, Nash D, Mathu-Muju K, Haney C, Mullins M, Bush H. Perspectives of board certified pediatric dentists on adding a pediatric oral health therapist to the dental team. Pediatric dentistry. 2010;32:505-512.
- 18. Litman T. Public perceptions of the physicians' assistant--a survey of the attitudes and opinions of rural Iowa and Minnesota residents. Am J Public Health. 1972;62:343-346.
- Breslau N, Novack A. Public attitudes toward some changes in the division of labor in medicine. Medical Care. 1979;17:859-867.
- Storms D, Fox J. The public's view of Physicians Assistants and Nurse Practitioners. Medical Care. 1979;17:526-535.
- Brands R. Acceptance of nurses as primary-care provides by retired people. Advances in Nursing Science. 1983;5:37-49.
- 22. Shamansky S, Schlung L, Holbrook T. Determining the market for Nurse Practitioner

services. Nursing Research. 1985;34:242-247.

- Roblin D, Becker E, Adams E, Howard D, Roberts M. Patient satisfaction with primary care. Medical Care. 2004;42:579-590.
- 24. Hooker R, Cipher D, Sekscenski E. Patient satisfaction with physician assistant, nurse practitioner, and physician care: a national survey of Medicare beneficiaries. Journal of clinical outcomes management. 2005;12:88-92.
- 25. Jeanmonod R, DelCollo J, Jeanmonod D, Dombchewsky O, Reiter M. Comparison of resident and mid-level provider productivity and patient satisfaction in an emergency department fast track. Emergency Medicine Journal. 2012;30:e12-e12.
- 26. Dyer T, Robinson P. Exploring the social acceptability of skill-mix in dentistry. International dental journal. 2008;58:173-180.
- 27. Dyer T, Robinson P. Public awareness and social acceptability of dental therapists. International Journal of Dental Hygiene. 2009;7:108-114.
- 28. Dyer T, Humphris G, Robinson P. Public awareness and social acceptability of dental therapists. BDJ. 2010;208:E2-E2.
- 29. American Academy of General Dentistry. White paper on increasing access to and utilization of oral health care services. Chicago, IL: Academy of General Dentistry; 2008 p. Accessed June 15 2014 at http://www.agd.org/media/54365/7025accesstocarewhitepaper7\_31\_08.pdf.
- 30. Drc.hhs.gov. CDC | Open Data Tools | Division of Oral Health [Internet]. 2015 [cited 15 June 2013]. Available from: http://drc.hhs.gov/SurveyQ/bydomain.htm
- 31. Baer L, Baldwin K, Sisk R, Watts P, Grinslade M, Brockschmidt B et al. Development of an instrument to measure community acceptance of Nurse Practitioners and Physician Assistants. Journal of nursing management. 1999;7:63-77.
- Bowling A. Mode of questionnaire administration can have serious effects on data quality. Journal of Public Health. 2005;27:281-291.

- 33. Kaiser Family Foundation. Kaiser Health Tracking Poll: September 2013 [Internet]. Kff.org. 2013 [cited 19 May 2014]. Available from: http://kff.org/health-reform/poll-finding/kaiserhealth-tracking-poll-september-2013/
- 34. The Pew Center on the States, Pew Children's Dental Campaign. A costly dental destination
   The Pew Charitable Trusts [Internet]. Pewtrusts.org. 2012 [cited 29 February 2012].
   Available from: http://www.pewtrusts.org/en/research-and-analysis/reports/2012/02/28/a-costly-dental-destination
- 35. Health Policy Resources Center Research Brief. Dental-related emergency department visits on the increase in the United States. American Dental Association; 2013.
- 36. Davis E, Deinard A, Maiga E. Doctor, my tooth hurts: the costs of incomplete dental care in the emergency room. Journal of Public Health Dentistry. 2010;70:205-210.
- 37. Shortridge E, Moore J. Use of emergency departments for conditions related to poor oral health care. Washington, DC: Office of Rural Health Policy; 2010.
- 38. The Pew Charitable Trusts. Expanding the dental team: studies of two private practices. [Internet]. pewtrusts.org. 2014 [cited 15 December 2014]. Available from: http://www.pewtrusts.org/~/media/legacy/uploadedfiles/pcs\_assets/2014/expandingdentalte amreportpdf.pdf
- 39. The Pew Charitable Trusts. Expanding the dental team: increasing access to care in public settings [Internet]. pewtrusts.org. 2014 [cited 15 December 2014]. Available from: http://www.pewtrusts.org/~/media/Assets/2014/06/27/Expanding\_Dental\_Case\_Studies\_Re port.pdf
- 40. Lake Research Partners. The dental access gap: findings from a national survey. WK Kellogg Foundation; 2011.
- 41. Dyer T, Owens J, Robinson P. What matters to patients when their care is delegated to dental therapists?. BDJ. 2013;214:E17-E17.
- 42. Street D, Cossman J. Does familiarity breed respect? Physician attitudes toward nurse practitioners in a medically underserved state. Journal of the American Academy of Nurse

Practitioners. 2010;22:431-439.

43. Rossignol S. Bill would create two-tier system of dental care. Kennebec Journal [Internet].
 2013 [cited 22 October 2013]; Available from: http://www.kjonline.com/opinion/bill-would-create-two-tier-system-of-dental-care\_2013-05-25.html


SGIG	Table 1:         Sample Demographic Characteristics
Male	39.1%
	(230/588)
Race <sup>a</sup>	
African-American	42.4%
Π	(241/569)
White	42.9%
	(244/569)
Ethnicity	
Latino	8.4%
	(39/466)
Arab/Middle Eastern	10.1%
	(47/466)
Age	
Senior (>=65)	20.1%
	(116/576)
Working Age (26-64)	71.2%
	(410/576)
Youth (18-25)	8.7%
	(50/578)

# Education

HS grad or less	30.3%	
	(177/585)	
Some college education	38.5%	
	(225/585)	
BA or more	31.3%	
$\bigcirc$	(183/585)	
Dental Insurance Status <sup>b</sup>		
Uninsured	37.4%	
$\mathbf{O}$	(219/585)	
Medicaid	29.9%	
U)	(175/585)	
Private	32.1%	
	(188/585)	
Have children under 18	40.5%	
<b>m</b>	(241/595)	

<sup>a</sup> Other possible race categories included: Indian-Native American (3), Asian-Pacific Islander (13); and other (60); 8 respondents checked more than one.


<sup>b</sup> Multiple responses were allowed.



Self-rated condition of teeth/gums	50.6%
fair or poor	(287/567)
Was there a time somebody skipped	43.1%
care? <sup>b</sup>	(258/599)
Cost	66.8%
	(159/238)
No time	15.6%
	(37/238)
Too afraid/nervous	14.3%
()	(34/238)
Unable to find dentist who took	13.9%
insurance	(33/238)

<sup>a</sup> Other possible responses included: about every two years, less often than every two years, and don't know.

<sup>b</sup> Multiple responses were allowed. These were the top four reasons; other choices included: didn't know who to call, couldn't get an appointment, transportation issue, lack of dependent care, some other reason, and don't know.



**Comfort with Seeing a Dental Therapist for Various Procedures** 

		Respondent
	Respondent	would take
Procedure	Would see a DT	child to a DT
	(n=600)	(n=241)

76.8%	73.4%
(461)	(177)
59.5%	66.0%
(357)	(159)
76.5%	66.8%
(459)	(161)
52.2%	47.3%
(313)	(114)
85.7%	82.2%
(514)	(198)
83.3%	83.0%
(500)	(200)
79.0%	76.4%
(474)	(184)
40%	39.8%
(240)	(96)
4.5%	6.6%
(27)	(16)
	<ul> <li>(461)</li> <li>59.5%</li> <li>(357)</li> <li>76.5%</li> <li>(459)</li> <li>52.2%</li> <li>(313)</li> <li>85.7%</li> <li>(514)</li> <li>83.3%</li> <li>(500)</li> <li>79.0%</li> <li>(474)</li> <li>40%</li> <li>(240)</li> <li>4.5%</li> </ul>

Note: Non-responses were taken as "no;" the number responding to each question ranged from 560 (caps/crowns) to 575 (fillings) for self, and from 202 (caps/crowns) to 229 (fillings) for children.



**Factors Potentially Affecting Discomfort with Dental Therapists** 

(asked of those uncomfortable with at least one procedure)

	Respondent	Respondent
Reason	would see a DT	would take
	(n=367)	child to DT
		(n=153)
If could get an appointment more	51.0%	49.0%
quickly	(187)	(75)
If it cost less	60.2%	56.2%
	(221)	(86)
If easier to get to	50.7%	50.3%
$\tilde{\mathbf{O}}$	(186)	(77)
If accepted insurance	56.1%	55.6%
	(206)	(85)
If dentist made plan and said ok	65.9%	62.1%
	(242)	(95)

Note: Non-responses were taken as "no;" the number responding to each question ranged from 313 (took insurance) to 320 (cost less) for self, and from 128 (dentist said ok) to 133 (quicker appointment; cost less) for children.

Table 5:

## **Comfort with Receiving Various Procedures from a Dental Therapist**

(Odds-ratio logistic regression results)

	Dependent Variable				
	<b>Fillings</b>	Extractions	Caps/Crowns	<b>Emergencies</b>	<u>All 7<sup>a</sup></u>
Male	1.21	1.40 +	1.37+	1.21	1.38+
	(0.27)	(0.27)	(0.26)	(0.28)	(0.26)
Black <sup>b</sup>	0.94	1.32	1.60*	0.64+	1.36

This article is protected by copyright. All rights reserved

	(0.22)	(0.27)	(0.31)	(0.16)	(0.27)
	(0.22)	(0.27)	(0.51)	(0.10)	(0.27)
Other race <sup>b</sup>	0.52*	1.04	0.92	0.36**	0.99
	(0.16)	(0.29)	(0.25)	(0.11)	(0.27)
Medicaid <sup>c</sup>	1.03	0.98	1.33	1.00	1.09
	(0.28)	(0.23)	(0.31)	(0.28)	(0.26)
Uninsured <sup>c</sup>	1.88*	1.94**	1.44 +	1.76*	1.43
	(0.51)	(0.44)	(0.31)	(0.50)	(0.32)
Unsure or other ins <sup>c</sup>	0.62	1.65	1.00	0.68	0.99
0	(0.25)	(0.65)	(0.38)	(0.29)	(0.39)
Youth (18-26) <sup>d</sup>	1.36	0.92	1.83+	2.15+	1.07
	(0.56)	(0.30)	(0.62)	(0.96)	(0.34)
Senior <sup>d</sup>	0.56*	0.50**	0.53**	1.15	0.51**
	(0.15)	(0.12)	(0.12)	(0.33)	(0.12)
HS or less <sup>e</sup>	0.52*	0.96	0.81	0.46**	0.83
Π	(0.14)	(0.22)	(0.18)	(0.13)	(0.18)
BA or more <sup>e</sup>	0.48**	0.44**	0.60*	0.78	0.52**
	(0.13)	(0.10)	(0.13)	(0.22)	(0.12)
Skipped care	0.99	1.30	1.57*	1.24	1.50*
	(0.22)	(0.25)	(0.29)	(0.29)	(0.28)
Self-administered	0.34**	0.44**	0.64*	0.33**	0.65*
	(0.08)	(0.09)	(0.12)	(0.09)	(0.12)
Constant	11.14**	2.08*	0.96	11.23**	0.71
	(4.22)	(0.62)	(0.27)	(4.40)	(0.20)
<b>—</b>					
Pseudo-R <sup>2</sup>	0.084	0.089	0.0656	0.085	0.057
Observations <sup>f</sup>	553	553	553	553	553
+ p < 10; * $p < 05$ ;	** p< 01				

+ p<.10; \* p<.05; \*\* p<.01

<sup>a</sup> also includes: injections, advice and info, and explanations

<sup>b</sup> reference category is white

<sup>c</sup> reference category is private insurance

This article is protected by copyright. All rights reserved

<sup>d</sup> reference category is working age

<sup>e</sup> reference category is some college education

<sup>f</sup> all dependent variables were coded 1 for a positive response and 0 for either a negative response or no response; 47 cases were excluded due to missing values on one or more explanatory variable

Note: Of the 553 respondents included in the regression sample, overall: 427 (77%) would get a filling; 330 (60%) would have a tooth pulled; 289 (52%) would get a cap/crown; 439 (79%) would be seen in an emergency; and 222 (40%) were comfortable with all 7 procedures.

Author Manus


Appendix 1

University of Michigan	For Office Use Number:
School of Social Work	S-A / O:
	Date:
We want your opinions on dental c	are!
are interested in learning about dental care. This survey is voluntary and y fidential. You do not even need to give us your name. The information we earchers studying dental care, but there will be no way for them to know w	gather will be used by ho said what. There are no
ht or wrong answers, and nothing you say will have any impact on your visi are just interested in your experiences and opinions.	it here today, or in the future.
1. How often do you go to the dentist? (check one)	
At least once a year	
About every 2 years	
Less often than every 2 years	
Only when needed, no regular schedule	
Don't know	
2. If you care for children (younger than age 18), on average, about how many to dentist? (check one)	imes a year do they go to the
No children (please go to the next question)	
At least once a year	
About every 2 years	
Less often than every 2 years	
Only when needed, no regular schedule	
Don't know	
Children are too young to go to the dentist (or today is child's first visit)	
<ol> <li>In the past 12 months, has there been a time when you or someone in your h but didn't go?</li> </ol>	ousehold needed to see a dentist
Yes No Don't know	
1	

 If you answered NO or DON'T KNOW to question 3 (there WASN'T or you WEREN'T SURE if there was a time when you, or someone in your household, needed to see a dentist but didn't), SKIP to question 5.

If you answered YES to question 3 (you, or someone in your household, DID need to see a dentist but didn't): Why did you, or someone in your household, not go to the dentist when needed? (check all that apply)

Didn't have time
Would cost too much
Couldn't find a dentist who took Medicaid or our insurance
Didn't know a dentist to call
Couldn't get an appointment
Would have to travel too far, or didn't have a way to get there
Didn't have anybody to care for children or other family member
Afraid, nervous, don't like needles
Some other reason:
Don't know
5. Why are you here today? (check all that apply)
5. Why are you here today: (check all that apply)
Came with somebody else (not seeing the dentist myself)
Regular checkup, or to get teeth cleaned
To have teeth filled
Trouble with gums
To have teeth pulled
Toothache
Loose tooth
To adjust, repair or have dentures made
Other:
Don't want to say
6. If you have an appointment here today, SKIP to question 7.
If you came with somebody who has an appointment, but you don't have one yourself: Do you feel you are currently in need of dental treatment?
bo you leel you are currency in need of dental deadheint.
Yes No Don't know
7. How would you describe the condition of your teeth and gums? (check one)
Poor
G Fair
Good
Very good     Don't know
2



This article is protected by copyright. All rights reserved

A dental therapist is like a nurse practitioner or a physician assistant, but for dentistry. They are in between a hygienist and a dentist because they can do all the things a dental hygienist can do, and some of the things a dentist can do. Dental therapists are allowed to assess the situation, do fillings, pull teeth, and a few other procedures. They are not allowed to do complicated procedures or treat more serious problems in your mouth. Only a dentist can do that. Dental therapists are taught to fill and pull teeth the same way dentists are, and they need to pass an exam to get licensed just like a dentist does. They work under the supervision of a dentist, and can talk with him in person or by phone or other electronic means if questions come up. Right now, dental therapists work in about 50 other countries. They also work in Minnesota. Other states are thinking about allowing them to work, to try to make it easier for people to get dental care.

Now we are going to ask some questions about dental therapists. We're interested in learning whether you think you would be comfortable seeing one in different kinds of situations. If you care for children, we're also interested in learning whether you think you would be comfortable taking your child to see a dental therapist.

14. If Michigan had dental therapists, do you think you personally would feel comfortable seeing one:

to have a simple cavity filled?	Yes	D No
to have a tooth taken out?	Yes	D No
to get a shot to make your teeth and gums go numb?	Yes	D No
to put on caps or crowns that stick permanently to your teeth?	Yes	D No
to get advice or information about your teeth?	Yes	D No
to explain treatment options for problems with your teeth?	Yes	D No
in an emergency for these things, if a dentist weren't available?	Yes	🗆 No

15. If you care for children (younger than age 18), do you think you would feel comfortable taking your child to see a dental therapist:

No children (please go to the next question)


to have a simple cavity filled?	Yes	D No
to have a baby tooth taken out?	Yes	D No
to get a shot to make their teeth and gums go numb?	Yes	D No
to put on caps or crowns that stick permanently to their teeth?	Yes	D No
to get advice or information about their teeth?	Yes	D No
to explain treatment options for problems with their teeth?	Yes	D No
in an emergency for these things, if a dentist weren't available?	Yes	No No

4

16. If you answered YES to all parts of questions 14 and 15 (you ARE comfortable with dental therapists), SKIP to 19.

If you answered NO to any part of questions 14 or 15 (you are NOT comfortable): What is the reason you think you would be uncomfortable with a dental therapist? (check all that apply) Worried about the quality of care Don't know enough about them Already have a dentist Other: 

Now, we're going to ask about some things that might, or might not, change your mind. We're going to ask about your opinion both for yourself, and then if you care for children, for them.




5

21. Are you:
C Male
□ Female
22. Are you:
C African American / Black
Caucasian / White
D American Indian / Alaskan Native
🗖 Asian / Pacific Islander
Other
23. Are you:
G Middle Eastern / Arabic
D Neither
24. What is your age:
25. What is the highest level of school you have completed? (check one)
Not a high school graduate (no diploma or GED). What was the last grade you completed?
High school graduate or GED
Some college, but less than a bachelor's degree
Bachelor's degree
Graduate or professional degree
26. What type of dental insurance do you, or others in your household, have? (check all that apply)
Medicaid (or other dental insurance from the state)
C Private (employer-provided) dental insurance
D Uninsured
D Other:
Don't know
27. What is your zip code:
You're done! Thank you for your time!
6

Appendix 2

The description of a dental therapist provided in the survey was: "A dental therapist is like a nurse practitioner or a physician assistant, but for dentistry. They are in between a hygienist and a dentist because they can do all the things a dental hygienist can do, and some of the things a dentist can do. Dental therapists are allowed to assess the situation, do fillings, pull teeth, and a few other procedures. They are not allowed to do complicated procedures or treat more serious problems in your mouth. Only a dentist can do that. Dental therapists are taught to fill and pull teeth the same way dentists are, and they need to pass an exam to get licensed just like a dentist does. They work under the supervision of a dentist, and can talk with him in person or by phone or other electronic means if questions come up. Right now, dental therapists work in about 50 countries. They also work in Minnesota. Other states are thinking about allowing them to work, to try to make it easier for people to get dental care." As a point of comparison, the mid-level medical provider introduction given to respondents by Breslau and Novack <sup>19</sup> was "One way of saving a doctor's time is by using people like nurses, paramedics, and physician's assistant who can perform some duties which a doctor might ordinarily do. Although they are not doctors, these people are well trained and work under the supervision of a physician." And Storms and Fox <sup>20</sup> described a PA as: "A physician assistant is not a doctor or nurse but a new health professional who has received special medical training in order to provide direct health care under the overall supervision of a doctor;" an NP was described as "A nurse practitioner is a nurse who has received additional training in order to provide direct health care in collaboration with a doctor."



Procedure	Self-	Read	Self-	Read
	Administered	Aloud	Administered	Aloud
	(n=382)	(n=218)	(n varies)	(n varies)
Filling	71.2%	86.7%	75.8%	87.5%
	(272/382)	(189/218)	(272/359)	(189/216)
Extraction	53.4%	70.2%	58.3%	70.5%
	(204/382)	(153/218)	(204/350)	(153/217)
Injection	69.9%	88.1%	75.9%	88.1%
	(267/382)	(192/218)	(267/352)	(192/218)
Cap/Crown	48.2%	59.2%	53.5%	59.7%
	(184/382)	(129/218)	(184/344)	(129/216)
Advice/Information	80.6%	94.5%	87.3%	94.5%
	(308/382)	(206/218)	(308/353)	(206/218)
Explanations	80.4%	88.5%	87.7%	88.5%
	(307/382)	(193/218)	(307/350)	(193/218)
Emergency	73.6%	88.5%	79.8%	88.9%
	(281/382)	(193/218)	(281/352)	(193/217)

Author N