

IN FOCUS

Passenger mutations and aberrant gene expression in congenic tissue plasminogen activator-deficient mouse strains

R. SZABO,* A. L. SAMSON,† D. A. LAWRENCE,‡ R. L. MEDCALF† and T. H. BUGGE*

*Proteases and Tissue Remodeling Section, Oral and Pharyngeal Cancer Branch, National Institute of Dental and Craniofacial Research, National Institutes of Health, Bethesda, MD, USA; †Australian Centre for Blood Diseases, Monash University, Alfred Medical Research and Education Precinct, Melbourne, Victoria, Australia; and ‡Internal Medicine, Division of Cardiovascular Medicine, University of Michigan Medical School, Ann Arbor, MI, USA

To cite this article: Szabo R, Samson AL, Lawrence DA, Medcalf RL, Bugge TH. Passenger mutations and aberrant gene expression in congenic tissue plasminogen activator-deficient mouse strains. *J Thromb Haemost* 2016; **14**:1618–28.

See also van Vlijmen BJM, Westrick RJ. Backseat drivers: passenger mutations take control of experimental phenotypes. This issue, pp 1615–7.

Essentials

- C57BL/6J-tissue plasminogen activator (tPA)-deficient mice are widely used to study tPA function.
- Congenic C57BL/6J-tPA-deficient mice harbor large 129-derived chromosomal segments.
- The 129-derived chromosomal segments contain gene mutations that may confound data interpretation.
- Passenger mutation-free isogenic tPA-deficient mice were generated for study of tPA function.

Summary. *Background:* The ability to generate defined null mutations in mice revolutionized the analysis of gene function in mammals. However, gene-deficient mice generated by using 129-derived embryonic stem cells may carry large segments of 129 DNA, even when extensively backcrossed to reference strains, such as C57BL/6J, and this may confound interpretation of experiments performed in these mice. Tissue plasminogen activator (tPA), encoded by the *PLAT* gene, is a fibrinolytic serine protease that is widely expressed in the brain. A number of neurological abnormalities have been reported in tPA-deficient mice. *Objectives:* To study genetic contamination of tPA-deficient mice. *Materials and methods:* Whole genome expression array analysis, RNAseq expression profiling, low- and high-density single nucleotide polymorphism

(SNP) analysis, bioinformatics and genome editing were used to analyze gene expression in tPA-deficient mouse brains. *Results and conclusions:* Genes differentially expressed in the brain of *Plat*^{-/-} mice from two independent colonies highly backcrossed onto the C57BL/6J strain clustered near *Plat* on chromosome 8. SNP analysis attributed this anomaly to about 20 Mbp of DNA flanking *Plat* being of 129 origin in both strains. Bioinformatic analysis of these 129-derived chromosomal segments identified a significant number of mutations in genes co-segregating with the targeted *Plat* allele, including several potential null mutations. Using zinc finger nuclease technology, we generated novel ‘passenger mutation’-free isogenic C57BL/6J-*Plat*^{-/-} and FVB/NJ-*Plat*^{-/-} mouse strains by introducing an 11 bp deletion into the exon encoding the signal peptide. These novel mouse strains will be a useful community resource for further exploration of tPA function in physiological and pathological processes.

Keywords: brain; congenic mice; gene targeting; mutation; tissue plasminogen activator.

Introduction

The development of methods to selectively disrupt genes in mice by homologous recombination in embryonic stem (ES) cells provided a potent new tool for analysis of gene function in mammals that has been of immeasurable value to a wide range of research fields [1]. Because germ line-competent ES cells were most successfully derived from various sub-strains of the 129 mouse strain, the large majority of gene disruptions initially were made in this strain. Due to the poor breeding characteristics, neurological and neuroanatomical abnormalities and high tumor incidence of 129 mice, gene-targeted mice developed using

Correspondence: Thomas H. Bugge, Proteases and Tissue Remodeling Section, Oral and Pharyngeal Cancer Branch, National Institute of Dental and Craniofacial Research, National Institutes of Health, 30 Convent Drive, Room 320, Bethesda, MD 20892, USA
Tel.: +1 301 435 1840; fax: +1 301 402 0823.
Email: thomas.bugge@nih.gov

Received 22 February 2016

Manuscript handled by: P. H. Reitsma

Final decision: P. H. Reitsma, 5 April 2016

129 ES cells typically were backcrossed to inbred reference strains, such as C57BL/6J. However, because of the low probability of meiotic crossover occurring close to the targeted gene, a large region of DNA is likely to remain of 129 origin, even in ‘congenic’ mice backcrossed for 10 generations ($P = 0.91$ for more than 1 centiMorgan of 129-derived DNA flanking each side of the targeted gene) [2]. Importantly in this respect, a recent direct comparison of the 129 and C57BL/6J genome sequences uncovered that the two strains differ by no less than 1395 deletions, insertions or nucleotide substitutions in protein coding regions, which resulted in 188 lost or gained stop codons, 875 frame shift variants and 332 splice donor or acceptor variants [2]. This finding led the authors of the above study to conclude that essentially all constitutively gene-targeted mouse strains generated using 129-derived ES cells carry ‘passenger mutations’ that confound data interpretation [2]. Indeed, several examples of erroneous attribution of observed phenotypes in gene-targeted mice generated using 129-derived ES cells have emerged recently [2–4]. With regards to extracellular proteases, the reported resistance of matrix metalloproteinase (MMP)-13-deficient mice and, most likely, MMP-7-deficient and MMP-8-deficient mice, to LPS-induced lethality was demonstrated not to be a consequence of the loss of the MMP. Rather, it was caused by a null mutation in the neighboring *Casp11* gene, which is present in 129 mice, but not in C57BL/6J mice [2]. A second issue with many ES cell-generated gene-disrupted mice relates to the use of homologous recombination to generate the null mutation. This strategy often involves the deletion of large segments of DNA within the targeted gene that could result in the deletion of cis-acting regulatory elements and microRNAs, while the insertion of a powerful neomycin selection cassette could similarly modulate the expression of genes flanking the targeted locus.

Tissue plasminogen activator (tPA) is a fibrinolytic serine protease [5] that is widely expressed in the brain [6–10], where it is involved in a diverse array of physiological and pathophysiological processes unrelated to fibrinolysis that were identified in large part through the analysis of tPA-deficient mice [11–25].

During transcriptomic profiling of genes differentially expressed in the brains of tPA-deficient mice extensively backcrossed to the C57BL/6J strain, we noted a highly significant over-representation of differentially expressed genes clustering around the *Plat* gene, encoding tPA, on chromosome 8. A similar anomalous clustering of differentially expressed genes was observed when we analyzed a second colony of tPA-deficient mice independently highly backcrossed to the C57BL/6J strain. Single nucleotide polymorphism (SNP) analysis provided evidence that at least 22 Mbp of DNA flanking *Plat* remained of 129 origin in each C57BL/6J-*Plat*^{-/-} strain. Furthermore, bioinformatic analysis identified several mutations, including potential null mutations, in protein coding genes located on this chromosomal segment.

Using zinc finger nuclease (Zfn) technology, we engineered a null mutation in the *Plat* gene of C57BL/6J and FVB/N mice. We demonstrate that isogenic C57BL/6J-*Plat*^{-/-} mice, free of 129 DNA, do not show anomalous clustering of differentially expressed genes in proximity to the targeted *Plat* gene. These novel isogenic *Plat*^{-/-} mouse strains constitute a valuable community resource for further exploration of tPA functions in the brain and other tissues.

Materials and methods

Mice

Mice were housed in standard barrier facilities under approved protocols.

Expression array analysis

Twelve- to 14-week-old male ‘Melbourne’ C57BL/6J-*Plat*^{-/-} and age-matched C57BL/6J mice (Jackson Laboratory, Bar Harbor, ME, USA) were transcardially transfused with ice-cold PBS (eight per genotype). The hippocampus was dissected out and placed in ‘RNA later’. RNA was extracted using the ‘RNeasy Lipid Tissue’ kit (Qiagen, Valencia, CA, USA) according to the manufacturer’s instructions. RNA was then pooled from pairs of mice to give four samples from each genotype for subsequent analysis. The extracted RNA was quality controlled, hybridized to Illumina microarray chips and analyzed by the SRC MicroRNA facility (University of Queensland, Australia) using MouseWG-6 v2.0 Expression BeadChip (Catalog ID: BD-201-0202). Genes were considered differentially expressed if the adjusted P value was less than 0.05. The original high-throughput microarray data are available through the following links: <https://cloudstor.aarnet.edu.au/plus/index.php/s/PYxAHTIkiH4zYy> and <https://cloudstor.aarnet.edu.au/plus/index.php/s/jPAXctPNeiIXKvQ>.

RNAseq

Six-week-old ‘Michigan’ C57BL/6J-*Plat*^{-/-} and ‘Bethesda’ C57BL/6J-*Plat*^{-/-} littermates (four mice per genotype), and age-matched wild-type C57BL/6J control mice (four mice per genotype), were euthanized by CO₂ inhalation. The brains were dissected out, the olfactory bulb and cerebellum removed, and the brains snap frozen in liquid nitrogen. cDNA library preparation and Illumina high-throughput sequencing were performed by The University of Michigan’s DNA Sequencing Core. Bioinformatic analysis of these data was conducted by the University of Michigan Bioinformatics Core. All data from these studies have been deposited in the gene expression omnibus (GEO) database with accession number GSE76093. Genes were considered differentially

expressed if presenting with an adjusted *P* value (*q*-value) of 0.05001 or less and fold-difference in expression of more than 0.3. Genes located on the Y chromosome were excluded from the analysis because the analyzed mice were both male and female.

SNP analysis

Low-density SNP analysis of the entire mouse genome was performed by the Jackson Laboratory using the 'JAX 150 SNP Panel' polymorphic between C57BL/6J and 129 (Table S1). The panel consists of 154 SNPs that cover the 19 autosomes and the X chromosome with a density of ~15–20 Mbp/SNP. High-density SNP analysis of mouse chromosome 8 was performed by the Jackson Laboratory using a chromosome 8-specific panel of 34 SNPs, polymorphic between C57BL/6J and 129, covering the length of the chromosome with a density of ~3.5 Mbp/SNP (Table S2).

Generation of *Plat*^{-/-} mice using *Zfns*

A pair of Zinc finger-FokI fragment fusion proteins (*Zfns*) binding, respectively, 5'-GACTGGCTTCCCAT-3' (nucleotides 9399–9413 of the *Plat* gene, NC_000074.6) and 5'-GACCAGGTGGGT-3' (nucleotides 9419–9430 of the *Plat* gene) were generated by Sigma-Aldrich (St Louis, MO, USA). mRNA encoding the *Zfn* pair was generated by *in vitro* transcription and microinjected into the male pronucleus of C57BL/6J and FVB/NJ zygotes, which were implanted into pseudo-pregnant mice. The ensuing founders were screened by PCR using the primer pair 5'-AAGAGT-CATTGCTGGATGGG-3' (forward) and 5'-GATCACTCCTGGGAACGTGT-3' (reverse), which amplifies the 325 bp fragment constituting nucleotides 9221–9545 of the *Plat* gene. Founders positive for mutations in the signal peptide-encoding exon 2 of *Plat* were further characterized by DNA sequencing and screened for germ line transmission by breeding to C57BL/6J or FVB/NJ.

tPA Western blot analysis

Non-reduced and non-boiled protein lysates from brains were separated by SDS-PAGE and transferred to poly(vinylidene difluoride) (PVDF) membranes. The membranes were blocked with 5% non-fat dry milk for 1 h. The blot was incubated with 2 µg/mL rabbit anti-human tPA (ASHTPA-GF; Molecular Innovation, Novi, MI, USA) for 1 h at room temperature. Bound antibody was visualized by incubation for 1 h with donkey anti-rabbit-HRP (Jackson Immuno Research, 711-036-152) 1 : 5000.

Plasminogen-casein zymography

Plasminogen-casein zymography was performed as previously described [26]. Lysis zones were visualized by staining of gels with Coomassie brilliant blue.

tPA activity assay

Active tPA was determined using a commercial sandwich tPA capture ELISA kit (Molecular Innovations, NTBIOCPAI) as recommended by the manufacturer.

Statistical analysis

Distribution of differentially expressed genes over all chromosomes: we used a chi-square statistic with a reference distribution calculated using simulation (because many of the expected hit counts are very small). Specifically, the expected number of 'hits' per chromosome, assuming uniform distribution, was calculated and compared with the observed number of hits per chromosome. This comparison utilized a chi-square statistic (but not a chi-square reference distribution). To obtain a reference distribution we generated Poisson counts from 100 000 randomizations with mean values equal to the expected number of hits per chromosome under uniformity, accounting for the differing lengths of the chromosomes.

Positions of differentially expressed genes on chromosome 8: to assess the probability of the observed hits on chromosome 8 being random, we divided the chromosome into 1 Mbp intervals and counted the number of hits per interval. We then used a chi-square statistic to compare this distribution of hit positions with what would be expected under a uniform distribution in which the hits were equally likely to occur at any point along chromosome 8. The reference distribution was calculated using simulation, with a uniform distribution to define random positions for hits along the length of chromosome 8.

Bioinformatic analysis

Bioinformatic analysis of 'passenger mutations' co-segregating with the targeted *Plat* allele was carried out using the Sanger 'Mouse Genomes Project – Query SNPs, indels or SVs' website (http://www.sanger.ac.uk/sanger/Mouse_SnpViewer/rel-1505) [27,28].

Results

Transcriptomic profiling of brains of two independently generated strains of congenic C57BL/6J-Plat^{-/-} *mice reveals anomalous clustering of differentially expressed genes around Plat on chromosome 8*

C57BL/6J-Plat^{tm1Mlg} mice [29] were housed at the Australian Centre for Blood Diseases, Monash University (hereafter termed Melbourne *C57BL/6J-Plat*^{-/-} mice). In 2009, when the expression analysis was carried out, these mice were estimated to be backcrossed to *C57BL/6J* mice for 13 generations. RNA was isolated from the brains of Melbourne *C57BL/6J-Plat*^{-/-} mice and from co-habiting

age- and sex-matched C57BL/6J mice and hybridized to whole genome expression arrays. Quadruplicate analysis was performed, with each sample representing pools of RNA from two mice. Overall, only 13 genes reporting in the array were identified as differentially expressed in the brain of C57BL/6J-*Plat*^{-/-} mice, when compared with the C57BL/6J control mice (Table 1). Surprisingly, 11 of these 13 genes (85%) were located on chromosome 8, which also harbors the *Plat* gene ($P < 0.000001$, see 'Materials and methods' for details on statistics). Furthermore, all of these 11 genes clustered around the *Plat* gene, on a chromosomal segment spanning approximately 19 Mbp upstream to 8 Mbp downstream of *Plat* (Fig. 1A). This clustering of differentially expressed genes within this 27 Mbp segment is also highly anomalous ($P < 0.000001$), considering that the total length of mouse chromosome 8 is approximately 129 Mbp and the total length of the mouse genome is more than 2600 Mbp.

We next set out to examine if the abnormal distribution of differentially expressed genes in Melbourne C57BL/6J-*Plat*^{-/-} mice was also observed in other congenic colonies of tPA-deficient mice. Specifically, we analyzed C57BL/6J-*Plat*^{tm1Mlg} mice housed at the University of Michigan (hereafter termed Michigan C57BL/6J-*Plat*^{-/-} mice), estimated to be backcrossed for > 10 generations to C57BL/6J in 2015 when the analysis was carried out. Brains of four Michigan C57BL/6J-*Plat*^{-/-} mice and four wild-type C57BL/6J mice from Jackson Laboratories were subjected to transcriptomic profiling by RNAseq. We again observed an abnormally high frequency of differentially expressed genes (eight of 32, 25%) located on chromosome 8 ($P < 0.003$) (Table 2). Furthermore, all eight

genes were located approximately 18 Mbp upstream to 5 Mbp downstream of *Plat* ($P < 0.000001$) (Fig. 1B).

Evidence for 129 origin of differentially expressed genes on chromosome 8 of tPA-deficient mice

To explore the basis of the anomalous clustering of differentially expressed genes around the *Plat* gene in the two independently backcrossed congenic strains of C57BL/6J-*Plat*^{-/-} mice, we next contracted Jackson Laboratories to perform SNP analysis of Michigan C57BL/6J-*Plat*^{-/-} mice and Melbourne C57BL/6J-*Plat*^{-/-} mice. We first employed a JAX 150 SNP Panel polymorphic between C57BL/6J and 129, which covers the 19 autosomes and the X chromosome with a density of ~15–20 Mbp. Strikingly, of 154 SNPs analyzed in Michigan C57BL/6J-*Plat*^{-/-} mice, only one was found to be 129 specific (rs3701395, Table S1) and this SNP was located 19.7 Mbp upstream of *Plat*, just upstream of a cluster of five differentially expressed genes. This SNP was also 129 specific in Melbourne C57BL/6J-*Plat*^{-/-} mice, as was the adjacent SNP (rs3701395 and rs3684251, Table S1). Melbourne C57BL/6J-*Plat*^{-/-} mice were also heterozygote for one chromosome 1 SNP (rs3697376, Table S1). Remarkably, this SNP was located very close to both the chromosome 1-located genes differentially expressed in Melbourne C57BL/6J-*Plat*^{-/-} mice (chromosome location-bp, rs3697376: 65042402. *C430010P07Rik*: 66719248-66817562. *Mtap2*: 66175273-66442583). Based on these findings, we next performed a high-density SNP analysis of chromosome 8, employing 34 SNPs, polymorphic between C57BL/6J and 129 (Fig. 2A and B and Table S2). In Michigan C57BL/6J-*Plat*^{-/-} mice, a continuous cluster of seven SNPs analyzed, located 19–0.3 Mbp upstream of *Plat*, were identified as being 129 specific (Fig. 2A). An identical cluster of 129-derived SNPs was found in Melbourne C57BL/6J-*Plat*^{-/-} mice, which also displayed two additional 129 SNPs located just downstream of *Plat* (Fig. 2B). Notably, the chromosomal segments identified in the SNP analysis as being 129 derived in the two mouse strains harbored most of the genes differentially expressed on chromosome 8 of the Michigan C57BL/6J-*Plat*^{-/-} mice and all of the identified genes differentially expressed in the Melbourne C57BL/6J-*Plat*^{-/-} mice (Fig. 2A and B, and Table S1).

Bioinformatic analysis identifies multiple passenger mutations in congenic C57BL/6J-*Plat*^{-/-} mice

Locating the specific regions of 129-derived DNA in Michigan C57BL/6J-*Plat*^{-/-} and Melbourne C57BL/6J-*Plat*^{-/-} mice allowed for the direct bioinformatic identification of 'passenger mutations' co-segregating with the targeted *Plat* allele by using the Sanger 'Mouse Genomes Project – Query SNPs, indels or SVs' website (http://www.sanger.ac.uk/sanger/Mouse_SnpViewer/rel-1505)

Table 1 Chromosomal location of genes differentially expressed in Melbourne C57BL/6J-*Plat*^{-/-} mice

Chromosome	Length (Mbp)	# Differentially expressed genes
1	195	2
2	182	0
3	160	0
4	157	0
5	152	0
6	150	0
7	145	0
8	129	11
9	125	0
10	131	0
11	122	0
12	120	0
13	120	0
14	125	0
15	104	0
16	91	0
17	95	0
18	91	0
19	61	0
X	161	0
Y	95	0




Fig. 1. The clustering of genes differentially expressed in brains of congenic tPA-deficient mice in the vicinity of *Plat* is unrelated to the loss of tPA. (A) RNA was extracted from congenic ‘Melbourne’ C57BL/6J-*Plat*^{-/-} mice and age- and gender-matched C57BL/6J wild-type controls, and whole-genome transcriptome analysis was performed by using expression arrays. The locations of *Plat* and of differentially expressed genes on chromosome 8 are shown. Eleven of 13 differentially expressed genes (excluding *Plat*) in tPA-deficient brains (85%) are located within a 27 Mbp interval, flanking *Plat*. (B) RNA was extracted from congenic ‘Michigan’ C57BL/6J-*Plat*^{-/-} mice and wild-type littermates and whole-genome transcriptome analysis was performed by RNAseq. The locations of *Plat* and of differentially expressed genes on chromosome 8 are shown. Eight of 32 differentially expressed genes (excluding *Plat*) in tPA-deficient brains (25%) are located within a 30 Mbp interval flanking *Plat*. (C) RNA was extracted from ‘Michigan’ C57BL/6J-*Plat*^{-/-} mice and ‘Bethesda’ C57BL/6J-*Plat*^{-/-} littermates and whole-genome transcriptome analysis was performed by RNAseq. The locations of *Plat* and of differentially expressed genes on chromosome 8 are shown. Seven of 19 differentially expressed genes (excluding *Plat*) (37%) are located within a 26 Mbp interval flanking *Plat*. (D) No anomalous clustering of differentially expressed genes in brains of isogenic ‘Bethesda’ C57BL/6J-*Plat*^{-/-} mice around *Plat* on chromosome 8. RNA was extracted from ‘Bethesda’ C57BL/6J-*Plat*^{-/-} mice and wild-type littermates and whole-genome transcriptome analysis was performed by RNAseq. The locations of *Plat* and of differentially expressed genes on chromosome 8 are shown.

[27,28]. Interestingly, the 129-derived regions of both Michigan C57BL/6J-*Plat*^{-/-} mice and Melbourne C57BL/6J-*Plat*^{-/-} mice harbored five protein-coding genes (*Cd209a*, *Cd209b*, *Ccl25*, *Defb34* and *Defb46*) predicted to be null in 129 mice and wild-type in C57BL/6J mice or *vice versa*, as evidenced by the presence of in-frame stop codons, frame shift-inducing insertions and deletions (Table 3). An additional eight genes (*Arhgef18*, *Cd209e*, *BC068157*, *Gm560*, *Tubgcp3*, *Mef2 l*, *Myom2* and *Mchp1*) harbored one or more mutations causing non-conservative amino acid substitutions. Owing to additional 129-derived regions upstream of *Plat*, Melbourne C57BL/6J-*Plat*^{-/-} mice also displayed mutations in *Adam3*.

C57BL/6J-Plat^{-/-} mice, generated by zinc finger nuclease targeting of C57BL/6J mice, do not show anomalous clustering of differentially expressed genes around *Plat*

We next used custom-designed zinc finger nucleases (Zfns) to generate tPA-deficient mice *de novo*. A Zfn was

engineered to introduce mutations into the signal peptide-encoding exon 2 of *Plat* (Fig. 3). mRNA encoding the Zfn was microinjected into fertilized C57BL/6J and FVB/NJ embryos that were implanted into pseudo-pregnant females. Analysis of born offspring identified a number of mice with mutations in exon 2 of *Plat*. This included C57BL/6J and FVB/NJ offspring with an identical 11 bp deletion in the signal peptide-coding exon 2, which were selected for further analysis. This deletion removes nucleotides 52–62 of the *Plat* open reading frame and introduces a frame shift mutation after Pro17 of the tPA signal peptide and a premature stop signal (Fig. 3A). This gave rise to a null allele, as confirmed by plasminogen-casein zymography (Fig. 3B), Western blot (Fig. 3C) and tPA activity assays (Fig. 3D) of brain tissue from mice bred to homozygosity for the targeted allele and their wild-type littermates. These novel mouse strains hereafter are termed, respectively, Bethesda C57BL/6J-*Plat*^{-/-} and Bethesda FVB/NJ-*Plat*^{-/-}. As expected, low-density SNP analysis of Bethesda C57BL/6J-*Plat*^{-/-} mice showed all

Table 2 Chromosomal location of genes differentially expressed in Michigan C57BL/6J-*Plat*^{-/-} mice

Chromosome	Length (Mbp)	# Differentially expressed genes
1	195	0
2	182	1
3	160	0
4	157	1
5	152	2
6	150	0
7	145	2
8	129	8
9	125	3
10	131	3
11	122	3
12	120	0
13	120	2
14	125	1
15	104	0
16	91	0
17	95	3
18	91	0
19	61	3
X	161	0

SNPs to be C57BL/6J derived (Table S1). High-density SNP analysis of chromosome 8 from Bethesda C57BL/6J-*Plat*^{-/-} mice (Table S2 and Fig. 2) showed all SNPs but one to be C57BL/6J specific. The single 129-specific SNP

(rs3709624, Table S2, indicated by red asterisk in Fig. 2) also genotyped as 129 specific in two C57BL/6J control mice purchased from Jackson Laboratories that were included in the analysis ('Control', columns L and M in Table S2), and we currently have no explanation for this finding.

We next set out to provide direct experimental evidence that the differential expression of chromosome 8 genes in brains of Melbourne and Michigan C57BL/6J-*Plat*^{-/-} mice was not a consequence of loss of tPA, but rather a result of retention of a 129-derived chromosomal segment. For this purpose, we interbred the new Bethesda C57BL/6J-*Plat*^{-/-} mice with Michigan C57BL/6J-*Plat*^{-/-} mice to generate littermate pairs of Bethesda C57BL/6J-*Plat*^{-/-} and Michigan C57BL/6J-*Plat*^{-/-} mice. These mice, both devoid of tPA, were then subjected to transcriptomic profiling by RNAseq using quadruplicate analysis. A conspicuous clustering of genes differentially expressed in Bethesda C57BL/6J-*Plat*^{-/-} mice and Michigan C57BL/6J-*Plat*^{-/-} littermates around chromosome 8 was again observed, with seven of 19 differentially expressed genes (37%) located on chromosome 8 ($P < 0.05$), clustering 19 Mbp upstream to 2.9 Mbp downstream from *Plat* ($P < 0.000001$) (Fig. 1C and Table 4). In sharp contrast, when the identical RNAseq analysis was performed comparing brains from Bethesda C57BL/6J-*Plat*^{-/-} mice with those from their wild-type littermates, no over-representation of differentially




Fig. 2. Congenic tPA-deficient mice contain large contiguous segments of 129-derived DNA. High-density single nucleotide polymorphism (SNP) analysis of congenic 'Michigan' C57BL/6J-*Plat*^{-/-} mice (A), congenic 'Melbourne' C57BL/6J-*Plat*^{-/-} mice (B) and isogenic 'Bethesda' C57BL/6J-*Plat*^{-/-} mice (C). Positions of SNPs specific for strain 129 mice vs. C57BL/6J mice are indicated with red asterisks, and the positions of SNPs specific for strain C57BL/6J mice vs. strain 129 mice are indicated with black circles.

Table 3 Passenger mutations associated with 129-DNA regions in 'Michigan' and 'Melbourne' *Plat*^{-/-} strains

Gene	Position on chr.8 (Mbp)	Distance from <i>Plat</i> (Mbp)	Coding region mutations in 129 vs. C57BL/6J			Mutated in Michigan	Mutated in Melbourne
			Stop gain/loss	Frame shift	Missense		
Arhgef18	3.40	19.30			T315M;S975P	+	+
Cd209a	3.74	18.96		After C64	R13G;T36P	+	+
Cd209e	3.85	18.85	W208Stop			+	+
Cd209b	3.92	18.78	R247Stop		C42S; I259V A175V; T369A; V389M; E428G; V439I; R524Q; P558L; T559A; S620P; P690S;	+	+
BC068157	4.21	18.49			F716L;P779T;S950A S82P; W86R; H116Y; E120K; P154S;	+	+
Ccl25	4.34	18.36		After P98	M169I	+	+
Tubgcp3	12.64	10.06			G190S	+	+
Mcf2l	12.95	9.75	Stop1102L			+	+
Myom2	15.10	7.60			R123Q;M131L;S234P;L977I; S1311N A300V; A303V; S469L; C580F; T615A;	+	+
McpH1	18.60	4.10			T615K	+	+
Defb34	19.12	3.58		After D25		+	+
Defb46	19.24	3.46	C32Stop		C21F	+	+
ENSMUSG00000097230	19.79	2.91				+	+
AY761185	20.94	1.76				+	+
Plat	22.70	0				N/A	N/A
ADAM3	24.70	2.00	Stop823S		G741D		+

N/A, not applicable.

expressed genes on chromosome 8 was found, with only two of 37 differentially expressed genes (5%) residing on chromosome 8 ($P = \text{NS}$), located, respectively, 50 and 61 Mbp from *Plat* (Fig. 1D and Table 5) ($P = \text{NS}$).

Discussion

This current study provides evidence that tPA-deficient mice highly backcrossed to C57BL/6J mice display differences in gene expression in the brain that are not related to the absence of tPA, but rather to the way the *Plat* gene was targeted. Two colonies of highly backcrossed C57BL/6J-*Plat*^{-/-} mice, made by targeting *Plat* 129-derived ES cells, showed the same anomalous clustering of differentially expressed genes in the vicinity of *Plat*, when independently analyzed by whole genome array analysis and RNAseq. In the two analyses, 86 and 25% of all differentially expressed genes were located within a segment of chromosome 8 roughly 19 Mbp upstream to 8 Mbp downstream of *Plat*. Considering the length of the mouse genome of more than 2600 Mbp, less than 1.5% of differentially expressed genes would be expected to be located on this short segment of chromosome 8, which constitutes just 1.03% of the mouse genome. Furthermore, an isogenic C57BL/6J-*Plat*^{-/-} mouse strain, generated in this study by Zfn-mediated genome editing in C57BL/6J mice, did not show this anomalous clustering of differentially

expressed genes. High-density SNP analysis showed that both Melbourne and Michigan C57BL/6J-*Plat*^{-/-} mice contained a surprisingly large contiguous 129-derived chromosomal segment, and we provide direct experimental evidence that the presence of this chromosomal segment is responsible for the anomalous clustering of differentially expressed genes in C57BL/6J-*Plat*^{-/-} mice.

Importantly, the 129-derived chromosomal segment that co-segregates with the targeted *Plat* allele displayed a number of differences in coding regions, when compared with the corresponding chromosomal segment of C57BL/6J mice, with at least four protein-coding genes potentially being null in the 129 mice and wild-type in C57BL/6J mice or *vice versa*, and eight additional genes displaying non-conservative amino acid substitutions. Notably, some of these genes have reported roles in human brain function: *ARHGGEF18* is involved in neurite retraction [30] and was linked in a genome-wide association study to sexual dysfunction in individuals treated for major depression [31]. *MCF2L* participates in the formation and stabilization of glutaminergic synapses of cortical neurons [32], and SNPs in *MCF2L* and *TUBGCP3* were identified as predictive for successful smoking cessation [33]. Homozygosity for mutations in *MCPHI* causes microcephaly [34], whereas homozygous loss of *ADAM3A* is associated with pediatric high-grade glioma and diffuse intrinsic pontine gliomas [35].




Fig. 3. Generation of isogenic C57BL/6J-*Plat*^{-/-} and FVB/NJ-*Plat*^{-/-} mice. (A) *De novo* generation of *Plat* null allele. Schematic structure of the proximal part of the mouse *Plat* gene (top), and the sequence of the signal peptide-encoding exon 2 (upper case letters) and flanking intron sequences (lower case letters) (bottom). The Zfn binding sites are in bold letters, with the Zfn cleavage site in red. The 11 bp deleted in the C57BL/6J and FVB/NJ strains carried forward for analysis are underlined. This deletion introduces a frame shift resulting in the production of a mRNA encoding amino acid 1-17 of tPA fused to a 19 amino acid nonsense peptide. (B) Plasminogen-casein zymography. Lanes 1-4: purified human tPA. Lanes 5-8: protein extracts from brains of a litter of FVB/NJ mice containing two wild-type (lanes 5 and 8) and 2 *Plat*^{-/-} littermates (lanes 6 and 7). (C) tPA Western blot. Lane 1: purified human tPA. Lanes 2-5: protein extracts from brains of a litter of mice containing two wild-type (lanes 2 and 5) and two *Plat*^{-/-} littermates (lanes 3 and 4). (D) Active tPA (middle column) and total tPA (right column) protein extracts from brains of a litter of mice containing two wild-type (top and bottom row) and two *Plat*^{-/-} littermates (middle rows).

Table 4 Chromosomal location of genes differentially expressed in Michigan and Bethesda C57BL/6J-*Plat*^{-/-} mice

Chromosome	Length (Mbp)	# Differentially expressed genes
1	195	2
2	182	2
3	160	0
4	157	0
5	152	0
6	150	1
7	145	2
8	129	7
9	125	0
10	131	1
11	122	1
12	120	1
13	120	1
14	125	0
15	104	0
16	91	0
17	95	1
18	91	0
19	61	0
X	161	0
Y	95	ND

ND, not determined.

Table 5 Chromosomal location of genes differentially expressed in Bethesda C57BL/6J-*Plat*^{-/-} mice

Chromosome	Length (Mbp)	# Differentially expressed genes
1	195	1
2	182	2
3	160	1
4	157	2
5	152	2
6	150	0
7	145	4
8	129	2
9	125	2
10	131	4
11	122	3
12	120	1
13	120	3
14	125	1
15	104	4
16	91	0
17	95	2
18	91	0
19	61	2
X	161	1
Y	95	ND

ND, not determined.

It should be noted that the 129-derived co-segregating chromosomal segments identified in the congenic C57BL/6J-*Plat*^{-/-} mouse strains have the potential to confound data interpretation irrespective of whether the wild-type mice used as controls are littermate-derived or age- and sex-matched C57BL/6J mice. Our study, thus, highlights the importance of reconstitution experiments in non-isogenic mice, where a wild-type phenotype can be restored via the introduction of exogenous tPA into tPA-deficient mice [11], or where tPA-overexpressing mice exhibit an opposite phenotype to that of tPA-deficient mice [36]. Similarly, with regards to plasmin-dependent functions of tPA, one can eliminate the confounding effects of co-inherited passenger mutations when tPA-deficient and plasminogen-deficient mice exhibit similar phenotypes [37]. It is important to stress, however, that the present study provides no evidence, directly or indirectly, that any of the published phenotypes in tPA-deficient mice are caused by 129-derived passenger mutations co-inherited with the targeted *Plat* allele. Nonetheless, it is also clear from the data presented in the current study that caution must be exerted when interpreting phenotypes observed in C57BL/6J-*Plat*^{-/-} mice. Related to this, it is interesting to note that this region of chromosome 8, which is syntenic with human chromosome 8, has been suggested to be a potential hub for genes associated with neuropsychiatric disorders and with cancer [38]. Undoubtedly, the C57BL/6J-*Plat*^{-/-} and FVB/NJ-*Plat*^{-/-} mice generated here should provide two novel mouse strains for the research community to further explore tPA function; strains which are free of co-inherited passenger mutations that are inevitable when using 129-derived ES cells. Nevertheless, the Zfn technology used to generate the two mouse strains is relatively new, and the degree to which 'off-targeting' of the germ line occurs has not been fully investigated [39,40]. Potential off-target mutations in the two novel mouse strains, however, would only constitute a potential long-term problem if located in proximity to *Plat*.

It should also be stressed that in conditionally-targeted strains, the potential of passenger mutations to confound data interpretation can largely be eliminated by the use of appropriate littermate controls, including Cre-negative littermates that carry two conditionally targeted alleles and, therefore, are expected to carry the same set of passenger mutations near the gene of interest [2].

The plasminogen activation system was among the first proteolytic systems to be analyzed by gene targeting in mice. Consequently, targeting of genes encoding most components of the system was carried out using 129-derived ES cells. This includes, besides *Plat*, *Plau*[29], *Plaur*[41,42], *Plg*[43,44], *Serpine1*[45], *SerpinF2*[46] and *Annat2*[47]. In light of the speed, efficiency and low cost of current genome editing technologies, the regeneration of isogenic strains carrying null mutations in these genes should be considered.

Addendum

A. L. Samson, D. A. Lawrence, T. H. Bugge, R. L. Medcalf and R. Szabo conceived and designed experiments. D. A. Lawrence and R. Szabo generated and analyzed *Plat*^{-/-} mice. A. L. Samson, T. H. Bugge, R. L. Medcalf and R. Szabo performed array analysis and analyzed data. A. L. Samson, D. A. Lawrence and T. H. Bugge performed RNAseq and analyzed data. T. H. Bugge wrote paper.

Acknowledgements

We thank K. Mann and M. Warnock for excellent technical assistance, R. McEachin of the University of Michigan Bioinformatics Core for Bioinformatics analysis of the RNAseq data, K. Shedden of the University of Michigan Center for Statistical Consultation and Research (CSCAR) for statistical analysis, and M. J. Danton for critically reviewing this manuscript. This study was supported by the NIDCR Intramural Research Program (T.H.B) and National Institutes of Health Grants HL055374 and NS079639 (D. A. L.). Bethesda C57BL/6J-*Plat*^{-/-} mice and Bethesda FVB/NJ-*Plat*^{-/-} mice can be obtained by contacting Thomas Bugge (Thomas.bugge@nih.gov).

Disclosure of Conflict of Interests

The authors state that they have no conflict of interest.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Table S1 Low-density single nucleotide polymorphism (SNP) analysis of all mouse chromosomes.

Table S2 High-density single nucleotide polymorphism (SNP) analysis of mouse chromosome 8.

References

- 1 Bronson SK, Smithies O. Altering mice by homologous recombination using embryonic stem cells. *J Biol Chem* 1994; **269**: 27155–8.
- 2 Vanden Berghe T, Hulpiau P, Martens L, Vandenbroucke RE, van Wouwerghem E, Perry SW, Bruggeman I, Divert T, Choi SM, Vuylsteke M, Shestopalov VI, Libert C, Vandenabeele P. Passenger mutations confound interpretation of all genetically modified congenic mice. *Immunity* 2015; **43**: 200–9.
- 3 Kayagaki N, Warming S, Lamkanfi M, Vande Walle L, Louie S, Dong J, Newton K, Qu Y, Liu J, Heldens S, Zhang J, Lee WP, Roose-Girma M, Dixit VM. Non-canonical inflammasome activation targets caspase-11. *Nature* 2011; **479**: 117–21.
- 4 Kenneth NS, Younger JM, Hughes ED, Marcotte D, Barker PA, Saunders TL, Duckett CS. An inactivating caspase 11

- passenger mutation originating from the 129 murine strain in mice targeted for c-IAP1. *Biochem J* 2012; **443**: 355–9.
- 5 Collen D, Lijnen HR. The fibrinolytic system in man. *Crit Rev Oncol Hematol* 1986; **4**: 249–301.
 - 6 Soreq H, Miskin R. Plasminogen activator in the rodent brain. *Brain Res* 1981; **216**: 361–74.
 - 7 Krystosek A, Seeds NW. Plasminogen activator secretion by granule neurons in cultures of developing cerebellum. *Proc Natl Acad Sci U S A* 1981; **78**: 7810–4.
 - 8 Qian Z, Gilbert ME, Colicos MA, Kandel ER, Kuhl D. Tissue-plasminogen activator is induced as an immediate-early gene during seizure, kindling and long-term potentiation. *Nature* 1993; **361**: 453–7.
 - 9 Sappino AP, Madani R, Huarte J, Belin D, Kiss JZ, Wohlwend A, Vassalli JD. Extracellular proteolysis in the adult murine brain. *J Clin Invest* 1993; **92**: 679–85.
 - 10 Salles FJ, Strickland S. Localization and regulation of the tissue plasminogen activator-plasmin system in the hippocampus. *J Neurosci* 2002; **22**: 2125–34.
 - 11 Tsirka SE, Gualandris A, Amaral DG, Strickland S. Excitotoxin-induced neuronal degeneration and seizure are mediated by tissue plasminogen activator. *Nature* 1995; **377**: 340–4.
 - 12 Siddiq MM, Tsirka SE. Modulation of zinc toxicity by tissue plasminogen activator. *Mol Cell Neurosci* 2004; **25**: 162–71.
 - 13 Wang YF, Tsirka SE, Strickland S, Stieg PE, Soriano SG, Lipton SA. Tissue plasminogen activator (tPA) increases neuronal damage after focal cerebral ischemia in wild-type and tPA-deficient mice. *Nat Med* 1998; **4**: 228–31.
 - 14 Frey U, Muller M, Kuhl D. A different form of long-lasting potentiation revealed in tissue plasminogen activator mutant mice. *J Neurosci* 1996; **16**: 2057–63.
 - 15 Huang YY, Bach ME, Lipp HP, Zhuo M, Wolfer DP, Hawkins RD, Schoonjans L, Kandel ER, Godfraind JM, Mulligan R, Collen D, Carmeliet P. Mice lacking the gene encoding tissue-type plasminogen activator show a selective interference with late-phase long-term potentiation in both Schaffer collateral and mossy fiber pathways. *Proc Natl Acad Sci U S A* 1996; **93**: 8699–704.
 - 16 Medina MG, Ledesma MD, Dominguez JE, Medina M, Zafra D, Alameda F, Dotti CG, Navarro P. Tissue plasminogen activator mediates amyloid-induced neurotoxicity via Erk1/2 activation. *EMBO J* 2005; **24**: 1706–16.
 - 17 Matys T, Pawlak R, Matys E, Pavlides C, McEwen BS, Strickland S. Tissue plasminogen activator promotes the effects of corticotropin-releasing factor on the amygdala and anxiety-like behavior. *Proc Natl Acad Sci U S A* 2004; **101**: 16345–50.
 - 18 Pawlak R, Magarinos AM, Melchor J, McEwen B, Strickland S. Tissue plasminogen activator in the amygdala is critical for stress-induced anxiety-like behavior. *Nat Neurosci* 2003; **6**: 168–74.
 - 19 Nagai T, Yamada K, Yoshimura M, Ishikawa K, Miyamoto Y, Hashimoto K, Noda Y, Nitta A, Nabeshima T. The tissue plasminogen activator-plasmin system participates in the rewarding effect of morphine by regulating dopamine release. *Proc Natl Acad Sci U S A* 2004; **101**: 3650–5.
 - 20 Maiya R, Zhou Y, Norris EH, Kreek MJ, Strickland S. Tissue plasminogen activator modulates the cellular and behavioral response to cocaine. *Proc Natl Acad Sci U S A* 2009; **106**: 1983–8.
 - 21 Noel M, Norris EH, Strickland S. Tissue plasminogen activator is required for the development of fetal alcohol syndrome in mice. *Proc Natl Acad Sci U S A* 2011; **108**: 5069–74.
 - 22 Yepes M, Sandkvist M, Moore EG, Bugge TH, Strickland DK, Lawrence DA. Tissue-type plasminogen activator induces opening of the blood-brain barrier via the LDL receptor-related protein. *J Clin Invest* 2003; **112**: 1533–40.
 - 23 Seeds NW, Basham ME, Haffke SP. Neuronal migration is retarded in mice lacking the tissue plasminogen activator gene. *Proc Natl Acad Sci U S A* 1999; **96**: 14118–23.
 - 24 Nicole O, Docagne F, Ali C, Margail I, Carmeliet P, MacKenzie ET, Vivien D, Buisson A. The proteolytic activity of tissue-plasminogen activator enhances NMDA receptor-mediated signaling. *Nat Med* 2001; **7**: 59–64.
 - 25 Pang PT, Teng HK, Zaitsev E, Woo NT, Sakata K, Zhen S, Teng KK, Yung WH, Hempstead BL, Lu B. Cleavage of proBDNF by tPA/plasmin is essential for long-term hippocampal plasticity. *Science* 2004; **306**: 487–91.
 - 26 Sawaya R, Highsmith R. Plasminogen activator activity and molecular weight patterns in human brain tumors. *J Neurosurg* 1988; **68**: 73–9.
 - 27 Yalcin B, Wong K, Agam A, Goodson M, Keane TM, Gan X, Nellaker C, Goodstadt L, Nicod J, Bhomra A, Hernandez-Pliego P, Whitley H, Cleak J, Dutton R, Janowitz D, Mott R, Adams DJ, Flint J. Sequence-based characterization of structural variation in the mouse genome. *Nature* 2011; **477**: 326–9.
 - 28 Keane TM, Goodstadt L, Danecek P, White MA, Wong K, Yalcin B, Heger A, Agam A, Slater G, Goodson M, Furlotte NA, Eskin E, Nellaker C, Whitley H, Cleak J, Janowitz D, Hernandez-Pliego P, Edwards A, Belgard TG, Oliver PL, *et al.* Mouse genomic variation and its effect on phenotypes and gene regulation. *Nature* 2011; **477**: 289–94.
 - 29 Carmeliet P, Schoonjans L, Kieckens L, Ream B, Degen J, Bronson R, De Vos R, van den Oord JJ, Collen D, Mulligan RC. Physiological consequences of loss of plasminogen activator gene function in mice. *Nature* 1994; **368**: 419–24.
 - 30 Tsuji T, Ohta Y, Hirose K, Ohashi K, Mizuno K. Involvement of p114-RhoGEF and Lfc in Wnt-3a- and dishevelled-induced RhoA activation and neurite retraction in N1E-115 mouse neuroblastoma cells. *Mol Biol Cell* 2010; **21**: 3590–600.
 - 31 Kurose K, Hiratsuka K, Ishiwata K, Nishikawa J, Nonen S, Azuma J, Kato M, Wakeno M, Okugawa G, Kinoshita T, Kurosawa T, Hasegawa R, Saito Y. Genome-wide association study of SSRI/SNRI-induced sexual dysfunction in a Japanese cohort with major depression. *Psychiatry Res* 2012; **198**: 424–9.
 - 32 Hayashi T, Yoshida T, Ra M, Taguchi R, Mishina M. IL1RAPL1 associated with mental retardation and autism regulates the formation and stabilization of glutamatergic synapses of cortical neurons through RhoA signaling pathway. *PLoS ONE* 2013; **8**: e66254.
 - 33 Rose JE, Behm FM, Drgon T, Johnson C, Uhl GR. Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. *Mol Med* 2010; **16**: 247–53.
 - 34 Jackson AP, Eastwood H, Bell SM, Adu J, Toomes C, Carr IM, Roberts E, Hampshire DJ, Crow YJ, Mighell AJ, Karbani G, Jafri H, Rashid Y, Mueller RF, Markham AF, Woods CG. Identification of microcephalin, a protein implicated in determining the size of the human brain. *Am J Hum Genet* 2002; **71**: 136–42.
 - 35 Barrow J, Adamowicz-Brice M, Cartmill M, MacArthur D, Lowe J, Robson K, Brundler MA, Walker DA, Coyle B, Grundy R. Homozygous loss of ADAM3A revealed by genome-wide analysis of pediatric high-grade glioma and diffuse intrinsic pontine gliomas. *Neuro Oncol* 2011; **13**: 212–22.
 - 36 Sashindranath M, Sales E, Daglas M, Freeman R, Samson AL, Cops EJ, Beckham S, Galle A, McLean C, Morganti-Kossmann C, Rosenfeld JV, Madani R, Vassalli JD, Su EJ, Lawrence DA, Medcalf RL. The tissue-type plasminogen activator-plasminogen activator inhibitor 1 complex promotes neurovascular injury in brain trauma: evidence from mice and humans. *Brain* 2012; **135**: 3251–64.

- 37 Tsirka SE, Bugge TH, Degen JL, Strickland S. Neuronal death in the central nervous system demonstrates a non-fibrin substrate for plasmin. *Proc Natl Acad Sci U S A* 1997; **94**: 9779–81.
- 38 Tabares-Seisdedos R, Rubenstein JL. Chromosome 8p as a potential hub for developmental neuropsychiatric disorders: implications for schizophrenia, autism and cancer. *Mol Psychiatry* 2009; **14**: 563–89.
- 39 Gabriel R, Lombardo A, Arens A, Miller JC, Genovese P, Kaeppel C, Nowrouzi A, Bartholomae CC, Wang J, Friedman G, Holmes MC, Gregory PD, Glimm H, Schmidt M, Naldini L, von Kalle C. An unbiased genome-wide analysis of zinc-finger nuclease specificity. *Nat Biotechnol* 2011; **29**: 816–23.
- 40 Pattanayak V, Ramirez CL, Joung JK, Liu DR. Revealing off-target cleavage specificities of zinc-finger nucleases by *in vitro* selection. *Nat Methods* 2011; **8**: 765–70.
- 41 Bugge TH, Suh TT, Flick MJ, Daugherty CC, Romer J, Solberg H, Ellis V, Dano K, Degen JL. The receptor for urokinase-type plasminogen activator is not essential for mouse development or fertility. *J Biol Chem* 1995; **270**: 16886–94.
- 42 Dewerchin M, Nuffelen AV, Wallays G, Bouche A, Moons L, Carmeliet P, Mulligan RC, Collen D. Generation and characterization of urokinase receptor-deficient mice. *J Clin Invest* 1996; **97**: 870–8.
- 43 Bugge TH, Flick MJ, Daugherty CC, Degen JL. Plasminogen deficiency causes severe thrombosis but is compatible with development and reproduction. *Genes Dev* 1995; **9**: 794–807.
- 44 Ploplis VA, Carmeliet P, Vazirzadeh S, van Vlaenderen I, Moons L, Plow EF, Collen D. Effects of disruption of the plasminogen gene on thrombosis, growth, and health in mice. *Circulation* 1995; **92**: 2585–93.
- 45 Carmeliet P, Kieckens L, Schoonjans L, Ream B, van Nuffelen A, Prendergast G, Cole M, Bronson R, Collen D, Mulligan RC. Plasminogen activator inhibitor-1 gene-deficient mice. I. Generation by homologous recombination and characterization. *J Clin Invest* 1993; **92**: 2746–55.
- 46 Lijnen HR, Okada K, Matsuo O, Collen D, Dewerchin M. Alpha2-antiplasmin gene deficiency in mice is associated with enhanced fibrinolytic potential without overt bleeding. *Blood* 1999; **93**: 2274–81.
- 47 Ling Q, Jacovina AT, Deora A, Febbraio M, Simantov R, Silverstein RL, Hempstead B, Mark WH, Hajjar KA. Annexin II regulates fibrin homeostasis and neoangiogenesis *in vivo*. *J Clin Invest* 2004; **113**: 38–48.