

# Anthropogenic N deposition increases soil organic matter accumulation without altering its biochemical composition

DONALD R. ZAK<sup>1,2</sup>, ZACHARY B. FREEDMAN<sup>1</sup>, RIMA A. UPCHURCH<sup>1</sup>, MARKUS STEFFENS<sup>3</sup> and INGRID KÖGEL-KNABNER<sup>3</sup>

<sup>1</sup>School of Natural Resources & Environment, University of Michigan, Ann Arbor, MI, USA, <sup>2</sup>Department of Ecology & Evolutionary Biology, University of Michigan, Ann Arbor, MI, USA, <sup>3</sup>Lehrstuhl für Bodenkunde, Department für Ökologie und Ökosystemmanagement, Wissenschaftszentrum Weihenstephan für Ernährung, Landnutzung und Umwelt, Technische Universität München, Freising-Weihenstephan 85350, Germany

## Abstract

Accumulating evidence indicates that future rates of atmospheric N deposition have the potential to increase soil C storage by reducing the decay of plant litter and soil organic matter (SOM). Although the microbial mechanism underlying this response is not well understood, a decline in decay could alter the amount, as well as biochemical composition of SOM. Here, we used size-density fractionation and solid-state <sup>13</sup>C-NMR spectroscopy to explore the extent to which declines in microbial decay in a long-term (*ca.* 20 yrs.) N deposition experiment have altered the biochemical composition of forest floor, bulk mineral soil, as well as free and occluded particulate organic matter. Significant amounts of organic matter have accumulated in occluded particulate organic matter (~20%; oPOM); however, experimental N deposition had not altered the abundance of carboxyl, aryl, alkyl, or O/N-alkyl C in forest floor, bulk mineral soil, or any soil fraction. These observations suggest that biochemically equivalent organic matter has accumulated in oPOM at a greater rate under experimental N deposition, relative to the ambient treatment. Although we do not understand the process by which experimental N deposition has fostered the occlusion of organic matter by mineral soil particles, our results highlight the importance of interactions among the products of microbial decay and the chemical and physical properties of silt and clay particles that occlude organic matter from microbial attack. Because oPOM can reside in soils for decades to centuries, organic matter accumulating under future rates of anthropogenic N deposition could remain in soil for long periods of time. If temperate forest soils in the Northern Hemisphere respond like those in our experiment, then unabated deposition of anthropogenic N from the atmosphere has the potential to foster greater soil C storage, especially in fine-texture forest soils.

**Keywords:** <sup>13</sup>C-NMR, anthropogenic N deposition, particulate organic matter, soil C storage, soil organic matter

Received 31 March 2016; revised version received 13 July 2016 and accepted 25 July 2016

## Introduction

By the end of this century, terrestrial ecosystems in eastern North America, as well as in other regions in the Northern Hemisphere, will receive quantities of biologically available nitrogen (N) that are unprecedented over the history of life on Earth (Lamarque *et al.*, 2005). This aspect of global change has the potential to constrain the accumulation of anthropogenic CO<sub>2</sub> in the Earth's atmosphere by increasing ecosystem C storage, thereby slowing the pace of climate warming. Increased net primary productivity, fostered by the anthropogenic supply of growth-limiting N to plants, was thought to be one mechanism that could elicit this

global biogeochemical response, although the extent of this mechanism is uncertain (Nadelhoffer *et al.*, 1999; Magnani *et al.*, 2007). Furthermore, a growing body of evidence reveals that anthropogenic N deposition can slow the microbial decay of plant detritus and increase soil C storage across a wide range of terrestrial ecosystems (Liu & Greaver, 2010; Frey *et al.*, 2015; Maaroufi *et al.*, 2015). However, the molecular and microbial mechanisms underlying this biogeochemical response are not well understood, and they are not a component of any coupled climate-biogeochemical model estimating ecosystem C storage (Thornton *et al.*, 2009; Burd *et al.*, 2016).

One plausible mechanism for reduced microbial decay under increased N deposition is a shift from basidiomycete fungal activity, some of which oxidize lignin in plant detritus and polyphenols in soil organic

Correspondence: Donald R. Zak, tel. 734 763 4991, fax 734 936 2195, e-mail: drzak@umich.edu

matter (SOM) to CO<sub>2</sub>, to a greater degree of metabolism by bacteria and ascomycete fungi, which only partially oxidize these organic substrates (Zak *et al.*, 2011; Freedman & Zak, 2014). Such a response should increase the amount of organic matter stored in soil, as well as alter its biochemical composition. In a widespread forest ecosystem in eastern North America, we have accumulated evidence consistent with many aspects of this microbial response to anthropogenic N deposition. For example, forest floor turnover time, an integrated measure of microbial decay, has significantly increased under experimental N deposition (+60%), while organic matter has rapidly accumulated in both forest floor (+51%) and surface mineral soil (+18%; Table 1). Additionally, extracellular phenol oxidase (−81%) and peroxidase (−30%) enzyme activities have consistently declined under experimental N deposition, whereas the production and leaching of phenolic DOC has substantially increased (+26%; Table 1). Although experimental N deposition has not altered the abundance of fungal laccase genes that encode the phenol oxidase enzyme (Table 1), it has reduced their expression by 50% (Table 1). In contrast, bacterial laccase-like multicopper oxidase (LMCO) genes, which weakly oxidize

polyphenols in plant detritus and SOM, were 26% more abundant in forest floor exposed to experimental N deposition (Freedman & Zak, 2014), suggesting that bacterial metabolism of these substrates is more prevalent. In combination, these observations are consistent with the proposed mechanism that experimental N deposition can diminish the complete metabolism of lignin in plant detritus and other polyphenols in SOM by soil fungi, while increasing their incomplete metabolism by soil bacteria and ascomycete fungi (Freedman & Zak, 2014; Freedman *et al.*, 2016).

We initially hypothesized that such a response should both increase the amount and alter the biochemical composition of SOM produced under experimental N deposition (Thomas *et al.*, 2012). This shift from complete metabolism of plant detritus to CO<sub>2</sub> by fungi to incomplete decay by bacteria and ascomycete fungi impacts not only the amount, but should also alter the biochemical composition of SOM (Thomas *et al.*, 2012; Freedman & Zak, 2014). Although the greater production of phenolic DOC under experimental N deposition is consistent with this expectation (Table 1; Pregitzer *et al.*, 2004), biochemical analyses (i.e., alkaline cupric oxide oxidation) of forest floor and mineral soil

**Table 1** Two decades of experimental N deposition have increased the storage and reduced the cycling of C in forest floor and surface mineral soil. Microbial responses are summarized for forest floor (Oe/Oa); mineral soil has exhibited similar directional trends. With the exception of laccase gene copy number, all of positive and negative responses are statistically significant ( $P < 0.05$ )


Response to chronic N deposition	% change	Citation
<b>Plant responses</b>		
Net primary production (NPP)	+10%	Pregitzer <i>et al.</i> (2008)
Leaf litter production	0%	Pregitzer <i>et al.</i> (2008)
Leaf litter N concentration	+25%	Xia <i>et al.</i> (2015)
Fine root litter	0%	Burton <i>et al.</i> (2004)
Fine root lignin concentration	0%	Xia <i>et al.</i> (2015)
Fine root N concentration	+8%	Xia <i>et al.</i> (2015)
Fine root respiration	0%	Burton <i>et al.</i> (2004)
AM fungal abundance (Soil)	−41%	Van Diepen <i>et al.</i> (2010)
Whole root system respiration	0%	Burton <i>et al.</i> (2012)
<b>Biogeochemical responses</b>		
Forest floor mass	+51%	Zak <i>et al.</i> (2008)
Forest floor turnover time	+60%	Zak <i>et al.</i> (2008)
Soil organic matter content	+18%	Zak <i>et al.</i> (2008)
Soil solution NO <sub>3</sub> <sup>−</sup> concentration	+288%	Zak <i>et al.</i> (2008)
NO <sub>3</sub> <sup>−</sup> leaching	+680%	Pregitzer <i>et al.</i> (2004)
DOC leaching	+26%	Pregitzer <i>et al.</i> (2004)
<b>Microbial responses</b>		
Soil respiration	−15%	Burton <i>et al.</i> (2004)
Active microbial biomass (PLFA)	−23%	DeForest <i>et al.</i> (2004)
Phenol oxidase activity	−81%	Freedman & Zak, (2014)
Peroxidase activity	−30%	DeForest <i>et al.</i> (2005)
Laccase copy number	−5 to −8%	Hassett <i>et al.</i> (2009)
Laccase expression	−50%	Edwards <i>et al.</i> (2011)
Bacterial LMCO copy number	+26%	Freedman & Zak (2014)

revealed that neither the amount of lignin-derived molecules nor their degree of oxidation have been altered by experimental N deposition (Thomas *et al.*, 2012); these prior observations did not support our hypothesis. However, it is plausible that other types of plant- and microbial-derived organic compounds (Amelung *et al.*, 2008; Kleber, 2010; Schmidt *et al.*, 2011) have accumulated under experimental N deposition due to the changes in microbial metabolism that we have documented above. As an alternative hypothesis, it also is plausible that experimental N deposition has altered the saprotrophic microbial community in a way that reduces the extent of decay (*sensu* Whittinghill *et al.*, 2012) without causing significant changes in the biochemical composition of the accumulating organic matter, possibly due to physical occlusion by silt and clay particles (Kleber *et al.*, 2015). To further test our initial hypothesis, as well as its alternative, we used solid-state  $^{13}\text{C}$ -NMR spectroscopy to characterize the degree of polyphenol oxidation in forest floor and SOM, as well as to characterize the abundance of other bonds that give insight into the biochemical composition of SOM (Baldock *et al.*, 1997). We further used this approach on free (fPOM) and occluded (oPOM) particulate organic matter in mineral soil to test our alternative hypothesis. Resolving our initial hypothesis from its alternative has direct implications for the amount and longevity of SOM accumulating under experimental N deposition and hence the extent to which these soils will function as future sinks for anthropogenic  $\text{CO}_2$  accumulating in the Earth's atmosphere.

## Materials and methods

### Study sites and sampling

Beginning in 1994, we have experimentally increased atmospheric  $\text{NO}_3^-$  deposition in a series of replicate northern hardwood forest stands spanning a 500-km distance (Fig. 1). Our study deliberately spans the north–south geographic range of the sugar maple-dominated (*Acer saccharum* Marsh.) northern hardwood forests of the Upper Great Lakes and eastern North America (Braun, 1950), enabling us to generalize our experimental results across this widespread and ecologically important ecosystem. Sites are floristically and edaphically matched (> 80% sugar maple on sandy Typic Haplorthods; see Burton *et al.*, 1991), but they differ in climate along the north–south latitudinal gradient (Fig. 1). The study sites also span a gradient of atmospheric N deposition, of which  $\text{NO}_3^-$  composes ~60% of wet-plus-dry N deposition. Ambient N deposition ranges from 6.8 kg N  $\text{ha}^{-1} \text{yr}^{-1}$  at Site A to 11.8 kg N  $\text{ha}^{-1} \text{yr}^{-1}$  at Site D. Soils in all sites are sandy, structureless (~85% sand) spodosols, with a well-developed Oe/Oa horizon (i.e., forest floor) that is permeated by a dense mat of fine roots. There are six 30-m  $\times$  30-m plots at each site, and every



**Fig. 1** Location of northern hardwood forest study sites in lower and upper Michigan, USA. Three plots (30 m  $\times$  30 m) in each stand receive ambient atmospheric N deposition, and three plots (30 m  $\times$  30 m) receive ambient plus 30 kg N  $\text{ha}^{-1} \text{yr}^{-1}$  of experimental N deposition. Treatments have been applied since 1994.

plot is surrounded on all sides by a 10-m-wide treated buffer. Three plots at each site receive ambient atmospheric N deposition; whereas the other three plots at each site receive ambient N deposition plus 30 kg  $\text{NO}_3^- \text{N} \text{ha}^{-1} \text{yr}^{-1}$ , a rate expected by 2050 across some portions of eastern North America (Galloway *et al.*, 2004). The additional  $\text{NO}_3^-$  is delivered over the growing season in six equal applications (5 kg N  $\text{ha}^{-1} \text{month}^{-1}$ ) of solid  $\text{NaNO}_3$  pellets, which are broadcast on the forest floor.

Forest floor and mineral soil sampling occurred in late May to early June 2013. In this way, samples from all four sites were collected during a phenologically similar period of decay. Within each 30-m by 30-m plot, 10 random 0.1-m by 0.1-m forest floor samples (Oe/Oa horizons; ~4 cm thick) were collected by hand to the mineral soil surface, after removing the Oi horizon. All forest floor samples were composited within each plot and homogenized by hand in the field. Mineral soil samples were collected directly beneath forest floor samples, using a 2.5-cm-diameter core extending to a depth of 10 cm. Mineral soil samples were passed through a 2-mm sieve in the field and also were composited within each plot. All samples were kept on ice during transport to the University of Michigan, where they were stored at  $-20^\circ\text{C}$ . After removing any fine roots, the entire air-dried, composite forest floor sample from each plot was ground using a ball mill; mineral soil samples were not ground. Air-dried forest floor and mineral soil samples then were shipped directly to the Technische Universität München for organic matter fractionation and biochemical analyses.

### Organic matter fractionation

We partitioned organic matter into five fractions based on density and size (Mueller & Kögel-Knabner, 2009). Air-dried

mineral soil samples (30 g; <2 mm) were saturated with sodium polytungstate ( $1.8 \text{ g cm}^{-3}$ ) and allowed to settle for 24 h. Free-floating particulate organic matter (fPOM) was removed from the solution surface by suction, and the remaining material was dispersed via ultrasonication at  $440 \text{ J mL}^{-1}$ . Occluded POM (oPOM) that was suspended in the sodium polytungstate was then separated from the mineral residue by centrifugation (30 min at  $3074 \text{ g}$ ). Sodium polytungstate was removed from fPOM and oPOM by washing each fraction with deionized water over a  $20\text{-}\mu\text{m}$  sieve, thereby sorting each POM fraction into material  $>20 \mu\text{m}$  and  $<20 \mu\text{m}$ . The aforementioned protocol created five organic matter fractions based on density ( $<1.8 \text{ g cm}^{-3}$ ) and size: fPOM  $>20 \mu\text{m}$ , fPOM  $<20 \mu\text{m}$ , oPOM  $>20 \mu\text{m}$ , and oPOM  $<20 \mu\text{m}$ ; the mineral residue fraction was  $>1.8 \text{ g cm}^{-3}$ . These fractions differ in their extent of decay, with fPOM  $>20 \mu\text{m}$  exhibiting the least modification by microbial decay and oPOM  $<20 \mu\text{m}$  the greatest, insights gained by  $^{14}\text{C}$  dating as well as the C : N of these fractions (von Lützow *et al.*, 2007). The mass of POM fractions ( $<1.8 \text{ g cm}^{-3}$ ) as well as organic matter remaining ( $>1.8 \text{ g cm}^{-3}$ ) in mineral soil was quantified, and the C and N content of each fraction was measured using a Vario EL elemental analyser (Elementar Analysensysteme, Hanau, Germany). From these data, we calculated the contribution of organic C and N in each fraction (Table S1).

#### Solid-state $^{13}\text{C}$ -NMR spectroscopy

To characterize the biochemical composition of forest floor and organic matter residing in mineral soil fractions, we employed solid-state  $^{13}\text{C}$ -NMR spectroscopy to determine the relative abundance of carboxyl, aryl, O/N-alkyl, and alkyl C (Baldock *et al.*, 1997). By doing so, we could directly and simultaneously test our hypothesis and its alternative. The cross-polarization magic angle spinning (CPMAS) technique was applied with a  $^{13}\text{C}$ -resonance frequency of 50.32 MHz and a spinning speed of 6.8 kHz using a Bruker DSX 200 NMR spectrometer (Karlsruhe, Germany; *sensu* Steffens *et al.*, 2009, 2011). A ramped  $^1\text{H}$ -pulse starting at 100% and decreasing to 50% of the initial power was used during a contact time of 1 ms in order to circumvent spin modulation during the Hartmann-Hahn contact. Pulse delays between 200 and 600 ms were used for all spectra. Depending on the C contents of the samples, between 2000 and 250 000 scans were accumulated, and a line broadening between 0 and 50 Hz was applied. The  $^{13}\text{C}$  chemical shifts were calibrated relative to tetramethylsilane (0 ppm). Relative contributions of the various C types were determined by integration of the signal intensity in their respective chemical shift regions according to Knicker *et al.* (2005). The region from 220 to 160 ppm was assigned to aldehyde, ketone, carboxyl, and amide bonds (i.e., carboxyl C). Olefinic and aromatic C were detected between 160 and 110 ppm and are noted as aryl C. Additionally, O-alkyl and N-alkyl C signals were found from 110 to 60 ppm and from 60 to 45 ppm are designated as O/N-alkyl C. Resonances of alkyl C were assigned to the region 45 to  $-10$  ppm. We calculated alkyl to O/N-alkyl ratio for forest floor and mineral soil fractions (Baldock *et al.*, 1997), because it is a

sensitive index for the extent of decay wherein a high ratio is indicative of a lower extent of decay.


#### Statistical analyses

To test our hypothesis, we used two-way ANOVA with a site by treatment interaction to compare the mean abundance of carboxyl, aryl, alkyl, O/N-alkyl C, as well as the alkyl to O/N-alkyl ratio in both forest floor and bulk mineral soil. We also used the same two-way ANOVA model to test our alternative hypothesis by comparing the mass, carbon concentration, nitrogen concentration, C : N, and abundance of the aforementioned organic bonds in both particulate and occluded organic matter fractions. To determine the extent of decay among fPOM and oPOM size fractions, we compared their alkyl to O/N-alkyl ratios as well as C : N ratios using a three-way ANOVA, with POM fraction, site and treatment as main effects. Means were compared using a Fisher's protected LSD, and statistical significance was accepted at  $\alpha = 0.05$ .

#### Results

Experimental N deposition did not significantly alter the abundance of carboxyl, aryl, O/N-alkyl, or alkyl C in forest floor (Fig. 2). Sites (main effect) differed from one another in the abundance of carboxyl and aryl C in forest floor, but the abundance of O/N-alkyl and alkyl C did not differ among them ( $P = 0.10\text{--}0.80$ ; data not shown). Moreover, we observed no interaction between site and treatment on the abundance of any C bond type in forest floor. In bulk mineral soil, although the abundance of carboxyl C in organic matter was significantly lower under experimental N deposition ( $P = 0.02$ ; Fig. 2), we found no difference in the proportion of aryl, O/N-alkyl, or alkyl C in mineral soil exposed to ambient and experimental N deposition (Fig. 2). Site was not a significant main factor in our analysis of bulk mineral soil, and we also observed no interaction between site and treatment. Further, experimental N deposition had no effect on the alkyl to O/N-alkyl C ratio in either forest floor (0.26 vs. 0.26; ambient vs. experimental N deposition, respectively) or mineral soil (0.59 vs. 0.60), indicating a similar degree of decay in both treatments. Taken together, these observations do not support our hypothesis that a reduction of microbial decay and oxidative enzyme activity has altered the biochemical composition of organic matter accumulating under experimental N deposition. Because the abundance of aryl C did not differ between ambient and experimental N deposition, we also have no evidence to support the notion that polyphenols in forest floor and mineral soil are less oxidized under experimental N deposition.

The mass ( $\text{g}_{\text{fraction}}/\text{g}_{\text{soil}}$ ) of large ( $>20 \mu\text{m}$ ) and small ( $<20 \mu\text{m}$ ) oPOM in mineral soil significantly increased




**Fig. 2** The abundance of C types (% of total) in forest floor and surface soil (0–10 cm) organic matter under ambient and experimental N deposition in a long-term study. Experimental N deposition had no effect on the proportion (%) of C types in forest floor (upper panel) and surface mineral soil (lower panel). Values are treatment means, and the half length of each bar is one standard error. Asterisks indicate a significant difference ( $P < 0.05$ ) in abundance between treatments.

under experimental N deposition, which contributed to an overall increase in POM under experimental N deposition (Fig. 3). This increase was driven by the greater contribution of small oPOM, relative to large oPOM (Fig. 3). Although experimental N deposition did not significantly alter either fPOM size fraction (Fig. 3), large fPOM was 23% greater under experimental N deposition, relative to the ambient treatment. Site had a significant influence on the mass of POM fractions, but site means for POM fractions did not display any consistent trend among our study sites (e.g., north to south, data not shown); site and treatment also did not interact to influence the mass of any POM fraction in mineral soil. Additionally, we experienced minimal loss of material during our size and density fractionation, wherein recovery ranged from 98.1% to 99.9%, averaging 99.1% across all samples. In combination, the

results described above support our alternative hypothesis that changes in microbial metabolism have interacted with the physical soil environment to store more organic matter in the oPOM fractions.

Consistent with our prior observations (Table 1), the C concentration in bulk mineral soil was 22% greater under experimental N deposition (15.6 vs. 19.1 mg C g<sup>-1</sup>; ambient vs. experimental N deposition, respectively;  $P = 0.02$ ), but the N concentration did not differ (0.36 vs. 0.28 mg N g<sup>-1</sup>;  $P = 0.52$ ). After POM fractions were removed from bulk mineral soil, the C (3.6 vs. 3.0 mg C g<sup>-1</sup>) and N (0.36 vs. 0.28 mg N g<sup>-1</sup>) in residual soil (>1.8 g cm<sup>-3</sup>) did not differ between ambient and experimental N deposition. However, the concentration of both C and N substantially increased under experimental N deposition (Fig. 4) in several POM fractions. For example, the C concentration of both oPOM size fractions was significantly greater under experimental N deposition, exhibiting an 80% increase in the large oPOM fraction and a 37% increase in the small oPOM fraction (Fig. 4). Similarly, experimental N deposition increased the C concentration of large fPOM by 33%, but it had no effect on the C concentration of small fPOM (Fig. 4). The N concentration in all POM fractions displayed the same trend as C concentration (Fig. 4), but increases in N concentration under experimental N deposition were only significant for large (+84%) and small (+26%) oPOM. Although site was a significant main effect in our analyses of the C and N concentration of POM fractions, means did not exhibit a consistent trend across our study sites; site and treatment also did not interact to influence the C or N concentration of any POM fraction or residual mineral soil. It appears that experimental N deposition has increased the accumulation of both C and N in bulk mineral soil as well as occluded POM fractions, which provides further support for our alternative hypothesis. Evidence for this is the significant increase in the mass (Fig. 3) and C concentration (Fig. 4) of several POM fractions, especially small oPOM.

In Fig. 5, we calculated the mass of organic C contained in each POM fraction as well as amounts contained in bulk and residual soil (>1.8 g cm<sup>-3</sup>). For each POM fraction, values are the product of its mass (g<sub>fraction</sub>/g<sub>soil</sub>) and C concentration (g C/g<sub>soil</sub>). The total amount of C contained in small and large oPOM significantly increased under experimental N deposition by 82% and 36%, respectively (Fig. 5). Although not significantly greater, the mass of organic C contained in large fPOM also increased by 33% under experimental N deposition. Interestingly, the difference in bulk soil C between ambient and experimental N deposition (3.5 mg C g<sup>-1</sup>) is equivalent to the additive increases in the mass of organic C in large fPOM (0.9 mg C), large



**Fig. 3** The percent of soil mass composed of POM fractions under ambient and experimental N deposition. Experimental N deposition has increased the mass of organic matter residing in oPOM fractions. Values are treatment means and the half length of each bar is one standard error. Asterisks indicate a significant difference ( $P < 0.05$ ) in concentrations between treatments.

oPOM ( $0.9 \text{ mg C g}^{-1}$ ) and small oPOM ( $1.9 \text{ mg C g}^{-1}$ ; combined increase =  $3.7 \text{ mg C g}^{-1}$ ). Clearly, the greater mass of C stored in soil under experimental N deposition results from increases in the amount of C that has accumulated in these POM fractions, especially small oPOM (Fig. 5).


The C : N ratio of bulk mineral soil (14.0 vs. 15.3) and residual mineral soil (10.1 vs. 10.7) was significantly greater under experimental N deposition (see Table S1). Site was not a significant main effect in our analysis of C : N in bulk soil or the residual soil after POM fractions were removed; there also was no interaction between site and treatment on the C : N of bulk soil or mineral residue. Interestingly, experimental N deposition had no influence on the C : N ratio of any POM fraction ( $P = 0.6\text{--}0.8$ ), but the C : N ratio differed dramatically among POM fractions, providing insight into their degree of decay. For example, both large fPOM (23.5) and large oPOM (27.7) had C : N ratios that were significantly greater than either small fPOM (13.4) or small oPOM (15.1), suggesting that these small size fractions have experienced a greater degree of microbial decay than the larger size fractions.

To determine whether experimental N deposition altered the biochemical composition of the four POM fractions, we quantified the abundance of carboxyl, aryl, O/N-alkyl, and alkyl C contained within them (Fig. 6). Similar to our analysis of forest floor and bulk mineral soil (Fig. 2), we have no evidence to support the hypothesis that experimental N deposition has altered the biochemical composition of soil organic matter in any POM fraction (Fig. 6). For example, there were no statistically significant differences in the abundance of any bond type under ambient and experimental N deposition, and this was true for both fPOM or oPOM size fractions (Fig. 6). Site was not a significant

main effect in our analysis of bond type in four POM fractions, and site did not significantly interact with treatment to influence the bond type abundance in any POM fraction. The alkyl to O/N-alkyl ratio did not significantly differ ( $P = 0.4\text{--}0.8$ ) for any fPOM or oPOM density fraction under ambient and experimental N deposition. However, averaged across N deposition treatments, there were significant differences in the alkyl to O/N-alkyl ratio among size and density fractions. For example, this ratio was greatest in small fPOM (0.73) and small oPOM (0.81), relative to large fPOM (0.54) and large oPOM (0.55). These observations indicate that organic matter residing in the small fPOM and oPOM fractions has undergone a greater degree of microbial decay. Although experimental N deposition has increased the amount of C stored in bulk mineral soil (Pregitzer *et al.*, 2008; Zak *et al.*, 2008) and the oPOM fractions, we have no evidence whatsoever that it has altered the biochemical composition of organic matter accumulating in soil.

## Discussion

Although we have accumulated evidence that declines in fungal polyphenol metabolism and increased bacterial metabolisms of these substrates have led to a significant accumulation of SOM under experimental N deposition (Table 1; Zak *et al.*, 2011; Freedman & Zak, 2014), our analyses of forest floor, mineral soil, and POM fractions using  $^{13}\text{C}$ -NMR provide no evidence that this change in microbial metabolism has altered the biochemical composition of organic matter to any extent. Rather, biochemically equivalent SOM has accumulated under experimental N deposition in oPOM fractions, which provides support for our alternative hypothesis. Moreover, the relatively low C : N of this



**Fig. 4** The concentration of organic C and N ( $\text{mg g}^{-1}$ ) in soil fractions under ambient and experimental N deposition. Values are treatment means, and the half length of each bar is one standard error. Asterisks indicate a significant difference ( $P < 0.05$ ) in concentration between treatments.

material, as well as the higher alkyl to O/N-alkyl ratio, indicates that it has undergone substantial decay by the microbial community (Baldock *et al.*, 1997). Because organic matter in oPOM decays at a slower pace relative to other fractions (Golchin *et al.*, 1994; Puget *et al.*, 1995; Besnard *et al.*, 1996), greater amounts of small oPOM under experimental N deposition could plausibly remain stored in soil for centuries (Rasmussen *et al.*, 2005; Yamashita *et al.*, 2006; McFarlane *et al.*, 2012) and perhaps longer (Mueller & Kögel-Knabner, 2009). If such a response occurs in other northern temperate forests experiencing anthropogenic N deposition, then it has important implications for the extent to which they

will function as future sinks for anthropogenic  $\text{CO}_2$  in the Earth's atmosphere.

Recently, Frey *et al.* (2015) synthesized the results of experimental N deposition in several temperate forest ecosystems, which revealed that significant increases in SOM resulted from the slowing of microbial decay, and not from the greater production of leaf or root litter; these observations are consistent with the evidence we have accumulated in our experiment (Table 1). Importantly, the biogeochemical changes (Table 1) in our experiment appear to be a general response of northern temperate forests to chronic anthropogenic N deposition, an aggregation of ecosystems that are a globally




Fig. 5 The amount of organic C contained in bulk soil and soil fractions under ambient and experimental N deposition. Values are the product of the mass of each soil fraction and their C concentration, providing an estimate of the mass of C (mg C g<sup>-1</sup>) contained in each soil fraction. Asterisks indicate a significant difference ( $P < 0.05$ ) between treatments.




Fig. 6 The abundance of C types in fPOM and oPOM size fractions. The proportion (%) of C types in free and occluded organic matter was not altered by experimental N deposition. Values are treatment means, and the half length of each bar is one standard error.

important sink for anthropogenic CO<sub>2</sub> in the Earth's atmosphere. If forests throughout the Northern Hemisphere respond in a manner similar to those in our

experiment, then it is plausible that the unabated deposition of anthropogenic N will increase the amounts and longevity of organic matter stored in their soils. Such a



biogeochemical response has the potential to counteract the accumulation of anthropogenic CO<sub>2</sub> in the atmosphere and therefore slow the pace of climate warming.

The formation of SOM is a complex process that is mediated by the amount and biochemical characteristics of detritus (i.e., plant and microbial) entering soil, the metabolism of this material by saprotrophic soil microorganisms, and the physical and chemical interaction of organic decay products with mineral particles (Kögel-Knabner, 2002; Lehmann & Kleber, 2015). Associations between organic matter and soil mineral particles involve occlusion of organic compounds into aggregates at the micrometer scale and interactions with mineral surfaces at the submicron scale (Chenu & Plante, 2006; Virto *et al.*, 2008; Kleber *et al.*, 2015). Several recent experiments using <sup>13</sup>C and <sup>15</sup>N consistently demonstrate the transfer of both organic C and N from litter into mineral soil fractions (Zeller & Dambrine, 2011; Hatton *et al.*, 2012; Bimüller *et al.*, 2013; Cotrufo *et al.*, 2015); this process is summarized in a conceptual model of C and N flow through soil mineral–organic associations by Hatton *et al.* (2012). Decaying plant residues enter the soil as coarse fragments that are progressively fragmented and decomposed. They are then transformed into fragments increasingly covered with fine mineral particles and form small aggregates, isolated as oPOM in our fractionation scheme. This physical transfer pathway (Cotrufo *et al.*, 2015) is associated with a progressive accumulation of microbial residues leading to low C : N ratios (Bimüller *et al.*, 2014), along with the progressive transfer of litter-derived materials from plant debris to microaggregates. As pointed out by Hatton *et al.* (2012), the progression from plant debris to aggregates induces increasing time lags prior to N reaching the dense organo-mineral associations. Mineral-stabilized organic matter is also derived from the direct association of dissolved organic matter released from decaying litter with mineral surfaces (Kaiser & Kalbitz, 2012; Cotrufo *et al.*, 2015), resulting in organo-mineral fractions that can be isolated due to their higher density (Chenu & Plante, 2006). Our results suggest that this pathway is not affected by experimental N deposition, because the fraction isolated with a density >1.8 g cm<sup>-3</sup> (i.e., residual soil) did not reveal differences in the amount of C or N contained within it (Table S1).

The results we present here amplify the importance of interactions among residual decay products and the soil mineral particles, especially silt and clay. For example, soils in our experiment are sandy (~85% sand), contain relatively small amounts of silt and clay (11 to 15%), and contain no macro-aggregates whatsoever (MacDonald *et al.*, 1991). However, despite the small proportion of fine soil particles in these soils, they had

a disproportionately large effect on stabilizing organic matter under experimental N deposition. Evidence supporting this assertion is the significant increase in the amount of organic matter residing in oPOM, especially the small oPOM fraction (Fig. 3). Organic matter contained in the small oPOM is thought to be stabilized by its association with mineral particles (von Lützow *et al.*, 2007; Hatton *et al.*, 2012; Lehmann & Kleber, 2015), which physically preclude attack by oxidative and hydrolytic extracellular enzymes, thereby reducing the extent of microbial decay. The higher alkyl to O/N-alkyl ratio of small oPOM, together with its lower C : N, further indicates that organic matter residing in this fraction has undergone a greater degree of microbial decay than organic matter found in the large fPOM and oPOM fractions.

Although we can demonstrate significant change in fungal and bacterial organic matter metabolism under experimental N deposition (Table 1), it appears that physical and chemical interactions between decay products and mineral surfaces are driving greater soil C storage under experimental N deposition. However, we do not understand why nearly 20 years of experimental N deposition has altered the soil environment to facilitate the greater occlusion of biochemically equivalent organic matter by silt and clay particles. Regardless, this observation is consistent with a recent modeling analysis indicating that SOM has accumulated in our experiment from a reduced extent of decay, rather than a slower rate of decay (Whittinghill *et al.*, 2012). Given the rapid and substantial accumulation of SOM that we have documented under experimental N deposition, developing this understanding appears central to anticipating the extent to which anthropogenic N deposition will foster greater soil C storage in northern temperate forests.

Several lines of prior evidence indicate that fine roots are the primary source of SOM under experimental N deposition, but we have no evidence whatsoever that broad changes in either leaf or root litter biochemistry have occurred under experimental N deposition (Xia *et al.*, 2015); this result is similar to our observations of the biochemical characteristics of organic matter in forest floor and mineral soil presented here and elsewhere (Thomas *et al.*, 2012). Foremost, fine roots in our experiment contain a threefold-greater lignin concentration (i.e., acid-insoluble fraction) and a twofold-greater concentration of condensed tannins than leaf litter (Xia *et al.*, 2015). Moreover, experimental N deposition has not substantially altered the concentrations of these decay-resistant molecules in either leaf or fine root litter (Xia *et al.*, 2015). For example, the relative difference in biochemical constituents (e.g., cellulose, hemicellulose, lipids, protein, soluble phenolics, tannins, and lignin)

between fine roots under ambient and experimental N deposition ranges from +6% for fine root N to -8% for condensed tannins. When combined with rates of leaf and fine root litter production (Table 1; Burton *et al.*, 2004), fine roots compose ~70% of the lignin and polyphenols entering soil, and that proportion does not differ between ambient and experimental N deposition (Xia *et al.*, 2015). Further, the biochemical composition of soil organic matter in our experiment is similar to that of fine roots, but not to leaf litter (Thomas *et al.*, 2012). Taken together, these observations indicate that fine roots are the dominant source of SOM in our experiment. Despite the change in microbial metabolism we have documented, it appears that the end products of fine root decay, which have been stabilized in oPOM, are the primary source of organic matter accumulating under experimental N deposition. Over the timescale of our experiment, it is implausible that relatively small biochemical differences (-8% to +6%) in fine root biochemistry under experimental N deposition have caused an 18% increase in SOM (Table 1). Rather, in some way, long-term experimental N deposition has increased the degree to which the products of fine root decay are stabilized by silt and clay particles, despite their relatively low abundance in the sandy soils of our experiment. Given the low amount of silt and clay in our experiment, this effect may be even more pronounced in soils of finer texture.

The insights we have gained from solid-state  $^{13}\text{C}$ -NMR spectroscopy provide no support for the hypothesis that declines in fungal polyphenol oxidation and the greater incomplete bacterial metabolism of these substrates have led to a change in the biochemical composition of soil organic matter. This result is unexpected for several reasons. First, the evidence we provide above demonstrates that experimental N deposition has not altered the amount or biochemical composition of either leaf or root litter entering soil (Xia *et al.*, 2015; Table 1). Given the consistent decline in the expression and activity of fungal extracellular enzymes that completely oxidize lignin and polyphenols in SOM (Edwards *et al.*, 2011; Freedman & Zak, 2015), one would expect to observe a greater proportion of aryl C (i.e., aromatic bonds) under experimental N deposition. However, the abundance of aryl C in bulk soil, as well as in all POM fractions, was equivalent under ambient and experimental N deposition. There are two plausible alternatives that could give rise to this observation. First, the insights provided by solid-state  $^{13}\text{C}$ -NMR spectroscopy provide an accurate assessment of the biochemical characteristics of SOM under ambient and experimental N deposition, and, indeed, changes in microbial metabolism have not altered the biochemistry of SOM. Alternatively, changes in saprotrophic litter

metabolism under experimental N have altered SOM biochemistry in manner that cannot be detected by this method of analysis. Presently, we are unable to determine which of these alternatives is correct and resolving them holds important insight into the mechanisms by which experimental N deposition has fostered the rapid accumulation of SOM.

One might argue that experimental N deposition has increased the abundance of arbuscular mycorrhizal fungi (AMF) and earthworms, both of which are known to aid in the formation of soil aggregates (i.e., oPOM), albeit through different mechanisms. However, several pieces of evidence suggest this is not the case. The young glacially derived soils (ca. 9000 yrs) in our study are ~85% sand and contain no visible macro-aggregates (D.R. Zak, *personal observation*). Approximately 85% of trees in our study are sugar maple (*Acer saccharum* Marsh.), whose roots are infected by AMF. These fungal symbionts produce glomalin-related soil proteins, which foster soil aggregation as well as its stability over time (Rillig, 2004). In contrast to this expectation, experimental N deposition has reduced the biomass of intraradical (-36%) and extraradical (-41%) AMF (Van Diepen *et al.*, 2010), making it implausible that greater glomalin production under experimental N deposition is responsible for the increase in oPOM we document here. Moreover, glaciation has removed many of the native earthworms across our sites, and they have yet to be invaded by either native or exotic earthworms (e.g., *Lumbricus terrestris*; Tiunov *et al.*, 2006; D.R. Zak, *personal observation*). Taken together, these observations suggest that interactions between the surface of fine soil particles and organic matter decay products are the most plausible mechanism increasing oPOM under experimental N deposition.

In conclusion, we have accumulated evidence that long-term experimental N deposition has not altered the production or biochemical composition of leaf or root litter entering soil and that the metabolic activities of lignolytic fungi have decreased, whereas the lignolytic activity of soil bacteria is potentially greater. These observations are consistent with a reduction in decay and the greater production of phenolic DOC. Despite these physiological changes and the biogeochemical responses (Table 1), the biochemical composition of forest floor and mineral soil was identical between ambient and experimental N deposition, at least as assayed by the methods used here and elsewhere (Thomas *et al.*, 2012). Rather, the accumulation of SOM has occurred primarily in oPOM, organic material that is stabilized by entrapment with silt and clay particles (Virto *et al.*, 2008; Hatton *et al.*, 2012). This result is unexpected and signals the importance of the interactions among physical and chemical

characteristics of fine soil particles and the biochemical residues from microbial decay. We do not understand the mechanisms by which *ca.* two decades of experimental N deposition has fostered the greater occlusion of organic matter by silt and clay particles. If other forests in the Northern Hemisphere respond similarly, then the unabated deposition of anthropogenic N has the potential to dramatically and rapidly increase soil C storage, especially in forests that occur on relatively fine-texture soil with high silt and clay contents.

## References

- Amelung W, Brodowski S, Sandhage-Hofmann A, Bol R (2008) Combining biomarker with stable isotope analyses for assessing the transformation and turnover of soil organic matter. *Advances in Agronomy*, **100**, 155–250.
- Baldock JA, Oades JM, Nelson PN *et al.* (1997) Assessing the extent of decomposition of natural organic materials using solid-state  $^{13}\text{C}$  NMR spectroscopy. *Australian Journal of Soil Research*, **35**, 1061–1083.
- Besnard E, Chenu C, Balesdent J *et al.* (1996) Fate of particulate organic matter in soil aggregates during cultivation. *European Journal of Soil Science*, **47**, 495–503.
- Bimüller C, Naumann PS, Buegger F *et al.* (2013) Rapid transfer of  $^{15}\text{N}$  from labeled beech leaf litter to functional soil organic matter fractions in a Rendzic Leptosol. *Soil Biology and Biochemistry*, **58**, 323–331.
- Bimüller C, Dannenmann M, Tejedor J *et al.* (2014) Prolonged summer droughts retard soil N processing and stabilization in organ-mineral fractions. *Soil Biology and Biochemistry*, **68**, 241–251.
- Braun E (1950) *Deciduous Forests of Eastern North America*. Macmillan Publishing Co, Inc, New York.
- Burd AB, Frey S, Cabre A *et al.* (2016) Terrestrial and marine perspectives on modeling organic matter degradation pathways. *Global Change Biology*, **22**, 121–136.
- Burton AJ, Ramm CW, Reed DD, Pregitzer KS (1991) Use of multivariate methods in forest research site selection. *Canadian Journal of Forest Research*, **21**, 573–1580.
- Burton AJ, Pregitzer KS, Crawford JN, Zogg GP, Zak DR (2004) Chronic  $\text{NO}_3^-$  additions reduce soil respiration in northern hardwood forests. *Global Change Biology*, **10**, 1080–1091.
- Burton AJ, Jarvey JJ, Jarvi MP, Zak DR, Pregitzer KS (2012) Chronic N deposition alters root respiration: tissue N relationships in northern hardwood forests. *Global Change Biology*, **18**, 258–266.
- Chenu C, Plante AF (2006) Clay-sized organo-mineral complexes in a cultivation chronosequence: revisiting the concept of the “primary organo-mineral complex”. *European Journal of Soil Science*, **57**, 596–607.
- Cotrufo MF, Soong JL, Horton AJ *et al.* (2015) Formation of soil organic matter via biochemical and physical pathways of litter mass loss. *Nature Geoscience*, **8**, 776–779.
- DeForest JL, Zak DR, Pregitzer KS, Burton AJ (2004) Experimental  $\text{NO}_3^-$  additions alter microbial community function in northern hardwood forests. *Soil Science Society of America Journal*, **68**, 132–138.
- DeForest JL, Zak DR, Pregitzer KS, Burton AJ (2005) Atmospheric  $\text{NO}_3^-$  deposition, declines in decomposition and increases in DOC: test of a potential mechanism. *Soil Science Society of American Journal*, **69**, 1233–1237.
- Edwards IP, Zak DR, Kellner H, Eisenlord SD, Pregitzer KS (2011) Simulated atmospheric N deposition alters fungal community composition and suppresses lignocellulolytic gene expression in forest floor of a northern hardwood forest. *PLoS ONE*, **6**, e20421.
- Freedman Z, Zak DR (2014) Atmospheric N deposition increases bacterial laccase-like multicopper oxidases: implications for organic matter decay. *Applied and Environmental Microbiology*, **80**, 4460–4468.
- Freedman Z, Zak DR (2015) Atmospheric N deposition alters co-occurrence in saprotrophic bacterial communities. *Molecular Ecology*, **24**, 3170–3180.
- Freedman Z, Upchurch R, Cline LC, Zak DR (2016) Anthropogenic N deposition slows decay by favoring bacterial metabolism: insights from metagenomic analyses. *Frontiers in Microbiology*, **7**, 259, doi:10.3389/fmicb.2016.00259.
- Frey SA, Ollinger S, Nadelhoffer KJ *et al.* (2015) Chronic nitrogen additions suppress decomposition and sequester carbon in temperate forests. *Biogeochemistry*, **121**, 305–316.
- Galloway JN, Dentener FJ, Capone DG *et al.* (2004) Nitrogen cycles; past, present and future. *Biogeochemistry*, **70**, 153–226.
- Golchin A, Oades JM, Skjemstad JO, Clarke P (1994) Study of free and occluded particulate organic matter in soils by solid-state  $\text{C-13}$  Cp/Mas nmr-spectroscopy and scanning electron-microscopy. *Australian Journal of Soil Science*, **32**, 285–309.
- Hassett JE, Zak DR, Blackwood CB, Pregitzer KS (2009) Are basidiomycete laccase gene abundance and composition related to reduced lignolytic activity under elevated atmospheric  $\text{NO}_3^-$  deposition in a northern hardwood forest? *Microbial Ecology*, **57**, 728–739.
- Hatton P-J, Kleber M, Zeller B *et al.* (2012) Transfer of litter-derived N to soil mineral-organic associations: evidence from decadal  $^{15}\text{N}$  tracer experiments. *Organic Geochemistry*, **42**, 1489–1501.
- Kaiser K, Kalbitz K (2012) Cycling downwards - dissolved organic matter in soils. *Soil Biology and Biochemistry*, **52**, 29–32.
- Kleber M (2010) What is recalcitrant soil organic matter? *Environmental Chemistry*, **7**, 320–332.
- Kleber M, Eusterhues K, Keiluweit M, Mikutta C, Mikutta R, Nico PS (2015) Mineral-organic associations: formation, properties, and relevance in soil environments. *Advances in Agronomy*, **130**, 1–140.
- Knicker H, Gonzalez-Vila FJ, Polvillo O, Gonzalez JA, Almendros G (2005) Fire-induced transformation of C- and N-forms in different organic soil fractions from a Dystric Cambisol under a Mediterranean pine forest (*Pinus pinaster*). *Soil Biology and Biochemistry*, **37**, 701–718.
- Kögel-Knabner I (2002) The macromolecular organic composition of plant and microbial residues as inputs to soil organic matter. *Soil Biology and Biochemistry*, **34**, 139–162.
- Lamarque JF, Kielhl JT, Brassuer GP *et al.* (2005) Assessing future nitrogen deposition and carbon cycle feedback using a multimodel approach: analysis of nitrogen deposition. *Journal of Geophysical Research*, **110**, D19303.
- Lehmann J, Kleber M (2015) The contentious nature of soil organic matter. *Nature*, **528**, 60–68.
- Liu LL, Greaver TL (2010) A global perspective on belowground carbon dynamics under nitrogen deposition. *Ecology Letters*, **13**, 819–828.
- von Lützwow M, Kögel-Knabner I, Klemens K *et al.* (2007) SOM fraction methods: relevance to functional pools and stabilization mechanisms. *Soil Biology and Biochemistry*, **39**, 321–332.
- Maaroufi NI, Nordin A, Hasselquist NJ, Bach LH, Palmqvist K, Gundale MJ (2015) Anthropogenic nitrogen deposition enhances carbon sequestration in boreal soils. *Global Change Biology*, **21**, 3169–3180.
- MacDonald NW, Burton AJ, Jurgensen MF *et al.* (1991) Variation in forest soil properties along a Great-Lakes air-pollution gradient. *Soil Science Society of America Journal*, **55**, 1709–1713.
- Magnani FM, Mencuccini M, Borghetti M *et al.* (2007) The human footprint in the carbon cycle of temperate and boreal forests. *Nature*, **477**, 848–850.
- McFarlane K, Torn M, Hanson P *et al.* (2012) Comparison of soil organic matter dynamics at five temperate deciduous forests with physical fractionation and radiocarbon measurements. *Biogeochemistry*, **8**, 1–20.
- Mueller CV, Kögel-Knabner I (2009) Soil organic carbon stocks, distribution and composition affected by historic land use changes on adjacent sites. *Biology and Fertility of Soils*, **45**, 347–359.
- Nadelhoffer KJ, Emmet BA, Gundersen P *et al.* (1999) Nitrogen deposition makes a minor contribution to carbon sequestration in temperate forests. *Nature*, **398**, 145–148.
- Pregitzer KS, Zak DR, Burton J, Ashby JA (2004) Chronic nitrate additions dramatically increase the export of carbon and nitrogen in northern hardwood forests. *Biogeochemistry*, **68**, 179–197.
- Pregitzer KS, Burton AJ, Zak DR, Talhelm AF (2008) Simulated chronic N deposition increases carbon storage in northern temperate forests. *Global Change Biology*, **14**, 142–153.
- Puget P, Chenu C, Balesdent J (1995) Dynamics of soil organic matter associated with particle-size fractions of water-stable aggregates. *European Journal of Soil Science*, **51**, 595–605.
- Rasmussen CG, Torn MS, Southard RJ (2005) Mineral assemblage and aggregates control carbon dynamics in a California conifer forest. *Soil Science Society of America Journal*, **69**, 1711–1721.
- Rillig MC (2004) Arbuscular mycorrhizae, glomalin and soil aggregation. *Canadian Journal of Soil Science*, **84**, 355–363.
- Schmidt MWI, Torn MS, Abiven S *et al.* (2011) Persistence of soil organic matter as an ecosystem property. *Nature*, **478**, 49–56.
- Steffens M, Kölbl A, Kögel-Knabner I (2009) Alteration of soil organic matter pools and aggregation in semi-arid steppe topsoils as driven by organic matter inputs. *European Journal of Soil Science*, **60**, 198–212.

- Steffens M, Kölbl A, Schörk GB, Kögel-Knabner I (2011) Distribution of soil organic matter fractions and aggregate size classes in grazed semiarid steppe soil profiles. *Plant and Soil*, **338**, 63–81.
- Thomas DC, Zak DR, Filley TR (2012) Chronic N deposition does not apparently alter the biochemical composition of forest floor and soil organic matter. *Soil Biology and Biochemistry*, **54**, 7–13.
- Thornton PE, Doney SC, Lindsay K *et al.* (2009) Carbon-nitrogen interactions regulate climate-carbon cycle feedbacks: results from an atmosphere-ocean general circulation model. *Biogeosciences*, **6**, 2099–2120.
- Tiunov AV, Hale CM, Holdsworth AR, Vesvolodov-Perel TS (2006) Invasion patterns of Lumbricidae into previously earthworm-free areas of northeastern Europe and the western Great Lakes Region of North America. *Biological Invasions*, **8**, 1223–1234.
- Van Diepen LTA, Lilleskov EA, Pregitzer KS, Miller RM (2010) Simulated nitrogen deposition causes a decline in the intra- and extraradical abundance of arbuscular mycorrhizal fungi and changes in microbial community structure in northern hardwood forests. *Ecosystems*, **13**, 683–695.
- Virto I, Barre P, Chenu C (2008) Microaggregation and organic matter storage at the silt-size scale. *Geoderma*, **146**, 326–335.
- Whittinghill KA, Currie WS, Zak DR, Burton AJ, Pregitzer KS (2012) Anthropogenic N deposition increases soil C storage by decreasing the extent of litter decay: analysis of field observations with a biogeochemical model. *Ecosystems*, **15**, 450–461.
- Xia M, Talhelm AF, Pregitzer KS (2015) Fine roots are the dominant source of recalcitrant plant litter in sugar maple-dominated northern hardwood forests. *New Phytologist*, **208**, 715–726.
- Yamashita T, Flessa H, John B, Helfrich M, Ludwig B (2006) Organic matter in density fractions of water-stable aggregates in silty soils: effect of land use. *Soil Biology and Biochemistry*, **38**, 3222–3234.
- Zak DR, Holmes WE, Burton AJ, Pregitzer KS, Talhelm AF (2008) Atmospheric NO<sub>3</sub>-deposition increases soil organic matter by slowing decomposition in a northern hardwood ecosystem. *Ecological Applications*, **18**, 2016–2027.
- Zak DR, Pregitzer KS, Burton AJ, Edwards IP, Kellner H (2011) Microbial responses to a changing environment: implications for the future functioning of terrestrial ecosystems. *Fungal Ecology*, **4**, 386–395.
- Zeller B, Dambrine E (2011) Coarse particulate organic matter is the primary source of mineral N in the topsoil of three beech forests. *Soil Biology and Biochemistry*, **43**, 542–550.

### Supporting Information

Additional Supporting Information may be found in the online version of this article:

**Table S1.** Results from physical fractionation of soils according to density and the subsequent analysis of these fractions.