

Patient-Reported Barriers are Associated with Lower HCC Surveillance Rates in Patients with Cirrhosis

Sherean Farvardin^{1*}, Jaimin Patel^{1*}, Maleka Khambaty¹, Olutola A. Yerokun¹, Hiram Mok¹, Jasmin A. Tiro^{2,3}, Adam C. Yopp^{2,4}, Neehar D. Parikh⁵, Jorge A. Marrero¹, Amit G. Singal^{1,2,3}

¹Department of Internal Medicine, UT Southwestern Medical Center and Parkland Health Hospital System, Dallas, TX

²Harold C. Simmons Cancer Center, UT Southwestern Medical Center, Dallas, TX

³Department of Clinical Sciences, UT Southwestern Medical Center, Dallas, TX

⁴Department of Surgery, UT Southwestern Medical Center, Dallas, TX

⁵Department of Internal Medicine, University of Michigan, Ann Arbor, MI

* Drs Farvardin and Patel contributed equally to this work and are co-first authors

Correspondence: Amit G. Singal, M.D., M.S.
Division of Digestive and Liver Diseases
UT Southwestern Medical Center
5959 Harry Hines Blvd, POB 1, Suite 420
Dallas TX 75390-8887
Tel: 214-645-6029
Fax: 214-645-6294
Email: amit.singal@utsouthwestern.edu

Running Title: HCC Surveillance Patient Barriers

Key Words: liver cancer, screening, knowledge, barriers, early detection

Financial Support: This work was conducted with support from the Agency for Healthcare Research and Quality (R24 HS022418), the University of Texas Southwestern Center for Patient-Centered Outcomes Research, and an ACG Junior Faculty Development Award awarded to Dr. Singal. The content is solely the responsibility of the authors and does not necessarily

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version record](#). Please cite this article as [doi:10.1002/hep.28770](https://doi.org/10.1002/hep.28770).

represent the official views of the University of Texas Southwestern Medical Center and its affiliated academic and health care centers or the Agency for Healthcare Research and Quality.

Conflicts of Interest: None of the authors have any relevant conflicts of interest to disclose

Author Contributions:

Sherean Farvardin - acquisition of data and critical revision of the manuscript for intellectual content

Jaimin Patel - acquisition of data and critical revision of the manuscript for intellectual content

Maleka Khambaty – acquisition of data and critical revision of the manuscript for intellectual content

Olutola Yerokun - acquisition of data and critical revision of the manuscript for intellectual content

Huram Mok – acquisition of data and critical revision of the manuscript for intellectual content

Jasmin Tiro – study concept and critical revision of the manuscript for intellectual content

Adam Yopp - critical revision of the manuscript for intellectual content

Neehar Parikh - critical revision of the manuscript for intellectual content

Amit Singal – study concept and design, acquisition of data, analysis and interpretation of the data, drafting of the manuscript, critical revision of the manuscript for intellectual content, obtained funding, and study supervision.

Abbreviations:

HCC – hepatocellular carcinoma; EMR – electronic medical record; HCV – hepatitis C virus; CT – computed tomography; MRI – magnetic resonance imaging; AFP – alpha fetoprotein; IQR – interquartile range; OR – odds ratio

ABSTRACT

Background: Over 20% of patients with cirrhosis are non-adherent with hepatocellular carcinoma (HCC) surveillance recommendations; however, few studies have evaluated the impact of patient-level factors on surveillance receipt. We characterized the association between HCC surveillance receipt and patient knowledge, attitudes, and perceived barriers in a racially diverse and socioeconomically disadvantaged cohort of patients with cirrhosis. Patients with cirrhosis followed at a large urban hospital were invited to complete a survey about HCC surveillance between August 2014 and December 2015. Multivariable logistic regression was performed to identify factors associated with HCC surveillance receipt during the 12-month period preceding and 6-month period after survey administration. We achieved a response rate of 71.8% (n=541/753). Patients demonstrated high levels of HCC-related knowledge (summary score 77.7%); however 48.6% believed eating a healthy diet precluded the need for HCC surveillance, and 34.0% believed HCC surveillance was not necessary if they had a normal physical exam and/or lacked clinical symptoms. Patients expressed worry about developing and dying from HCC, but nearly half (49.9%) of patients reported barriers to receiving HCC surveillance including difficulty with the scheduling process (30.5%), costs of surveillance testing (25.3%), and transportation difficulties (17.3%). HCC surveillance receipt was significantly higher in patients who knew cirrhosis is a risk factor for developing HCC (OR 3.09, 95%CI 1.25-7.62) and significantly lower in those reporting barriers to surveillance (OR 0.42, 95%CI 0.25-0.70).

Conclusions: Patients with cirrhosis are knowledgeable and interested in HCC surveillance; however, patient-reported barriers are associated with lower surveillance rates in clinical practice and represent potential intervention targets to improve HCC surveillance effectiveness.

INTRODUCTION

Hepatocellular carcinoma (HCC) is the third leading cause of cancer-related death worldwide and one of the leading causes of death in patients with cirrhosis.(1) Prognosis for patients with HCC depends on tumor stage at diagnosis, with curative options only available for patients diagnosed at an early stage.(2) Despite improvement in overall survival over time, most patients continue to be diagnosed at advanced stages and 5-year survival rates remain below 20%.(3) Racial/ethnic minorities have higher rates of late stage tumor detection as well as significantly lower stage-adjusted survival rates than non-Hispanic Caucasians.(4, 5)

Poor HCC clinical outcomes are in part attributed to underuse of surveillance among at-risk patients with cirrhosis. Several cohort studies have demonstrated HCC surveillance is associated with higher rates of early tumor detection, curative treatment receipt, and overall survival;(6) however, fewer than 20% of patients with cirrhosis in the United States undergo HCC surveillance.(7) HCC surveillance rates are even lower among racial-ethnic minorities and patients of low socioeconomic status, likely related to a combination of medical, socioeconomic, and cultural issues.(8)

Although provider recommendation is reported to be a strong predictor for HCC surveillance,(9) patient-level factors may also play an important role in surveillance underuse. In prospective cohort studies, nearly 1 in 5 patients were non-adherent with provider recommendations for HCC surveillance.(10, 11) The cancer screening literature has applied theoretical models of behavior change and demonstrated that patients' poor knowledge, negative attitudes, or perceived barriers to cancer screening may be associated with lower screening rates.(12, 13) However, most studies evaluating HCC surveillance correlates have primarily focused on patient socio-demographic and clinical characteristics.(7, 14) The effect of theory-

based patient-reported psychosocial factors on HCC surveillance completion has not been quantified. A better understanding of patient knowledge, attitudes and barriers, particularly among low income minorities, is crucial to develop and implement effective interventions to improve HCC surveillance rates.(15) Therefore, the aim of our study was to characterize patient-level knowledge, attitudes, and barriers regarding HCC surveillance and their association with receipt of HCC surveillance in a racially diverse and socioeconomically disadvantaged cohort of patients with cirrhosis.

METHODS

Study Population

Electronic medical records of patients being seen at Parkland Health and Hospital System (Parkland), the safety net health system of Dallas County, were screened to identify adult patients with cirrhosis. Parkland is an integrated healthcare system including 12 primary care provider clinics, outpatient Hepatology and multidisciplinary Liver Tumor clinics, and a tertiary hospital—all sharing a single electronic medical record (EMR) system.

Patients were initially identified using a validated set of ICD-9 codes for cirrhosis (571.2, 571.5) or cirrhosis complications (456.0, 456.1, 456.2, 456.21, 567.23, 572.2, 572.3, 572.4).(16) The presence of cirrhosis was then confirmed by chart review prior to survey administration by one of the authors (A.S., M.K. or J.P.), with diagnosis based on consistent histology, non-invasive markers of fibrosis including Fibrosure, or imaging showing a cirrhotic-appearing liver with associated signs of portal hypertension including splenomegaly, varices, or thrombocytopenia. We excluded patients with a) grade 3-4 encephalopathy, b) patient self-reported low literacy precluding survey completion, c) language other than English or Spanish,

d) personal history of HCC, or e) history of liver transplantation. The study was approved by the Institutional Review Board of UT Southwestern Medical Center.

Survey Development and Administration

Eligible patients were recruited to complete the survey at the time of a clinic appointment or via telephone between August 2014 and December 2015. A convenience sample of patients were approached in the waiting room by one of the study investigators (S.F. or J.P.) prior to their clinic appointment, consented, and asked to complete a self-administered paper survey. Study investigators (S.F or M.K) also consented and administered the survey via telephone to a random sample of patients with cirrhosis who did not have an upcoming clinic appointment.

We used a theoretical model of patient behavior for HCC surveillance (Figure 1), based on the Health Behavior Framework, to guide selection of relevant variables for survey development.(17, 18) The survey took approximately 20 minutes to complete and was divided into 4 sections: 1) knowledge about HCC, 2) potential barriers to HCC surveillance completion, 3) patient attitudes, and 4) demographic information. The first section assessed level of knowledge and consisted of 12 true-false questions with 3 domains: 5 questions regarding HCC risk and need for surveillance, 4 questions about surveillance logistics and effectiveness, and 3 questions about HCC treatment and prognosis. The second section assessed the presence of potential patient-, provider-, and system-level barriers to HCC surveillance completion including patient refusal, financial or transportation concerns, lack of provider recommendation, and scheduling difficulties using yes/no response options. The section assessing patient attitudes included 2 domains: worry about HCC and information-seeking behavior. The patient apprehension domain estimated patient level of concern about developing HCC and/or dying from HCC as well as any impact of surveillance on level of concern using a 5-point Likert scale.

Information-seeking questions assessed confidence in current level of HCC-related knowledge and desire for more information. Demographic information included age, gender, race and ethnicity, marital status, living situation, level of education, employment status and household income. Questions were adapted from earlier validated surveys when available.(19, 20) After initial development of the survey, we pretested it using cognitive interviewing methods with 10 English-speaking and 10 Spanish-speaking patients.(21, 22)

Electronic Medical Record Data Collection

Additional correlates including patient demographics, clinical history, and laboratory data were obtained through standardized abstraction of the EMR. We did not collect provider- or clinic-level characteristics, although we previously demonstrated there are no differences in clinic practices supporting HCC surveillance.(19) Patient age, gender, race, ethnicity, preferred language, and insurance type as reported on the survey were verified using EMR data. We detailed drug, alcohol and smoking history, with active alcohol abuse defined as drinking more than 40 grams/day. Patients were classified according to etiology of liver disease using laboratory data and clinical notes as follows: hepatitis C virus (positive HCV antibody or viral load), hepatitis B virus (positive HBsAg or viral load), alcohol-related liver disease (as determined by clinic provider), nonalcoholic steatohepatitis (presence of metabolic syndrome in absence of other causes of chronic liver disease or as determined by clinic provider), and other. Data regarding presence of decompensation (ascites or hepatic encephalopathy) were abstracted from clinical notes and classified as none, mild or controlled, and severe or uncontrolled. Laboratory data of interest at time of survey administration included platelet count, creatinine, bilirubin, albumin, and international normalized ratio (INR).

Study Outcomes

Our primary outcome of interest was receipt of abdominal imaging for surveillance purposes within the 12-month period preceding and 6-month period after survey administration. We evaluated surveillance completion instead of patients' intent to complete surveillance because it could be collected in a standardized and objective manner using EMR data. We only included imaging for surveillance purposes because patient knowledge and attitudes about HCC surveillance would not affect the likelihood of having exams for diagnostic purposes. We used a study period surrounding the time of survey administration to minimize potential for changes in patient knowledge or attitudes. We recorded all imaging (ultrasound, contrast-enhanced CT, or contrast-enhanced MRI) and alpha fetoprotein (AFP) results during the study period. Imaging performed at outside institutions were included if results were recorded in the EMR. Intent of imaging (surveillance vs. non-surveillance purposes) was determined through review of imaging orders and associated clinical notes. Indications including "surveillance", "screening", "rule out HCC", and "cirrhosis" were classified as surveillance indications, whereas exams performed for any symptoms or signs, e.g. abdominal pain or elevated liver enzymes, were classified as diagnostic exams.

Surveillance receipt was first characterized as an ordinal outcome: consistent surveillance (≥ 1 imaging study during each 6-month period), inconsistent surveillance (≥ 1 imaging study during each study period but not meeting criteria for semiannual surveillance), and no surveillance. Given the small number of patients with consistent surveillance, patients with consistent and inconsistent surveillance were combined for correlates analyses.

A secondary outcome of interest was overall patient knowledge. Each of the 12 true-false questions in the survey was awarded 1 point if correct and 0 if incorrect, "don't know", or left

blank. A summary scale of the items was created, with higher scores indicating more accurate knowledge.

Statistical Analysis

Distributions of patient characteristics and perceptions of surveillance were reported with descriptive statistics. Univariate and multivariable logistic regression was performed to identify factors (patient knowledge, attitudes, barriers, and EMR-derived clinical variables) associated with our dichotomous measure of HCC surveillance receipt (any surveillance vs. no surveillance). The multivariable model included variables of *a priori* clinical importance (e.g., race/ethnicity, Child Pugh score) and factors significant on univariate analyses. Predictor variables with $p < 0.10$ in univariate analyses were included in multivariable models to minimize type II error. Statistical significance was defined as $p < 0.05$ for multivariable analyses. We required 480 patients to have 90% power to detect a difference of 15% in HCC surveillance rates¹⁸, assuming a baseline surveillance rate of 50% and pre-specified alpha of 0.05. To identify factors associated with overall patient knowledge, we performed a secondary analysis using multivariable ordinal logistic regression. The multivariable model included variables of *a priori* clinical importance (e.g., race/ethnicity, educational attainment) and factors significant on univariate analyses. All data analysis was performed using Stata 11 (StataCorp, College Station, TX).

RESULTS

Patient Characteristics

Of 753 patients approached, 541 (71.8%) agreed to complete the survey including 384 patients who completed the survey in clinic and 157 via telephone. Reasons for non-response

include non-working phone number or did not answer telephone (n=109), patient refusal (n=52 via telephone and 32 in clinic), and being admitted to the hospital, rehab facility or jail (n=19). Patient characteristics are detailed in Table 1. Median age was 56 (IQR 51 – 61) years and 56.4% were men. The cohort was racially/ethnically diverse with 43.4% Hispanics, 27.4% non-Hispanic Caucasians, and 25.7% Blacks. Most (73.0%) patients had Medicaid or were covered by a county medical assistance plan, 23.3% had Medicare, and 3.7% had private insurance. The most common etiologies of cirrhosis were HCV infection (50.7%), alcohol-induced (26.0%), and NASH/cryptogenic (10.2%). Most patients had compensated cirrhosis, with 48.7% having Child Pugh A cirrhosis, 36.5% Child Pugh B cirrhosis, and 14.8% Child Pugh C cirrhosis. Ascites was present in 48.2% of patients, and 30.5% had grade I-II hepatic encephalopathy.

Patient Knowledge of HCC Surveillance

The percent of correct responses for each of the 12 knowledge questions ranged from 51.4% to 92.6% (Table 2). Patients demonstrated lower levels of knowledge about HCC risk factors than HCC surveillance and treatment logistics and effectiveness ($p < 0.001$). Although over 90% of patients knew that patients with cirrhosis are higher risk for HCC than those without cirrhosis, there were several notable knowledge deficiencies. Nearly half believed eating a healthy diet could sufficiently lower risk of HCC to no longer require HCC surveillance. Similarly, 20% of patients erroneously believed HCC surveillance is not necessary in the absence of abdominal pain, 28.5% believed HCC surveillance can be stopped after a couple normal ultrasound exams, and 34% believed HCC surveillance is not necessary if their physical exam is normal.

Factors associated with higher levels of overall knowledge on univariate analysis (Table 3) included Caucasian race, English as primary language, HCV-related cirrhosis, Child Pugh A

cirrhosis, receipt of Hepatology subspecialty care, higher educational attainment, and high perceived likelihood of dying from HCC. Being married, living alone, and lack of insurance were associated with lower levels of knowledge. Due to collinearity between language and race/ethnicity, language was not included in the multivariable model. Age, gender, and self-reported household income were examined but were not significant on univariate analysis. On multivariable analysis, increased level of knowledge was significantly associated with Caucasian race, Medicare insurance status, receipt of Hepatology subspecialty care, and higher educational attainment (Table 3).

Patient Attitudes about HCC Surveillance

Patients demonstrated high levels of worry regarding HCC (42.3% worried and 24.0% extremely worried), with 282 (52.1%) expressing concern that they are likely to develop HCC in their lifetime and 250 (46.3%) expressing concern they are likely to die from HCC. Only 40 (7.4%) patients believed they are extremely unlikely to die from HCC. Patients expressed reassurance from undergoing HCC surveillance, with 375 (69.4%) reporting HCC surveillance is likely to find HCC at an early stage allowing curative therapy and 221 (41.0%) reported HCC surveillance decreases their risk of dying from HCC. Accordingly, 89.5% of patients rated having knowledge about HCC surveillance as important, and 88.8% believed it is important to undergo HCC surveillance.

Patient-Reported Barriers to HCC Surveillance

The prevalence of patient-reported barriers to HCC surveillance is illustrated in Figure 2. The most common perceived barriers to HCC surveillance were difficulty with the scheduling process for ultrasound exams (30.5%), costs of surveillance tests (25.3%), uncertainty of where

to get surveillance ultrasound performed (19.6%), and difficulty with transportation (17.3%).

Half (49.9%) of patients reported at least one of these barriers, with 22.8% reporting a multitude of these barriers. Perceived poor surveillance test efficacy (13.9%), fear of finding cancer (11.3%), time commitment (10.2%), and fear of pain from surveillance tests (9.5%) were uncommon barriers to HCC surveillance.

HCC surveillance Receipt

Surveillance imaging was performed at least once during the study period in 356 (65.8%) patients, with only 15 (2.8%) having consistent surveillance every 6 months. An additional 91 patients received ultrasound imaging for non-surveillance purposes, and were classified as having no surveillance. AFP was performed at least once during the study period in 92.4% of 356 patients with imaging-based surveillance and 70.2% of the 185 patients without imaging-based surveillance.

In univariate analysis, receipt of any surveillance was directly associated with Hepatology subspecialty care (70.8% vs. 36.7%, $p < 0.001$), HCV etiology of cirrhosis (69.6% vs. 62.4%, $p = 0.08$), reporting lower levels of HCC-related worry because of surveillance (69.3% vs. 60.9%, $p = 0.04$), belief that HCC surveillance reduces chance of dying from HCC (71.5% vs. 63.7%, $p = 0.10$), and knowledge that cirrhosis is a primary risk factor for HCC development (67.0% vs. 54.2%, $p = 0.08$). Receipt of HCC surveillance was inversely associated with the presence of patient-reported barriers to surveillance (53.6% vs. 71.1%, $p = 0.001$). Receipt of HCC surveillance was not associated with patient age ($p = 0.74$), gender ($p = 0.33$), Caucasian race ($p = 0.60$), or English as primary language ($p = 0.76$). Similarly, we did not find an association between receipt of surveillance and overall level of knowledge ($p = 0.89$), perceived importance of undergoing surveillance ($p = 0.62$), or perceived likelihood of developing HCC ($p = 0.36$).

In multivariable analysis, receipt of any surveillance was positively associated with receipt of Hepatology subspecialty care (OR 2.65, 95%CI 1.30 – 5.41) HCV etiology of cirrhosis (OR 2.08, 95%CI 1.19 – 3.63) and patient knowledge that cirrhosis is a high-risk state for developing HCC (OR 3.09, 95%CI 1.25-7.62); presence of any of the commonly reported barriers was inversely associated with receipt of HCC surveillance (OR 0.42, 95%CI 0.25 – 0.70) (Table 4).

In a post-hoc analysis, receipt of imaging for any reason over the 18-month period was directly associated with receipt of Hepatology care (OR 4.77, 95%CI 2.63 – 8.64) and inversely associated with the presence of any of the commonly reported barriers to surveillance (OR 0.47, 95%CI 0.30 – 0.74). We did not find an association between receipt of any imaging and patient age ($p=0.64$), gender (0.36), Caucasian race ($p=0.94$), HCV etiology ($p=0.75$), overall level of knowledge ($p=0.58$), or knowledge that cirrhosis is a high-risk state for developing HCC ($p=0.77$).

DISCUSSION

To the best of our knowledge, our study is the first survey assessing patient knowledge, attitudes, and barriers for HCC surveillance in a racially diverse and socioeconomically disadvantaged cohort of patients. We found patients demonstrated high levels of HCC-related knowledge and expressed concern for developing and dying from HCC; however, they reported several barriers to receiving HCC surveillance including difficulty with the scheduling process, costs of surveillance testing, and transportation difficulties. Surveillance rates were significantly higher in patients who knew cirrhosis was a risk factor for HCC development and significantly lower in those who reported barriers to HCC surveillance, suggesting patient knowledge deficits

and surveillance barriers may be important intervention targets to increase HCC surveillance rates in the United States.

Although a prior survey study from the University of Michigan among 160 patients with cirrhosis reported high levels of knowledge about HCC risk, surveillance logistics, and treatment, this patient population represented a highly selected patient population (>80% Caucasian, >63% college educated, and highly insured) and may not be applicable to other populations.(20) Our current survey was administered in a racially diverse and socioeconomically disadvantaged cohort of patients followed in a large safety net health system. Patient knowledge, attitudes, and barriers of this population is particularly important to understand given increased HCC risk, lower surveillance rates, and worse prognosis among racial/ethnic minorities and patients of low socioeconomic status.(4, 23, 24)

Despite differences in patient populations, patients demonstrated high levels of basic knowledge regarding HCC risk, surveillance logistics, and prognosis. Most importantly, 90% of patients understood that cirrhosis is a high-risk state for HCC and that ultrasound-based surveillance should be performed every 6-12 months. However, HCC surveillance rates were significantly lower among the subset of patients who did not know patients with cirrhosis were at high risk for developing HCC, highlighting the importance of discussing this issue with all patients with cirrhosis. We also found several other knowledge deficiencies that warrant targeted patient education efforts. Nearly half of patients believed eating a healthy diet could preclude the need for HCC surveillance, and over one-third believed HCC surveillance was not necessary if they had a normal physical exam and/or were asymptomatic. Further, over one-fourth of patients believed HCC surveillance could be stopped after two normal ultrasound exams, potentially

partly explaining “surveillance fatigue” and decreasing adherence rates as patients are followed over time.

Overall, patients expressed high levels of concern for developing and/or dying from HCC. Although they expressed reassurance in terms of HCC surveillance improving early tumor detection, less than half of patients believed HCC surveillance reduces mortality. This uncertainty regarding HCC surveillance’s survival benefit could contribute to patients placing a lower priority on HCC surveillance and lead to non-adherence with surveillance recommendations. The reason for the disparity in reassurance for early tumor detection and mortality reduction is unclear but may relate to how providers communicate benefits of HCC surveillance. A similar pattern was seen in a survey study among primary care providers caring for patients with cirrhosis.(19) Provider communication style and their level of enthusiasm for cancer screening benefits has been a strong predictor of adherence in colon and breast cancer screening.(25) If provider communication also impacted HCC surveillance adherence, this would highlight the need for higher quality data characterizing the survival benefit of HCC surveillance in patients with cirrhosis. These data would also support studies evaluating the impact of interventions such as provider communication training or audit-feedback on patient attitudes and surveillance adherence.(26)

Nearly half of all patients reported issues with ultrasound scheduling, costs, transportation, or uncertainty where to get surveillance ultrasound performed. While these barriers are likely more common among socioeconomically disadvantaged patients, they highlight important intervention targets that can increase HCC surveillance rates. Effective intervention strategies can be adapted from breast and colon cancer screening programs, where similar barriers have been reported.(27-30) Simple reassurance about insurance coverage may be

sufficient to allay fears of surveillance costs, and having clinic staff schedule ultrasounds the same day as clinic visits can likely mitigate patient issues with ultrasound scheduling, transportation, and uncertainty regarding potential surveillance sites. Comparative effectiveness trials evaluating intervention strategies to address patient-reported barriers are needed.

It is important to note that our study had limitations. Our study was performed in a single large safety-net hospital and may not be generalized to other practice settings. For example, our results might not be generalizable to patients associated with transplant centers given most patients in safety net health systems are not eligible for liver transplantation given social and/or financial barriers.⁽³¹⁾ However, we feel this study population is particularly important to evaluate given higher incidence and mortality rates among racial/ethnic minorities and socioeconomically disadvantaged patients. A national survey is needed to see if our data are representative of patients receiving care in other settings. Survey studies are inherently limited by response and recall biases. Although we had a high response rate (>70%), there is also the possibility of non-response bias, in which patients knowledgeable and/or interested in HCC surveillance were more likely to respond. However, we feel these limitations are outweighed by its notable strengths including its large racially diverse sample size and the fact that it is one of the first studies to evaluate the combination of socio-demographic, clinical, and patient-reported psychosocial factors affecting HCC surveillance utilization, providing insight into potential patient-level intervention targets to increase HCC surveillance rates.

Overall, our study provides important insights into patient knowledge, attitudes and barriers regarding HCC surveillance among patients with cirrhosis. Despite having overall high levels of HCC-related knowledge, they have misconceptions about HCC surveillance and report several barriers to implementation. These misconceptions and perceived barriers might partly

explain the gap between the efficacy of HCC surveillance and its effectiveness in clinical practice. Overall, these findings highlight the need for patient education and systems-level interventions to decrease barriers for increasing HCC surveillance rates and optimizing HCC surveillance effectiveness.

Accepted Article

References

1. El-Serag HB, Rudolph KL. Hepatocellular carcinoma: epidemiology and molecular carcinogenesis. *Gastroenterology* 2007 Jun;132(7):2557-76.
2. Singal AG, Marrero JA. Recent advances in the treatment of hepatocellular carcinoma. *Curr Opin Gastroenterol* 2010 May;26(3):189-95.
3. Altekruse SF, McGlynn KA, Reichman ME. Hepatocellular carcinoma incidence, mortality, and survival trends in the United States from 1975 to 2005. *J Clin Oncol* 2009 Mar 20;27(9):1485-91.
4. Ha J, Yan M, Aguilar M, Bhuket T, Tana M, Liu B, Gish R, Wong RJ. Race/Ethnicity-Specific Disparities in Cancer Incidence, Burden of Disease, and Overall Survival among Patients with Hepatocellular Carcinoma in the U.S. *Cancer* 2016;122.
5. Sloane D, Chen H, Howell C. Racial disparity in primary hepatocellular carcinoma: tumor stage at presentation, surgical treatment and survival. *J Natl Med Assoc* 2006 Dec;98(12):1934-9.
6. Singal AG, Pillai A, Tiro J. Early Detection, Curative Treatment, and Survival Rates for Hepatocellular Carcinoma Surveillance in Patients with Cirrhosis: A Meta-analysis. *PLoS Med* 2014 Apr;11(4):e1001624.
7. Singal AG, Yopp A, C SS, Packer M, Lee WM, Tiro JA. Utilization of hepatocellular carcinoma surveillance among American patients: a systematic review. *J Gen Intern Med* 2012 Jul;27(7):861-7.
8. Singal AG, X. L, Tiro J, Kandunoori P, Adams-Huet B, Nehra M, Yopp A. Racial, Social, and Clinical Determinants of Hepatocellular Carcinoma Surveillance. *Am J Med* 2014.
9. Singal AG, Yopp AC, Gupta S, Skinner CS, Halm EA, Okolo E, Nehra M, Lee WM, Marrero JA, Tiro JA. Failure Rates in the Hepatocellular Carcinoma Surveillance Process. *Cancer Prev Res (Phila)* 2012 Aug 7;5:1124-30.
10. Collier J, Sherman M. Screening for hepatocellular carcinoma. *Hepatology* 1998 Jan;27(1):273-8.
11. Henrion J, Libon E, De Maeght S, Deltenre P, Schapira M, Ghilain JM, Maisin JM, Heller FR. [Screening for hepatocarcinoma in a cohort with cirrhosis mainly of alcoholic origin]. *Gastroenterol Clin Biol* 2003 May;27(5):534-9.

12. Klein WM, Bloch M, Hesse BW, McDonald PG, Nebeling L, O'Connell ME, Riley WT, Taplin SH, Tesauro G. Behavioral research in cancer prevention and control: a look to the future. *Am J Prev Med* 2014 Mar;46(3):303-11.
13. Rakowski W, Breslau ES. Perspectives on behavioral and social science research on cancer screening. *Cancer* 2004 Sep 1;101(5 Suppl):1118-30.
14. Davila JA, Morgan RO, Richardson PA, Du XL, McGlynn KA, El-Serag HB. Use of surveillance for hepatocellular carcinoma among patients with cirrhosis in the United States. *Hepatology* 2010 Jul;52(1):132-41.
15. Glanz K, Bishop DB. The role of behavioral science theory in development and implementation of public health interventions. *Annu Rev Public Health* 2010;31:399-418.
16. Nehra MS, Ma Y, Clark C, Amarasingham R, Rockey DC, Singal AG. Use of administrative claims data for identifying patients with cirrhosis. *J Clin Gastroenterol* 2013 May-Jun;47(5):e50-4.
17. Bastani R, Glenn BA, Taylor VM, Chen MS, Jr., Nguyen TT, Stewart SL, Maxwell AE. Integrating theory into community interventions to reduce liver cancer disparities: The Health Behavior Framework. *Prev Med* 2010 Jan-Feb;50(1-2):63-7.
18. Maxwell AE, Bastani R, Chen MS, Jr., Nguyen TT, Stewart SL, Taylor VM. Constructing a theoretically based set of measures for liver cancer control research studies. *Prev Med* 2010 Jan-Feb;50(1-2):68-73.
19. Dalton-Fitzgerald E, Tiro J, Kandunoori P, Halm EA, Yopp A, Singal AG. Practice Patterns and Attitudes of Primary Care Providers and Barriers to Surveillance of Hepatocellular Carcinoma in Patients with Cirrhosis. *Clin Gastroenterol Hepatol* 2014 Jul 11.
20. Singal A, Volk M, Rakoski M, Fu S, Su G, McCurdy H, Marrero J. Patient Involvement is Correlated with Higher HCC Surveillance in Patients with Cirrhosis. *J Clin Gastroenterol* 2011;45(8):727-32.
21. Beatty P, Willis G. Research synthesis: the practice of cognitive interviewing. *Public Opinion Quarterly* 2007;71:287-311.
22. Willis G. Cognitive interviewing: A tool for improving questionnaire design. Thousand Oaks, CA: Sage Publications; 2004.

23. Artinyan A, Mailey B, Sanchez-Luege N, Khalili J, Sun CL, Bhatia S, Wagman LD, Nissen N, Colquhoun SD, Kim J. Race, ethnicity, and socioeconomic status influence the survival of patients with hepatocellular carcinoma in the United States. *Cancer* 2010 Mar 1;116(5):1367-77.
24. Shebl FM, Capo-Ramos DE, Graubard BI, McGlynn KA, Altekruse SF. Socioeconomic status and hepatocellular carcinoma in the United States. *Cancer Epidemiol Biomarkers Prev* 2012 Aug;21(8):1330-5.
25. Fox SA, Heritage J, Stockdale SE, Asch SM, Duan N, Reise SP. Cancer screening adherence: does physician-patient communication matter? *Patient Educ Couns* 2009 May;75(2):178-84.
26. Price-Haywood EG, Harden-Barrrios J, Cooper LA. Comparative effectiveness of audit-feedback versus additional physician communication training to improve cancer screening for patients with limited health literacy. *J Gen Intern Med* 2014 Aug;29(8):1113-21.
27. Green AR, Peters-Lewis A, Percac-Lima S, Betancourt JR, Richter JM, Janairo MP, Gamba GB, Atlas SJ. Barriers to screening colonoscopy for low-income Latino and white patients in an urban community health center. *J Gen Intern Med* 2008 Jun;23(6):834-40.
28. Sabatino SA, Lawrence B, Elder R, Mercer SL, Wilson KM, DeVinney B, Melillo S, Carvalho M, Taplin S, Bastani R, Rimer BK, Vernon SW, Melvin CL, Taylor V, Fernandez M, Glanz K. Effectiveness of interventions to increase screening for breast, cervical, and colorectal cancers: nine updated systematic reviews for the guide to community preventive services. *Am J Prev Med* 2012 Jul;43(1):97-118.
29. Singal AG, Tiro JA, Gupta S. Improving hepatocellular carcinoma screening: applying lessons from colorectal cancer screening. *Clin Gastroenterol Hepatol* 2013 May;11(5):472-7.
30. **Singal AG, Gupta S.** Tiro JA, Skinner CS, et al. Outreach invitations for FIT and colonoscopy improve colorectal cancer screening rates: A randomized controlled trial in a safety net health system. *Cancer* 2016; 122(3): 456-63.
31. Singal AG, Chan V, Getachew Y, Guerrero R, Reisch JS, Cuthbert JA. Predictors of liver transplant eligibility for patients with hepatocellular carcinoma in a safety net hospital. *Dig Dis Sci* 2012 Feb;57(2):580-6.

Accepted Article

Figure Legends

Figure 1: Model of Patient Behavior for HCC Surveillance

Figure 2: Patient Reported Barriers to HCC Surveillance

Accepted Article

Table 1: Patient Characteristics

Characteristic	Receipt of any surveillance (n=356)	No surveillance (n=185)
Age (years)	55.4 ± 8.6	55.1 ± 10.1
Gender (% male)	206 (57.9%)	99 (53.5%)
Race/Ethnicity		
Non-Hispanic Caucasian	100 (28.1%)	48 (25.9%)
Black	84 (23.6%)	55 (29.7%)
Hispanic	157 (44.1%)	78 (42.2%)
Other/Unknown	15 (4.2%)	4 (2.2%)
Primary language (% English)	273 (76.7%)	144 (77.8%)
Insurance status		
Uninsured	160 (44.9%)	94 (50.8%)
Medicaid	102 (28.7%)	39 (21.1%)
Medicare	80 (22.5%)	46 (24.9%)
Private Insurance	14 (3.9%)	6 (3.2%)
Educational attainment		
Did not graduate high school	92 (25.8%)	38 (20.5%)
High school degree	168 (47.2%)	86 (46.5%)
Attended but did not graduate college	74 (20.8%)	51 (27.6%)
College degree	20 (5.6%)	10 (5.4%)
Annual household income		
Less than \$10,000	233 (65.4%)	126 (68.1%)
\$10,000 - \$20,000	70 (19.7%)	31 (16.8%)
Greater than \$20,000	38 (10.7%)	21 (11.4%)
Not reported	15 (4.2%)	7 (3.8%)
Marital status (% married)	112 (31.5%)	53 (28.7%)
Living situation (% alone)	80 (22.5%)	46 (24.9%)
Etiology of Liver Disease		
Hepatitis C	190 (53.4%)	83 (44.9%)

Hepatitis B	8 (2.2%)	6 (3.2%)
Alcohol-related	91 (25.6%)	49 (26.5%)
Non-alcoholic steatohepatitis	34 (9.5%)	21 (11.4%)
Other	33 (9.3%)	26 (14.0%)
Child Pugh class		
Child Pugh A	166 (48.4%)	74 (49.3%)
Child Pugh B	122 (35.6%)	58 (38.7%)
Child Pugh C	55 (16.0%)	18 (12.0%)
Presence of hepatic encephalopathy	120 (33.7%)	45 (24.3%)
Presence of ascites	175 (49.2%)	86 (46.5%)
Receipt of Hepatology Care	327 (91.9%)	135 (73.0%)

Accepted Article

Table 2: Proportion of Patient Knowledge Items Answered Correctly

Knowledge Item	Proportion correct (n=541)
Patients with cirrhosis are at high risk for developing HCC	491 (90.8%)
Eating a healthy diet does not sufficiently lower HCC risk to preclude a need for surveillance	278 (51.4%)
Lack of abdominal pain does not preclude need for HCC surveillance	433 (80.0%)
Patients with two consecutive normal surveillance ultrasound exams require continued HCC surveillance	387 (71.5%)
Normal physical exam and labs does not preclude need for HCC surveillance	357 (66.0%)
Ultrasound is the primary recommended modality for HCC surveillance	462 (85.4%)
Patients with cirrhosis should have HCC surveillance performed at least once per year	478 (88.4%)
Surveillance does not prevent development of HCC	418 (77.3%)
Surveillance does not preclude diagnosis with advanced HCC	392 (72.5%)
HCC can be cured if diagnosed at an early stage	429 (79.3%)
Patients with HCC will die without cancer treatment	419 (77.5%)
HCC does not spontaneously resolve without cancer treatment	501 (92.6%)
Summary score for all questions (n=12)	77.7%
Summary score for risk factors questions (n=5)	71.9%
Summary score for surveillance logistics and efficacy questions (n=4)	80.9%
Summary score for treatment questions (n=3)	83.1%

HCC – hepatocellular carcinoma

Table 3: Predictors of Higher Patient Knowledge

Variable	Univariate Models OR (95% CI)	Multivariable Model AOR (95% CI) *
Race/Ethnicity		
Non-Hispanic Caucasian	Reference	Reference
Black	0.46 (0.30 – 0.70)	0.53 (0.33 – 0.85)
Hispanic	0.20 (0.14 – 0.30)	0.38 (0.24 – 0.61)
Other/Unknown	0.37 (0.15 – 0.91)	0.53 (0.21 – 1.31)
English language as preferred language	3.83 (2.67 – 5.48)	Collinear
Insurance status		
Uninsured	Reference	Reference
Medicaid	1.49 (1.04 – 2.15)	1.25 (0.84 – 1.87)
Medicare	1.73 (1.18 – 2.53)	1.75 (1.15 – 2.65)
Private Insurance	0.96 (0.45 – 2.04)	1.0 (0.42 – 2.38)
Educational attainment		
Did not graduate high school	Reference	Reference
High school degree	3.03 (2.07 – 4.43)	2.14 (1.41 – 3.26)
Attended but did not graduate college	3.84 (2.46 – 6.00)	2.68 (1.61 – 4.44)
College degree	5.79 (2.77 – 12.11)	3.20 (1.42 – 7.20)
Marital status (being married)	0.55 (0.40 – 0.76)	NS
Living alone	1.43 (1.01 – 2.03)	NS
HCV etiology of cirrhosis	1.79 (1.33 – 2.42)	NS
Child Pugh A cirrhosis	1.42 (1.03 – 1.94)	NS
Perceived likelihood of dying from HCC	1.57 (1.16 – 2.12)	NS
Receipt of Hepatology care	1.49 (1.10 – 2.00)	1.45 (1.05 – 2.01)

OR - Odds Ratio; AOR - Adjusted Odds Ratio; CI = Confidence Interval; HCV – hepatitis C virus; HCC – hepatocellular carcinoma; NS – not significant

* Significant associations in bold


Table 4: Predictors of HCC Surveillance Receipt

Variable	Univariate Models	Multivariable Model*
	OR (95% CI)	AOR (95% CI)
Age (continuous)	1.00 (0.98 – 1.02)	**
Female gender	0.84 (0.59 – 1.20)	**
Caucasian race	1.11 (0.74 – 1.67)	**
English as primary language	0.94 (0.61 – 1.43)	**
HCV etiology of cirrhosis (vs. other etiologies)	1.37 (0.96 – 1.97)	2.09 (1.20 – 3.65)
Receipt of Hepatology care	4.18 (2.53 – 6.88)	2.64 (1.30 – 5.40)
Knowledge: Cirrhosis is risk factor for HCC	1.72 (0.94 – 3.13)	3.02 (1.23 – 7.39)
Attitude: Surveillance reduces HCC mortality	1.43 (0.94 – 2.19)	NS
Attitude: Surveillance reduces worry about HCC	1.45 (1.01 – 2.07)	NS
Patient reported barriers to HCC surveillance	0.47 (0.30 – 0.73)	0.42 (0.25 – 0.70)

* Adjusted for race/ethnicity and Child Pugh class A cirrhosis given a priori importance

**Not entered into multivariable model due to lack of significance in univariate analyses

OR - Odds Ratio; AOR - Adjusted Odds Ratio; CI = Confidence Interval; HCV – hepatitis C virus; HCC – hepatocellular carcinoma; NS – not significant




* Not measured in current study.

Model of Patient Behavior for HCC Surveillance

Figure 1

26x29mm (300 x 300 DPI)

Accept



Patient Reported Barriers to HCC Surveillance
Figure 2
352x264mm (72 x 72 DPI)

Accep