

**Title: Characteristics of Children with Cancer Discharged or Admitted from the Emergency Department**

Emily L Mueller MD, MSc<sup>1,2</sup>

Matt Hall PhD<sup>3</sup>

Samir S Shah, MD, MSCE<sup>4</sup>

Keith J August, MD, MS<sup>5</sup>

Mohamed Radhi, MD<sup>5</sup>

Michelle L Macy, MD, MS<sup>6,7</sup>

1. Section of Pediatric Hematology Oncology, Department of Pediatrics, Indiana University, Indianapolis, IN 46202
2. Center for Pediatric and Adolescent Comparative Effectiveness Research, Indiana University, Indianapolis, IN 46202
3. Children's Hospital Association, Overland Park, KS 66202
4. Division of Hospital Medicine, Department of Pediatrics, University of Cincinnati, Cincinnati, OH, 45229
5. Section of Pediatric Hematology/Oncology/Stem Cell Transplantation, University of Missouri, Kansas City, MO 64108
6. Child Health Evaluation and Research (CHEAR) Unit, Division of General Pediatrics, Department of Pediatrics and Communicable Diseases, University of Michigan, Ann Arbor, MI, 48109
7. Department of Emergency Medicine, University of Michigan, Ann Arbor, MI 48109

Correspondence:

Emily L Mueller, MD, MSc

410 West 10th Street, Suite 4099C

Indianapolis, IN 46202

Cell: 312-399-0245 Fax: 317-321-0128

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/pbc.25872](https://doi.org/10.1002/pbc.25872).

This article is protected by copyright. All rights reserved.

Admit Peds Ca

elmuelle@iu.edu

Abstract word count: 240

Main text word count: 2892

Number of tables: 3

Number of figures: 1

Short Title: ED Utilization for Children with Cancer

Key words: Child, Adolescent, Oncology, Supportive Care, Febrile Neutropenia

Abbreviations:

|          |  |
|----------|--|
| ALL      | Acute lymphoblastic leukemia   |
| CNS      | Central nervous system tumors  |
| CI       | Confidence interval  |
| ED       | Emergency department   |
| FN       | Febrile neutropenia  |
| ICD-9-CM | International Classification of Diseases, Ninth Edition, Clinical Modification |
| NHL      | Non-Hodgkin lymphoma   |
| OR       | Odds ratio   |
| PHIS     | Pediatric Health Information System  |
| SCT      | Stem cell transplant   |
| U.S.     | United States  |

### Abstract

**Background:** Emergency department (ED) utilization by children with cancer is poorly understood. Among children with cancer, we explored reasons for ED visits and factors associated with admission within U.S. children's hospitals.

**Methods:** A retrospective study of the 2011-2013 Pediatric Health Information System (PHIS) was conducted. Eligible ED visits included those within 365 days from the first inpatient encounter with

This article is protected by copyright. All rights reserved.

Admit Peds Ca

an ICD-9-CM code for cancer. Patient characteristics and reasons for ED visits were assessed. Factors associated with admission from the ED were examined with multivariable regression.

Results: There were 26,770 ED visits by 17,943 children with cancer at 39 children's hospitals during the study period. Half of children with cancer visited the ED within one year after their first cancer hospitalization in PHIS. Fifty-six percent of ED visits resulted in admission. Fever or neutropenia accounted for the largest proportion of reasons for visits (34.6%). Risk factors for admission were: "Other" race/ethnicity as compared to white, non-Hispanic (OR=1.4, 95% CI 1.2-1.6), history of transplant (OR=1.7, 95% CI 1.4-2.1), and ED visits reasons including neutropenia (OR=43.4, 95% CI 36.0-52.3), blood stream infection (OR=3.3, 95% CI 2.8-3.9), pancytopenia (OR=28.8, 95% CI 18.1-45.9), dehydration (OR=2.3, 95% CI 1.9-2.9) or pneumonia (OR=3.8, 95% CI 2.8-5.1).

Conclusion: Children with cancer have high ED usage within one year after their first cancer hospitalization. Age, demographic factors and reasons for ED visits significantly impacted admission from the ED. Further research should focus on ED utilization among children with cancer.

Admit Peds Ca

## Background

Children with cancer represent a high-risk population for disease and treatment related complications that require urgent or emergent medical care.[1-4] These children are frequently referred to emergency departments (ED) for evaluation of high risk, time-sensitive conditions such as febrile neutropenia (FN) and sepsis.

Improved understanding of the ED utilization of children with cancer could lead to optimization of ED referral practices and ED management through targeted interventions aimed at the most common reasons for ED visits and admissions. While admission rates for children with cancer vary by reason for the ED visit,[5] the role of patient level factors on the decision to hospitalize are poorly defined.

The objectives of this study were to (a) assess ED utilization by pediatric patients with cancer and evaluate the reasons why children with cancer visit the ED, and (b) assess factors associated with admission versus discharge from the ED at children's hospitals in the United States, including patient specific factors unavailable in previously analyzed datasets.

## Methods

### *Study Design and Setting*

We conducted a retrospective cohort study using data from the Pediatric Health Information System (PHIS), an administrative and resource utilization database from 45 freestanding children's hospitals. Participating hospitals are located in non-competing markets of 27 states plus the District of Columbia and account for ~15% of all pediatric hospitalizations in the United States. Hospitals were included if they provided inpatient and ED data between January 1, 2011, and September 30, 2013, without interruption, resulting in the inclusion of 39 hospitals. The PHIS database contains the

This article is protected by copyright. All rights reserved.

Admit Peds Ca

following: patient medical record data (demographic characteristics, up to 41 diagnoses, and up to 41 procedures) and billing data (all medication, diagnostic imaging, laboratory, and supply charges to individual patients). Diagnoses are coded after discharge, so we are unable to separate ED versus inpatient diagnoses for those patients who were admitted. Data are de-identified before inclusion in the database; however, encrypted medical record numbers allow for tracking of individual patients across multiple encounters at the same hospital. The Children's Hospital Association (Overland Park, KS) and participating hospitals jointly monitor the quality and integrity of data, described previously.[6] The analysis was based on de-identified data and was therefore considered exempt from institutional review board approval by Indiana University School of Medicine.

#### *Study Population/Identification of Cases*

Children with cancer were defined based on the identification of an inpatient encounter that included an International Classification of Diseases, Ninth Edition, Clinical Modification (ICD-9-CM) code for cancer (140.x-209.x, 235.x-239.x), as described previously.[5] The index inpatient encounter was defined as the first inpatient encounter with an ICD-9-CM code for cancer in the study period only if there were no previous hospitalizations with cancer diagnoses within 365 days prior. We then identified all ED visits that occurred in the 365 days following the index hospitalization. We restricted our analysis to those patients between the ages of 0-19 years.

#### *Outcome and Exploratory Variables*

Patient characteristics that were evaluated included: gender, age (<1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years – based on clinical differences in types of cancers and development differences), race/ethnicity (non-Hispanic white, non-Hispanic black, Hispanic, Asian, or other), type of cancer (acute lymphoblastic leukemia (ALL), acute myelogenous leukemia, central nervous system tumors (CNS), solid tumors (non-CNS), Hodgkin lymphoma, and non-Hodgkin lymphoma (NHL)),

This article is protected by copyright. All rights reserved.

Admit Peds Ca

transplant status flag (defined using Feudtner's complex chronic condition codes, including stem cell transplant)[7], primary payer (public/governmental, private, other), median income quartile per ZIP code and urban/rural patient residence (based on the Rural Urban Commuting Area (RUCA) code of the patient's home ZIP code).[8] We also included an indicator variable for frequent ED visitor, which was defined as 4 or more visits (greater than the 90<sup>th</sup> percentile among the study population of children with cancer) during the year (365 days) following the index admission.

For the logistic regression, the primary outcome of interest was whether an ED visit resulted in admission to the same institution versus discharged to home (patient treated and released from the ED). Deaths occurring in the ED were excluded from this analysis; there were only 7 ED deaths in the time period analyzed.

#### *Reason for ED visit*

The reason prompting an ED visit was defined by the primary ICD-9-CM discharge diagnosis associated with the encounter, unless the primary diagnosis was a cancer diagnosis (ICD-9-CM Codes 140-239). In cases where a cancer diagnosis was the primary diagnosis, the second listed ICD-9-CM diagnosis was considered to be the reason prompting the ED visit. Cancer diagnoses were not considered the reason prompting an ED visit in order to focus on symptoms or complications associated with ED visits made by pediatric patients with cancer. A rank list of diagnoses was generated by frequency. Diagnoses with similar codes were collapsed into single categories, for example "neutropenia, unspecified" (288.00) and "drug induced neutropenia" (288.03) were collapsed into a single "neutropenia" variable. After collapsing the similar codes, the top 10 diagnoses were analyzed.

Admit Peds Ca

*Statistical Analyses*

Hospital and patient characteristics were summarized using frequencies and percentages, and compared across groups (admitted vs. discharged from the ED) using chi-square tests. The number of ED visits and corresponding disposition status (admitted or discharged) were analyzed. The top 10 reasons for ED visits by disposition status (admitted or discharged) were calculated by frequencies and proportions. Multivariable regression was performed using generalized linear mixed effects models to estimate factors associated with admission for pediatric patients with cancer while accounting for clustering of patients within hospitals through the inclusion of a random hospital intercept. Variables included were based on our defined model: gender, patient's age category, type of cancer, presence of a transplant flag, primary expected payer, median income quartile per ZIP code, urban/rural patient residence, a dichotomous indicator variable for frequent ED visitors, and variables for the presence or absence of each of the top 10 most common visit diagnoses. All analyses were performed using SAS v9.4 (SAS Institute, Cary, NC), and p-values <0.05 were considered statistically significant.

## Results

### *Characteristic of the study population*

We identified 17,943 children with cancer who received inpatient care with associated ICD-9-CM codes for cancer at 39 children's hospitals in the United States, from January 2011-September 2013. There were a total of 26,770 ED visits in the year following the index hospitalization among 9,114 (50.8%) of the identified children with cancer. The range for the number of ED visits per patient was 0-24, the distribution as shown in Figure 1. Among all of the ED visits, 56.1% (n=15,022) were admitted and 43.9% (n=11,748) were discharged to home; there were no transfers to other institutions.

Table 1 demonstrates the encounter characteristics of children with cancer who presented to the ED, stratified by admission versus discharge from the ED. Overall, most ED visits were for children ages 0-9 years (65.9%), white, non-Hispanic (53.5%), and residing in urban areas (86.1%). There were significant differences between encounters resulting in admission vs. discharge for all factors investigated, except for primary payer.

The ten most common reasons prompting an ED visit for the children with cancer (Table 2) accounted for one half (50.8%) of all non-cancer primary diagnoses for ED visits among children with cancer. A diagnosis of fever or neutropenia, common complications of cancer therapy, accounted for the largest percentage of visits (34.6%).

Variation in ED disposition was found across diagnoses. Children with cancer were admitted the majority of the time if their reasons for ED visit was neutropenia (97.5%), pancytopenia (96.4%), blood stream infections (75.5%), pneumonia (78.1%), and dehydration (69.6%). The lowest admission rates were for principal diagnoses of fever (23.3%), complication of a vascular device (20.0%), and headache (20.0%).



Admit Peds Ca

*Factors affecting admission versus discharge among children with cancer*

In a multivariable analysis, patient factors associated with significantly increased odds of admission included “Other” race/ethnicity as compared to white, non-Hispanic (Odds Ratio [OR]=1.4, 95% Confidence Interval [CI] 1.2-1.6) or having a history of transplant (OR=1.7, 95% CI 1.4-2.1), as shown in Table 3. Conversely, all age groups had significantly decreased odds of admission when compared with the patient encounters for those ages 15-19 years. Among the cancer diagnoses, none of the categories had significantly increased odds of admission when compared to the reference of ALL. Meeting criteria for a frequent ED user also did not increase the odds of admission.

Patients who lived in urban settings had decreased odds of admission (OR=0.7, 95% CI 0.6-0.8) compared with those living in a rural residence.

ED visit diagnoses associated with increased odds of admission included neutropenia (OR=43.4, 95% CI 36.0-52.3), blood stream infection (OR=3.3, 95% CI 2.8-3.9), pancytopenia (OR=28.8, 95% CI 18.1-45.9), dehydration (OR=2.3, 95% CI 1.9-2.9) or pneumonia (OR=3.8, 95% CI 2.8-5.1).

## Discussion

In this study using administrative data from 39 U.S. children's hospitals, half of children with cancer visited the ED within one year of their index inpatient encounter in the study period. Over half of these ED visits resulted in hospital admission, which is a higher rate than the general pediatric population (about 10%).<sup>[9,10]</sup> Consistent with our previous analysis of a nationally representative sample of ED visits, we found that there was significant variation in admission rates across the wide range of reasons for ED visits in this patient population.<sup>[5]</sup> Though the variation cannot be fully explained using administrative data, we have confirmed that admission rates are highest for those with neutropenia and lowest for those with headaches. This analysis highlights two potential areas for future intervention. For ED visit reasons that had high admission rates, research endeavors could focus understanding the patient experience and improving patient-centered outcomes. Conversely, research concentrated on those reasons with low admission rates could aim to improve anticipatory guidance given to caregivers to decrease reliance on the ED.

Consistent with our prior research, there was a wide range of reasons for ED visits among children with cancer.<sup>[5]</sup> We chose to focus on only the primary diagnosis, or the second diagnosis code if the first was a cancer diagnosis, in order to ascertain the reason for the ED visit. Yet, it is likely that children with cancer can present to the ED with more than one medical issue. Febrile neutropenia (FN) is a clinically important entity, but is not a single diagnostic code. We also found similar admission rates between patients with a primary ED visit diagnosis of neutropenia and FN (when we identified a combination of fever and neutropenia among all diagnostic fields). To truly understand ED utilization for children with cancer who experience fever in the setting of neutropenia, datasets will need to include chief complaints and laboratory data.

While children with cancer are a small subset of all children who visit the ED (0.2%),<sup>[5]</sup> there are important distinctions regarding the care for this unique population. Counter to ED admission patterns for common pediatric conditions, children with cancer who were younger (as compared to the

Admit Peds Ca

15-19 year olds) had decreased odds of admission. This finding lends itself to future research regarding reasons for these differences. Possible explanations include adolescents waiting to visit the ED only when they have reached the point where their condition requires inpatient admission or, alternatively, young children may be taken to the ED early in the course of an acute illness due to their limited ability to explain their symptoms. Also, ALL occurs more commonly in younger children and requires the longest duration of active chemotherapy, but the majority of therapy is maintenance chemotherapy with low risk for substantial immunosuppression. Further research that includes greater patient detail regarding the reason for seeking ED care will be needed to improve our understanding of this dynamic.

Socioeconomic factors appeared to impact the disposition of children with cancer from the ED. Patients who lived in an urban location, had decreased odds of admission, which may be because patients who live in urban areas may live closer to the free-standing children's hospitals that contributed to this analysis. Therefore, patients living in closer urban areas are more easily able to return for re-evaluation if their condition worsens. Those who live further away, in rural areas, may experience hardship related to travel including lack of adequate means of transportation, the cost of traveling or lodging nearby. This may result in longer hospital admissions to ensure stability prior to discharge or visits to other hospitals with subsequent transfer only if admission is deemed necessary. Of note, risk stratification schemas for the management of febrile neutropenia (FN) among children with cancer do not take into account the differences in patient proximity to the treating hospital.[11-13]

The PHIS database has several unique features that revealed important information in this study. First, we had the ability to identify patients with a comorbid condition of transplant. Within the population of patients with cancer diagnoses, this most commonly represents a stem cell transplant (SCT).[7] We found that children with cancer with SCT had an increased odds of admission when evaluated in the ED. Patients who receive a SCT are a uniquely vulnerable population of patients with

Admit Peds Ca

high rates of complications that can require emergent evaluation.[14,15] Stem cell transplant patients tend to be followed very closely as outpatients and remain within close proximity to the treating institution (for those who do not live nearby) for a period of time after discharge. Due to the prolonged immunocompromised status and high readmission rates of SCT patients, further analysis of the ED utilization of this specific population of patients may yield information that could be used to implement improved outpatient or phone triaging practices or provide targets for anticipatory guidance.

A second unique feature of the PHIS database is the ability to identify children with cancer who visited the ED frequently in the year following their index admission. Frequent ED users among children with cancer deserve additional study to gain an understanding of the reasons for their higher levels of ED utilization. Our analysis determined that meeting the frequent ED user criterion was not associated with increased odds of admission. Further research to characterize why this population is seeking care in the ED could lead to targetable interventions to decrease reliance on the ED.

While this analysis contributes to the knowledge of ED utilization by children with cancer, it has also highlighted areas that deserve further evaluation. We did not have the clinical information that would allow us to ascertain the appropriateness of ED use by this population. The care of urgent medical issues among children with cancer occurs along a continuum, from outpatient triage regarding the need for in-person evaluation to urgent evaluation to inpatient admission. Most research among children with cancer to date has focused on what happens after admission,[16-18] but we poorly understand the events or processes that occur prior to the ED visit and admission decision. The decision making process for patients and their caregivers is complex and includes many facets that need to be better understood including decisions on when to seek care, where to seek care and who to contacted regarding these issues. Lastly, both patient/caregiver and provider perspectives play a role in triaging and decision making regarding ED use or disposition and should be addressed. All of these

Admit Peds Ca

areas will need to be further explored in order to improve the care of children with cancer within the ED.

### **Limitations**

While this study adds insight into ED utilization by children with cancer, there are several important limitations. For our analysis, we captured children with cancer by identifying the first inpatient encounter with an associated ICD-9-CM code for cancer in the study period and then followed these patients for 365 days. It is possible that we are not accurately evaluating all patients with cancer at these institutions since a proportion of patients with cancer are diagnosed and initiated on therapy as outpatients. For example, many of our Hodgkin lymphoma patients are evaluated and diagnosed as outpatients, though this is a small population. Unfortunately, PHIS does not include information related to disease status, type of therapy, or time since last treatment for the patient at the time they visit the ED. This information should be included in prospective data collection regarding ED utilization in order to provide context and help to delineate associations between types of therapy and reasons for seeking urgent evaluation in an ED.

Our analyses relied on discharge ICD-9-CM codes. This limits our understanding of the chief complaint that brought the child to seek care in the ED and therefore we are missing key information for future interventions aimed at anticipatory guidance. Also, we extrapolated the reason for the ED visit from ICD-9-CM codes rather than the initial chief complaint, which was not available in this dataset. For those patients who were admitted to the inpatient hospital from the ED, the diagnosis codes evaluated are from the entirety of the encounter and cannot be differentiated between diagnoses associated with the ED versus diagnoses that arose during the inpatient stay. Only a single diagnosis was evaluated per patient encounter, yet children with cancer may present to the ED with multiple complications or concerns.

This article is protected by copyright. All rights reserved.

Admit Peds Ca

### **Conclusion**

In conclusion, half of children with cancer will visit the emergency department at a tertiary care institution within one year after the index inpatient admission for cancer. Factors that significantly impacted whether a child with cancer was admitted versus discharged from the ED included age, demographic factors and reasons for the ED visits. Knowledge of reasons for visits and patient characteristics may be used to target common diagnoses or design health system modifications to improve the care of children with cancer who visit the ED.

Funding Source: None

Financial Disclosure: The authors have no financial disclosures

Conflict of Interest: The authors have no conflicts of interest to disclose

## References

1. Behl D, Hendrickson AW, Moynihan TJ. Oncologic Emergencies. *Crit Care Clin* 2010;26(1):181–205.
2. Bodey GP, Buckley M, Sathe YS, Freireich EJ. Quantitative relationships between circulating leukocytes and infection in patients with acute leukemia. *Ann Intern Med* 1966;64(2):328–340.
3. Henry M, Sung L. Supportive Care in Pediatric Oncology: Oncologic Emergencies and Management of Fever and Neutropenia. *Pediatr Clin North Am* 2015;62(1):27-46.
4. McCurdy MT, Shanholtz CB. Oncologic emergencies. *Crit Care Med* 2012;40(7):2212–2222.
5. Mueller EL, Sabbatini A, Gebremariam A, Mody R, Sung L, Macy ML. Why pediatric patients with cancer visit the emergency department: United States, 2006-2010. *Pediatr Blood Cancer* 2015;62(3):490-495.
6. Mongelluzzo J, Mohamad Z, Ten Have TR, Shah SS. Corticosteroids and mortality in children with bacterial meningitis. *JAMA* 2008;299(17):2048-2055.
7. Feudtner C, Feinstein JA, Zhong W, Hall M, Dai D. Pediatric complex chronic conditions classification system version 2: updated for ICD-10 and complex medical technology dependence and transplantation. *BMC Pediatr* 2014;14:199.
8. Hart LG, Larson EH, Lishner DM. Rural definitions for health policy and research. *Am J Public Health* 2005;95(7):1149-1155.
9. Kharbanda AB, Hall M, Shah SS, Freedman SB, Mistry RD, Macias CG, Bonsu B, Dayan PS, Alessandrini EA, Neuman MI. Variation in resource utilization across a national sample of pediatric emergency departments. *J Pediatr* 2013;163(1):230-236.
10. O'Mahony L, O'Mahony DS, Simon TD, Neff J, Klein EJ, Quan L. Medical complexity and pediatric emergency department and inpatient utilization. *Pediatrics* 2013;131(2):e559-565.
11. Alexander SW, Wade KC, Hibberd PL, Parsons SK. Evaluation of risk prediction criteria for episodes of febrile neutropenia in children with cancer. *J Pediatr Hematol Oncol* 2002;24(1):38-42.
12. Ammann RA, Bodmer N, Hirt A, Nigli FK, Nadal D, Simon A, Ozsahin H, Kontny U, Kuhne T, Popovic MB, Luthy AR, Aebi C. Predicting adverse events in children with fever and chemotherapy-induced neutropenia: the prospective multicenter SPOG 2003 FN study. *J Clin Oncol* 2010;28(12):2008-2014.

Admit Peds Ca

13. Ammann RA, Hirt A, Luthy AR, Aebi C. Identification of children presenting with fever in chemotherapy-induced neutropenia at low risk for severe bacterial infection. *Med Pediatr Oncol* 2003;41(5):436-443.
14. Miano M, Faraci M, Dini G, Bordigoni P, Party EPW. Early complications following haematopoietic SCT in children. *Bone Marrow Transplant* 2008;41 Suppl 2:S39-42.
15. Shulman DS, London WB, Guo D, Duncan CN, Lehmann LE. Incidence and causes of hospital readmission in pediatric patients after hematopoietic cell transplantation. *Biol Blood Marrow Transplant* 2015;21(5):913-919.
16. Basu SK, Fernandez ID, Fisher SG, Asselin BL, Lyman GH. Length of Stay and Mortality Associated With Febrile Neutropenia Among Children With Cancer. *J of Clin Oncol* 2005;23(31):7958–7966.
17. Costa VC, Ferraz MB, Petrilli AS, Pereira CA, Rogerio JW. Resource utilization and cost of episodes of febrile neutropenia in children with acute leukemias and lymphomas. *Support Care Cancer* 2003;11(6):356–361.
18. Mueller EL, Walkovich KJ, Mody R, Gebremariam A, Davis MM. Hospital discharges for fever and neutropenia in pediatric cancer patients: United States, 2009. *BMC Cancer* 2015;15:388.



Admit Peds Ca

Figure 1. Distribution of the number of ED encounters




TABLE I. Encounter Level Characteristics of Children with Cancer Presenting to the Emergency Department, by Admission Status – January 2011 to September 2013

|                                | Overall<br>n=26,770 | Admitted<br>n=15,022 | Discharged<br>n=11,748 | p-value |
|--------------------------------|---------------------|----------------------|------------------------|---------|
|                                | N (% Column)        | N (% Row)            |                        |         |
| <b>Patient Characteristics</b> |                     |                      |                        |         |
| <b>Gender</b>                  |                     |                      |                        | 0.230   |
| Female                         | 12,491 (46.7)       | 5,433 (43.5)         | 7,058 (56.5)           |         |
| Male                           | 14,279 (53.3)       | 6,315 (44.2)         | 7,964 (55.8)           |         |
| <b>Age</b>                     |                     |                      |                        | <0.001  |
| <1 year                        | 4,306 (16.1)        | 2,065 (48.0)         | 2,241 (52.0)           |         |
| 1-4 years                      | 7,370 (27.5)        | 3,400 (46.1)         | 3,970 (53.9)           |         |
| 5-9 years                      | 5,979 (22.3)        | 2,659 (44.5)         | 3,320 (55.5)           |         |
| 10-14 years                    | 5,021 (18.8)        | 2,045 (40.7)         | 2,976 (59.3)           |         |
| 15-19 years                    | 4,094 (15.3)        | 1,579 (38.6)         | 2,515 (61.4)           |         |
| <b>Race/Ethnicity</b>          |                     |                      |                        | <0.001  |
| White, Non-Hispanic            | 14,311 (53.5)       | 6,299 (44)           | 8,012 (56)             |         |

Admit Peds Ca

|   |               |               |               |        |
|---|---------------|---------------|---------------|--------|
| Black, Non-Hispanic                         | 2,728 (10.2)  | 1,332 (48.8)  | 1,396 (51.2)  |        |
| Hispanic                                    | 6,196 (23.1)  | 2,727 (44.0)  | 3,469 (56.0)  |        |
| Asian                                       | 817 (3.1)     | 371 (45.4)    | 446 (54.6)    |        |
| Other                                       | 2,718 (10.2)  | 1,019 (37.5)  | 1,699 (62.5)  |        |
| <b>Type of Cancer</b>                       |               |               |               | <0.001 |
| Acute Lymphoblastic Leukemia                | 9,510 (35.5)  | 4,259 (44.8)  | 5,251 (55.2)  |        |
| Acute Myelogenous Leukemia                  | 1,324 (4.9)   | 608 (45.9)    | 716 (54.1)    |        |
| Solid Tumor                                 | 8,621 (32.2)  | 3,601 (41.8)  | 5,020 (58.2)  |        |
| Central Nervous System Tumor                | 4,712 (17.6)  | 2,212 (46.9)  | 2,500 (53.1)  |        |
| Hodgkin Lymphoma                            | 784 (2.9)     | 326 (41.6)    | 458 (58.4)    |        |
| Non-Hodgkin Lymphoma                        | 1,819 (6.8)   | 742 (40.8)    | 1,077 (59.2)  |        |
| <b>Transplant</b>                           |               |               |               | <0.001 |
| Yes   | 693 (2.6)     | 255 (36.8)    | 438 (63.2)    |        |
| No  | 26,077 (97.4) | 11,493 (44.1) | 14,584 (55.9) |        |
| <b>Frequent ED User</b>                     |               |               |               | <0.001 |
| Yes   | 15,519 (58.0) | 6,943 (44.7)  | 8,576 (55.3)  |        |
| No  | 11,251 (42.0) | 4,805 (42.7)  | 6,446 (57.3)  |        |
| <b>Socioeconomic Characteristics</b>        |               |               |               |        |
| <b>Primary Payer</b>                        |               |               |               | 0.095  |
| Public/ Governmental                        | 13,308 (49.7) | 5,833 (43.8)  | 7,475 (56.2)  |        |
| Private                                     | 12,038 (45.0) | 5,251 (43.6)  | 6,787 (56.4)  |        |
| Other                                       | 1,424 (5.3)   | 664 (46.6)    | 760 (53.4)    |        |
| <b>Median Household Income per Zip Code</b> |               |               |               | 0.001  |
| 1 <sup>st</sup> quartile                    | 5,546 (21.3)  | 2,341 (42.2)  | 3,205 (57.8)  |        |
| 2 <sup>nd</sup> quartile                    | 6,160 (23.7)  | 2,688 (43.6)  | 3,472 (56.4)  |        |
| 3 <sup>rd</sup> quartile                    | 6,585 (25.3)  | 2,864 (43.5)  | 3,721 (56.5)  |        |
| 4 <sup>th</sup> quartile                    | 7,723 (29.7)  | 3,528 (45.7)  | 4,195 (54.3)  |        |
| <b>Urban vs Rural Patient Residence</b>     |               |               |               | <0.001 |
| Urban                                       | 23,051 (86.1) | 10,354 (44.9) | 12,697 (55.1) |        |
| Rural                                       | 3,719 (13.9)  | 1,394 (37.5)  | 2,325 (62.5)  |        |

TABLE II. Top Reasons Prompting ED Visits Among Pediatric Cancer Patients – Rank and Disposition Status by Frequent ED Visitor Status

| Rank | Diagnosis | Overall      | Disposition Status   |                    |
|------|-----------|--------------|----------------------|--------------------|
|      |           | n=26,770     | Admitted to Hospital | Discharged to Home |
|      |           | N (% Column) | N (% Row)            |                    |

Admit Peds Ca

|    |                                 |              |              |              |
|----|---------------------------------|--------------|--------------|--------------|
| 1  | Neutropenia                     | 5,189 (19.4) | 5,061 (97.5) | 128 (2.5)    |
| 2  | Fever                           | 4,057 (15.2) | 940 (23.2)   | 3,117 (76.8) |
| 3  | Blood Stream Infection          | 883 (3.3)    | 667 (75.5)   | 216 (24.5)   |
| 4  | Nausea or vomiting              | 611 (2.3)    | 169 (27.7)   | 442 (72.3)   |
| 5  | Pancytopenia                    | 557 (2.1)    | 537 (96.4)   | 20 (3.6)     |
| 6  | Acute URI                       | 501 (1.9)    | 196 (39.1)   | 305 (60.9)   |
| 7  | Dehydration                     | 497 (1.9)    | 346 (69.6)   | 151 (30.4)   |
| 8  | Complication of vascular device | 485 (1.8)    | 97 (20)      | 388 (80)     |
| 9  | Headache                        | 450 (1.7)    | 90 (20)      | 360 (80)     |
| 10 | Pneumonia                       | 310 (1.2)    | 242 (78.1)   | 68 (21.9)    |

URI=Upper respiratory infection

TABLE III. Multivariate Logistic Regression to Evaluate Factors Associated with Admission Versus Discharge from the ED Among Pediatric Cancer Patients

| Factors                        | Adj. Odds Ratio (OR) | 95% CI         | p-value          |
|--------------------------------|----------------------|----------------|------------------|
| <b>Patient Characteristics</b> |                      |                |                  |
| <b>Gender</b>                  |                      |                |                  |
| Female                         | 1.1                  | 1.0-1.1        | 0.114            |
| <b>Age</b>                     |                      |                |                  |
| 15-19 years                    | Ref                  |                |                  |
| 10-14 years                    | 0.9                  | 0.8-1.0        | 0.012            |
| 5-9 years                      | 0.7                  | 0.6-0.8        | <0.001           |
| 1-4 years                      | 0.6                  | 0.6-0.7        | <0.001           |
| <1 year                        | 0.7                  | 0.6-0.8        | <0.001           |
| <b>Race/Ethnicity</b>          |                      |                |                  |
| White, Non-Hispanic            | Ref                  |                |                  |
| Black, Non-Hispanic            | 0.9                  | 0.8-1.0        | 0.049            |
| Hispanic                       | 1.1                  | 1.0-1.2        | 0.089            |
| Asian                          | 1.2                  | 1.0-1.5        | 0.121            |
| Other                          | <b>1.4</b>           | <b>1.2-1.6</b> | <b>&lt;0.001</b> |
| <b>Type of cancer</b>          |                      |                |                  |
| Acute Lymphoblastic Leukemia   | Ref                  |                |                  |
| Acute Myelogenous Leukemia     | 0.9                  | 0.8-1.1        | 0.304            |
| Solid Tumors                   | 0.9                  | 0.9-1.0        | 0.173            |
| Central Nervous System Tumor   | 1.1                  | 1.0-1.2        | 0.221            |

Admit Peds Ca

|  |             |                  |                  |
|--|-------------|------------------|------------------|
| Hodgkin Lymphoma                                       | 0.7         | 0.6-0.9          | 0.005            |
| Non-Hodgkin Lymphoma                                   | 0.9         | 0.8-1.1          | 0.191            |
| <b>Transplant</b>                                      |             |                  |                  |
| Yes  | <b>1.7</b>  | <b>1.4-2.1</b>   | <b>&lt;0.001</b> |
| <b>Frequent ED Visitor</b>                             |             |                  |                  |
| Yes  | 0.9         | 0.9-1.0          | 0.124            |
| <b>Socioeconomic Characteristics</b>                   |             |                  |                  |
| <b>Primary Payer</b>                                   |             |                  |                  |
| Public/ Governmental                                   | Ref         |                  |                  |
| Private  | 0.9         | 0.9-1.0          | 0.103            |
| Other  | 0.8         | 0.6-0.9          | 0.001            |
| <b>Median Household Income per ZIP Code</b>            |             |                  |                  |
| 4 <sup>th</sup> quartile                               | Ref         |                  |                  |
| 3 <sup>rd</sup> quartile                               | <b>1.1</b>  | <b>1.0-1.2</b>   | <b>0.017</b>     |
| 2 <sup>nd</sup> quartile                               | 1.1         | 1.0-1.2          | 0.267            |
| 1 <sup>st</sup> quartile                               | 1.0         | 0.9-1.2          | 0.400            |
| <b>Urban vs Rural Patient Residence</b>                |             |                  |                  |
| Urban  | 0.7         | 0.6-0.8          | <0.001           |
| <b>ED Visit Characteristics</b>                        |             |                  |                  |
| <b>Top 10 ED Visit Diagnoses (in descending order)</b> |             |                  |                  |
| Neutropenia  | <b>43.4</b> | <b>36.0-52.3</b> | <b>&lt;0.001</b> |
| Pancytopenia   | <b>28.8</b> | <b>18.1-45.9</b> | <b>&lt;0.001</b> |
| Pneumonia  | <b>3.8</b>  | <b>2.8-5.1</b>   | <b>&lt;0.001</b> |
| Blood Stream Infection                                 | <b>3.3</b>  | <b>2.8-3.9</b>   | <b>&lt;0.001</b> |
| Dehydration  | <b>2.3</b>  | <b>1.9-2.9</b>   | <b>&lt;0.001</b> |
| Acute URI  | 0.7         | 0.6-0.9          | 0.001            |
| Nausea or vomiting                                     | 0.4         | 0.3-0.4          | <0.001           |
| Fever  | 0.3         | 0.3-0.3          | <0.001           |
| Complication of vascular device                        | 0.2         | 0.2-0.3          | <0.001           |
| Headache   | 0.2         | 0.2-0.3          | <0.001           |

ED=Emergency Department, URI=Upper respiratory infection

**Bold** indicates significant factors associated with admission versus discharge