

TITLE: Clinical practice guidelines on the evidence-based use of integrative therapies during and following breast cancer treatment

RUNNING TITLE: Integrative therapies during and following breast cancer treatment

AUTHORS AND AFFILIATIONS

Heather Greenlee, ND, PhD, MPH; Melissa J. DuPont-Reyes, MPH, MPhil; Lynda G. Balneaves, RN, PhD; Linda E. Carlson, PhD; Misha R. Cohen, OMD, LAc; Gary Deng, MD, PhD; Jillian A. Johnson, MSc; Matthew Mumber, MD; Dugald Seely, ND, MSc; Suzanna Zick, ND, MPH; Lindsay Boyce, MLIS; Debu Tripathy, MD

Department of Epidemiology, Mailman School of Public Health (HG, MD), Herbert Irving Comprehensive Cancer Center (HG), Columbia University, New York, NY; College of Nursing, Rady Faculty of Health Sciences, Winnipeg, MB, Canada (LGB); Department of Oncology, University of Calgary, Calgary, AB, Canada (LEC); Institute for Health and Aging, University of California San Francisco, CA (MC); Chicken Soup Chinese Medicine, San Francisco, CA (MC); Department of Medicine (GD), Memorial Sloan Kettering Library (LB), Memorial Sloan Kettering Cancer Center, New York, NY (GD); Department of Biobehavioral Health, The Pennsylvania State University, University Park, PA (JJ); Harbin Clinic, Rome, GA (MM); Ottawa Integrative Cancer Center, Ottawa, ON, Canada (DS); Canadian College of Naturopathic Medicine, Toronto, ON, Canada (DS); Department of Family Medicine, University of Michigan Health System (SMZ), Department of Nutritional Sciences, School of Public Health (SMZ), and Department of Family Medicine (SMZ), University of Michigan, Ann Arbor, MI; Department of Breast Medical Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX (DT)

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version record](#). Please cite this article as [doi:10.3322/caac.21397](https://doi.org/10.3322/caac.21397).

Correspondence to: Heather Greenlee, ND, PhD, MPH, Department of Epidemiology, Mailman School of Public Health, Columbia University, 722 W. 168th Street, 7th Floor, New York, NY 10032 (email: hg2120@columbia.edu)

Date: February 14, 2017

Accepted Article

ABSTRACT (Words: 211; 200 max)

Breast cancer patients commonly use complementary and integrative therapies as supportive care during cancer treatment and to manage treatment-related side effects. However, evidence supporting the use of such therapies in the oncology setting is limited. This report provides updated clinical practice guidelines from the Society for Integrative Oncology on the use of integrative therapies for specific clinical indications during and after breast cancer treatment, including anxiety/stress, depression/mood, fatigue, quality of life/physical functioning, chemotherapy-induced nausea and vomiting (CINV), lymphedema, chemotherapy-induced peripheral neuropathy (CIPN), pain and sleep disturbance. Clinical practice guidelines are based upon a systematic literature review from 1990 through 2015. Music therapy, meditation, stress management and yoga are recommended for anxiety/stress reduction. Meditation, relaxation, yoga, massage and music therapy are recommended for depression/mood disorders. Meditation and yoga are recommended to improve quality of life. Acupressure and acupuncture are recommended for reducing CINV. Acetyl-L-carnitine is not recommended to prevent CIPN due to harm. No strong evidence supports the use of ingested dietary supplements to manage breast cancer treatment-related side effects. In summary, there is a growing body of evidence supporting the use of integrative therapies, especially mind-body therapies, as effective supportive care strategies during breast cancer treatment. Many integrative practices, however, remain understudied with insufficient evidence to be definitively recommended or avoided.

KEYWORDS (4-10 words)

Breast cancer, integrative therapies, complementary therapies, implementation strategies, integrative oncology, integrative medicine, music therapy, meditation, yoga, stress management, relaxation, massage, acupressure, acupuncture

COI: The authors report no conflicts of interest

FINANCIAL DISCLOSURES: none

INTRODUCTION

Breast cancer patients and survivors are frequent users of complementary and integrative therapies and there are a growing number of formal integrative oncology programs within cancer centers.¹⁻⁶ A variety of terms are used to describe such use and it is helpful at the outset to define terms. *Complementary and alternative therapies* are generally defined as any medical system, practice, or product that is not part of conventional medical care.^{7,8} Other relevant terminology includes “*complementary medicine*”, which comprises therapies used as a complement alongside conventional medicine; “*alternative medicine*”, which comprises therapies used in place of conventional medicine; and “*integrative medicine*”, which is the coordinated use of evidence-based complementary practices and conventional care. *Integrative oncology* refers to the use of complementary and integrative therapies in collaboration with conventional oncology care. In oncology, individuals employ complementary and integrative therapies with the intent of enhancing wellness, improving quality of life (QOL), and relieving symptoms of disease and side effects of conventional treatments. However, the evidence supporting the use of complementary and integrative therapies in the oncology setting is limited.

In November 2014, the Society for Integrative Oncology (SIO) published clinical practice guidelines to inform both clinicians and patients on the use of integrative therapies during breast cancer treatment and to treat breast cancer treatment-related symptoms.⁹ SIO adapted methods established by the U.S. Preventive Services Task Force¹⁰ to develop graded recommendations on the use of specific integrative therapies for defined clinical indications based on strength of available evidence concerning associated benefits and harms. The 2014 clinical practice guidelines were derived from a systematic review of randomized clinical trials published between 1990 and 2013 and organized by specific clinical conditions (e.g., anxiety/stress, fatigue). This review provides an updated set of clinical practice guidelines based upon a current systematic literature review of randomized controlled trials published through December 2015. In addition, this review provides detailed definitions of integrative therapies and clinical outcomes of interest, a detailed summary of the literature upon which the clinical practice

guidelines are based, and suggestions for how appropriate therapies may be integrated into clinical practice.

Of note, it is important to define the use of the term “recommendation” in these clinical practice guidelines. In many settings, a clinical guideline “recommendation” infers that it be used as the standard of care and is favorable or equal compared to all other options based on best clinical evidence for benefit/risk ratio. Here, in the setting of integrative oncology, we use the term “recommendation” to infer that the therapy should be considered as a viable but not singular option for the management of a specific symptom or side effect. Few studies have conducted a head to head comparison of a given integrative therapy against a conventional treatment, and most integrative therapies are used in conjunction with standard therapy and have been studied in this manner. Moreover, combination-based approaches and the interactions of the numerous permutations of integrative and conventional treatments have not been formally investigated, such that recommendations must account for this limitation of our knowledge. Despite these limitations to the evaluation of the use of integrative therapies in the oncology setting, there is a body of well-conducted trials of specific therapies for specific conditions that provide sufficient evidence to warrant recommendations on the therapies as viable options for treating specific conditions.

This review provides clinicians and patients with updated SIO clinical practice guidelines on the use of integrative therapies to manage symptoms and side effects during and after breast cancer treatment. The clinical practice guidelines do not address breast cancer recurrence and survival endpoints as there are very few adequately powered randomized controlled trials examining the effect of integrative therapies on these outcomes. This review provides a definition of each integrative therapy that had a sufficiently large body of evidence to formulate specific recommendations. Information is also provided on how to implement the recommendations into the clinical setting, with caveats for specific clinical situations. In addition, this review summarizes pertinent meta-analyses and identifies promising areas for future investigation. The information that arose from other published reviews and meta-analyses did not change the interpretation of the findings or quality of specific trials, but the information was used to influence the establishment of specific recommendation grades based on consistency, reproducibility and

assessment of potential harms and benefits. The goal of this current review is to provide clinicians and patients with practical information and tools to evaluate whether there is an evidence-base to support the use of a defined integrative therapy for a specific clinical application in the context of breast cancer.

METHODS

Systematic Review Methodology

To update the previously published clinical practice guidelines that were based upon the a systematic review of the literature from January 1, 1990 through December 31, 2013⁹, we conducted a systematic review of published randomized controlled trials from January 1, 2014 through December 31, 2015 using the same search criteria and process. The process followed the methods set forth by the Institute of Medicine on clinical guideline development.¹¹ The following databases were searched: Embase, MEDLINE, PsychINFO, and CINAHL. As previously reported⁹, trials were selected for inclusion in the systematic review if they met the following criteria: (1) peer-reviewed published randomized controlled trial; (2) available in English; (3) included $\geq 50\%$ breast cancer patients and/or reported results separately for breast cancer patients; (4) used an integrative therapy as an intervention during standard treatment with surgery, chemotherapy, radiation therapy, and/or hormonal therapy, or addressed symptoms and side effects resulting from diagnosis and/or treatment; and (5) addressed an endpoint of clinical relevance to breast cancer patients and survivors (Supplemental Table 1).⁹ Several lifestyle and psychological interventions were excluded from current as well as previous guidelines because they have already been well summarized by other groups (e.g., diet^{12, 13} and physical activity¹²⁻¹⁴ recommendations for cancer survivors) and/or because they have a strong evidence base and are often considered to be mainstream rather than integrative or complementary (e.g., cognitive-behavioral therapy¹⁵, psychoeducation¹⁶, counseling¹⁷, and support groups¹⁶). Other interventions excluded were in early or pilot stages of research (e.g., attention restoration therapy) or were not considered to be an integrative oncology therapy for the purposes of the SIO guidelines (e.g., prayer, spirituality). Each article was scored according to quality of design and reporting based on the Jadad scoring scale and a modified scale

adapted from the Delphi scoring system.^{18, 19} Finally, grades of evidence were determined for each therapy as applied to a specific clinical outcome using a modified version of the U.S. Preventive Services Task Force Grading System.¹⁰ Grades were based upon strength of evidence, determined by the number of trials, quality of trials, magnitude of effect, statistical significance, sample size, consistency of results across studies, and whether the outcomes were primary or secondary. The highest grades (A and B) indicate that a specific therapy is recommended for a particular clinical indication. Grade A indicates there is high certainty that the net benefit is substantial, while grade B indicates there is high certainty that the net benefit is moderate or there is moderate certainty that the net benefit is moderate to substantial. Grade C indicates that the evidence is equivocal, or that there is at least moderate certainty that the net benefit is small. The lowest grades (D, H, and I) indicate no demonstrated effect, suggest harm, or indicate that the current evidence is inconclusive, respectively.

Following the clinical guideline development process outlined by the Institute of Medicine,¹¹ drafts prepared by the SIO Guideline Working group were distributed to an interdisciplinary group of SIO internal and external reviewers. Reviewer comments, suggestions and critiques were incorporated into the final version of these guidelines.

It is important to note that as we reviewed the literature, we recognized that there is a difference between statistical and clinical significance. The graded recommendations reflect our assessment of the clinical significance based upon our assessment of the body of literature, including the importance of statistical significance with respect to the primary endpoint. We did not report on specific magnitudes of effect because of the range of outcome measures and statistical methods used across the trials, making it difficult to describe detailed data on effect sizes across all trials. Though some of the trials with small sample sizes ($n < 100$) may have been methodologically sound, we downplayed their contribution to the graded recommendation because larger trials provided more information on generalizability of results to larger populations. Due to space limitations, p -values are reported and citations are provided to reference the primary reports for additional details.

Definitions of Complementary and Integrative Therapies^{20,21}

Below are definitions listed alphabetically for each of the complementary and integrative therapies that received a Grade A, B, C, D or H in the updated clinical practice guidelines presented here. Table 1 displays the graded recommendations. Table 2 provides background information on the specific training, licensure, and professional organizations associated with each therapy. If a therapy is known to have a specific contraindication or caution, it is noted in the description. The descriptions include statements on how the therapies are often used by cancer patients and survivors, but do not indicate the level of evidence supporting such use. The guideline recommendations provide the summary of the evidence on the use for specific conditions. In addition to the information provided below, there are continuously updated well-referenced websites that can provide additional details on the range of therapies, including Natural Medicines (<https://www.naturalmedicines.therapeuticresearch.com>), Memorial Sloan Kettering Cancer Center's *About Herbs* website (<https://www.mskcc.org/cancer-care/treatments/symptom-management/integrative-medicine/herbs>), and the National Cancer Institute's Office of Cancer Complementary and Alternative Medicine CAM Therapies: A-Z website (https://cam.cancer.gov/health_information/cam_therapies_a-z.htm).

Acetyl-L-carnitine. Acetyl-L-carnitine is a dietary supplement that some patients use to treat cancer-related fatigue by enhancing energy and lowering inflammation in the body²². It has shown to be effective at preventing and treating diabetic neuropathy and, therefore, was of interest to examine in the context of chemotherapy-induced peripheral neuropathy. It is a substance made in muscle and liver tissue and is found in foods including meats, poultry, fish, and some dairy products.

Acupuncture. Acupuncture involves the stimulation of specific points, (i.e., acupoints) by penetrating the skin with thin, solid, metallic needles.^{23,24} A variation of acupuncture includes electroacupuncture, in which a small electric current is passed along acupuncture needles to provide a stronger stimulus than acupuncture alone, with distinct effects suggested by functional MRI^{25,26}. Acupuncture has been practiced in Asia for thousands of years as a component of traditional medicine systems (e.g., traditional forms of Chinese, Japanese and Korean medicine) and is thought to stimulate the

flow of a form of energy called *qi* (chee) throughout the body. Traditional Chinese acupuncture, commonly used in North America, requires needle manipulation to produce a *de qi* sensation (a soreness, fullness, heaviness, or local area distension^{27, 28}), along with a period of time of rest with the placed needles.²⁹ This is posited to remove energetic blockages, thus reestablishing homeostasis. The mechanisms for acupuncture's effects are not well understood but are thought to function in part through modulation of specific neuronal/cortical pathways.³⁰ Acupuncture practice typically requires formal education through schools, training programs, and certifications (see Table 2). Acupuncture is often used in the oncology setting for chemotherapy-induced nausea/vomiting (CINV), pain management, musculoskeletal complaints, hot flashes, fatigue, stress, anxiety and sleep disorders. The practice of acupuncture in North American is regulated by some U.S. states and Canadian provinces and territories (see Table 2)

Acupressure. Acupressure draws on the same knowledge and philosophical system as acupuncture. A trained therapist, or the patient, uses his/her hands and fingers, or possibly a device to apply pressure to specific points on the body (acupoints), in contrast to metallic needles.³¹ Practices can range from stimulating a single point or a combination of points to achieve the intended outcome. In the oncology setting, acupressure is often used for CINV pain, stress management and fatigue.

Aloe Vera. *Aloe vera* gel is derived from the leaves of the perennial succulent plant, *Aloe vera* (Liliaceae). It is typically applied topically or ingested in the form of a clear thick gel.²² *Aloe vera* gel is found in multiple skin products, such as lotions, creams, and sunblock. *Aloe vera* is used as a topical ointment to heal wounds, sunburn, insect bites, and skin conditions including psoriasis and frostbite.³² In oncology, it is typically used with the goal of healing surgical wounds or preventing or treating radiation induced dermatitis.

Ginger. Ginger (*Zingiber officianale*) comes from the rhizome or root of a tropical plant with green-purple flowers and an aromatic stem.^{22, 33} Ginger can be used as a food in cooking and for medicinal purposes. In Asian medicine, ginger is used to treat stomach aches, nausea, and diarrhea. For cancer patients, it has been studied for the treatment of chemotherapy-induced nausea and vomiting

(CINV). Ginger is available in capsule form, fresh as a root, as a tea, as a candy, or in very diluted quantities in ginger ale. Ginger supplementation should not be used in perioperative settings or in patients with bleeding disorders due to potential risk for increased bleeding.³²

Ginseng. Ginseng is derived from a plant root and has been used to treat certain medical problems.²² There are two common types of ginseng used, Asian ginseng (*Panax ginseng*) and American ginseng (*Panax quinquefolius*), while the herb called Siberian ginseng or eleuthero is not a true ginseng.³² Asian and American ginsengs are used to boost the immune system and promote well being and stamina. Ginseng comes in capsule form made of ground ginseng, extracts, and teas as well as creams and other products for topical use. Taken as an herbal supplement, ginseng is often used to treat cancer-related fatigue.³² Side effects of taking ginseng may include headaches, breast tenderness and menstrual irregularities, sleep problems, restlessness, rapid heart rate, low blood sugar, allergic reactions, and gastrointestinal problems.³²

Glutamine. Glutamine is a non-essential amino acid used in the biosynthesis of proteins and is primarily synthesized in skeletal muscle.³² Most of the glutamine synthesized in the body is utilized by the intestinal tract. Glutamine has numerous biological functions including protein and lipid synthesis, regulation of acid-base balance in the kidney, and as an important mitochondrial cellular energy source. Normally the body can synthesize its own glutamine; however, during a critical illness such as cancer not enough glutamine is made, leading to problems such as fatigue and muscle wasting. Glutamine has been used as an oral supplement in cancer patients to reverse cachexia in patients with advanced cancer. It has also been used for CINV in cancer patients. Glutamine can be obtained from food or supplements, with important food sources being beef, pork, chicken, fish, eggs, milk, dairy products, wheat, cabbage, beets, beans, spinach, and parsley.³²

Guarana. Guarana is an herbal supplement from the guarana plant (*Paullinia cupana*) native to the Amazon basin.³² Guarana supplements contains various phytochemicals, including caffeine, theobromine, theophylline, tannins, saponins, catechins, epicatechins, proanthocyanidols and other

compounds in minor concentrations. Guarana has been used as a stimulant since pre-Columbian times.³² In the oncology setting, guarana is often used to decrease fatigue.

Healing touch. Healing touch (also known as therapeutic touch) is based on the belief that vital energy flows through and around the human body and may be transferred or modified.^{34,35} A healing touch practitioner (often a nurse trained in the practice) passes his/her hands over, or gently touches, a patient's body to balance or increase their energy. Healing touch is often used among cancer patients to improve quality of life, pain, fatigue and depression.

Hyaluronic acid cream. Hyaluronic acid cream is a topical cream containing hyaluronic acid that is used to heal wounds through repair-promoting, skin moisturizing, and potential radioprotective properties.²² When the cream is applied, the hyaluronic acid adheres to injured tissue, provides hydration to the skin, and protects against dehydration, chemical and mechanical irritation. Hyaluronic acid cream is often used by cancer patient to prevent and treat radiation induced dermatitis.

Hypnosis. Hypnosis is facilitated by a specially trained therapist or is practiced on one's own (self-hypnosis). It is characterized by a trance-like state, which allows a patient to be more aware, focused and open to suggestion. A person in a hypnotic state can concentrate more clearly on specific feelings, thoughts, images, sensations, or behaviors without distraction.³⁶ The hypnotic state is obtained by first relaxing the body, then shifting attention toward a narrow range of object or ideas given by the hypnotist or hypnotherapist. A person under hypnosis may feel more calm, relaxed, and open to suggestion. In cancer patients, hypnosis is often used to help relieve stress, anxiety, and pain.

Laser therapy. Low-level laser treatment (wave lengths 650-1000 nm) has been approved in some countries, including the United States, to treat lymphedema after breast cancer surgery and is administered by trained users. Low-level laser therapy is thought to stimulate macrophages and the immune system, and break down scar tissue, thus improving lymphatic flow.³⁷

Manual lymphatic drainage and compression bandaging. Manual lymphatic drainage and compression bandaging are used singly and in combination for the treatment of lymphedema following breast cancer surgery.³⁷ Manual lymph drainage is a specific type of therapeutic massage delivered by a

health professional ideally certified in the technique. Manual lymph drainage can decrease lymphedema when administered early, before symptoms advance.³⁷ Compression bandages or garments, including sleeves, stockings, bras, compression shorts, gloves, bandages, or neck compression wraps, are also used to treat lymphedema and can be worn during the day or night depending on the garment and individual.³⁷

Massage. There are many different forms of this form of physical therapy that involves a therapist stroking, kneading, applying friction, and stretching specific muscles and other connective tissues at an even tempo with varying levels of pressure.³⁸ In cancer patients, the goal of massage is to promote relaxation, address muscle stiffness and pain, and to resolve musculoskeletal complaints. There are multiple forms of massage, including but not limited to, Swedish, shiatsu, and deep tissue massage. The National Cancer Institute urges massage therapists to take precautions with all cancer patients and avoid massaging specific vulnerable areas of the body including open wounds, bruises, skin breakdown, a blood clot in a vein, a tumor site, near a medical device (e.g., drain), or sensitive skin following radiation therapy.^{39, 40} In addition, certain patients with multiple bone metastases may be at risk for fracture during deep massage. The practice of massage therapy in North American is regulated by some U.S. states and Canadian provinces and territories (see Table 2).

Meditation. Meditation is a group of self-regulation practices that focus on training attention and awareness on bringing mental processes under greater voluntary control.⁴¹ In cancer patients, these practices are intended to foster general mental well-being, calmness, clarity, and concentration. The ultimate goal of meditation varies depending on the type of practice, its history, and application. Most meditation practices have four elements in common: a quiet location with few distractions; a specific, comfortable posture achieved by sitting or lying down; a focus of attention; and an open attitude of letting thoughts come and go naturally without judgment.⁴² The focus of attention may be towards a specific target, such as on the breath, repeating a sound or mantra (known as *concentration meditation*), on all mental events that enter the field of awareness (called *mindfulness meditation*), or a combination of both. There has been a growing interest in cancer care on the practice of a secular form of mindfulness meditation called Mindfulness-Based Stress Reduction (MBSR), based on the work of Jon Kabat-Zinn.⁴³

MBSR is typically delivered in an 8-week structured group program consisting of a range of meditation practices including a sensate focus body scan, sitting meditation, walking meditation, loving-kindness practice and gentle Hatha yoga postures. All formal practices are designed to cultivate increasing levels of mindfulness in day-to-day life. Participants engage in home practice daily throughout the program, and each session involves teaching of relevant concepts, discussion of progress and barriers to practice, and introduction and practice of new meditation modalities. A number of recent papers and meta-analyses have reviewed the literature on the use of MBSR in the oncology setting, some specific to breast cancer.⁴⁴⁻

47

Mistletoe. The use of mistletoe in cancer care is based on the premise that injections of specially prepared extracts of the plant during chemotherapy and radiation therapy can create a host response that is immune stimulatory, preferentially cytotoxic to cancer cells and protective of host cells.⁴⁸ Mistletoe is a parasitic plant from the Santalacea family that attaches to and penetrates the branches of a tree or shrub in order to absorb water and nutrients from the host plant. Use of mistletoe as a medicine extends back centuries, whereas modern use for cancer care was promoted in Europe and particularly in Germany in the early part of the 20th century.⁴⁹ There are three main types of mistletoe: European Mistletoe (*Viscum album*), Korean Mistletoe (*Viscum album var. coloratum*) and American Mistletoe (*Phoradendron leucarpum*), but multiple methods of preparation and formulation exist. Preparations from European Mistletoe are some of the most commonly prescribed substances internationally in out-patient clinics for cancer, where it is provided most often as a subcutaneous injection or occasionally intravenously by infusion.⁵⁰ While this therapy is often used clinically for its antineoplastic potential, the clinical trial evidence on the use of mistletoe is based on trials where it is co-administered with conventional treatments to improve quality of life.⁵⁰

Music therapy. Music therapy is the clinical use of music to accomplish individualized goals within a therapeutic relationship by a credentialed professional.⁵¹ In cancer care, music therapy is used to address a variety of physical, emotional, cognitive, and social needs. Qualified music therapists assess patient's strengths and needs and provide indicated treatment such as creating, singing, moving to, and listening to

music. Music therapy interventions can be described as either *passive* (e.g., listening to music prior to a medical intervention) or *active* (e.g., therapist instructing a patient to engage in the creation of live music) depending on the level of engagement required. Although the exact mechanisms by which music therapy works are not well understood, the most commonly accepted theories are through neurologic, psychological, behavioral and physiological pathways^{52, 53} The practice of music therapy in North American is regulated by some U.S. states (see Table 2).

Reflexology. In reflexology, a trained practitioner applies pressure to the feet, hands, ears, and face with specific thumb, finger, and hand techniques with the goal of stimulating the reflex areas to promote physiological change in the body. The theory behind reflexology states that specific areas on the feet and hands correspond to specific glands, organs, and other parts of the body, which are stimulated in order to help numerous health problems.⁵⁴ Reflexology is used to cause relaxation and healing in those specific stimulated parts of the body. In oncology, reflexology is often used to promote relaxation and improve quality of life.

Relaxation techniques. A variety of techniques are employed to promote relaxation in cancer patients. The National Cancer Institute (NCI) defines relaxation techniques as including progressive muscle relaxation (PMR), guided imagery, autogenic training, biofeedback, self-hypnosis, and deep breathing exercises.⁵⁵ PMR focuses on the tightening and relaxation of specific successive muscle groups and is usually combined with breathing and imagery exercises.⁵⁶ Guided imagery can be self-directed or led by a practitioner or a recording and often involves focusing on pleasant imagery to replace negative or stressful feelings.⁵⁶ Autogenic training involves concentrating on physical sensations of warmth, heaviness, and relaxation in different parts of the body.⁵⁶ Biofeedback uses electronic devices to monitor and teach control of certain bodily functions, such as breathing or heart rate, in order to facilitate relaxation.⁵⁶ Self-hypnosis refers to the training of patients to learn to induce a hypnotic state, a natural state of aroused, attentive focal concentration along with a relative suspension of peripheral awareness, on their own or when prompted by a phrase or a cue.⁵⁶ Deep breathing exercises involve the use of slow, deep, and even breaths, sometimes called diaphragmatic or belly breathing.⁵⁶

Qigong. The word qigong consists of two Chinese words: ‘*qi*’ (chee) meaning life force or vital energy that flows through all things in the universe; and ‘*gong*’ (gung) meaning accomplishment or skill that is cultivated through steady practice.⁵⁷ Qigong is a form of ancient and traditional Chinese medicine that integrates movement (physical postures), meditation (focused attention), and controlled breathing. Qigong aims to enhance vital energy or life force that balances a patient’s spiritual, emotional, mental and physical health. Qigong practices are used to increase the *qi*, circulate it, use it to cleanse and heal the body, store it, or emit *qi* to help heal others. Practices range in intensity from the gentle movements of Tai Chi to the more vigorous practice of Kung Fu.⁵⁷ In cancer patients, qigong is often used to reduce anxiety, fatigue, and pain, support the immune system, and improve physical and emotional balance.

Stress management. Acute stress is a normal physical and emotional reaction that people experience as they encounter changes in life⁵⁸, including following a cancer diagnosis, during cancer treatment, and during cancer survivorship. Long-term chronic stress may contribute to or worsen a range of health problems including digestive disorders, headaches, sleep disorders, depression, anxiety, and other mental health problems.⁵⁸ To address stress and induce the relaxation response, stress management programs teach such techniques as progressive muscle relaxation (PMR), guided imagery, and breathing exercises. Stress management also typically incorporates elements of cognitive-behavioral therapy, such as understanding the effects of appraisal and perception on the experience of subjective stress.^{59, 60} Participants are taught coping skills and practice various techniques for cognitive re-appraisal. One common structured group stress reduction program studied in oncology is called Cognitive Behavioral Stress Management (CBSM).^{61, 62} There are overlaps in some techniques used in stress management, relaxation, and meditation therapies. For example, meditation, guided imagery, and yoga may be practiced as techniques in isolation or combined. In this review, we distinguish between stress management, relaxation, and meditation interventions. Stress management interventions include psychoeducation on stress and coping, and emphasize cognitive-behavioral therapy and coping skills training; relaxation interventions typically consist of PMR and guided imagery; and meditation interventions use some form of meditation practice as the focal point of the training.

Soy. Soy is a plant in the pea family that has been common in Asian diets for thousands of year and more recently in the American diet.³² Soybeans are the seeds of the soy plant and contain isoflavones and soy protein. Soy is available in dietary supplements in tablet or capsule form, and contains isoflavones and/or soy protein. Soybeans can be cooked or eaten or used to make tofu, soy milk, and other food products. Soy is also used as an additive to other processed foods such as baked goods. Soy is used to treat menopausal symptoms, osteoporosis, memory problems, high blood pressure, and high cholesterol levels.³² In cancer patients, soy is often used to treat hot flashes.

Yoga. Yoga is a mind-body practice with origins in ancient South Asian philosophy and practice.⁶³ The term “yoga” is derived from the Sanskrit word *yug*, meaning “yoke” or “union”.⁶⁴ This, according to traditional yoga philosophy, is the ultimate intent of a yoga practice—to unite the individual with the totality of the universe. The techniques of yoga include ethical daily living (*yamas and niyamas*), physical postures (*asanas*), breathing techniques (*pranayama*), and meditation training (*dhyana*). There are a wide range of yoga forms and styles. The most commonly practiced form of yoga in the U.S. and Canada is Hatha yoga, which emphasizes postures (*asanas*) and often breathing exercises (*pranayama*). In cancer patients, yoga is used for a variety of conditions, including stress, anxiety, depression, and fatigue, as well as a method to increase physical activity.

LITERATURE REVIEW ON USE OF COMPLEMENTARY AND INTEGRATIVE THERAPIES FOR CLINICAL OUTCOMES IN BREAST CANCER PATIENTS

The clinical outcomes addressed here are common symptoms and side effects that breast cancer patients experience during treatment or as sequelae of treatment. The outcomes of interest include: anxiety/stress, pain, depression/mood, fatigue, sleep disturbances, quality of life and physical functioning, CINV, radiation dermatitis, vasomotor outcomes, lymphedema, chemotherapy-induced peripheral neuropathy (CIPN), pain and sleep disturbance. Guidelines outlining conventional approaches to managing these symptoms and outcomes have been issued by national organizations such as NCCN and the American Society of Clinical Oncology (ASCO), but many of the prior guidelines and reviews have

not included thorough reviews of complementary and integrative medicine approaches.⁶⁵⁻⁷¹ This review fills that gap for breast cancer patients.

Below, for each therapy and clinical outcome of interest that received an A or B grade, we summarize the trials that contributed to the graded recommendation in order for the reader to have an understanding of the specific interventions that were tested. To provide additional context we provide a brief review of the literature on the use of the integrative therapy for conditions other than breast cancer. We also provide a risk/benefit assessment of each therapy as well as suggestions for future research. For therapies and clinical outcomes of interest that received C, D or H grades, we provide a brief overview of the rationale for the graded recommendation. Supplemental tables provide detailed information on each trial that informed an A or B graded recommendations (Supplemental Tables 2-5). Table 3 lists the clinical outcome and integrative therapy combinations that had insufficient evidence to make an A, B, C, D or H graded recommendation.

Updated Recommendations

While the majority of graded recommendations remained the same from the previously published guidelines,⁹ there are five noteworthy changes. For the outcome of anxiety and stress reduction, the use of meditation moved from grade B to grade A due to results of a fifth trial⁷², in addition to the previously published four trials.⁷³⁻⁷⁶, showing beneficial effects. The use of yoga for depression and mood disturbance was downgraded from grade A to grade B due to four new published studies demonstrating conflicting results.⁷⁷⁻⁸⁰ The use of yoga for improving quality of life changed from grade C to grade B due to two added trials showing beneficial effects.^{81, 82} Finally, new trials on the use of yoga^{77-79, 82} and hypnosis^{83, 84} for fatigue upgraded previous recommendations from grade I to grade C.

Use of Integrative Therapies for Anxiety/Stress Reduction

Description of anxiety/stress

Cancer patients may experience stress related to the life changes associated with a cancer diagnosis, both during and after treatment. Anxiety is categorized under CTCAE psychiatric disorders from grade 1 (mild symptoms and no intervention required) to grade 4 (life-threatening). Stress is often the result of life challenges that exceed the individual's perceived ability to cope, and is a common and normal reaction during cancer diagnosis and treatment. This stress is associated with symptoms of anxiety and somatic complaints that can significantly diminish quality of life.⁸⁵ Patients with anxiety may worry more frequently, have difficulty relaxing, or feel tense. Patients with cancer-related anxiety also may have elevated heart rate, myalgias, headaches, sleep disturbances, changes in appetite, nausea, diarrhea, and difficulty concentrating. The percentage of breast cancer patients reporting anxiety ranges from 12% to 47%, with approximately 11-16% of patients experiencing combined symptoms of anxiety and depression.⁸⁶⁻⁸⁸ Evidence suggests that effective anxiety management is associated with improvements in quality of life, psychological adjustment, understanding of the disease, decision making, and adherence to treatment.⁸⁹⁻⁹¹

Meditation (A grade)

Overview of meditation interventions for anxiety/stress reduction

Meditation is recommended for reducing anxiety in breast cancer patients including during radiation therapy (Grade A). Many uncontrolled trials have been published, but this recommendation is based on five RCTs completed between 2009 and 2013 that used meditation to reduce anxiety symptoms (Supplemental Table 2).⁷²⁻⁷⁶ Anxiety was the primary outcome for four of the trials. In all five studies, a meditation intervention was compared with a usual-care control condition. Study participants included women undergoing radiation or chemotherapy, breast cancer survivors who had completed treatment, and older adult breast cancer survivors ages 50 years and older. The study sample sizes ranged from 49 to 336 participants. Among these trials, three types of meditation interventions were tested. Three trials implemented an intensive, integrated MBSR program, customized for breast cancer patients, in which participants were trained in mindfulness meditation and gentle yoga for body awareness.^{72, 75, 76} A fourth

intervention was called the Mindful Movement Program and was also an intensive, integrated program, customized for breast cancer patients, that included mindful walking/moving, group discussion, exploration of body parts, specific and deliberate movements, moving with intentional effort, active energetic movement, and partner work.⁷³ The fifth trial assessed a Brain Wave Vibration meditation⁷⁴, or a mind/body training technique that combines simple, rhythmic movements with music, action, and positive messages.⁷²

A systematic review and meta-analysis examined meditation in terms of its ability to reduce general psychological distress and stress-related health problems in adult clinical populations with a variety of health conditions; this analysis included 47 trials with 3,515 participants.⁹² Overall, mindfulness meditation programs showed moderate evidence of improved anxiety at 8 weeks and at 3-6 months and showed low evidence of improved stress/distress and mental health-related quality of life. The findings of these reviews across other patient populations and disease types support our recommendations.

The earliest work in MBSR interventions specifically showed sustained benefits for individuals with anxiety disorders, and more recent research has continued to show a benefit for generalized anxiety.⁹³⁻⁹⁵ The first study conducted with cancer patients, a RCT of 89 patients with a variety of cancer types, found substantial decreases in anxiety for the group receiving MBSR compared with results for a usual-care control group; results for the MBSR interventions were maintained at 6-month follow-up.^{96, 97} The reduction in anxiety observed in the above trials, specifically those that used more traditional forms of MBSR, provide support for the recommendation that meditation can be beneficial for the management of anxiety in women with breast cancer.

A recent systematic review and meta-analysis of 22 studies examined the effect of mindfulness-based therapy specifically on symptoms of anxiety and depression in adult cancer patients and survivors where 21 studies included a substantial percentage or all breast cancer patients.⁹⁸ Overall, this review included 12 nonrandomized studies and RCTs. In the nonrandomized studies, mindfulness-based therapy was associated with significantly reduced symptoms of anxiety post-intervention with a moderate effect

size, while the pooled effects sizes of RCTs, including the one listed above,⁷⁵ resulted in a larger effect size ($P < 0.001$). Although the review reported that overall study quality varied among studies included, there appears to be sound evidence from carefully conducted RCTs ($n=9$) supporting the use of mindfulness-based therapies for the management of anxiety in breast cancer patients and survivors. Another review of nine studies (including two RCTs, one quasi-experimental case-control study, and six single-group pre-post intervention studies) investigated the efficacy of MBSR on mental health specifically in breast cancer patients and found an overall decrease in anxiety scores following MBSR ($P < 0.01$).⁹⁹ This review provides further support for the use of MBSR to manage anxiety, specifically in women with breast cancer. Many other reviews reached similar conclusions, reporting positive moderate effect sizes of mind-body interventions on anxiety and distress in cancer.^{47, 100}

Risk/Benefit assessment of meditation interventions

Meditation therapies pose very little risk to participants in this type of intervention. Few adverse events have been reported in any trials, but there has been recent interest within the meditation research community in exploring adverse reactions to intensive meditation practice, particularly in vulnerable individuals.^{101, 102} Typically, potential participants are screened through individual orientation interviews prior to joining meditation group programs, and participants who have serious mental health issues are often re-directed to individual counseling or psychiatric intervention prior to or concomitant with mind-body intervention participation. Group facilitators are typically mental health care professionals trained to identify and manage psychological symptoms and reactions that may occur during the training.

Typically, participants in these meditation therapy groups report that the sessions are enjoyable, and dropout rates are often low and comparable to other psychosocial group programs. Because they are offered in group settings, meditation interventions are more cost-effective than traditional individual counseling or psychotherapy are and can often achieve similar results. However, the literature on meditation therapy is lacking in head-to-head comparisons with other forms of therapy, including individual counseling, cognitive-behavioral therapy, or other mind-body interventions. Hence, the

specificity of meditation therapy is not yet known. With the exception of Carlson et al.,⁷² studies have not compared mindfulness-based interventions with other effective interventions. Other research suggests that the benefit is related to the degree of the participant's engagement in and commitment to the practice,^{96, 103} in that participants who practice more at home often benefit more, but this area is still being investigated.¹⁰⁴ Drawbacks of these types of group interventions are the requirement for highly trained facilitators and the need for and ability of participants to attend in person, usually in large cities with tertiary cancer centers. In response to these issues, online and home-based adaptations of MBIs are being developed. For example, Zernicke et al.¹⁰⁵ demonstrated that an online live MBI group in which rural and remotely located cancer patients participated weekly over eight weeks had similar benefit to the on-site, in-person version, and patients were highly satisfied with the remote MBI adaptations.

Future research in meditation interventions for anxiety reduction

Future research on the use of meditation interventions for anxiety can similarly test novel interventions in populations who may not have ready access to in-person meditation programs.

Music therapy (B grade)

Overview of music therapy interventions for anxiety/stress reduction

Passive music therapy is recommended to reduce anxiety during radiation therapy, chemotherapy sessions, and post-surgery (Grade B), based upon results from five randomized clinical trials (RCTs) comparing music therapy interventions to standard care (Supplemental Table 2).¹⁰⁶⁻¹¹⁰ Study participants included breast cancer patients undergoing mastectomy, chemotherapy, and/or radiation therapy. The sample sizes of these studies ranged from 30 to 170 participants. The music therapy interventions were described as either *passive* or *active* music therapy, depending on the level of engagement required by the individual. Four trials examined the effect of passive music therapy, which was found to decrease anxiety scores in the intervention group^{106, 107, 109, 110}, including reducing sedation requirements during radiation therapy (measured as a secondary outcome).^{106, 107, 109} The fifth trial, conducted by Hanser et al.,¹⁰⁸

examined active music therapy (which required active engagement of the participant) and yielded substantially different results from the four trials of passive music therapy. In this study, active music therapy did not result in decreases in anxiety. The discrepancy in trial results between passive and active music therapy might have occurred because the participant engagement required for active therapy does not include the potential relaxing components of passive music therapy (e.g., listening to music).

A recent systematic review and meta-analysis by Boehm et al.¹¹¹ assessed the effect of different expressive therapies, including passive and active music therapy, on improving anxiety, depression, and quality of life in breast cancer patients. The review included three of the RCTs that formed the basis for our recommendation of passive music therapy for anxiety/stress reduction, of which two trials tested passive music therapy^{106, 109} and one trial tested active music therapy.¹⁰⁸ Another recent meta-analysis included one additional RCT evaluating art therapy.¹¹² Boehm et al. found a clinically and statistically significant mean difference ($P < 0.01$) in the anxiety scores of patients receiving music therapy compared to the control group,¹¹¹ thus further supporting our recommendation of passive music therapy for reducing anxiety. Passive music therapy has also been shown to reduce anxiety among patients undergoing mammographic screening indicating that the recommendation may apply broadly to an adult women in a clinical cancer setting.¹¹³

Risk/benefit assessment of music therapy

Passive music therapy is noninvasive, does not interfere with a patient's privacy, and has no reported deleterious effects. Furthermore, it does not require costly, technologically advanced equipment and can be implemented in a variety of locations. As such, passive music therapy can be safely and effectively implemented in clinical settings with breast cancer patients to help reduce short-term anxiety associated with receiving medical care for their cancer.

Future research in music therapy for anxiety reduction

Future research should assess long-term effects of passive music therapy on anxiety, as the trials reviewed here only assessed short-term reductions in anxiety. Although active music therapy may also have benefit, more than one existing trial will be needed for a comprehensive risk/benefit assessment. Trials that directly compare the effect of passive music therapy and active music therapy on anxiety are needed to clarify whether benefit is due to listening to music in a relaxed state or participating in the creation of music.

Stress Management (B grade)

Overview of stress management interventions for anxiety/stress reduction

Stress management is recommended to reduce anxiety in patients during breast cancer treatment, (Grade B) but long-term stress management group programs appear to be better than self-administered home programs. This recommendation is based on four RCTs completed in 2008-2013 with breast cancer patients that tested a stress management intervention compared with usual care in improving anxiety as the primary outcome.¹¹⁴⁻¹¹⁷ Study participants included only breast cancer patients in two^{114, 117} of the four trials and mixed cancer patients with a significant proportion of breast cancer patients in the other two.^{115, 116} For all four trials, breast cancer patients were included from defined periods along the continuum of care, including pre-surgery, during chemotherapy, and after cancer treatment. Patients were thus either undergoing or had undergone surgery, chemotherapy, or radiation therapy, either alone or in combination. The study sample sizes for these trials ranged from 85 to 286 participants. Of the four trials, two studies found that the improvement was statistically significantly different compared to the control group.^{115, 117} One trial found similar improvement in both the intervention and control group with no statistically significant differences across groups,¹¹⁴ while the fourth trial found no improvement.¹¹⁶ (see Supplemental Table 2, Stress Management).

Stress management interventions varied among the studies (Supplemental Table 2).¹¹⁴⁻¹¹⁷ One trial¹¹⁴ implemented a self-administered stress management intervention before receiving chemotherapy. The intervention included video and booklet information specific to stress management and exercise and

was delivered by a doctoral level psychologist.¹¹⁴ A second trial¹¹⁴ offered stress management modalities that included guided imagery techniques, relaxation, meditative exercises, and counseling that aimed to promote active coping, alert relaxation, and a positive attitude toward change. Instruction was given in person and was complemented by audio CDs for use at home. Face-to-face sessions lasted 45-60 minutes each and took place in the hospital.¹¹⁴ The stress management portion included information, demonstrations and instructions for paced breathing, progressive muscle relaxation with guided imagery, and use of coping strategies to manage stress. A third trial¹¹⁵, a multi-center community clinical oncology program involving 20 clinical sites, was designed to determine whether a self-administered stress management intervention, previously found to be beneficial to cancer patients, would improve quality of life and decrease psychological distress (including anxiety and depression) in patients receiving chemotherapy at community clinical centers. The study used a factorial design to test stress management training, exercise training, the combination of stress management and exercise training, and usual care. Patients assigned to stress management training were provided written materials and videos on three techniques, including progressive muscle relaxation and guided imagery, abdominal breathing, and coping skills training.¹¹⁵ A fourth trial¹¹⁷ examined the effects of a cognitive behavioral stress management intervention (comprised of relaxation training including meditation, guided imagery, PMR, and breathing techniques, cognitive restructuring, and coping skills training) on late-afternoon serum cortisol and relaxation indicators in women undergoing treatment for non-metastatic breast cancer.

A systematic review by Trijsburg et al. evaluated the effects of stress management techniques on mental health outcomes, including anxiety, across 22 studies using samples of heterogeneous cancer populations that included six studies of breast cancer-only samples.¹¹⁸ The interventions assessed were all structured and included counseling and coping components in addition to some interventions including progressive muscle relaxation, guided imagery, self-hypnosis, and deep breathing. Overall, the review found positive effects for mental health outcomes, including anxiety scores.

Risk/benefit assessment of stress management interventions

Stress management therapy is noninvasive, nontoxic, and has no appreciable risk to patients. Stress management does not require specialized equipment and can be implemented in a variety of locations. Resources for providing instruction on self-management practices are broadly available. As the majority of the trials discussed above tested long-term interventions, the guideline recommendation is specific to long-term stress management programs. This may, however, be complicated by other factors, including the potential increase to the cost of delivery for providers/facilitators as a result of the implementation of longer term programs and could also present a barrier to patients with access-to-care and other health care disparity issues. Overall, it is recommended that stress management therapy can be safely and effectively implemented in clinical settings for breast cancer patients.

Future research in stress management interventions for anxiety reduction

Future research in this area should compare in-person, professionally led stress management group programs to home-based self-study and internet-based stress management intervention options that are less costly and more broadly accessible. Research should also examine long-term outcomes of each type of delivery modality.

Yoga (B grade)

Overview of yoga interventions for anxiety/stress reduction

Yoga is recommended for reducing anxiety in breast cancer patients (Grade B). This recommendation is based on nine RCTs, completed between 2007 and 2014, in which a yoga intervention was implemented to reduce physical symptoms and psychological distress, including anxiety, which assessed as the primary outcome in four of the studies (Supplemental Table 2).^{79, 119-126} These nine trials tested five different yoga interventions. The first intervention was an intensive, integrated yoga program, customized for breast cancer patients, in which participants were led through slow stretching and loosening exercises, various postures (*asanas*), guided imagery specific to cancer, positive thought provocation, chanting exercises, various breathing exercises (*pranayama*), and soothing sound vibrations

and guided imagery (*yoga nidra*).¹¹⁹⁻¹²² A second intervention implemented *Iyengar* yoga, a traditional form of Hatha yoga that emphasizes postures and breathing techniques that target symptom-specific concerns using passive inversions (upside-down postures with the head lower than the heart) and passive backbends (supported spinal extensions).¹²³ A third intervention used Patanjali's Yoga Sutras that included warm-up movements synchronized with breathing, selected postures, deep relaxation techniques, alternate-nostril breathing (*pranayama*), and meditation.¹²⁶ The fourth intervention implemented meditation and breathing exercises that focused attention on internal body sensations, as well as yoga exercises (modified *asanas*) composed of gentle stretching and strengthening exercises.⁷⁹ *Pranayama* or yoga breathing practices was tested in the remaining trials.^{124, 125} Study participants included women who had been recently diagnosed with breast cancer, those who were currently undergoing radiation or chemotherapy or a combination of both, and those who were experiencing significant fatigue post treatment. The study sample sizes ranged from 23 to 98 participants.

Several recent reviews of yoga interventions for individuals with cancer have been published.¹²⁷⁻¹³⁰ The first meta-analysis investigating the impact of yoga interventions on psychological health outcomes¹³¹ identified 10 articles that examined outcomes in breast cancer patients (n=7 articles), lymphoma patients (n=1), and mixed cancer populations (n=2). Analyses involving eight trials that assessed anxiety found lower anxiety associated with the yoga groups relative to controls ($P = 0.009$); similar results with yoga interventions were observed for distress ($P = 0.003$) and stress ($P = 0.006$). However, since this was a heterogeneous sample of patients, it was unclear whether the results could be generalized specifically to breast cancer patients. More recently, a systematic review and meta-analysis assessing the impact of yoga interventions on quality of life and psychological health specifically in breast cancer patients and survivors was conducted.¹³² Overall, 12 randomized clinical trials were included in the analysis with a total of 742 participants. Analyses revealed short-term positive effects of yoga interventions on improved psychological health, including anxiety ($P < 0.01$), perceived stress ($P = 0.03$), and psychological distress ($P < 0.01$). It is, however, notable that these effects were applicable only to those who engaged in yoga during active cancer treatment and not in the post-treatment period. The

authors state that based on these positive preliminary results, a wide variety of yoga practices could be recommended for this population.¹³² Finally, although the methodological quality across the trials is variable, yoga has been shown to be beneficial for reducing psychological distress in a wide variety of non-cancer clinical populations, including adult patients with anxiety and depression.^{133, 134}

Risk/Benefit assessment of yoga interventions

Yoga interventions are noninvasive and with proper instruction can be adapted to people with functional and other impairments without harm. Yoga interventions are low cost and can be practiced at home with instructional videos but in the context of breast cancer are best undertaken under the guidance of certified yoga instructors with specific training in teaching cancer patients and survivors. Such programs also have the ability to be adapted and modified for people with medical conditions or limited mobility.¹³⁵ Older adults, individuals with limited mobility, and those with chronic medical conditions should proceed with yoga therapy only under the guidance of a certified instructor to minimize potential risk of harms such as strained muscles and dizziness if yoga postures are attempted incorrectly or prematurely.

Future research in yoga interventions for anxiety reduction

Many of the studies investigating the impact of yoga on psychological outcomes in individuals with cancer should be interpreted with caution given that many of these studies are small and preliminary. Furthermore, the studies summarized above have some inconsistency of results; the trials from India¹¹⁹⁻¹²² reported consistently positive results while the trials from North America^{79, 123-126} reported less consistent positive results. Nevertheless, many of the studies and reviews that examined yoga interventions reported overall positive outcomes in several physical, psychological, and quality of life measures. Future trials of yoga interventions for anxiety/stress reduction should focus on testing forms of yoga that can be more easily applied, and in order to improve the generalizability of the results, test the interventions in larger

sample size and in minority and underserved populations who may not have easy access to yoga programs.

C graded therapies for anxiety and stress reduction

Acupuncture¹³⁶⁻¹³⁸, massage¹³⁹⁻¹⁴² and relaxation¹⁴³⁻¹⁴⁷ can be considered for use for anxiety and stress reduction (Grades C). Three high-quality trials assessed acupuncture for anxiety and stress reduction, which were secondary outcomes; two of the three positive trials were small^{136, 137} and one trial showed no effect¹³⁶. Four studies examined massage for anxiety, however, three studies with positive findings included fewer than 40 participants¹³⁹⁻¹⁴¹ and one study found no effect¹³⁹. Results were inconclusive regarding relaxation for anxiety and stress reduction due to inconsistent results and small sample sizes. Future directions in research can focus on assessing these modalities in a large high-quality trial assessing anxiety as the primary outcome.

Use of Integrative Therapies for Depression/Mood Disturbances

Description of depression/mood disturbances

Both during and after cancer therapy, patients may experience symptoms of depression, as they often feel a sense of loss of health and the life they had before their cancer diagnosis.⁸⁵ The CTCAE categorizes depression as a psychiatric disorder on a scale of 1 (mild) to 5 (death). Symptoms of depression in cancer patients may include persistent feelings of sadness, numbness, nervousness, guilt, worthlessness, helplessness or hopelessness, difficulty concentrating or behavior that includes being short-tempered or moody, crying for long periods of time or many times each day, lacking interest or pleasure in performing activities, and having suicidal thoughts. Other symptoms may include weight change, sleep disturbances, tachycardia, dry mouth, increased perspiration, gastrointestinal symptoms, diarrhea, changes in energy level, persistent fatigue, headaches, or myalgias. The percentage of breast cancer patients reporting depression ranges from 3% to 34% and 11-16% of patients experienced combined depression and anxiety symptoms, depending on the population studied.⁸⁶⁻⁸⁸ Effectively

managing depression may improve quality of life, psychological adjustment, understanding of the disease, decision making, adherence with cancer treatment, and response to cancer treatment.⁸⁹⁻⁹¹

Meditation (A grade)

Overview of meditation interventions for depression/mood disturbances

Meditation, particularly MBSR, is recommended for treating mood disturbance and depressive symptoms in breast cancer patients (Grade A). This recommendation is based on 10 RCTs, completed between 2009 and 2015 that used meditation to help reduce depressive symptoms (Supplemental Table 3).^{72-76, 148-152} Depression was the primary or secondary outcome for all of the trials. In eight^{73-76, 148-150, 152} of the 10 trials, a meditation intervention was compared with a usual care group, a waitlist control, or another active intervention; two other trials used a three-arm trial design.^{72, 151} Study participants included women undergoing current radiation therapy or chemotherapy for breast cancer, breast cancer survivors who had completed treatment, and adult breast cancer survivors 55 years and older. The study sample sizes ranged from 33 to 336 participants and tested six different types of meditation interventions, including an intensive, integrated MBSR program customized for breast cancer patients^{72, 75, 76, 149, 151}, the Mindful Movement Program⁷³, Brain Wave Vibration meditation⁷⁴, Tibetan sound meditation¹⁴⁸, Cognitively-Based Compassion Training¹⁵², and Transcendental Meditation¹⁵⁰.

In the meta-analysis examining the effect of mindfulness-based therapy on psychological outcomes in adult cancer populations,⁹⁸ compared with results in controls, mindfulness-based therapy was associated with significantly reduced depression post-intervention, with a moderate effect size in the nonrandomized studies and the RCTs, including one trial listed above⁷⁵ ($P < 0.001$). Using evidence from nine well-conducted studies, the review and meta-analysis by Zainal et al. confirms the use of mindfulness-based therapies for the management of depression in breast cancer patients and survivors despite the heterogeneity in the reviewed studies; the results of the meta-analysis found a significant pooled effect size for MBSR on depression scores (effect size = 0.575; 95% CI = 0.429, 0.722; $P < 0.01$).⁴⁶ In their study, Teasdale et al.¹⁵³ modified traditional MBSR by combining it with principles of

cognitive behavioral therapy (CBT) to create mindfulness-based cognitive therapy (MBCT), which was designed specifically to prevent recurrence of depressive symptoms in individuals with relapsed major depression.¹⁵⁴ The intervention proved effective for preventing depression relapse¹⁵³ in a sample of >75% women with cancer (though type of cancer is not specified), and has since been widely applied and adapted to treat depression symptoms in a range of clinical samples.¹⁵⁵ In cancer patients, other RCTs of meditative interventions showed decreasing depressive symptoms,⁹⁶ with results maintained after 6-months' follow-up.⁹⁷ The reduction in depressive symptoms observed in the above trials comprising these reviews and meta-analyses provide further support for the recommendation that meditation can be beneficial for the management of depressive symptoms in women with breast cancer.

Risk/benefit assessment of meditation interventions

The risk/benefit assessment of the effects of meditation intervention on depression and mood disturbance outcomes is similar to the assessment of anxiety outcome. Despite these limitations, the evidence suggests that meditation and MBSR could be added to treatment plans or added in the post-treatment period, provided that these interventions are facilitated by appropriately trained instructors and can be adapted and modified if needed for individuals with cancer.

Future research in meditation interventions for depression/mood

Future research on meditation interventions to improve mood disturbances/depression should focus on understanding the type, duration and lasting effect of specific meditation techniques on specific mental health outcomes.

Relaxation (A grade)

Overview of relaxation interventions for depression/mood disturbances

Relaxation therapy is recommended for improving mood disturbances and depressive symptoms when added to standard care (Grade A). The recommendation put forth in the guidelines is based on

results from six RCTs, completed between 1999 and 2007, which implemented a relaxation program with or without guided imagery as an intervention to improve mood disturbances and treat depression (Supplemental Table 3).^{143, 146, 147, 156-158} Five of the six trials measured depression as the primary outcome. In all six trials, a relaxation therapy intervention group was compared with a standard care group. The interventions comprised some form of relaxation therapy that included progressive muscle relaxation and guided imagery or visualization techniques. One trial¹⁴³ assessed autogenic training that provided training in relaxation that is meant to induce heaviness and warmth of limbs, calming of the heart and breathing, abdominal warmth, and cooling of the forehead. Limitations of all of these trials included the potential for expectation bias attributable to the inability to blind participants and a lack of attention control groups, although one study used a health education attention control.¹⁵⁸ Participants in these studies included breast cancer patients who had or were currently undergoing surgery, chemotherapy, or radiation therapy. The study sample sizes ranged from 31 to 183 participants.

The majority of pertinent systematic reviews combine relaxation techniques with stress management, psychosocial, and psychological interventions for breast cancer patients. Thus, a review specific to relaxation interventions that includes progressive muscle relaxation and guided imagery for depression/mood or other psychological outcomes in breast cancer patients and survivors is warranted. A review study of guided imagery as adjuvant cancer therapy broadly assessed six RCTs and found the methodological quality inconsistent.¹⁵⁹ Across trials, the results provided few details, and the studies were implemented with heterogeneous cancer populations, interventions, and outcome measures, which ultimately precluded statistical pooling of the results. Despite these limitations, guided imagery as a sole adjuvant cancer therapy was found to be supportive and to increase comfort in patients with few risks.

Beyond decreasing depression, relaxation therapy may also have a beneficial impact on other symptoms important to women with breast cancer. For instance, progressive muscle relaxation has been found effective at ameliorating sleep problems and fatigue in women undergoing chemotherapy for breast cancer.¹⁶⁰ Other likely benefits attributable to this therapy in this breast cancer population include reduced nausea and anxiety.¹⁶¹⁻¹⁶⁴

Risk/benefit assessment of relaxation interventions

Relaxation therapy is non-invasive and positively engages the patient with very little potential for harm. The goal of this treatment approach is to use principles of psychoneuroimmunology to better regulate the hypothalamic pituitary-adrenal axis, modulate cortisol production, and decrease stress, which may have other health benefits including psychological outcomes that may be affected by stress and distress (such as anxiety and depression/mood disturbances). Some of the appeal of relaxation therapy includes its low cost, safety, and portability. With adequate training, patients themselves can apply this therapy when and where they want without the need for supervision. However, the durability of relaxation therapy and the frequency required to sustain a long-term positive effect on depression remain unclear. It seems that relaxation therapy is not only applicable for those with breast cancer and depression but also has been shown to be beneficial in other contexts, including to decrease depression in adults^{165, 166} and depression associated with cardiac disease.¹⁶⁷ The minimal cost and low potential for harm with relaxation therapy in conjunction with its evidence of benefit supports its A grade recommendation for depression.

Future research in relaxation interventions for depression/mood

In addition to the gaps in the literature described above, future research on the use of relaxation interventions to improve mood disturbances should focus on how to use novel electronic communication strategies to deliver low-cost relaxation techniques to diverse patient populations.

Yoga (B grade)

Overview of yoga interventions for depression/mood disturbances

Yoga is recommended for improving mood disturbances and depressive symptoms in women with breast cancer (Grade B). This recommendation is based on the results of 15 RCTs completed

between 2006 and 2015 that employed a yoga intervention to reduce physical symptoms and psychological distress, including depression (Supplemental Table 3).^{77-80, 119-121, 123-126, 168-171} Depression or depressive symptoms were the primary or secondary outcome for all studies included in this review. Study participants included women recently diagnosed with or having a recurrence of breast cancer; women who were undergoing radiation therapy, chemotherapy, or a combination of both; an ethnically diverse and underserved sample of female patients, and women who had completed breast cancer treatment. Five different yoga interventions were tested in study samples ranging from 23 to 200 participants. Five trials assessed an intensive, integrated yoga program, customized for breast cancer patients, including *asanas*, *pranayama*, and *yoga nidra*.^{80, 119-121, 171} Five trials implemented *Iyengar* yoga, a traditional form of Hatha yoga, and passive backbends.^{78, 123, 168-170} Two trials assessed Patanjali's Yoga Sutras that included warm-up movements synchronized with breathing, selected postures, deep relaxation techniques, meditation, and alternate-nostril breathing (*Pranayama*),^{77, 126} while two trials evaluated only the *Pranayama* practices.^{124, 125} Finally, a yoga exercise intervention implemented meditation and breathing exercises that focused attention on internal body sensations, as well as yoga exercises (modified *asanas*) composed of gentle stretching and strengthening exercises.⁷⁹

Yoga has been shown to be beneficial for reducing depression in a wide variety of clinical populations.^{134, 172-174} and specifically for individuals with cancer.¹²⁷⁻¹³⁰ One meta-analysis investigating the impact of yoga interventions on psychological health outcomes¹³¹ analyzed eight trials of yoga interventions for depression and showed improvement in depressive symptoms in the yoga groups when compared with patients in the control groups ($P = 0.002$) among a heterogeneous sample of patients. Specifically in breast patients and survivors, a meta-analysis¹³² of 12 RCTs, representing a total of 742 participants, revealed short-term effects of yoga interventions on improved psychological health, including depression ($P < 0.01$). A caveat was that the observed efficacy was only applicable for yoga practiced during active cancer treatment and not necessarily in the post-treatment period. Overall, the authors state that based on these positive preliminary results, yoga therapy should be used in this population.

Risk/benefit assessment of yoga interventions

The risk/benefit assessment of the effects of yoga on depression and mood disturbance outcomes is similar to the assessment for anxiety outcomes. Studies assessing yoga for psychological outcomes in individuals with cancer are typically small, not well controlled, and preliminary in nature. Furthermore, comparability across studies is difficult as findings differ between populations from India^{119-121, 171} and North America.^{123, 124, 126, 168-170} Despite these limitations, the evidence suggests that yoga interventions could be added to treatment plans or added in the post-treatment period, provided these interventions are facilitated by appropriately trained yoga instructors and can be adapted and modified for people with medical conditions or limited mobility.¹³⁵

Future research in yoga interventions for depression/mood

Future trials of yoga interventions to improve mood disturbances/depression should test the effects of different types, doses and durations of yoga on patient populations with varying degrees and types of mood disturbances and depression in addition to including larger sample sizes and testing across active controls.

Massage (B Grade)*Overview of massage therapy interventions for depression/mood disturbances*

Massage therapy is recommended to improve mood disturbance in breast cancer survivors following active treatment (i.e., surgery, chemotherapy, radiation) (Grade B). This recommendation is based on results from six trials completed between 2004 and 2012.^{140-142, 175-177} In the trial reported by Listing et al.,¹⁷⁶ the primary outcomes focused on physical discomfort and fatigue, with mood disturbance being a secondary outcome. The other five trials assessed depression as the primary outcome and had other secondary outcomes, including perceived stress, quality of life, pain, heart rate variability, and emesis. In the five trials, the effect of massage therapy was compared with the effect of standard care. The

studies, which took place in the United States,¹⁴⁰ Germany^{141, 175, 176}, England¹⁴², and Spain,¹⁷⁷ all included breast cancer survivors who had completed active cancer treatment at least three months prior to study enrollment. The study sample sizes ranged from 20 to 288 participants. In three of the six trials, the massage therapy intervention was a variation of classic massage involving rhythmic stroking, kneading, and acupressure at select areas on the body.^{141, 175, 176} The study of Fernandez-Lao used an experienced and trained physical therapist to administer manual massage therapy.¹⁷⁷ Wilkinson et al.¹⁴² included aromatherapy as part of the massage therapy, which was individualized across study participants. Hernandez-Reif et al.¹⁴⁰ combined massage with acupressure and Trager®, which uses hundreds of small, rocking and elongating movements that release muscle tension. The number and frequency of massage therapy sessions varied across the studies.

All of the trials assessing classic massage therapy used the same protocol of bi-weekly, 30-minute massages for five weeks^{141, 175, 176} with the exception of the Fernandez-Lao et al. trial, which assessed a one-time 40-minute massage session.¹⁷⁷ Across these studies, compared with control groups, women in the intervention group reported significant improvement in mental health outcomes, including reduced depression and mood disturbance scores particularly in relation to anxious depression, anger, and tiredness. These differences between groups, however, were not consistently sustained across all studies. For example, in the study of Fernandez-Lao et al., the improvements in depression outcomes with one-session massage intervention depended on the participant's individual level of enthusiasm for that intervention. The classic massage studies^{141, 175, 176} and one trial with massage combined with aromatherapy¹⁴² found that the massage interventions significantly decreases depression only immediately following the intervention but not long term. The remaining study that combined massage therapy with acupressure and Trager®¹⁴⁰ demonstrated a positive effect of the interventions on anxiety and depression with these effects sustained at long term follow-up.

A systematic review and meta-analysis examined the role of massage interventions in the management of treatment-related side effects associated with breast cancer in terms of improvement in overall quality of life, including measures of depression.¹⁷⁸ A total of 18 RCTs were included in this

review. Overall regular massage therapy was shown to have positive effects on other outcomes such as anger and fatigue symptoms, but the meta-analysis of eight trials, three of which were summarized in this section,^{140, 141, 176} did not find significant beneficial effects of massage therapy for depression. This meta-analysis, however, was limited by significant heterogeneity across the studies assessing depression as the outcome ($P = 0.002$) as well as by small and possibly underpowered sample sizes that may have accounted for the non-significant results. An earlier systematic review of massage therapy in breast cancer populations that found no positive effect of massage on depression also concluded that few rigorous trials have been conducted and that the risk of bias in such trials is high.¹⁷⁹

Risk/benefit assessment of massage therapy interventions

Classic massage is a non-invasive therapy that has limited adverse effects. For cancer patients, massage therapy by a trained massage professional appears to have few risks and may reduce pain, promote relaxation, and boost mood, at least in the short term.¹⁸⁰ Since trained, licensed therapists delivered the interventions tested in these studies, the recommendation of massage for depression applies specifically to massage by trained therapists.^{181, 182} Some caution is needed, however, for individuals receiving anti-coagulation therapy due to their risk of bruising. Women with breast cancer who have undergone radiation therapy or surgery or have implanted medical devices also may need to be cautious in having massage to the affected regions. In addition, some women may be reluctant to disrobe due to altered body image, modesty, or ethnocultural issues. Thus, important considerations include appropriate draping of the individual and ensuring that the massage therapist is sensitive to the treatment experiences of women with breast cancer and develops a therapeutic relationship with the individual. Ensuring that the massage therapist is the same gender as the patient is considered best practice in the massage profession and may also increase an individual's comfort with receiving massage therapy. However, this practice was not tested in the studies included in this review.

Future research in massage interventions for depression/mood

A 2008 systematic review of massage therapy for depression in the general public concluded that evidence to support massage as an effective treatment for this indication and population was lacking.¹⁸³ However, a more recent (2010) meta-analysis of 17 RCTs concluded that massage therapy had a positive effect on individuals suffering from depression.¹⁸⁴ That meta-analysis also highlighted the heterogeneity across massage therapy trials in terms of therapy protocols, outcomes measurement, and populations and underscored the need for standardization across future massage therapy trials. These issues are also true for the trials assessing massage interventions for depression/mood in breast cancer patients. Future research on massage therapy interventions to improve depression/mood disturbances should focus on understanding how best to disseminate cost-effective massage interventions in routine clinical settings.

Music therapy (B grade)

Overview of music therapy interventions

Passive music therapy is recommended to improve depression/mood disturbances in newly diagnosed breast cancer patients (Grade B). This recommendation is based on four RCTs completed between 200 and 2011 that tested a music therapy intervention to improve mood/depression (Supplemental Table 3).^{108, 110, 185, 186} Depression/mood disturbances was assessed as either the primary outcome or secondary outcome to anxiety. In all four trials, a music therapy intervention group was compared with either a waitlist or a standard care control group. Study participants included women with a breast cancer diagnosis who had completed mastectomy, chemotherapy, or radiation therapy or who had metastatic disease. The study sample sizes ranged from 8 to 170 participants. Three trials examined the effect of passive music therapy, which was found to decrease depression scores compared to control^{110, 185, 186}, while the fourth trial examining active music therapy¹⁰⁸ yielded no clinically meaningful long-term effects between groups or over time. Active music therapy resulted in immediate effects on happiness within the intervention group, which was not sustained over time. Therefore, the guideline recommendation is specific to passive music therapy.

Out of five studies combined in the meta-analysis described above that assessed multiple types of art therapies for improving anxiety, depression, and quality of life among breast cancer patients¹¹¹ (four music therapy interventions including two trials identified in our review^{108, 112}, and one art therapy trial), a clinically and statistically significant mean difference ($P = 0.05$) was found across all depression scores in the music therapy intervention groups compared with control groups. Further, a systematic review of music therapy specifically for depression in the Cochrane Database of Systematic Reviews found that although only five RCTs have tested music therapy interventions for depression and met the review study inclusion criteria, this therapy is widely accepted and beneficial to a broad range of individuals with depression and is associated with improvements in mood disturbances.¹⁸⁷ Taken together, this literature supports our recommendation of passive music therapy for reducing depression and improving mood.

Risk/benefit assessment of music therapy interventions

The risk/benefit assessment of passive and active music therapy interventions for depression/mood among breast cancer patients is the same as for anxiety depression (see above).

Future research in music therapy interventions for depression/mood

Future research in this area should assess long-term effects of passive music therapy on mood disturbances/depression because the reviewed trials are limited to assessing short-term improvement of mood/depression following breast cancer treatments or during metastatic cancer diagnosis. In addition, studies should attempt to replicate the null findings of active music therapy interventions in order to formulate a comprehensive risk/assessment regarding active music therapy. Suggestions for future research in music therapy for improving depression/mood are similar to those for anxiety, as stated in the anxiety section above.

C graded therapies for depression/mood

As trials have been conducted on the effects of acupuncture^{136-138, 188, 189}, healing touch^{34, 35}, and stress management^{114-116, 190, 191} on depression and improvement in mood, these therapies can be considered (Grades C). Five trials assessed acupuncture as a treatment for depression/mood (but as a secondary outcome) and only one trial was large,¹³⁸ with mixed findings in terms of effect. Two studies assessed the effect of healing touch on depression as a primary outcome, with one small study showing no effect³⁵ and a larger study showing a positive effect³⁴. Though five large studies were reviewed for the effect of stress management on depression/mood, findings were inconclusive due to inconsistent results across trials. Future research directions include conducting trials with larger samples sizes and replicating trials with these modalities to examine impact on depression symptoms and improving mood as the primary outcome.

Use of Integrative Therapies for Fatigue

Description of fatigue

Among cancer patients, fatigue is commonly referred to as cancer-related fatigue. Fatigue is a multi-factorial condition marked by extreme tiredness and inability to function due to lack of energy.¹⁹² According to National Comprehensive Cancer Network (NCCN) guidelines, ratings of fatigue of 4 or higher on a scale of 0 to 10 (where 10 is very severe fatigue) are further evaluated for known contributing factors such as pain, emotional distress, anemia, sleep, nutrition, and level of activity. These comorbidities are then treated. The CTCAE measures fatigue from grade 1 (relieved by rest) to grade 3 (not relieved by rest, limiting activities of daily living). Fatigue is the most frequent and distressing side effect of common antineoplastic therapies, including chemotherapy, radiation therapy, surgery, and selected biologic response modifiers.⁷⁰ Although cancer-related fatigue typically improves upon the conclusion of treatment, it can last for months or years in up to one-third of patients and become a chronic condition that leads to a variety of physical and psychological effects long into survivorship. Symptoms of fatigue include feeling tired, weak, worn-out, heavy, slow, or having “no energy or get-up-and-go”. Cancer-related fatigue is different from common tiredness in both its magnitude and quality. Cancer-

related fatigue is not typically relieved by rest and is much more profound than simply feeling tired. About 33% of patients with breast cancer experience moderate to severe fatigue.¹⁹³⁻¹⁹⁵ Fatigue has disruptive consequences and can have a negative impact on a patient's quality of life, mood, and self-esteem.⁶⁶ Cancer-related fatigue can prevent patients from taking part in daily activities, relationships, social events, community activities, as well as work or school, which can have financial consequences, such as loss of employment and health insurance.

C and D graded therapies for fatigue

There are no A or B graded therapies to report for fatigue. Trials evaluating hypnosis^{83, 84}, ginseng^{196, 197}, Acetyl-L-carnitine¹⁹⁸, and guarana^{199, 200} have examined their effects on fatigue during treatment. Two trials from a single research group reported beneficial effects of hypnosis on fatigue during treatment, therefore hypnosis can be considered for use for fatigue during treatment (Grade C). Neither Acetyl-L-carnitine nor guarana are recommended for improving fatigue during treatment due to a lack of effect in clinical trials among cancer patients (Grades D). Ginseng received a Grade C for fatigue during treatment based upon a single high quality trial with a large sample size.¹⁹⁶ A previous large, high quality dose finding study by the same group indicated that a higher dose was more effective.¹⁹⁷ In subset analyses in the subsequent trial, which tested use in patients reporting fatigue either during or after treatment, results showed that ginseng was more effective in patients undergoing active treatment. Trials testing the effects of acupuncture^{138, 201-203} and yoga^{78, 82, 123, 170, 204, 205} for post treatment fatigue yielded modest results (Grades C). Four high quality acupuncture trials assessed fatigue as the primary outcome; three of the four trials included a large sample size.^{138, 201-203} Acupuncture for post treatment fatigue received a C grade mainly due to inconsistent comparison groups across the trials, including sham acupuncture,^{201, 203} standard of care,^{138, 202} self-administered acupuncture,²⁰² and wait-list control.²⁰³ Four trials evaluated yoga for post treatment fatigue as a secondary outcome,^{82, 170, 204, 205} only two studies had a large sample size⁷⁸ where each reported contrasting results.

Use of Integrative Therapies for Quality of Life

Description of quality of life

Quality of life is a multi-dimensional construct typically measuring the functioning of emotional, physical, role, and social domains using validated questionnaires.²⁰⁶ The majority of breast cancer patients report some level of diminished quality of life during cancer treatment and/or survivorship.²⁰⁷ The physical domain includes common physical side effects of cancer and cancer treatment, including constipation and diarrhea, fatigue, hair loss, fever, hot flashes and night sweats, lymphedema, nausea and vomiting, poor nutrition, oral complications, pain, and skin changes, as well as the ability to function physically in everyday life. The emotional domain includes psychological functioning with indicators of anxiety, depression, distress, confusion, and memory problems. The sexual functioning domain refers to patients' perception of sexuality and sexual functioning, attractiveness, and fertility. Finally, the social domain refers to the patients' social functioning, their social role, and level of social support. Each of the domains, either individually or in combination, influences a patient's quality of life.²⁰⁸⁻²¹⁰ Decrements in quality of life may persist upon the conclusion of treatment, on chronic/long-term hormonal and biotherapies and into survivorship.²¹¹

Meditation (A grade)

Overview of meditation interventions for quality of life

Meditation is recommended for improving quality of life in breast cancer patients (Grade A). This recommendation is based on seven RCTs, completed between 2009 and 2013, which used meditation for this indication (Supplemental Table 4).^{73-75, 149-151, 212} Quality of life was the primary outcome in four of the seven trials.^{149-151, 212} In four trials, a meditation intervention group was compared with a usual care or waitlist control condition group, and three other trials^{149, 151, 212} used a three-arm design. Study participants included women undergoing radiation therapy for breast cancer, newly diagnosed breast cancer patients of stages 0-IV, breast cancer survivors who had completed treatment, and older adult breast cancer survivors aged 55 years and older. The study sample sizes ranged from 47 to 180 participants. These trials

overlap with the trials described under meditation interventions for anxiety/stress reduction and depression outcomes (see above).

Previous research has supported the role of MBSR interventions for improved quality of life in heterogeneous samples of cancer patients and survivors, with effect sizes ranging from small to large on quality of life scales.^{103, 213, 214} One review in particular assessed the impact of MBSR on quality of life in breast cancer patients.⁴⁷ In that review, the authors identified only three studies that met their criteria and also measured quality of life as an outcome.^{75, 212, 215} Of the three studies that measured quality of life, only one study (also included in our review) reported significant improvements post-MBSR intervention⁷⁵ relative to results in control or comparator groups. The other two trials reported no significant improvements in quality of life after MBSR intervention or at follow-up. A second systematic review and meta-analysis was conducted to investigate the effect of MBSR in the cancer care setting on several psychological outcomes, including quality of life.¹⁰⁰ The analyses specific to the quality of life outcome included 248 patients in six studies and reported a small effect size ($P < 0.01$). The authors suggest that although these findings support the use of MBSR for improving quality of life, more well-conducted RCTs are required that implement adequate controls, longer follow-up periods, larger samples sizes, and obtainment of patients' psychological profiles.

Risk/benefit assessment of meditation interventions

There is very little risk to participants who use meditation therapies to improve quality of life and/or physical functioning. Few adverse events have been reported in any trials involving meditation, with participants typically reporting positive feedback about meditation, resulting in low dropout rates from the programs. Group meditation formats and online and home-based adaptations of MBIs¹⁰⁵ are cost-effective and beneficial therapies that can be used as adjuncts to traditional individual counseling or psychotherapy.

Future research in meditation interventions for quality of life

To improve specificity of the effective components of meditation and to compare meditation interventions to other mind-body interventions, research assessing meditation as a treatment for improved quality of life and/or physical functioning should be extended to directly compare meditation with other forms of therapy, including individual counseling, cognitive behavioral therapy, and other mind-body interventions, similar to the therapies compared in the trial by Carlson et al.⁷²

Yoga (B grade)

Overview of yoga interventions for quality of life

Yoga is recommended for improving the quality of life in breast cancer patients (Grade B). This recommendation is based on 12 RCTs, completed between 2006 and 2015, which tested a variety of yoga programs (Supplemental Table 4).^{81, 82, 121, 124, 126, 168-171, 204, 205, 216} Quality of life was the primary outcome for six of the 12 trials.^{81, 168, 169, 204, 205, 216} In 10 trials, a yoga intervention group was compared with a usual care or waitlist control condition group, and two others used brief supportive therapy as a comparison group.^{121, 171} Study participants included women undergoing radiation therapy or chemotherapy for breast cancer, newly diagnosed breast cancer patients, and breast cancer survivors who had completed treatment. The study sample sizes ranged from 15 to 128 participants. Several of these trials overlap with those described under yoga interventions for anxiety and depression outcomes (see above). A total of four different types of yoga interventions were investigated, including: *Iyengar* or Hatha yoga,^{82, 168-170, 204, 205} an integrated yoga program,^{81, 121, 171} *Pranayama* or yoga breathing,^{124, 216} and Patanjali's Yoga Sutras¹²⁶.

A systematic review and meta-analysis assessed the impact of yoga interventions on quality of life and psychological health specifically in breast cancer patients and survivors.¹³² Overall, 12 RCTs were included in the analysis with a total of 742 participants. Analyses revealed short-term effects on improved psychological health, including anxiety ($P < .01$), perceived stress ($P = .03$), and psychological distress ($P < .01$). It is, however, notable that these effects were only applicable to those who engaged in yoga during active cancer treatment and not in the post-treatment period. The authors state that with these positive preliminary results, yoga should be used in this population.

Risk/benefit assessment of yoga interventions

As stated above regarding yoga for anxiety/stress and for depression/mood disturbances, yoga can be adapted and modified for use in this population with low risk to the patient. Overall, yoga has shown preliminary efficacy in improving quality of life, and is recommended for use with breast cancer patients.

Future research in yoga interventions for quality of life

As stated earlier, future trials of yoga to improve quality of life/physical functioning should examine the effects of different types, doses, and durations of yoga on quality of life outcomes. Higher quality trials, including trials with larger and more diverse samples, should be conducted.

C and D graded therapies for quality of life

Trials in acupuncture^{136, 138, 202, 217, 218}, mistletoe²¹⁹⁻²²², Qigong^{223, 224}, reflexology²²⁵⁻²²⁷, and stress management^{61, 114-116, 190, 191, 228} and have assessed the effect of these therapies on quality of life and received a C Grade indicating that they can be considered for use. The five trials evaluating acupuncture had mixed findings and small sample sizes; future studies should replicate the trials of acupuncture that compared real and sham acupuncture, which were the study designs showing no effect.^{136, 202} Two trials^{223, 224} found qigong to have beneficial effects on quality of life, however, the studies were fairly small and should be replicated in larger and more diverse patient populations. Three large high-quality trials²²⁵⁻²²⁷ of reflexology for improving quality of life reported mixed findings. The trials of stress management had conflicting results and employed a broad range of control groups.^{61, 113-115, 189, 190, 227}

There is some evidence that mistletoe may improve the quality of life in breast cancer patients²¹⁹⁻²²². However, while the trials have study quality and sample sizes that could merit a “B” grade, the final decision to assign a “C” grade is due to two areas of uncertainty. First, while several different preparations and formulations have been found to be effective in trials of moderate size, the assessment does not result in a higher grade because of the non-specificity and variability of formulations of agents

tested. Secondly, mistletoe is an injected bioactive compound with a potential for a differential risk benefit ratio due to toxicities and drug interaction with standard cancer therapies that may not be detected in smaller studies; a similar stringency need not to be applied to mind-body interventions as they have lower risk profiles. The Grade C recommendation is based on four RCTs, completed between 2004 and 2014, which tested the use of mistletoe for improving quality of life.²¹⁹⁻²²² Quality of life was the primary outcome in all four trials where a mistletoe product group was compared with placebo^{219, 220} or standard care^{221, 222} control group. The mistletoe products tested, all injected subcutaneously, included PS76A2^{219, 220}, Helixor A²²², and Iscador²²¹. Study participants included women receiving chemotherapy treatment for breast cancer and sample sizes ranged from 61 to 352 participants. To improve specificity of the effectiveness of mistletoe as a treatment for improved quality of life in breast cancer patients, double blind trials need to directly evaluate and compare the different products available and also assess long-term benefit and safety from the use of mistletoe products. Trials of bioactive agents carry the additional requirement of adequate size and statistical power to exclude drug interactions and attenuation of cancer outcome benefits of concurrently administered adjuvant treatments. Two systematic literature reviews^{50, 229} of controlled clinical trials of mistletoe, including a Cochrane database analysis, did find an improvement in survival in the adjuvant setting. Though this outcome is outside the scope of this current review, the two reviews found a suggestion of benefit for quality of life and called for further confirmatory trials.^{50, 229}

Use of Integrative Therapies for Chemotherapy Induced Nausea and Vomiting (CINV)

Description of chemotherapy-induced nausea and vomiting

Chemotherapy-induced nausea and vomiting (CINV) is experienced by some cancer patients following chemotherapy administration.^{71, 230, 231} Acute CINV is typically defined as occurring during the first 24-hour period following chemotherapy administration. Delayed or late CINV is felt to be mediated by different mechanisms compared to acute CINV²³² and occurs more than 24 hours following chemotherapy administration. In a large prospective study of breast cancer patients receiving

chemotherapy treatments, 37% reported any nausea and 13% reported any vomiting at the first 24-hour period.²³³ In the 2-5 days following chemotherapy administration: 70% reported any nausea and 15% reported any vomiting.^{69, 233} Consequences of CINV include dehydration, serious metabolic derangements, nutritional depletion and anorexia, deterioration of physical and mental status, withdrawal from potentially useful and curative antineoplastic treatment, and decrease in self-care and functional ability. CINV is considered to be one of the most severe and feared adverse effects of cancer treatment by patients and can have a significant impact on quality of life.^{230, 234-236} Standard of care anti-emetics for managing CINV have changed considerably in the last 5 years, thus many of the trials evaluating integrative approaches are not tested with the newest and most effective standard treatment regimen.^{71, 231} Most contemporary studies use the endpoint of proportion of patients achieving a complete response defined as no emesis or use of rescue medication. Additionally, anti-emetics themselves have side effects such as headaches, constipation, and neuropsychiatric effects and therefore merit study designs that replace medications with integrative approaches and use equivalence or non-superiority designs for the CINV endpoints and medication side effect endpoints.

Acupressure (B grade)

Overview of acupressure interventions for CINV

For breast cancer patients receiving chemotherapy, acupressure can be considered as an addition to anti-emetics to help control nausea and vomiting (Grade B). This recommendation is based on results from three RCTs, reported between 2000 and 2007, of an acupressure intervention used in conjunction with antiemetics to treat CINV (Supplemental Table 5).²³⁷⁻²³⁹ Acute and delayed nausea and vomiting were the primary outcomes for all three trials. In two trials^{237, 239}, the acupressure plus usual care intervention group was compared with a usual care group. The third trial²³⁸ was a three-arm trial comparing 1) true acupressure at the P6 and SI3 points in addition to usual care; 2) sham acupressure, or placebo acupressure on a different acupressure point, in addition to usual care; and 3) usual care only. (Of note, the use of sham controls in acupressure and acupuncture studies is to attempt to control for the

experience of receiving the treatment and if implemented well, participants will not be able to discern between the true and sham techniques.) Study participants included breast cancer patients undergoing the first, second, or third cycle of chemotherapy. The study sample sizes in the trials ranged from 17 to 160 participants. The acupressure interventions included self-acupressure^{237, 238} using a finger and wearing acupressure wristbands.²³⁹ Across the three trials, acupressure therapy produced significant decreases in nausea, retching, and vomiting ($P < 0.05$ for multiple outcomes assessing CINV; see Supplemental Table 5 for details).

A review assessing acupressure as a nonpharmacologic adjunctive intervention for CINV control across all cancers concluded that acupressure should be strongly recommended as an effective intervention along with standard care for CINV control.²⁴⁰ Other studies of acupressure to reduce nausea and vomiting have shown efficacy in other populations, including pregnant women and postoperative patients, including after thyroidectomy.²⁴¹ All of these studies were conducted with acupressure wristbands placed on both the patient's arms at the PC-6 acupoint.²⁴² A review of acupuncture and acupressure for CINV control among breast cancer patients concludes that the therapies are both safe and effective.²⁴³ A secondary data analysis of the Dibble et al.²³⁸ multi-center study concluded that breast cancer patients whose nausea intensity started higher from the acute phase continued to experience higher symptom intensity during the 11 days after chemotherapy administration and required more frequent acupressure on acupressure point PC6 even after the peak of nausea.²⁴⁴ However, a recent publication by Molassiotis, a lead author of one of the included trials in our review,²⁴⁵ suggests an overall placebo effect in the study of acupressure for control of CINV, although this interpretation included a mix of cancer populations and was not limited to breast cancer patients.

Risk/benefit assessment of acupressure interventions

Self-administered acupressure is easy to perform, safe, cost-effective, noninvasive, does not interfere with a patient's privacy, and has no deleterious effects on patients. Acupressure can be performed anywhere with little or no equipment.

Future research in acupressure interventions for CINV

Future research in this area could assess how to identify the best patients that can be instructed to perform self-administered acupressure, when acupressure can be performed, and whether additional points can be administered along with PC6 to increase the effectiveness of self-administered acupressure to reduce nausea and vomiting.

Electroacupuncture (B grade)*Overview of electroacupuncture interventions for CINV*

Electroacupuncture or acustimulation can be considered as an addition to antiemetics to control CINV in breast cancer patients during chemotherapy (Grade B). This recommendation is based on two RCTs, published in 2000 and 2012 (Supplemental Table 5)^{246, 247}, as well as the 1997 NIH Consensus Conference on acupuncture.²⁴⁸ However, most of these trials predated the use of newer agents including 5-HT₃ or NK1 receptor antagonists, which have become standard anti-emetic therapies for patients receiving highly emetogenic chemotherapy. We do not have a body of evidence to evaluate whether the addition of acupuncture to contemporary anti-emetics yields added benefit. Participants in the trials in this analysis included breast cancer patients who had received moderately high or highly emetogenic chemotherapy without a 5-HT₃ or neurokinin-1 (NK1) receptor antagonist.⁷¹ Both trials used PC-6 and ST-36 acupoints and sham controls, and both trials evaluated the effects of these acupoints on acute and delayed CINV. One trial additionally used acupoint LI-4 and found that electroacupuncture was no better than sham electroacupuncture and that a likely contributor to the lack of effect of electroacupuncture in CINV is the fact that the study evaluated only feasibility with minimal electroacupuncture intervention and without a no-acupuncture arm.²⁴⁶ The second trial found that patients receiving electroacupuncture experienced significantly fewer emesis episodes over the five days of the acupressure intervention than did patients receiving mock therapy or antiemetics alone ($P < 0.001$).²⁴⁷ Between-group differences in the

number of emesis episodes were also significant for electroacupuncture versus minimal needling ($P < .001$) and for minimal needling versus antiemetics alone ($P = 0.01$).

A multicenter study by Yang et al.²⁴⁹ that compared ST-36 electroacupuncture plus anti-emetics with anti-emetics alone in 246 patients with heterogeneous cancers indicated an additive effect with the use of electroacupuncture ($P < 0.01$) with greater decreases in nausea and vomiting scores ($P < 0.001$) compared to use of anti-emetics alone. Two early studies by Dundee et al.^{250, 251} reported significantly less CINV with a PC-6 electroacupuncture intervention, although investigators noted that the brevity of emetic action was a major problem. In a recent review of acupuncture studies including studies that evaluated CINV, investigators concluded that only the electroacupuncture study of Shen et al.²⁴⁷ had a low risk of bias.³⁰ In addition, an earlier meta-analysis by Ezzo et al.²⁵² determined that electroacupuncture, not manual acupuncture, was beneficial for first-day vomiting and that needle insertion as part of manual and electroacupuncture provides greater intensity of stimulation and produces more beneficial effects than surface electrostimulation.

Risk/benefit assessment of electroacupuncture interventions for CINV

With proper administration, electroacupuncture has been shown to be both safe and effective.²⁵³⁻²⁵⁸ In addition to possibly reducing CINV, PC-6 stimulation has been associated with other positive benefits, including analgesic,²⁵³ sedative,²⁵⁹ and anxiolytic effects.²⁶⁰ Further, because CINV is drug-specific rather than disease-specific, these benefits should extend to CINV in other cancer patient populations, as suggested in the study of Yang et al.²⁴⁹ with participants who had a variety of cancers. Practical issues to consider are that electroacupuncture should not be used in patients with a pacemaker or implantable defibrillators and that special attention is required when treating patients who are pregnant, have seizure disorders, or are disoriented.³⁰

Future research in electroacupuncture interventions for CINV

Future trials on the use of electroacupuncture interventions for CINV in breast cancer patients can focus on the testing the use of electroacupuncture with new standard of care treatment regimens, as well as the dissemination and implementation of this technique in the clinical setting. In addition, nausea that is unrelated to chemotherapy is also a common problem in cancer patients and this modality could be tested in comparison to less potent anti-emetics or to best supportive care when other therapies are ineffective.

C and D graded therapies for CINV

Three trials of ginger²⁶¹⁻²⁶³ and two trials of relaxation^{146, 156} have examined their effects on CINV as a primary outcome. Though there were a limited number of trials, results suggested enough of an effect to result in Grade C recommendations stating that ginger and relaxation can be considered as an addition to antiemetics for the control of acute CINV. Future directions in research should focus on replicating trials of these modalities for CINV as the primary outcome. Glutamine is not recommended for improving CINV due to a lack of effect from two trials^{264, 265} in which CINV was assessed as a secondary outcome (Grade D).

Use of Integrative Therapies for Acute Radiation Dermatitis

As radiation is a localized, targeted therapy, side effects are most often locoregional within the targeted area. However, damage to normal local tissues and adjacent organs at risk can result in fatigue as the body expends energy in normal tissue repair. The most common side effect in breast cancer patients is acute skin irritation. Late changes to normal tissue can occur years following therapy and can include soft tissue fibrosis, lymphedema, lung, heart and chest wall damage.²⁶⁶ Not all patients experience one or all of the side effects of radiation, but patients who do experience acute side effects of radiation typically see the effects go away after several weeks following treatment.²⁶⁶ Acute radiation dermatitis can occur with radiation therapy and may result in reactions ranging from faint erythema to dry, itchy, and peeling skin and ultimately to moist desquamation and ulceration.²⁶⁷ The CGCTC is the most common scale to

measure acute radiation skin toxicity that is used by cooperative groups during cancer clinical trials and grades skin reactions from 0 to 4 with grade 1 being erythema and grade 4 being ulceration. Most breast cancer patients develop mild to moderate acute skin reactions of grade 1-3 during and shortly after a course of radiation therapy. These reactions usually resolve quickly but can cause significant symptoms, especially with higher grade toxicity.²⁶⁷

D graded therapies for acute radiation dermatitis

There are no A, B or C graded therapies to report for acute radiation dermatitis post treatment. Aloe vera gel and hyaluronic cream are both not recommended as a standard therapy to prevent or treat acute radiation dermatitis simply due to lack of effect (Grade D). Our review consisted of two quality studies for each product with large sample sizes for both the aloe vera trials^{268, 269} and hyaluronic cream trials^{270, 271}. Each trial assessed the acute skin reaction from radiation therapy as its primary outcome.

Use of Integrative Therapies for Vasomotor Outcomes

Vasomotor symptoms are common in breast cancer patients and include hot flashes, intense sweating, flushing on the face and chest, and may come with heart palpitations and anxiety.²⁷² These symptoms occur episodically, including nocturnally when night sweats can significantly disrupt women's sleep. According to NCI, about two thirds of postmenopausal women with a history of breast cancer experience hot flashes. These symptoms may occur naturally or as a consequence of surgery, chemotherapy or endocrine therapy.²⁷³ While vasomotor symptoms may resolve on their own, 20% of affected women suffer from persistent hot flashes four years later after last menses.²⁷² Together, vasomotor symptoms can significantly impact women's quality of life.²⁷³

C and D graded therapies for vasomotor outcomes

There are no A or B graded therapies to recommend for vasomotor outcomes. Acupuncture can be considered as a therapy for hot flashes based upon nine trials assessing acupuncture for hot flashes

(Grade C).^{136, 188, 189, 274-279} Seven trials assessed hot flashes as the primary outcome and only one trial²⁷⁹ had more than 100 participants. Overall, the literature showed mixed findings; however, the single large trial showed significant reductions in hot flashes in their electroacupuncture group compared to sham and control groups. The use of soy as a therapy for hot flashes is not recommended due to lack of effect (Grade D). Three large trials²⁸⁰⁻²⁸² assessed soy for the treatment of hot flashes as the primary outcome and showed a lack of effect.

Use of Integrative Therapies for Lymphedema

Lymphedema

Lymphedema is a condition following treatment such as surgery or radiation therapy where parts of the lymph system become damaged or blocked, leading to an accumulation of lymph fluid that does not drain properly, builds up in tissues, and causes swelling.³⁷ The CTCAE grades edema of the limbs from grade 1 (5-10% interlimb discrepancy) up to grade 3 (>30% interlimb discrepancy). Lymphedema commonly affects the arm or leg but can also impact other parts of the body. For breast cancer patients and survivors, lymphedema is most common in the upper extremities and sometimes in the breast and/or chest wall, and can occur up to 30 years following treatment. Owing to difference in diagnosis and characteristics of patients studied, and inadequate follow-up the overall incidence of arm lymphedema after breast cancer has been found to range from 8% to 56%.²⁸³ Breast cancer survivors with arm lymphedema in particular have been found to have decreased quality of life and increased psychological distress and disability compared to survivors without lymphedema.²⁸³⁻²⁸⁵

C graded therapies for lymphedema

There are no A or B graded therapies to report for lymphedema. Two trials assessed laser therapy^{286, 287} and seven trials assessed manual lymphatic drainage²⁸⁸⁻²⁹⁴ for the treatment of lymphedema as a primary outcome. The two trials evaluating laser therapy were small in sample size and showed mixed findings. Only two of the seven trials assessing manual lymphatic drainage had a sample size

greater than 100 participants^{289, 294}. Overall, the literature suggests that manual lymphatic drainage and compression bandaging are equivalent. Thus, either therapy can be considered as treatment options for lymphedema, with manual lymphatic drainage being considered for those who have sensitivity to bandaging (Grade C).

Use of Integrative Therapies for Chemotherapy Induced Peripheral Neuropathy (CIPN)

Chemotherapy induced peripheral neuropathy (CIPN)

Cancer treatments, including chemotherapy, may cause damage to the peripheral nerves resulting in neuropathy.²⁹⁵ The CGCTC categorizes neuropathy under neurologic sensory and grades from 0-3 with 3 being severe objective sensory loss or paresthesias that interfere with function. Sensory neuropathy can include symptoms of pain, tingling, numbness, or a pins-and-needles feeling, the inability to feel a hot or cold sensation, or the inability to feel pain. Motor neuropathy can include problems with balance, weak or achy muscles, twitching, cramping or wasting muscles, or swallowing or breathing difficulties. Autonomic nerve damage can cause dizziness or faintness, and digestive, sexual, sweating and urination problems.²⁹⁵

H graded therapies for CIPN

There are no A or B graded therapies to report for the prevention or treatment of CIPN. Acetyl-L-carnitine is not recommended as a standard therapy to prevent or treat CIPN due to harm (Grade H). A single large and high-quality study¹⁹⁸ assessing the use of acetyl-L-carnitine capsules to prevent CIPN following taxane therapy as a primary outcome found that acetyl-L-carnitine administered during taxane chemotherapy was associated with worse CIPN symptoms.

Use of Integrative Therapies for Pain

Pain

According to the International Association for the Study of Pain, pain can be defined as “an unpleasant sensory and emotional experience associated with actual or potential tissue damage.”²⁹⁶ The CTCAE grades pain from 1 (mild) to 3 (severe, limiting self care). Pain can be caused by cancer therapies, including surgery, radiation therapy, chemotherapy, targeted therapy, supportive care therapies and/or diagnostic procedures.²⁹⁷ Pain is commonly experienced by breast cancer patients with a prevalence ranging from 40-89%.²⁹⁶ Pain management requires proper assessment including measurement of intensity.²⁹⁷ It is also important to evaluate the impact of pain on the patients physical, mental, and social health as pain can negatively impact a patient’s functional status and quality of life. Pain management can include both pharmacologic and non-pharmacologic modalities. Proper education about treatment and longitudinal follow-up are essential.

C graded therapies for pain

There are no A or B graded therapies to report for pain. Healing touch³⁴ for pain following chemotherapy; music therapy^{106, 109}, hypnosis^{298, 299}, and acupuncture²⁵³ for pain following surgery, and acupuncture³⁰⁰⁻³⁰⁴ for pain associated with aromatase inhibitor associated musculoskeletal symptoms were examined and each received a C grade indicating that they can be recommended as a therapy for pain. A single large trial assessed healing touch³⁴ for pain following chemotherapy as a secondary outcome showing small positive effects favoring the therapy. Similarly, trials assessed music therapy^{106, 109}, hypnosis^{298, 299} and acupuncture²⁵³ for pain following surgery as a primary outcome showing small positive effects favoring the therapy. However, there is a lack of multiple large trials to support each therapy. Finally, five trials evaluated acupuncture for pain associated with aromatase inhibitor-associated musculoskeletal symptoms as the primary outcome.³⁰⁰⁻³⁰⁴ All trials had small sample sizes and reported mixed findings.

Use of Integrative Therapies for Sleep Disturbance

Sleep disturbances

Studies have shown that nearly half of all breast cancer patients have sleep-related problems due to a range of causes including side effects of antineoplastic medications, long hospital stays, or stress.³⁰⁵ In addition, a growing number of breast cancer patients are obese,³⁰⁷ which increases the incidence of sleep apnea, a major cause for insomnia. Insomnia, a specific sleep disorder of initiating and maintaining sleep, is most common in cancer patients and often occurs along with anxiety and depression.³⁰⁸ The CTCAE measures insomnia under psychiatric disorders from grade 1-3, with grade 3 being severe difficulty falling asleep, staying asleep or waking up early.

C graded therapies for sleep disturbance

There are no A or B graded therapies to report for sleep disturbance. Yoga can be considered for sleep disturbance (Grade C). Five trials assessed yoga for sleep disturbance^{77, 123, 126, 170, 309} and in four of these trials sleep was a secondary outcome. Two trials were of high quality with more than 100 participants^{77, 309}. Overall, the body of literature showed no greater effect in sleep quality than health education classes, stretching groups, and wait list control.

CONCLUSION

In this review, we closely examined and described the RCTs that provide support for the highest graded therapy recommendations for the use of integrative therapies during the patient experience of breast cancer and for side effects related to breast cancer treatment. High levels of evidence support the routine use of mind-body practices, such as yoga, meditation, relaxation techniques and passive music therapy to address common mental health concerns among breast cancer patients, including anxiety, stress, depression, and mood disturbances. Additionally, meditation has been shown to improve quality of life and physical functioning, and yoga has been found to improve quality of life and fatigue. Massage has been shown to improve mood and acupressure and electroacupuncture decrease CINV. Given the high level of evidence of benefit coupled with the and the relative low level of risk, these therapies can be incorporated as an option care for patients, especially when there is poor symptom control. As is the case

with many standard therapies, the impact of integrative approaches on symptom management is highly individualized. Therefore, a patient-centered trial and evaluation approach may be needed and can be guided by the grade of recommendations and altered as needed along with the incorporation of patient preferences. In addition to the modalities discussed in this review that were given lower grade (C or D), patients are using many other forms of integrative therapies with little or no supporting evidence; this serves as a compelling call for further research to support patients and health care providers in making more informed decisions that avoid harm. In the meantime, while further clinical evaluation is underway, clinicians and patients need to be cautious about the use of therapies that received a Grade of C or D and need to fully understand the potential risks and benefits of use, including the risk associated with not using a conventional therapy that may effectively prevent or treat the condition. For example, in a patient with incurable disease who has marked symptoms not adequately managed with conventional therapies, carefully monitored use of a Grade C therapy could be medically reasonable even though more research clearly would be needed to apply this broadly across a patient population. This review and others support referral or provision of clinical services to include both evidence-based conventional and integrative therapy options.

The limited number of integrative modalities with A or B grades emphasizes the need for all cancer care providers to initiate a dialogue early in their relationship with cancer patients to develop a framework for how evidence forms the basis for all clinical decisions. Patients and clinicians should engage in shared decision making based on the best available evidence on the benefits and harms, while reflecting patient values and preferences. A careful appraisal of the evidence base for integrative therapies can help allay a patient's concern that their care team is informed and not overlooking options that may be of interest to them. In addition, such an appraisal of the evidence will offer those modalities that do merit consideration and allow for better personalization of care and shared decision making.

This systematic review with grades of evidence adds to a growing literature base including reviews of integrative therapy for breast cancer and other cancer populations. For example, there are numerous reviews that support the use of integrative therapies including passive music therapy¹¹¹, stress

management programs¹¹⁸, various yoga practices¹³¹, meditation and MBSR⁹⁸, massage¹⁷⁸, and relaxation techniques¹⁵⁹ as adjunctive therapies for psychological outcomes, specifically the anxiety/stress and depression/mood outcomes assessed in this review. Acupressure for CINV is also well supported in the review literature across all cancer patients.²⁴⁰ There are mixed findings in the areas of meditation and MBSR for quality of life^{47, 100}, and electroacupuncture for CINV,³⁰ which suggests overall low quality or too few studies. Thus, future research on the impact of these integrative therapies on the relevant clinical outcomes is warranted. A limitation to the generalizability of our findings is that the majority of participating subjects in the clinical trials evaluated were non-Hispanic white women with high socioeconomic status relative to the general population. In addition, none of the trials examined age-related responses, and or differential responses in pre- vs. post-menopausal women. There is a clear need to design well-powered controlled trials using the best standard treatment control or an appropriate placebo.

Challenges of implementing integrative therapies in breast oncology

Cancer patients face a number of psychological and physical challenges as they move through cancer diagnosis, treatment and survivorship. According to NCCN guidelines, comprehensive clinical programs should systematically screen for cancer-related symptoms and side effects in the process of mandated screening for distress. This review and others support subsequent referral or provision of clinical services to include both evidence-based conventional and integrative therapy options.^{68, 310}

Based on recent estimates from the U.S. National Health Interview Survey, 75% of individuals with a history of cancer use one or more complementary and integrative therapies.³¹¹ Many North American cancer centers now operate formal integrative oncology programs. As most of these services are not reimbursable by insurance, the methods and models of funding and implementing these programs vary; some programs and services are fully funded and provided free of charge to patients, some are entirely paid for out-of-pocket by patients, and some are a combination of the two. Often mind-body therapies are available from trained clinical staff already employed at cancer centers, such as oncology

nurses or social workers, and hence, are more readily accessible at low or no cost. Others such as massage therapy and acupuncture may be covered by some forms of insurance, varying by country, province/territory and state.

Implementing complementary and integrative therapies in a clinical setting not only requires funding and infrastructure, but also requires well-trained, knowledgeable providers. Many of the integrative therapies do not have a one-size-fits-all approach and need to be provided and administered by appropriately trained practitioners who can evaluate which are the best forms and techniques to use with a specific patient. Though training and credentialing for many integrative providers varies by jurisdictions, best practices suggest that providers be trained to the highest standard of their profession, even if that exceeds the state or province-based standards, such as a requirement for institutional credentialing that may include proctoring. As the fields of integrative therapies are expanding, there are now new professional associations that specialize in oncology, for example the Society for Oncology Massage and the Oncology Association of Naturopathic Physicians.^{312, 313}

Future research

Rigorous clinical research that appropriately reflects integrative care as it is used in the community and at integrative cancer centers is needed to responsibly move this field forward. Integrative modalities can be tested in addition to standard supportive treatments, or even in place of them, if the standard therapy is associated with side effects or significant costs and the trial design allows for early discontinuation in the event of futility. Clinical trials designed to test efficacy in tightly controlled academic research settings are often testing protocols that are not realistically implementable in the community setting. Implementation and dissemination research designs to consider include pragmatic trials that involve multi-modal therapies that are applied in the manner in which they are typically offered in clinical settings. This approach, while unable to pinpoint clear causal relationships between specific interventions and outcomes, allows an exploration and evaluation of clinical impact that is more truly generalizable. Head-to-head comparisons of different integrative therapies and conventional symptom

management therapies would help provide some specificity and direction for health care providers making recommendations to patients. Comparative effectiveness research testing integrative modalities in relationship to pharmacological and other approaches would also be informative in providing options as well as comparisons of toxicities and cost effectiveness. Studies that examine mechanism of action are also needed, however, the emphasis here is on trials of agents that are actively in use, unlike novel pharmacological therapies. Importantly, interventions need to be tested in economically and culturally diverse patient populations in order to understand the applicability of an intervention to the growing population of cancer survivors.

Future studies need to include systematic assessments of treatment toxicities, including toxicities from both the integrative and the conventional therapies. In this review, when possible, the NCI Common Terminology Criteria for Adverse Events (CTCAE) are used to describe cancer treatment side effects. However, many of the trials did not report toxicities or adverse events, and among the trials that did, the majority of the trials did not assess toxicities and adverse events systematically. The CTCAE are a set of criteria for the standard grading and classifications of adverse effects of drugs used in cancer therapy and the FDA is increasingly using CTCAE patient reported outcomes (CTCAE-PROs) to monitor treatment side effects.^{314, 315} Ideally, trials will include systematic evaluation of both provider (i.e., CTCAE) and patient (i.e., CTCAE-PROs) assessments of adverse events. If future trials do not use these methods, at minimum, validated measurement tools need to be used to allow for ongoing quantitative assessments of adverse events using robust statistical analyses.

Ongoing challenges include the inability to blind subjects to most of the integrative modalities studied, as most measures are subjective and thus susceptible to suggestive biases in which patients perceive benefit to an intervention simply because they are receiving it. Using a mixed methods model of research including both qualitative inquiry that explores the patient's experience of their treatments and quantitative data will be helpful to validate and better justify the use of integrative therapies. Additionally, the use of both subjective and objective patient-reported outcomes should be employed within a mixed methods model. This approach can be used in both clinical trials or in prospective observational studies.

To better enable real clinical uptake and change, knowledge translation experts, cancer patients, policy makers and decision makers should be involved in both study design and interpretation to better enable integration of these therapies into clinical practice.

In conclusion, awareness of the base of evidence for complementary and integrative therapies based on the recently published SIO guidelines and the emerging literature should be a core competence for the cancer care provider and should be applied in decision making with breast cancer patients requiring supportive care. Billions of dollars are spent each year on complementary and integrative health therapies, with unknown benefits and for those that have thus far been shown to be ineffective.³¹⁶ Research in this area could save large amounts of healthcare dollars and resources and more importantly can redirect patients to treatments with known benefits and better safety profiles. This article provides greater depth of discussion of these interventions, such that clinicians and patients can begin the process of integration, based upon patient needs in their specific setting and context.

ACKNOWLEDGEMENTS

We thank the following internal and external reviewers for their insightful comments and critiques:

Internal reviewers: Executive Committee of the Society for Integrative Oncology; Board of Trustees of the Society for Integrative Oncology; Donald Abrams, MD (University of California, San Francisco), Ting Bao, MD (Memorial Sloan Kettering Cancer Center), Gustav Dobos, MD (Duisberg-Essen University), Petra Klose, MD (Duisberg-Essen University), Omer Kucuk, MD (Emory University), Jodi MacLeod (University of Pennsylvania), Gregory Plotnikoff, MD (Minnesota Personalized Medicine) and Santosh Rao, MD (Banner MD Anderson). External SIO reviewers included Gabriel Hortobagyi, MD (MD Anderson Cancer Center), Shelley Hwang, MD (Duke University), and Anna Wu, PhD (University of Southern California).

TABLES

Accepted Article

Table 1: Graded integrative therapies for use in breast cancer patients according to clinical outcomes*

Clinical Outcomes	Recommended therapy	Strength of Evidence Grade
Acute Radiation Skin Reaction	Aloe vera ^{268, 269} and Hyaluronic acid cream ^{270, 271} should not be recommended for improving acute radiation skin reaction.	D
Anxiety/ Stress Reduction	Meditation is recommended for reducing anxiety. ⁷²⁻⁷⁶	A
	Music therapy is recommended for reducing anxiety. ¹⁰⁶⁻¹¹⁰	B
	Stress management is recommended for reducing anxiety during treatment, but longer group programs are likely better than self-administered home programs or shorter programs. ¹¹⁴⁻¹¹⁷	B
	Yoga is recommended for reducing anxiety. ^{79, 119-126}	B
	Acupuncture ¹³⁶⁻¹³⁸ , Massage ¹³⁹⁻¹⁴² , Relaxation ¹⁴³⁻¹⁴⁷ can be considered for reducing anxiety.	C
Chemotherapy Induced Nausea and Vomiting	Acupressure can be considered as an addition to anti-emetics drugs to control nausea and vomiting during CT. ²³⁷⁻²³⁹	B
	Electroacupuncture can be considered as an addition to anti-emetics drugs to control vomiting during CT. ^{246, 247}	B
	Ginger ²⁶¹⁻²⁶³ , Relaxation ^{146, 156} can be considered as an addition to anti-emetics drugs to control nausea and vomiting during CT.	C
	Glutamine ^{264, 265} should not be recommended for improving nausea and vomiting during CT.	D
Depression/ Mood Disturbance	Meditation, particularly MBSR, is recommended for treating mood disturbance and depressive symptoms. ^{72-76, 148-152}	A
	Relaxation is recommended for improving mood disturbance and depressive symptoms. ^{143, 146, 147, 156-158}	A
	Yoga is recommended for improving mood and depressive symptoms. ^{77-80, 119-121, 123-126, 168-171}	B
	Massage is recommended for improving mood disturbance. ^{140-142, 175-177}	B
	Music therapy is recommended for improving mood. ^{108, 110, 185, 186}	B
	Acupuncture ^{136-138, 188, 189} , Healing Touch ^{34, 35} , Stress Management ^{114-116, 190, 191} can be considered for improving mood disturbance and depressive symptoms.	C
Fatigue	Hypnosis ^{83, 84} and ginseng ^{196, 197} can be considered for improving fatigue during treatment.	C
	Acupuncture ^{138, 201-203} and Yoga ^{78, 82, 123, 170, 204, 205} can be considered for improving fatigue post treatment.	C

	Acetyl-L-carnitine ¹⁹⁸ and Guarana ^{199, 200} should not be recommended for improving fatigue during treatment.	D
Lymphedema	Laser therapy ^{286, 287} , Manual lymphatic drainage and compression bandaging ²⁸⁸⁻²⁹⁴ can be considered for improving lymphedema.	C
Neuropathy	Acetyl-L-carnitine is not recommended for prevention of chemotherapy-induced peripheral neuropathy in breast cancer patients due to harm. ¹⁹⁸	H
Pain	Acupuncture ^{253, 300-304} , Healing Touch ³⁴ , Hypnosis ^{298, 299} , Music therapy ^{106, 109} can be considered for the management of pain.	C
Quality of life	Meditation is recommended for improving quality of life. ^{73-75, 149-151, 212}	A
	Yoga is recommended for improving quality of life. ^{81, 82, 121, 124, 126, 168-171, 204, 205, 216}	B
	Acupuncture ^{136, 138, 202, 217, 218} , mistletoe ²¹⁹⁻²²² , Qigong ^{223, 224} , Reflexology ²²⁵⁻²²⁷ , Stress management ^{61, 114-} ^{116, 190, 191, 228} can be considered for improving quality of life.	C
Sleep disturbance	Gentle yoga ^{77, 123, 126, 170, 309} can be considered for improving sleep.	C
Vasomotor / Hot flashes	Acupuncture ^{136, 188, 189, 274-279} can be considered for improving hot flashes.	C
	Soy ²⁸⁰⁻²⁸² is not recommended for hot flashes in BC patients due to lack of effect.	D
<p>Abbreviations: BC, Breast Cancer; CT, Chemotherapy; MBSR, Mindfulness-Based Stress Reduction Notes: The clinical population is breast cancer (BC) patients during treatment, including surgery, chemotherapy, hormonal/biological therapy, and radiation therapy. The clinical question is "What integrative therapies can be used to prevent, treat and manage symptoms and side effects encountered during breast cancer treatment?" Definitions of the Grade of Recommendations are listed below.¹⁰ Grade A: Recommends the modality. There is high certainty that the net benefit is substantial. Offer/provide this modality. Grade B: Recommends the modality. There is high certainty that the net benefit is moderate or there is moderate certainty that the net benefit is moderate to substantial. Offer/provide this modality. Grade C: Recommends selectively offering or providing this service to individual patients based on professional judgment and patient preferences. There is at least moderate certainty that the net benefit is small. Offer/provide this modality for selected patients depending on individual circumstances. Grade D: Recommends against the service. There is moderate or high certainty that the modality has no net benefit. Discourage the use of this modality. Grade H: Recommends against the service. There is moderate or high certainty that the harms outweigh the benefits. Discourage the use of this modality.</p>		

Table 2: Training and licensure information for A and B grade integrative therapies

Modality or Therapy	Training	Licensure and Regulation	Professional Societies and Organizations
Acupuncture, electro-acupuncture, and acupressure	<p>Licensed acupuncturists generally have attended formal schools of Asian medicine and have passed national certification examinations in order to sit for state or provincial licensing exams. The Accreditation Commission for Acupuncture and Oriental Medicine (ACAOM) accredits schools of Asian medicine in the U.S. Degrees include the Masters of Acupuncture requiring a minimum of 3 years and 1900 hours of training and the Masters of Acupuncture and Oriental Medicine requiring a 3-4 year program, averaging 2,700-3,465 hours of training.</p> <p>Advanced degrees include Doctor of Acupuncture and Oriental Medicine (DAOM), Oriental Medicine Doctor (OMD), Doctor of Acupuncture and Chinese Medicine (DACM) and Doctor of Oriental Medicine (DOM) who provide advanced evaluations, integrative medicine as well as acupuncture and herbal therapy and can provide access to the whole breadth of Chinese medicine. Advanced degrees require a Masters level degree with additional 1000+ hours of training. The National Certification Commission for Acupuncture and Oriental Medicine (NCCAOM) offers acupuncture, herbal and Oriental medicine certifications on a national level.²⁴</p>	<p>US: Acupuncture and Chinese herbology and Oriental medicine are regulated via state licensing bodies. NCCAOM reports 44 states plus the District of Columbia have acupuncture practice acts. 43 states currently require passage of NCCAOM national examinations for initial licensure. The NCCAOM certifies acupuncture, Chinese herbology and Oriental medicine. Currently, California requires passage of its own state examinations for licensure and does not accept the NCCAOM. A subset of states regulate acupressure under acupuncture or massage regulations.</p> <p>CANADA: Acupuncture is regulated via provincial licensing bodies in the provinces of British Columbia, Alberta Ontario, Quebec and Newfoundland. Chinese medicine is currently only regulated in British Columbia and Ontario.</p>	<p>The Accreditation Commission for Acupuncture and Oriental Medicine (ACAOM): http://www.acaom.org</p> <p>National Certification Commission for Acupuncture and Oriental Medicine (NCCAOM): http://mx.nccaom.org</p> <p>College of Traditional Chinese Medicine Practitioners and Acupuncturists of British Columbia (CTCMA): http://www.ctcma.bc.ca</p> <p>College of Traditional Chinese Medicine Practitioners and Acupuncturists of Ontario (CTCMPAO): http://www.ctcmpao.on.ca/</p> <p>College and Association of Acupuncturists of Alberta (CAA): http://acupuncturealberta.ca/</p> <p>Association of Acupuncturists of Quebec (AAQ): http://www.acupuncture-quebec.com/en/home.html</p> <p>Newfoundland and Labrador Council of Health Professionals (nlchp): http://www.nlchp.ca/</p>

<p>Hypnosis</p>	<p>Mental health and medical professionals typically practice hypnosis as a specialty or sub-specialty. Certified hypnotherapists, in general, hold a graduate-level or bachelor's-level degree in a broad range of specialties including: MD, registered nurse, dentist, social worker, licensed counselor or psychologist, pastoral counselor, ordained minister, chiropractor, among many others, prior to obtaining training in hypnosis.</p> <p>At present, there are no accredited schools offering standard college or university degrees in hypnosis; therefore, training in one of the above professions is typically required prior to acceptance into one of many training or certification programs. These programs have a wide range of training requirements, but in general require anywhere from 50 to 200 hours of classroom and clinical training prior to certification.</p>	<p>US: To be certified as a clinical hypnotherapist, applicants typically require anywhere from 50 to 200 hours or training, and often years of experience. There are a number of certification programs with a range of requirements (see links). Regulations for the practice of hypnosis vary on a state-by-state basis (http://www.aihcp.org/hypnosis-regulation.htm).</p> <p>Typically clinical hypnotherapists must renew their certification every 2 to 4 years and have completed 20+ hours of approved training during that time.</p> <p>CANADA: Certification is similar to the US however requirements are much higher, between 225 and 1100 hours: https://www.archcanada.ca/</p> <p>The Council of Professional Hypnosis encompasses many professional societies and organizations listed: http://www.copho.com/org.htm.</p>	<p>Association of Registered Clinical Hypnotherapists (ARCH): https://www.archcanada.ca/</p> <p>American College of Hypnotherapy at the American Institute of Health Care Professionals: http://aihcp.net/american-college-of-hypnotherapy/</p> <p>American Society of Clinical Hypnosis (ASCH): http://www.asch.net/</p> <p>Hypnosis Motivation Institute (HMI): https://hypnosis.edu/</p> <p>National Board for Certified Clinical Hypnotherapists (NBCCH): http://www.natboard.com</p> <p>The International Society of Hypnosis (ISH): http://www.ishhypnosis.org/</p> <p>Society for Clinical and Experimental Hypnosis (SCEH): http://www.sceh.us/</p> <p>The Milton H. Erickson Foundation: http://www.erickson-foundation.org/</p>
-----------------	---	--	--

Accepted Article

Massage	<p>Associated Bodywork and Massage Professional (ABMP) members at the certified or professional levels must possess a valid massage license from a regulated state/province/territory, have completed 500 approved educational hours, or be certified through the National Certification Board for Therapeutic Massage and Bodywork (NCTMB). Licensed nurse and physical therapists may qualify for membership at either the certified or professional level with a minimum of 50 hours of additional massage therapy training. Board Certification is the highest voluntary credential attainable to massage therapists and bodyworkers in the profession today. For the requirements of Board Certification, see: http://www.ncbtmb.org/board-certification.</p>	<p>US: Massage therapy is regulated by some U.S. states. See link for state-by-state massage licensure information. http://www.massagetherapy.com/careers/stateboards.php</p> <p>CANADA: Massage therapists can only be registered not licensed in Canada. Currently only four provinces regulate massage therapists: Ontario, British Columbia, New Brunswick and Newfoundland. The website for the Ontario Massage Therapists has links to the four provinces registration procedures and links to other provincial massage organizations.</p>	<p>American Massage Therapy Association: http://www.amtamassage.org</p> <p>Associated Bodywork and Massage Professionals (ABMP): www.abmp.com</p> <p>The National Certification Board for Therapeutic Massage and Bodywork (NCTMB): http://www.ncbtmb.org/</p> <p>Society for Oncology Massage: http://www.s4om.org/</p> <p>Massage Therapists' Association of Ontario: https://secure.rmtao.com/massage_therapy/regulation_of_mt/massage_therapy_in_canada.htm</p>
Meditation	<p>Mindfulness-Based Stress Reduction: The Centre for Mindfulness in Massachusetts provides meditation specialist training –: http://www.umassmed.edu/cfm/training/detailed-training-information/teacher-certification-review/</p> <p>University of California San Diego (UCSD) Center for Mindfulness provides extensive training: http://mbpti.org/</p> <p>University of Bangor in Wales awards a Masters in Mindfulness and also has a range of teacher training programs: http://www.bangor.ac.uk/mindfulness/courses.php.en</p> <p>Centre for Mindfulness Studies in Toronto, Canada offer one-day workshops as well as intensive professional training in Mindfulness-Based Cognitive Therapy (MBCT). http://www.mindfulnessstudies.com/</p>	<p>In both the US and Canada board certification is the highest credential attainable for meditation teachers in the healthcare profession today. There is no registration or licensure currently available.</p> <p>Completing mindfulness teacher training courses at one of the recognized training programs, in addition to professional certification, would be the optimal level of training for providing these interventions.</p>	<p>American Mindfulness Research Association: https://goamra.org/</p>

Music therapy	A professional music therapist holds a bachelor's degree or higher in music therapy from one of over 70 American Music Therapy Association (AMTA) approved college and university programs. Music therapists who currently hold professional designation are listed on the National Music Therapy Registry (NMTR) and are qualified to practice music therapy	Music therapist is regulated by some U.S. states (e.g., North Dakota, Nevada). See link for state-by-state massage licensure information: http://www.cbmt.org/examination/state-licensure/ Currently no Canadian provinces/territories license or regulate music therapists.	The Certification Board for Music Therapists (CBMT): http://www.cbmt.org/ Canadian Association for Music Therapy (CAMT): http://www.musictherapy.ca/en/
Relaxation and Stress Management	Relaxation and stress management are usually provided by trained mental health professionals such as Registered/Chartered/Counseling Psychologists, Psychiatrists, Marital and Family Counselors or, Clinical Social Workers and Nurses. Any of these professions could have specific training in these two modalities.	Typically the aforementioned professionals would take courses in relaxation training or stress management as part of their training.	Every jurisdiction has a college of psychologists and social workers. American Psychological Association (APA): http://www.apa.org/ Canadian Psychological Association (CPA): http://www.cpa.ca
Yoga	There are very stringent criteria to be a yoga instructor and most jurisdictions have their own requirements. Most jurisdictions have criteria in place in order to become a "certified yoga instructor". There are also requirements for specific genres of yoga such as Bikram, Iyengar, Ashtanga, restorative yoga, etc. Most states and provinces/territories have a designation as a certified yoga instructor. In the U.S. and Canada, the Yoga Alliance is the biggest body that registers teachers: https://www.yogaalliance.org/Credentialing/Credentials_for_Teachers ; and https://www.yogaalliance.org/Credentialing/CredentialsforSchools ; http://www.canadianyogicalliance.com	There are currently no states or Canadian provinces/territories that regulate or license yoga.	Most states and provinces have a professional association (or more than one) for certified yoga instructors.

Table 3: Clinical outcomes and integrative therapies with insufficient evidence to form a clinical recommendation (I-statement)	
Outcome	Integrative Therapies
Adherence	Acupressure ³¹⁷ , Multi-Modal ³¹⁸
Anemia	LCS101 Combination botanical ³¹⁹ , RG-CMH Combination botanical ³²⁰ , Shenqi Fuzheng Injection ³²¹
Anxiety/Stress Reduction	Art Therapy ^{112, 322} , Comprehensive Coping Strategy ³²³ , Electrical Nerve Stimulation ³²⁴ , Healing Touch ^{35, 325} , Hypnosis ³²⁶ , Myofascial Release ³²⁷ , Multi-Modal ^{328, 329} , Reflexology ^{226, 227, 330} , Reiki ³³¹ , Tai-Chi ³³²
Cardiomyopathy	N-Acetylcysteine ³³³
Chemotherapy Induced Nausea and Vomiting	Acupressure ³³⁴ , Aromatherapy ³³⁵ , <i>Agaricus sylvaticus</i> ³³⁶ , Cocculine (complex homeopathic Rx) ³³⁷ , Comprehensive Coping Strategy ³²³ , Massage ¹³⁹ , Nevasic Audio Program ³³⁸ , Yoga ¹²¹
Cognition	Natural Environment ³³⁹ , <i>Ginkgo biloba</i> ¹⁹⁶ , Meditation ¹⁴⁸ , Yoga ³⁴⁰
Constipation	Self-Management Program ³⁴¹
Depression/ Mood	Art Therapy ¹¹² , Biofield Healing ³⁴² , Comprehensive Coping Strategy ³²³ , CoQ10 ³⁴³ , Electrical Nerve Stimulation ³²⁴ , <i>Gandoderma lucidium</i> ³⁴⁴ , Guarana ¹⁹⁹ , Hypnosis ³⁴⁵ , Multi-Modal ^{318, 328, 346, 347} , Myofascial Release ³²⁷ , Qi-gong ²²³ , Reflexology ^{226, 227} , Tai-Chi ³³²
Fatigue	Acupressure ³¹⁷ , Acupuncture ^{137, 348, 349} , Biofield Healing ³⁴² , Comprehensive Coping Strategy ³²³ , CoQ10 ³⁴³ , <i>Gandoderma lucidium</i> ³⁴⁴ , Light treatment ³⁵⁰ , Massage ¹⁷⁶ , Meditation ^{148, 152} , Mind-Body Cognitive Therapy ³⁵¹ , Movement ³⁵² , Multi-Modal ^{347, 353, 354} , Multivitamin ³⁵⁵ , Polarity Therapy ^{356, 357} , Stress Management ³⁵⁸ , Qigong ²²³ , Reflexology ²²⁷ , Relaxation ^{359, 360} , Stress Management ¹¹⁴ , Yoga ^{77, 79, 124-126, 169}
Lymphedema	CYCLO 3 FORT ³⁶¹ , Electro-therapy ³⁶² , Ginkgo forte ³⁶³ , Pentoxifylline & Vitamin E ^{364, 365} , Yoga ³⁶⁶
Neuropathy	Omega 3 Fatty Acids ³⁶⁷ , Vitamin E ^{368, 369} , Acupuncture ²¹⁸
Neutropenia/ Leukopenia	Cat's Claw ³⁷⁰ , LCS101 Combination botanical ³¹⁹ , RG-CMH Combination botanical ³²⁰ , Mistletoe ^{221, 222} , Shenqi Fuzheng Injection ³²¹
Pain	Comprehensive Coping Strategy ³²³ , Stress Management ¹¹⁴ , Vitamin D2 ³⁷¹ , Electrical Nerve Stimulation ³²⁴ , Cognitive and Behavioral Therapy ³⁷² , Hypnosis ³⁷³ , Massage ¹⁷⁷ , Myofascial Release ³²⁷ , Reflexology ^{227, 330}

Quality of life	Acupressure ³³⁴ , Biofield Healing ³⁴² , Calendula cream ³⁷⁴ , Cannabis ³⁷⁵ , Chlorella extract ³⁷⁶ , CoQ10 ³⁴³ , Curcuminoids ³⁷⁷ , Electrical Nerve Stimulation ³²⁴ , Electro-therapy ³⁶² , Flaxseed ³⁷⁸ , <i>Ganoderma lucidum</i> ³⁴⁴ , Ginkgo forte ³⁶³ , Guided Imagery ³⁷⁹ , Healing Touch ³⁵ , Homeopathy ³⁸⁰⁻³⁸² , Hypnosis ⁸³ , Laser therapy ²⁸⁷ , Manual Lymphatic Draining ²⁸⁹ , Massage ¹⁷⁶ , Meditation ^{72, 148} , Movement ^{352, 383} , Music Therapy ¹⁸⁵ , Multi-modal ^{318, 328, 347, 353, 354, 384} , Multivitamin ³⁵⁵ , Polarity therapy ^{356, 357} , Relaxation ^{156, 157} , Shengji Fuzheng ³²¹ , Shark cartilage ³⁸⁵ , Soy ²⁸⁰ , Supportive-Expressive Group therapy ³⁸⁶ , Tai-Chi ^{332, 387-389}
Physical Functioning	Mind-Body Cognitive Therapy ³⁵¹ , Music Therapy ¹⁰⁸ , Multi-modal ³¹⁸ , Myofascial Release ³²⁷ , Reflexology ²²⁷ , Stress management ¹¹⁵ , Tai-Chi ^{387, 389} , Yoga ^{77, 123, 170, 390}
Radiation therapy induced toxicity outcomes	Adlay Bran Extract ³⁹¹ , Alpha Ointment with Henna ³⁹² , Aquaphor-Biafine-RadiaCare ³⁹³ , Boswellia cream ³⁹⁴ , Calendula Cream ^{374, 395} , Chamomile ³⁹⁶ , Curcumin ³⁹⁷ , Glutamine ³⁹⁸ , Homeopathic Pills ³⁹⁹ , Honey ^{400, 401} , Hydration ⁴⁰² , Massage ⁴⁰³ , Oil-In-Water Emulsion ⁴⁰⁴ , Ray Gel ⁴⁰⁵ , Wheat Grass Extract ⁴⁰⁶ , Pentoxifylline & Vitamin E ^{407, 408}
Sleep disturbance	Acupuncture techniques ^{136, 137, 217, 114} , Calendula cream ³⁷⁴ , Meditation ^{148, 215, 409, 410} , Qigong ²²³ , Stress management
Vasomotor Outcomes	Black Cohosh ^{411, 412} , Flaxseed ³⁷⁸ , Homeopathy ^{381, 382} , Hypnosis ⁴¹³ , Magnetic therapy ⁴¹⁴ , Meditation ¹⁴⁹ , Peppermint ⁴¹⁵ , Vitamin E ⁴¹⁶ , Yoga ^{82, 417}

Notes: Definition of the I Statement: Concludes that the current evidence is insufficient to assess the balance of benefits and harms of the service. Evidence is lacking, of poor quality, or conflicting, and the balance of benefits and harms cannot be determined. Read the Clinical Considerations section of the USPSTF Recommendation Statement. If the service is offered, patients should understand the uncertainty about the balance of benefits and harms.

Accepted Article

SUPPLEMENTAL TABLES

Supplemental Table 1: List of clinical outcomes of interest (in alphabetical order)

0
1
2
3
4
5
6
7
8
9
0
1
2
3
4
5
6
7
8
9
0
1
2
3
4
5
6
7
8
9
0
1
2
3
4
5
6
7
8
9

Fatigue
Gastrointestinal
Gynecological
Hematological
Lymphedema
Neurological
Neuromuscular
Pain
Psychological
Quality of life
Renal
Skin
Sleep
Vasomotor symptoms

Supplemental Table 2: Summary of randomized controlled trials of A and B grade integrative therapies for anxiety

Modality/ Therapy	Author, Year	Population, Sample Size	Intervention and Control Groups	Study Conclusions and Limitations
Meditation (A grade)	Carlson, 2013 ⁷²	Stage I-III; N=271	INTERVENTION: A mindfulness intervention patterned after Jon Kabat Zinn's MBSR program consisted of 8 weekly 90-minute group sessions and a 6-hour workshop. INTERVENTION: Supportive Expressive Therapy of 12 weekly 90-minute group sessions aiming to facilitate family support, enhance emotional expression, integrate a changed body image into the new view of the self, improve coping skills and doctor patient relationships, and detoxify feelings around death CONTROL: 6-hour didactic seminar on stress management	Greater decrease in stress levels in MBSR intervention compared to control
	Crane-Okada, 2012 ⁷³	50+ yo post-treatment; N=49	INTERVENTION: A Mindful Movement Program of 12 weekly 2-hour sessions led by a trained instructor included mindful walking/moving, group discussion, exploration of body parts, specific and deliberate movements, moving with intentional effort, active energetic movement, and partner work consisted CONTROL: SC	No group differences in anxiety symptoms
	Kim, 2013 ⁷⁴	Post-surgery receiving RT; N=102	INTERVENTION: Brain Wave Vibration meditation of two 60-minute sessions per week for 6 weeks included simple, rhythmic movements with music, action, and positive messages CONTROL: SC	Greater decrease in anxiety in intervention compared to control from implementation of meditation program following surgery
	Lengacher, 2009 ⁷⁵	Within 18 months post-treatment transitioning back to daily life; N=84	INTERVENTION: Led by a trained psychologist, MBSR included sitting and walking meditation, body scan, yoga, and home practice for 2-hour sessions weekly for 6 weeks CONTROL: SC	Compared to control, greater improvements in anxiety in intervention group, and in trait anxiety and perceived stress in those who engaged in more hours of body scan and sitting meditation, suggesting those as more beneficial practices
	Würtzen, 2013 ⁷⁶	Stage I-III, 3-18 months post-surgery; N=336	INTERVENTION: Led by a trained clinical psychologist, a MBSR program of 8 weekly 2-hour group sessions and a 5-hour silent retreat included guided meditation, yoga, psycho-educational advice, group discussion, and home practice CONTROL: SC	Greater decrease in anxiety in intervention compared to control after 12 months from implementation of meditation program following surgery
Music Therapy (B grade)	Binns-Turner, 2011 ¹⁰⁶	Undergoing surgery; N=30	INTERVENTION: Passive music therapy where participants chose a music genres (classical, easy listening, inspirational, and new age), and listened with earphones and an iPod in a carrying case that covered the display to 4 hours of non-repeating continuous music immediately after sedation CONTROL: SC; iPod and earphones that stored no music	Greater decrease in anxiety in intervention compared to control from implementation of passive music therapy following sedation for surgery

	Bulfone, 2009 ¹⁰⁷	Stage I-II, 40-60 yo post-surgery entering hospital for scheduled CT; N=60	INTERVENTION: Passive music therapy where participants chose a music genre (new age, nature melodies, film soundtrack, Celtic melodies, and classical music) and listened for 15 minutes on a Walkman with earphones. CONTROL: SC; in the waiting room with no music	Greater decrease in anxiety in intervention compared to control from implementation of passive music therapy while waiting for CT
	Hanser, 2006 ¹⁰⁸	Stage IV; N=70	INTERVENTION: Led by a certified music therapist during treatment or in a consultation room, participants created live music through improvisation, song writing, and playing simple percussion instruments for three 45-minute sessions CONTROL: SC	Significant immediate effects of music therapy were observed on relaxation, but not on anxiety
	Li, 2012 ¹⁰⁹	Post-surgery; N=120	INTERVENTION: Passive music therapy where participants chose volume and a music genre (Chinese classical folk music, famous world music, music recommended by the American Association of Music Therapy, and Chinese relaxation music), and listened on earphones and an MP3 player twice a day for 30 minutes per session following surgery, and during the second and third time of hospital admission for CT CONTROL: SC; routine nursing care	Greater decrease in anxiety in intervention compared to control from implementation of passive music therapy at all three time points, following surgery and waiting for CT
	Zhou, 2015 ¹¹⁰	Post-surgery; N=170	INTERVENTION: Within 48 hours after surgery, researchers trained in music therapy and PMR taught patients to do PMR and listen to music on MP3 players using headphones, twice a day in the early morning and late evening for 30-minute sessions; music was delivered through MP3 players with 230 songs in Chinese relaxation music, classical folk music, religious music, and music recommended by the American Association of Music Therapy. CONTROL: SC	While anxiety symptoms decreased over time for both groups, a larger and faster decrease in anxiety was seen in the music therapy and PMR training group compared to control.
Stress Management (B grade)	Aguado Loi, 2012 ¹¹⁶	Newly diagnosed, pre-CT; N=220; 59.5% BC patients	INTERVENTION: Patients received instructions on PMR, guided imagery, abdominal breathing and coping skills training CONTROL: SC	No significant improvements in either group on anxiety, although patients presented with lower stress scores at the outset than in previous studies
	Garssen, 2013 ¹¹⁴	Stage I-III post-surgery; N=85	INTERVENTION: Four 45-60 minute sessions on days 1 and 5 pre-surgery and 2 and 30 post-surgery included guided imagery techniques, relaxation, meditative exercises and counseling CONTROL: SC	Anxiety similarly decreased in both groups. Patients were very satisfied with the meetings with the clinical psychologist and gave high-perceived value to the information delivered
	Jacobsen, 2013 ¹¹⁵	With treatment plan including CT; N=460; 33% BC patients	INTERVENTION: Self-directed stress management training singly and in combination with exercise CONTROL: SC	Greater decrease in anxiety in intervention with combined stress management and exercise compared to either alone and control
	Phillips, 2008 ¹¹⁷	Recently diagnosed; N=128	INTERVENTION: Led by 2 trained female facilitators, 10 2-hour weekly sessions consisted of cognitive restructuring, social support, assertiveness, anger management, meditation, coping skills training, abdominal breathing, guided imagery and PMR CONTROL: A 6-hour psychoeducational group seminar of a condensed demonstration of several of the above techniques	Greater increases in ability to relax and reduced cortisol levels in cognitive-behavioral stress management intervention compared to controls across time

Yoga (B grade)	Banerjee, 2007 ¹¹⁹	Post-surgery with treatment plan of RT or RT/CT; N=68	INTERVENTION: Led by expert yoga teachers, a custom, intensive, integrated yoga program of 90-minute sessions over 6-weeks included slow stretching and loosening exercises, various postures (asanas), guided imagery specific to cancer, positive thought provocation, chanting exercises, various breathing exercises (Pranayama), and soothing sound vibrations and guided imagery (yoga nidra) CONTROL: Brief weekly supportive counseling and advised to engage in light exercise	Decrease in perceived stress and anxiety in yoga group after 6 weeks, and no change in control. Unclear if yoga intervention was weekly or daily, or if it was individual or in groups
	Bower, 2012 ¹²³	Stage 0-II, 40-65 yo post-treatment with fatigue; N=31	INTERVENTION: Led by a trained instructor, a 90-minute twice a week for 12 weeks Iyengar and Hatha yoga in small group classes CONTROL: A 120-minute health education class facilitated by a psychologist, held once a week for 12 weeks	Perceived stress decreased over the assessment period in both groups
	Chandwani, 2010 ¹²⁶	Stage 0-III scheduled for RT; N=61	INTERVENTION: A trained yoga instructor provided two 60-minute yoga classes per week for 6-weeks based on Patanjali's Yoga Sutras during RT treatment with up to two participants, and included home practice. CONTROL: WLC	No changes in anxiety in either group
	Dhruva, 2012 ¹²⁴	Receiving CT; N=16	INTERVENTION: Pranayama or yogic breathing consisted of a weekly 60-minute class with 4 breathing practices with an instructor and 1-2 participants during 2 consecutive CT cycles, and included home practice CONTROL: SC during cycle 1 of CT and intervention during cycle 2	Anxiety decreased more in yoga group; greater improvements with more practice of pranayama
	Pruthi, 2012 ¹²⁵	Receiving treatment; N=30	INTERVENTION: Led by a registered yoga instructor, 60-minute weekly individual therapeutic gentle hatha yoga sessions for 8 weeks focused on breathing techniques (pranayama) and included home practice CONTROL: SC	Perceived stress improved similarly in both groups
	Raghavendra, 2007 ¹²¹	Stage II or III receiving CT; N=62	INTERVENTION: Individual 30-minute bedside sessions prior to receiving each CT injection of yogic relaxation, meditation using breath awareness or chanting a mantra from a Vedic text, and included home practice CONTROL: Individual 60-minute coping preparation and psychodynamic supportive-expressive therapy sessions was given at the bedside prior to first CT cycle	Greater improvements in state and trait anxiety in yoga group than control
	Rao, 2009 ¹²²	Stage II-III RT with or without CT; N=98	INTERVENTION: Facilitated by 2 instructors, yoga program included asanas (postures), breathing exercises, pranayama (voluntarily regulated nostril breathing), meditation, yogic relaxation techniques with imagery, and home practice for 4 individual sessions during the pre-post-operative period and 3 individual sessions per week for 6 weeks during RT and CT CONTROL: Brief 15-minute supportive therapy sessions with individual counseling by a trained social worker once every 10 days, during hospital visits over 24-weeks	Decrease in both state and trait anxiety in yoga group compared to controls

Taso, 2014 ⁷⁹	Stage I-III, nonmetastatic, undergoing CT; N=60	INTERVENTION: Yoga exercise program facilitated by an experienced instructor twice a week for 60 minute classes over an eight week period (16 sessions) during chemotherapy treatment CONTROL: SC	No improvement of anxiety following intervention
Vadiraja, 2009 ¹²⁰	Stage II and III post-surgery receiving RT; N=88	INTERVENTION: Same as Banerjee et al. 2007 program for a minimum of 3 60-minute sessions per week over a period of 6 weeks during RT CONTROL: Brief weekly supportive counseling and advised to engage in light exercise	Greater decreases in anxiety in the yoga compared with control groups

0
1 Abbreviations: BC, Breast Cancer; CBT, Cognitive Behavioral Therapy; CT, Chemotherapy; MBSR, Mindfulness-Based Stress Reduction; PMR, Progressive Muscle Relaxation;
2 RT, Radiation Therapy; SC, Standard Care; WLC, Wait List Control; yo, years old
3
4
5
6
7
8
9
0
1
2
3
4
5
6
7
8
9
0
1
2
3
4
5
6
7
8
9
0
1
2
3
4
5
6
7
8
9

Supplemental Table 3: Summary of randomized controlled trials of A and B grade integrative therapies for mood disturbances/depression

Modality/Therapy	Author, Year	Population, Sample Size	Intervention and Control Groups	Study Conclusions and Limitations
Meditation (A grade)	Crane-Okada, 2012 ⁷³	50+ yo post-treatment; N=49	INTERVENTION: A Mindful Movement Program that included mindful walking/moving, group discussion, exploration of body parts, specific and deliberate movements, moving with intentional effort, active energetic movement, and partner work consisted of 12 weekly 2-hour sessions led by a trained instructor CONTROL: SC	No group differences on depression symptoms
	Carlson, 2013 ⁷²	Stage I-III; N=271	INTERVENTION: A mindfulness intervention patterned after Jon Kabat Zinn's MBSR program consisted of 8 weekly group sessions of 90 minutes and a 6 hour workshop. INTERVENTION: Supportive Expressive Therapy of 12 weekly group sessions of 90 minutes that aims to facilitate mutual and family support, enhance emotional openness and expression, integrate a changed body image into the new view of the self, improve coping skills and doctor patient relationships and detoxify feelings around death CONTROL: 6-hour didactic seminar on stress management	Women in MBSR improved more over time in mood
	Dodds, 2015 ¹⁵²	Survivors; N=33	INTERVENTION: The Cognitively-Based Compassion Training was delivered in eight weekly, 2-hour group classes and one booster session 4 weeks later by a certified instructor. The program is a secular adaptation of traditional Tibetan Buddhist methods for cultivating compassion (<i>lojong</i>). The classes included developing meditative concentration (<i>shamatha</i>) and mindfulness, and incorporated a series of contemplative exercises. CONTROL: WLC	CBCT group showed improvements in depression scores
	Henderson, 2013 ¹⁵¹	Stage I or II receiving RT; N=172	INTERVENTION: Facilitated by trained graduate-level instructor, MBSR included an introductory and 8 weekly 2.5-3.5 hour larger group sessions, a 7.5-hour retreat, and 3 2-hour sessions facilitated by a psychiatrist at monthly intervals following the conclusion of the MBSR intervention. CONTROL: Nutrition education program (NEP) led by a registered dietician and matched the MBSR intervention for contact time and homework assignments CONTROL: SC	At 4 months, depression scores decreased more in MBSR than nutrition control and SC but this did not persist over follow-up
	Hoffman, 2012 ¹⁴⁹	Stage 0-III post-surgery post-treatment 2 months - 2 years prior; N=229	INTERVENTION: Administered by a trained instructor, MBSR consisted of 8 weekly 2-hour classes, plus a 6-hour retreat and included body scan and sitting meditation, gentle lying and standing yoga-based stretches, group discussion, didactic teaching, and home practice CONTROL: SC, WLC	MSBR improved mood and well-being more than SC, which persisted at three months

	Kim, 2013 ⁷⁴	Post-surgery receiving RT; N=102	INTERVENTION: Brain Wave Vibration meditation combines simple, rhythmic movements with music, action, and positive messages and consisted of two 60-minute sessions per week for 6 weeks CONTROL: SC	No group differences on depression symptoms
	Lengacher, 2009 ⁷⁵	Within 18 months post-treatment; N=84	INTERVENTION: Led by a trained psychologist, MBSR included sitting and walking meditation, body scan, yoga, and home practice for 2-hour sessions weekly for 6 weeks CONTROL: SC	MBSR had significantly lower levels of depression at 6 weeks
	Milbury, 2013 ¹⁴⁸	Stage I-III post-treatment; N=47	INTERVENTION: Tibetan sound meditation includes breathing, awareness, concentration, visualization and sound exercises, and composed of 2 weekly instructor led 60-minute classes for 6 weeks, and included home practice CONTROL: WLC	Improved depression symptoms at end of treatment in meditation over control but not one month later
	Nidich, 2009 ¹⁵⁰	Stage II-IV over 55 yo; N=130	INTERVENTION: Transcendental Meditation in a 7-session course led by a qualified instructor and included home practice for the duration of the study (32 months) CONTROL: SC; provided with basic educational materials	Significant improvements in the intervention group compared with controls in emotional well-being and overall mental health
	Würtzen, 2013 ⁷⁶	Stage I-III 3-18 months post-surgery; N=336	INTERVENTION: Led by a trained clinical psychologist, a MBSR program including guided meditation, yoga, psycho-educational advice, group discussion, and home practice composed of 8 weekly 2-hour group sessions and a 5-hour silent retreat CONTROL: SC	After 12 months MBSR had clinically meaningful improvements on depression symptoms compared to control
Relaxation (A grade)	Gudenkauf, 2015 ¹⁵⁸	Stage 0-III; N=183	INTERVENTION: two groups, relaxation training (RT) and cognitive-behavioral training (CBT) RT: In-session experiential training including abdominal breathing, guided imagery, meditation, and PMR, and out-of-session information and assignments for home practice administered over 5-weeks CBT: structured group cognitive-behavioral sessions that included awareness of stressors, coping skills, skills for social support, and anger management administered over 5-weeks CONTROL: Health Education control	Both CBT and RT groups reported reduced depressive affect compared to control. No differences between CBT and RT groups for depressive affect were found.
	Hidderley, 2004 ¹⁴³	Early stage post-surgery receiving RT; N=31	INTERVENTION: Autogenic training or Western meditation that is meant to induce heaviness and warmth of limbs; calming of the heart and breathing; abdominal warmth; and cooling of the forehead; timing was not clearly reported but appears to have occurred weekly following lumpectomy and during RT over a 2-month period CONTROL: Home visit	Significant reductions in depression symptoms compared to control
	Molassiotis, 2002 ¹⁴⁶	Recently diagnosed scheduled to receive CT; N=71	INTERVENTION: PMR followed by imagery limited to the hour prior to and for five days after the first CT round ever. CONTROL: SC	Mood disturbance score decreased in intervention group, while it increased significantly in the control
	Nunes, 2007 ¹⁴⁷	Stage I or II receiving RT; N=34	INTERVENTION: Relaxation and visualization therapy for 24 days total CONTROL: SC	Significant reductions in depression symptoms compared to control

	Yoo, 2005 ¹⁵⁶	Stage II-III post-surgery within 2 months; N=60	INTERVENTION: PMR and guided imagery applied the hour before 6 CT administrations, and included home practice CONTROL: SC	Significant reductions in depression symptoms compared to control
	Walker, 1999 ¹⁵⁷	Newly diagnosed; N=96	INTERVENTION: PMR and imagery daily across 6 cycles of CT CONTROL: SC	Slight improvement in mood in intervention compared to control
Yoga (B grade)	Banerjee, 2007 ¹¹⁹	Post-surgery with treatment plan of RT or RT/CT; N=68	INTERVENTION: Led by expert yoga teachers, a custom, intensive, integrated yoga program of 90-minute sessions over 6-weeks included slow stretching and loosening exercises, various postures (asanas), guided imagery specific to cancer, positive thought provocation, chanting exercises, various breathing exercises (Pranayama), and soothing sound vibrations and guided imagery (yoga nidra) CONTROL: Brief weekly supportive counseling and advised to engage in light exercise	Greater decrease in depression symptoms in yoga group
	Bower, 2012 ¹²³	Stage 0-II 40-65 yo post-treatment; N=31	INTERVENTION: Led by a trained instructor, a 90-minute twice a week for 12 weeks Iyengar and Hatha yoga in small group classes CONTROL: A 120-minute health education class facilitated by a psychologist, held once a week for 12 weeks	Both groups had reduced depressive symptoms from baseline to post-treatment, with a greater decline in the yoga group. No group differences at 3-month follow-up
	Chandwani, 2014 ⁷⁷	Stage 0-III scheduled for RT; N=178	INTERVENTION: A trained yoga instructor provided up to three 60-minute yoga classes based on Patanjali's Yoga Sutras each week during 6-week RT treatment including up to two participants at a time, with the majority of sessions being one-on-one, and included home practice CONTROL: A 60-minute stretching program was provided up to three times a week during 6-week RT treatment by physiotherapists. Stretches were introduced in a stepped approach and approximated the gross movements of the yoga intervention CONTROL: WLC	No group differences on depression scales
	Chandwani, 2010 ¹²⁶	Stage 0-III scheduled for RT; N=61	INTERVENTION: A trained yoga instructor provided up to 2 60-minute yoga classes based on Patanjali's Yoga Sutras each week during 6-week RT treatment including up to two participants at a time, with the majority of sessions being one-on-one, and included home practice CONTROL: WLC	No group differences on depression symptoms
	Culos-Reed, 2006 ¹⁶⁸	Post-treatment; N=38	INTERVENTION: Led by a certified yoga instructor, weekly 75-minute small group classes over a 7-week period consisted of gentle Hatha yoga including breathing, stretching, strengthening exercises, and relaxation CONTROL: WLC	Yoga group showed trend to reduced depression
	Danhauer, 2009 ¹⁷⁰	Post-surgery; N=44	INTERVENTION: Led by a registered yoga instructor, 10 weekly 75-minute group yoga classes incorporated physical postures, breathing exercises, and deep relaxation, with no requirement for home practice CONTROL: WLC	Yoga group improved more on mental health, depression, positive affect scores than control; benefits were greater for those with greater psychological morbidity at baseline

Dhruva, 2012 ¹²⁴	Receiving CT; N=16	INTERVENTION: Pranayama or yogic breathing consisted of a weekly 60-minute class with 4 breathing practices with an instructor and 1-2 participants during 2 consecutive CT cycles, and included home practice CONTROL: SC during cycle 1 of CT and intervention during cycle 2	No group differences on depression symptoms
Kiecolt-Glaser, 2014 ⁷⁸	Stage 0-IIIa post-treatment; N=200	INTERVENTION: Certified instructors facilitated two 90-minute group hatha yoga classes per week for 12 weeks. The classes consisted of 4 to 20 women. Home practice was encouraged. CONTROL: WLC	No group differences on depression
Moadel, 2007 ¹⁶⁹	Ethnically diverse, recently diagnosed; N=128	INTERVENTION: 12 weekly 1.5-hour group classes based on Hatha yoga including physical stretches and poses, breathing exercises, meditation, and home practice, all completed in a seated or reclined position CONTROL: WLC	Among patients not receiving chemotherapy, yoga improved emotional well-being and mood
Pruthi, 2012 ¹²⁵	Receiving treatment; N=30	INTERVENTION: Led by a registered yoga instructor, 60-minute weekly individual therapeutic gentle hatha yoga sessions for 8 weeks focused on breathing techniques (pranayama) and included home practice CONTROL: SC	Mood improved equally in both groups
Rao, 2015 ⁸⁰	Stage II and III undergoing surgery followed by adjuvant RT and/or CT; N=98	INTERVENTION: Individual integrated yoga program provided during hospital visits and at-home practice on remaining days. Yoga practices consisted of a set of asanas, breathing exercises, voluntarily regulated nostril breathing (pranayama), meditation, and relaxation with imagery. CONTROL: Individual, unstructured supportive-expressive therapy with an education component provided during hospital visits and extended over the course of RT and CT cycles (i.e., once in 10 days for 30 minute sessions)	Improvements in depression symptoms in yoga compared to control, though both groups improved in time.
Raghavendra, 2007 ¹²¹	Stage II or III receiving CT; N=62	INTERVENTION: Individual 30-minute bedside sessions prior to receiving each CT injection of yogic relaxation, meditation using breath awareness, or chanting a mantra from a Vedic text and included home practice CONTROL: Individual 60-minute coping preparation and psychodynamic supportive-expressive therapy sessions was given at the bedside prior to first CT cycle	Improvements in depression symptoms in yoga compared to control.
Taso, 2014 ⁷⁹	Stage I-III, nonmetastatic, undergoing CT; N=60	INTERVENTION: Yoga exercise program facilitated by an experienced instructor twice a week for 60 minute classes over an eight week period (16 sessions) during chemotherapy treatment CONTROL: SC	No improvement of mood disturbance following intervention
Vadiraja, 2009 ^{120, 171}	Stage II and III post-surgery receiving RT; N=88	INTERVENTION: Same as Banerjee et al. 2007 program for a minimum of three 60-minute sessions per week over a period of 6 weeks during RT CONTROL: Brief weekly supportive counseling and advised to engage in light exercise	Significant decreases in depression and improvement in positive affect in yoga compared to control

Massage (B grade)	Fernandez-Lao, 2012 ¹⁷⁷	Stage I-IIIa with fatigue between 25 and 65 yo post-treatment; N=20	INTERVENTION: Led by a trained physical therapist with BC survivors, a 40-minute myofascial intervention focused on the neck–shoulder area, longitudinal stroke, J stroke, sub-occipital sustained pressure, frontalis bone spread and ear pull techniques CONTROL: SC plus special attention to the patient for 40 minutes consisting of a clinician speaking about nutrition, physical activity, and relaxation techniques for improving quality of life after BC	Massage improved overall mood and subscales or tension–anxiety, depression–dejection and anger–hostility more than control
	Hernandez-Reif, 2004 ¹⁴⁰	Stage I or II post-surgery; N=34	INTERVENTION: Classic massage with acupressure and Trager®, which uses hundreds of small, rocking and elongating movements that release muscle tension, was provided by a registered massage therapist for three 30-minute massages each week for 5 weeks CONTROL: SC	Immediate massage therapy effects included reduced depression
	Krohn, 2011 ¹⁷⁵	Early Stage post-treatment; N=34	INTERVENTION: Classic massage therapy by licensed, female massage therapists in a quiet and private room CONTROL: SC	Massage significantly decreased depression
	Listing, 2009 ¹⁷⁶	Stage 0-II post-treatment; N=86	INTERVENTION: Classic massage therapy by licensed, female massage therapists in a quiet and private room CONTROL: SC and informed that they would receive PMR after the completion of the study	Classical massage improved short-term mood disturbance though study suffered from unbalanced study arms reducing power of the study
	Listing, 2010 ¹⁴¹	Post-treatment; N=34	INTERVENTION: Classic massage therapy by the same licensed, female massage therapist for each session in a quiet and private room CONTROL: WLC	Massage improved mood compared to control
	Wilkinson, 2007 ¹⁴²	Diagnosed with clinical depression and/or anxiety; N=288; 55% BC	INTERVENTION: Standard supportive care and a 1-hour individually tailored massage provided weekly for 4 weeks CONTROL: SC	Greater improvement in self report depression at week 6 in intervention; both groups improved in depression at week 10, but no group differences
Music Therapy (B grade)	Burns, 2001 ¹⁸⁵	Completed RT and CT, under 65 yo; N=8	INTERVENTION: Led by a board-certified music therapist who implemented the Bonny Method of Guided Imagery and Music and played compact discs from the collection “Music for the Imagination”, music genres included Western art music and classical music for 10 weekly 90-110-minute sessions. CONTROL: WLC	More improvement in mood in music group than WLC.
	Hanser, 2006 ¹⁰⁸	Stage IV; N=70	INTERVENTION: Led by a certified music therapist during treatment or in a consultation room, participants created live music through improvisation, song writing, and playing simple percussion instruments for 3 45-minute sessions CONTROL: SC	Significant immediate effects of active music therapy for happiness, but no significant differences between conditions over time on depression.
	Zhou, 2015 ¹¹⁰	Post-surgery; N=170	INTERVENTION: Within 48 hours after surgery, researchers trained in music therapy and PMR taught patients to do PMR and listen to music on MP3 players using headphones, twice a day in the early morning and late evening for 30-minute sessions; music was delivered through MP3 players with 230 songs in Chinese relaxation music, classical folk music, religious music, and music recommended by the American Association of Music Therapy.	While depression symptoms decreased over time for both groups, a larger and faster decrease in depression was seen in the music therapy and PMR training group compared to control.

			CONTROL: SC	
	Zhou, 2011 ¹⁸⁶	Post-surgery under 65 yo; N=120	INTERVENTION: Passive music therapy where participants chose volume and a music genre (Chinese classical folk music, famous world music, music recommended by the American Association of Music Therapy, and Chinese relaxation music), and listened on earphones and an MP3 player twice a day for 30 minutes per session following surgery, and during the second and third time of hospital admission for CT CONTROL: SC; routine nursing care	Music therapy group had lower depression scores than control at all post-tests (2 months)

3 Abbreviations: BC, Breast Cancer; CBT, Cognitive Behavioral Therapy; CT, Chemotherapy; MBSR, Mindfulness-Based Stress Reduction; NS, Not Specified; PMR, Progressive
4 Muscle Relaxation; RT, Radiation Therapy; SC, Standard Care; UC, Usual Care; WLC, Wait List Control; yo, years old

Supplemental Table 4: Summary of randomized controlled trials of A and B grade integrative therapies to improve quality of life

Modality/Therapy	Author, Year	Population, Sample Size	Intervention and Control Groups	Study Conclusions and Limitations
Meditation (A grade)	Crane-Okada, 2012 ⁷³	50+ yo post-treatment; N=49	INTERVENTION: A Mindful Movement Program that included mindful walking/moving, group discussion, exploration of body parts, specific and deliberate movements, moving with intentional effort, active energetic movement, and partner work consisted of 12 weekly 2-hour sessions led by a trained instructor CONTROL: SC	Mindful Movement Program better than SC
	Henderson, 2012 ²¹²	Stage I-II; N=172	INTERVENTION: Facilitated by a trained graduate-level instructor, MBSR included an introductory and 8 weekly 2.5-3.5 hour larger group sessions, a 7.5-hour retreat, and 3 2-hour sessions facilitated by a psychiatrist at monthly intervals following the conclusion of the MBSR intervention. CONTROL: Nutrition education program (NEP) led by a registered dietician and matched the MBSR intervention for contact time and homework assignments CONTROL: SC	Stress management better than nutrition control and SC
	Henderson, 2013 ¹⁵¹	Stage I or II receiving RT; N=110	Same as Henderson, 2012 above	MBSR better than control groups
	Hoffman, 2012 ¹⁴⁹	Stage 0-III post-surgery post-treatment 2 months - 2 years prior; N=229	INTERVENTION: MBSR administered by a trained instructor, followed the traditional program design, with women attending 8 weekly 2-hour classes, plus a 6-hour retreat; formal mindfulness practices included: body scan meditation, gentle lying and standing yoga-based stretches, sitting meditation, group discussion, didactic teaching, and home practice; participants were provided with four 45-minute guided meditation CDs and a manual to facilitate the recommended 40-45 minute daily at home practice (6 or 7 days per week). CONTROL: SC, WLC	MBSR better than WLC
	Kim, 2013 ⁷⁴	Post-surgery receiving RT; N=102	INTERVENTION: Brain Wave Vibration meditation combines simple, rhythmic movements with music, action, and positive messages and consisted of two 60-minute sessions per week for 6 weeks CONTROL: SC	Meditation better than SC
	Lengacher, 2009 ⁷⁵	Within 18 months post-treatment; N=84	INTERVENTION: Led by a trained psychologist, MBSR included sitting and walking meditation, body scan, yoga, and home practice for 2-hour sessions weekly for 6 weeks CONTROL: SC	MBSR better than SC
	Nidich, 2009 ¹⁵⁰	Stage II-IV over 55 yo; N=130	INTERVENTION: Transcendental Meditation in a 7-session course led by a qualified instructor and included home practice for the duration of the study (32 months)	Meditation better than SC

			CONTROL: SC; provided with basic educational materials	
	Banasik, 2011 ²⁰⁴	Stage II-IV, at least 2 months post-treatment; N=18	INTERVENTION: An 8-week traditional Iyengar yoga class was delivered in 90-minute sessions, two times per week. CONTROL: WLC	Yoga better than WLC
	Chandwani, 2010 ¹²⁶	Stage 0-III scheduled for RT; N=61	INTERVENTION: A trained yoga instructor provided up to 2 60-minute yoga classes based on Patanjali's Yoga Sutras each week during 6-week RT treatment including up to two participants at a time, with the majority of sessions being one-on-one, and included home practice CONTROL: WLC	Yoga better than WLC and SC
	Cramer, 2015 ⁸²	Stage I-III post-treatment; N=40	INTERVENTION: A 12-week traditional Hatha yoga and meditation intervention was facilitated weekly by certified a instructor in 90-minute sessions. Classes focused on guided relaxation, breathing techniques, yoga postures, and meditation practices. Participants were encouraged to practice at home. CONTROL: WLC	Yoga better than WLC
	Culos-Reed, 2006 ¹⁶⁸	Post-treatment; N=38	INTERVENTION: Led by a certified yoga instructor, weekly 75-minute small group classes over a 7-week period consisted of gentle Hatha yoga including breathing, stretching, strengthening exercises, and relaxation CONTROL: WLC	Yoga better than WLC
	Danhauer, 2009 ¹⁷⁰	Post-surgery; N=44	INTERVENTION: Led by a registered yoga instructor, 10 weekly 75-minute group yoga classes incorporated physical postures, breathing exercises, and deep relaxation, with no requirement for home practice CONTROL: WLC	Yoga better than WLC
	Dhruva, 2012 ¹²⁴	Receiving CT; N=16	INTERVENTION: Pranayama or yogic breathing consisted of a weekly 60-minute class with 4 breathing practices with an instructor and 1-2 participants during 2 consecutive CT cycles, and included home practice CONTROL: SC during cycle 1 of CT and intervention during cycle 2	Yoga better than SC
	Littman, 2012 ²⁰⁵	Stage 0-III post-treatment with BMI ≥ 24 kg/m ² ; N=63	INTERVENTION: –Intervention was based on viniyoga, a Hatha style of yoga that involves physical stretches and poses, breath control, and meditation. Adapted for use with overweight or obese BC survivors without yoga experience. Practiced five times per week, including at least one 75-minute facility-based class. CONTROL: WLC	Yoga better than WLC
	Moadel, 2007 ¹⁶⁹	Recently diagnosed; N=128	INTERVENTION: 12 weekly 1.5-hour group classes based on Hatha yoga including physical stretches and poses, breathing exercises, meditation, and home practice, all completed in a seated or reclined position CONTROL: WLC	Yoga better than SC
	Pruthi, 2012 ²¹⁶	Receiving treatment; N=15	INTERVENTION: Led by a registered yoga instructor, 60-minute weekly individual therapeutic gentle hatha yoga sessions for 8 weeks focused on breathing techniques (pranayama) and included home practice CONTROL: SC	Yoga better than SC

Raghavendra, 2007 ¹²¹	Stage II or III receiving CT; N=62	INTERVENTION: Individual 30-minute bedside sessions prior to receiving each CT injection of yogic relaxation, meditation using breath awareness, or chanting a mantra from a Vedic text and included home practice CONTROL: Individual 60-minute coping preparation and psychodynamic supportive-expressive therapy sessions was given at the bedside prior to first CT cycle	Yoga better than SC
Siedentopf, 2013 ⁸¹	Post-surgery; N=93	INTERVENTION: 75-minute classes facilitated by certified instructors were provided at the gym in the hospital twice a week for five weeks. The program focused on yoga poses (asanas), breathing exercises, eye exercises, and concentration exercises. CONTROL: WLC 5 weeks after surgery	Yoga better than WLC
Vadiraja, 2009 ¹⁷¹	Stage II and III post-surgery receiving RT; N=88	INTERVENTION: Same as Banerjee et al. 2007 program for a minimum of 3 60-minute sessions per week over a period of 6 weeks during RT CONTROL: Brief weekly supportive counseling and advised to engage in light exercise	Yoga better than control group

Abbreviations: BC, Breast Cancer; CT, Chemotherapy; MBSR, Mindfulness-Based Stress Reduction; RT, Radiation Therapy; SC, Standard Care; WLC, Wait List Control; yo, years old;

Supplemental Table 5: Summary of randomized controlled trials of A- and B-graded integrative therapies in breast cancer patients with chemotherapy-induced nausea and vomiting outcomes

Modality/Therapy	Author, Year	Population, Sample Size	Intervention and Control Groups	Study Conclusions and Limitations
Acupressure (B grade)	Dibble, 2000 ²³⁷	Receiving CT; N=17	INTERVENTION: Self-applied acupressure immediately prior to CT session during one cycle of CT. CONTROL: SC	Acupressure better than SC
	Dibble, 2007 ²³⁸	Receiving CT; N=160	INTERVENTION: Daily self-treatment with verum acupressure to the P6 acupressure point CONTROL: Daily self-treatment with placebo acupressure to the SI3 acupressure point CONTROL: SC	Acupressure better than sham and SC to decrease incidence of delayed nausea and vomiting; Acupressure no different than sham or SC for acute nausea and vomiting

	Molassiotis, 2007 ²³⁹	Stage I-III receiving CT; N=54	INTERVENTION: Acupressure was applied using Sea-Band™ wristbands worn bilaterally with the stud pressing on the P6 acupressure point that they wore for the 5 days following CT administration CONTROL: WLC; participants received antiemetics and were told that they would receive the acupressure instructions and be given the wristbands to use starting in their next cycle of CT	Sea-bands better than SC
0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 6 7 8 9	Beith, 2012 ²⁴⁶	Early stage receiving CT; N=32	INTERVENTION: Electroacupuncture for 20 minutes at least 2 hours prior to CT on the first two days of the first two cycles using acupoints PC-6, LI-4, and ST-36 where prior to electrical stimulation, manual needling techniques were performed to induce the de qi sensation CONTROL: Sham electroacupuncture receiving identical treatment using placebo needles and non-functioning electrodes.	Electroacupuncture no better than sham likely due to the fact that the study was underpowered, only evaluating feasibility with minimal electroacupuncture intervention, and without a no-acupuncture arm
	Shen, 2000 ²⁴⁷	BC patients; N=104	INTERVENTION: Bilateral stimulation at PC-6 and ST-36 for 20 minutes once daily over 5 days where prior to electrical stimulation, manual needling techniques were performed to induce the de qi sensation CONTROL: Minimal needling with mock stimulation plus SC CONTROL: SC	Electroacupuncture better than minimal needling and SC

Abbreviations: BC, Breast Cancer; CINV, Chemotherapy-Induced Nausea Vomiting; CT, Chemotherapy; RT, Radiation Therapy; SC, Standard Care; WLC, Wait List Control; yo, years old

REFERENCES

1. Boon HS, Olatunde F, Zick SM. Trends in complementary/alternative medicine use by breast cancer survivors: comparing survey data from 1998 and 2005. *BMC Womens Health*. 2007;7: 4.
2. Greenlee H, Kwan ML, Ergas IJ, et al. Changes in vitamin and mineral supplement use after breast cancer diagnosis in the pathways study: a prospective cohort study. *BMC Cancer*. 2014;14: 382-397.
3. Link AR, Gammon MD, Jacobson JS, et al. Use of self-care and practitioner-based forms of complementary and alternative medicine before and after a diagnosis of breast cancer. *Evid Based Complement Alternat Med*. 2013;2013: 1-16.
4. Matsuno RK, Pagano IS, Maskarinec G, Issell BF, Gotay CC. Complementary and alternative medicine use and breast cancer prognosis: a pooled analysis of four population-based studies of breast cancer survivors. *J Womens Health (Larchmt)*. 2012;21: 1252-1258.
5. Greenlee H, Kwan ML, Ergas IJ, et al. Complementary and alternative therapy use before and after breast cancer diagnosis: the Pathways Study. *Breast Cancer Res Treat*. 2009;117: 653-665.
6. Bright-Gbejy M, Makambi KH, Rohan JP, et al. Use of multivitamins, folic acid and herbal supplements among breast cancer survivors: the Black Women's Health Study. *BMC Complement Altern Med*. 2011;11: 30-35.
7. Office of Cancer Complementary and Alternative Medicine. CAM Definitions. Available from URL: http://cam.cancer.gov/health_definitions.html [accessed 06/18/2015].
8. National Center for Complementary and Integrative Health. Complementary, alternative, or integrative health: what's in a name? Available from URL: <http://nccam.nih.gov/health/whatisacam> [accessed 6/18/15].

9. Greenlee H, Balneaves LG, Carlson LE, et al. Clinical Practice Guidelines on the Use of Integrative Therapies as Supportive Care in Patients Treated for Breast Cancer. *JNCI Monographs*. 2014;2014: 346-358.
10. U.S. Preventive Services Task Force. Grade Definitions. Available from URL: <http://www.uspreventiveservicestaskforce.org/uspstf/grades.htm> [accessed 5/10/14].
11. Institute of Medicine Committee on Standard for Developing Trustworthy Clinical Practice Guidelines. *Clinical Practice Guidelines We Can Trust*. Washington DC: The National Academies Press, 2011.
12. World Cancer Research Fund / American Institute for Cancer Research. *Food, nutrition, physical activity, and the prevention of cancer: a global perspective*. Washington, D.C.: AICR, 2007.
13. Rock CL, Doyle C, Demark-Wahnefried W, et al. Nutrition and physical activity guidelines for cancer survivors. *CA Cancer J Clin*. 2012;62: 243-274.
14. Schmitz KH, Courneya KS, Matthews C, et al. American College of Sports Medicine roundtable on exercise guidelines for cancer survivors. *Med Sci Sports Exerc*. 2010;42: 1409-1426.
15. Duijts SF, Faber MM, Oldenburg HS, van Beurden M, Aaronson NK. Effectiveness of behavioral techniques and physical exercise on psychosocial functioning and health-related quality of life in breast cancer patients and survivors--a meta-analysis. *Psychooncology*. 2011;20: 115-126.
16. Faller H, Schuler M, Richard M, Heckl U, Weis J, Kuffner R. Effects of psycho-oncologic interventions on emotional distress and quality of life in adult patients with cancer: systematic review and meta-analysis. *J Clin Oncol*. 2013;31: 782-793.
17. Galway K, Black A, Cantwell M, Cardwell CR, Mills M, Donnelly M. Psychosocial interventions to improve quality of life and emotional wellbeing for recently diagnosed cancer patients. *Cochrane Database Syst Rev*. 2012;11: CD007064.
18. Jadad AR, Moore RA, Carroll D, et al. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Control Clin Trials*. 1996;17: 1-12.

19. Verhagen AP, de Vet HC, de Bie RA, et al. The Delphi list: a criteria list for quality assessment of randomized clinical trials for conducting systematic reviews developed by Delphi consensus. *J Clin Epidemiol.* 1998;51: 1235-1241.
20. National Center for Complementary and Integrative Health. Health Topics A-Z. Available from URL: <https://nccih.nih.gov/health/atoz.htm> [accessed 6/18/15].
21. National Cancer Institute. Complementary and Alternative Medicine. Available from URL: <http://www.cancer.gov/cancertopics/cam> [accessed 6/18/15].
22. National Cancer Institute. NCI Dictionary of Cancer Terms. Available from URL: <http://www.cancer.gov/publications/dictionaries/cancer-terms> [accessed 6/27/16].
23. National Center for Complementary and Integrative Health. Acupuncture. Available from URL: <https://nccih.nih.gov/health/acupuncture> [accessed 6/18/15].
24. Cohen MR. *The New Chinese Medicine Handbook: An Innovative Guide to Integrating Eastern Wisdom with Western Practice for Modern Healing.* Fairwinds Press, 2015.
25. Shen Y, Liu L, Chiang JS, et al. Randomized, placebo-controlled trial of K1 acupoint acustimulation to prevent cisplatin-induced or oxaliplatin-induced nausea. *Cancer.* 2015;121: 84-92.
26. Napadow V, Makris N, Liu J, Kettner NW, Kwong KK, Hui KK. Effects of electroacupuncture versus manual acupuncture on the human brain as measured by fMRI. *Hum Brain Mapp.* 2005;24: 193-205.
27. Gottschling S, Reindl TK, Meyer S, et al. Acupuncture to alleviate chemotherapy-induced nausea and vomiting in pediatric oncology - a randomized multicenter crossover pilot trial. *Klin Padiatr.* 2008;220: 365-370.
28. Rithirangsiroj K, Manchana T, Akkayagorn L. Efficacy of acupuncture in prevention of delayed chemotherapy induced nausea and vomiting in gynecologic cancer patients. *Gynecol Oncol.* 2015;136: 82-86.

29. Melchart D, Ihbe-Heffinger A, Leps B, von Schilling C, Linde K. Acupuncture and acupressure for the prevention of chemotherapy-induced nausea--a randomised cross-over pilot study. *Support Care Cancer*. 2006;14: 878-882.
30. Garcia MK, McQuade J, Haddad R, et al. Systematic review of acupuncture in cancer care: a synthesis of the evidence. *J Clin Oncol*. 2013;31: 952-960.
31. Zick SM, Sen A, Wyatt GK, Murphy SL, Arnedt JT, Harris RE. Investigation of 2 Types of Self-administered Acupressure for Persistent Cancer-Related Fatigue in Breast Cancer Survivors: A Randomized Clinical Trial. *JAMA Oncol*. 2016;2: 1470-1476.
32. Memorial Sloan Kettering Cancer Center. Integrative Medicine: About Herbs, Botanicals and Other Products. Available from URL: <https://www.mskcc.org/cancer-care/treatments/symptom-management/integrative-medicine/herbs> [accessed June 27, 2016].
33. National Center for Complementary and Integrative Health. Ginger. Available from URL: <https://nccih.nih.gov/health/ginger> [accessed 6/27/16].
34. Post-White J, Kinney ME, Savik K, Gau JB, Wilcox C, Lerner I. Therapeutic massage and healing touch improve symptoms in cancer. *Integr Cancer Ther*. 2003;2: 332-344.
35. FitzHenry F, Wells N, Slater V, Dietrich MS, Wisawatapnimit P, Chakravarthy AB. A randomized placebo-controlled pilot study of the impact of healing touch on fatigue in breast cancer patients undergoing radiation therapy. *Integrative cancer therapies*. 2014;13: 105-113.
36. National Cancer Institute. Topics in Integrative, Alternative, and Complementary Therapies (PDQ)-Patient Version. Available from URL: https://www.cancer.gov/about-cancer/treatment/cam/patient/cam-topics-pdq-link/34_toc [accessed 11/29/16].
37. National Cancer Institute. Lymphedema (PDQ) Health Professional Version. Available from URL: <http://www.cancer.gov/about-cancer/treatment/side-effects/lymphedema/lymphedema-hp-pdq> [accessed 6/27/16].

38. National Center for Complementary and Integrative Health. Massage therapy for health purposes: what you need to know. Available from URL: <https://nccih.nih.gov/health/massage/massageintroduction.htm> [accessed 6/18/15].
39. Geccedi RA. Massage therapy for patients with cancer. *Clin J Oncol Nurs*. 2002;6: 52-54.
40. National Cancer Institute. Pain-for health professionals (PDQ): Physical, Integrative, Cognitive-Behavioral and Psychosocial Interventions. Available from URL: <http://www.cancer.gov/about-cancer/treatment/side-effects/pain/pain-hp-pdq - section/ 231> [accessed 6/18/15].
41. Walsh R, Shapiro SL. The meeting of meditative disciplines and Western psychology - A mutually enriching dialogue. *American Psychologist*. 2006;61: 227-239.
42. Bond K, Ospina MB, Hooton N, et al. Defining a Complex Intervention: The Development of Demarcation Criteria for "Meditation". *Psychology of Religion and Spirituality*. 2009;1: 129-137.
43. Kabat-Zinn JNH, T. *Full Catastrophe Living*. New York, New York: Bantam Dell, 1990.
44. Shennan C, Payne S, Fenlon D. What is the evidence for the use of mindfulness-based interventions in cancer care? A review. *Psychooncology*. 2011;20: 681-697.
45. Matchim Y, Armer JM, Stewart BR. Mindfulness-based stress reduction among breast cancer survivors: a literature review and discussion. *Oncol Nurs Forum*. 2011;38: E61-71.
46. Zainal NZ, Booth S, Huppert FA. The efficacy of mindfulness-based stress reduction on mental health of breast cancer patients: a meta-analysis. *Psychooncology*. 2013;22: 1457-1465.
47. Cramer H, Lauche R, Paul A, Dobos G. Mindfulness-based stress reduction for breast cancer-a systematic review and meta-analysis. *Curr Oncol*. 2012;19: e343-352.
48. Marvibaigi M, Supriyanto E, Amini N, Abdul Majid FA, Jaganathan SK. Preclinical and clinical effects of mistletoe against breast cancer. *Biomed Res Int*. 2014;2014: 785479.
49. Bussing A. *Mistletoe: The Genus Viscum* The Netherlands: CRC Press, 2000.
50. Horneber MA, Bueschel G, Huber R, Linde K, Rostock M. Mistletoe therapy in oncology. *Cochrane Database Syst Rev*. 2008: CD003297.

51. American Music Therapy Association. What is music therapy? Available from URL: <http://www.musictherapy.org/about/musictherapy/> [accessed 6/22/15].
52. Nilsson U. The anxiety- and pain-reducing effects of music interventions: a systematic review. *AORN J*. 2008;87: 780-807.
53. Guetin S, Charras K, Berard A, et al. An overview of the use of music therapy in the context of Alzheimer's disease: a report of a French expert group. *Dementia (London)*. 2013;12: 619-634.
54. National Center for Complementary and Integrative Health. Reflexology. Available from URL: <https://nccih.nih.gov/health/reflexology> [accessed 6/27/16].
55. National Center for Complementary and Integrative Health. Relaxation Techniques for Health: What you need to know. Available from URL: <https://nccih.nih.gov/health/stress/relaxation.htm> [accessed 6/18/15].
56. National Center for Complementary and Integrative Health. Relaxation Techniques for Health: What you need to know. Available from URL: <https://nccih.nih.gov/health/stress/relaxation.htm> [accessed 3/10/15].
57. National Center for Complementary and Integrative Health. Tai Chi and Qi Gong: In Depth. Available from URL: <https://nccih.nih.gov/health/taichi/introduction.htm> [accessed 6/27/16].
58. National Center for Complementary and Integrative Health. Stress. Available from URL: <https://nccih.nih.gov/health/stress> [accessed 6/18/15].
59. National Cancer Institute. Depression- for health professionals (PDQ). Available from URL: http://www.cancer.gov/about-cancer/coping/feelings/depression-hp-pdq/-link/_74 [accessed 8/10/15].
60. National Cancer Institute. Cancer-related post-traumatic stress- for health professionals (PDQ). Available from URL: http://www.cancer.gov/about-cancer/coping/survivorship/new-normal/ptsd-hp-pdq/-link/stoc_h2_4 [accessed 8/8/15].
61. Antoni MH, Lechner SC, Kazi A, et al. How stress management improves quality of life after treatment for breast cancer. *J Consult Clin Psychol*. 2006;74: 1143-1152.

62. Antoni MH, Lehman JM, Kilbourn KM, et al. Cognitive-behavioral stress management intervention decreases the prevalence of depression and enhances benefit finding among women under treatment for early-stage breast cancer. *Health Psychol.* 2001;20: 20-32.
63. National Center for Complementary and Integrative Health. Yoga. Available from URL: <https://nccih.nih.gov/health/yoga> [accessed 6/18/15].
64. Iyengar B. *Light on pranayama: the yogic art of breathing.* Crossroad Publishing Company, 2005.
65. Andersen BL, DeRubeis RJ, Berman BS, et al. Screening, assessment, and care of anxiety and depressive symptoms in adults with cancer: an American Society of Clinical Oncology guideline adaptation. *J Clin Oncol.* 2014;32: 1605-1619.
66. Bower JE, Bak K, Berger A, et al. Screening, assessment, and management of fatigue in adult survivors of cancer: an American Society of Clinical oncology clinical practice guideline adaptation. *J Clin Oncol.* 2014;32: 1840-1850.
67. Hershman DL, Lacchetti C, Dworkin RH, et al. Prevention and management of chemotherapy-induced peripheral neuropathy in survivors of adult cancers: American Society of Clinical Oncology clinical practice guideline. *J Clin Oncol.* 2014;32: 1941-1967.
68. Runowicz CD, Leach CR, Henry NL, et al. American Cancer Society/American Society of Clinical Oncology Breast Cancer Survivorship Care Guideline. *J Clin Oncol.* 2016;34: 611-635.
69. Jordan K, Jahn F, Aapro M. Recent developments in the prevention of chemotherapy-induced nausea and vomiting (CINV): a comprehensive review. *Ann Oncol.* 2015;26: 1081-1090.
70. Berger AM, Abernethy AP, Atkinson A, et al. NCCN Clinical Practice Guidelines Cancer-related fatigue. *J Natl Compr Canc Netw.* 2010;8: 904-931.
71. Hesketh PJ, Bohlke K, Lyman GH, et al. Antiemetics: American Society of Clinical Oncology Focused Guideline Update. *J Clin Oncol.* 2016;34: 381-386.
72. Carlson LE, Doll R, Stephen J, et al. Randomized controlled trial of Mindfulness-based cancer recovery versus supportive expressive group therapy for distressed survivors of breast cancer. *J Clin Oncol.* 2013;31: 3119-3126.

73. Crane-Okada R, Kiger H, Sugerma F, et al. Mindful movement program for older breast cancer survivors: a pilot study. *Cancer Nurs*. 2012;35: E1-13.
74. Kim YH, Kim HJ, Ahn SD, Seo YJ, Kim SH. Effects of meditation on anxiety, depression, fatigue, and quality of life of women undergoing radiation therapy for breast cancer. *Complement Ther Med*. 2013;21: 379-387.
75. Lengacher CA, Johnson-Mallard V, Post-White J, et al. Randomized controlled trial of mindfulness-based stress reduction (MBSR) for survivors of breast cancer. *Psycho-Oncology*. 2009;18: 1261-1272.
76. Wurtzen H, Dalton SO, Elsass P, et al. Mindfulness significantly reduces self-reported levels of anxiety and depression: results of a randomised controlled trial among 336 Danish women treated for stage I-III breast cancer. *Eur J Cancer*. 2013;49: 1365-1373.
77. Chandwani KD, Perkins G, Nagendra HR, et al. Randomized, controlled trial of yoga in women with breast cancer undergoing radiotherapy. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2014;32: 1058-1065.
78. Kiecolt-Glaser JK, Bennett JM, Andridge R, et al. Yoga's impact on inflammation, mood, and fatigue in breast cancer survivors: a randomized controlled trial. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2014;32: 1040-1049.
79. Taso CJ, Lin HS, Lin WL, Chen SM, Huang WT, Chen SW. The effect of yoga exercise on improving depression, anxiety, and fatigue in women with breast cancer: a randomized controlled trial. *The journal of nursing research: JNR*. 2014;22: 155-164.
80. Rao RM, Raghuram N, Nagendra HR, et al. Effects of an integrated Yoga Program on Self-reported Depression Scores in Breast Cancer Patients Undergoing Conventional Treatment: A Randomized Controlled Trial. *Indian J Palliat Care*. 2015;21: 174-181.
81. Siedentopf F, Utz-Billing I, Gairing S, Schoenegg W, Kentenich H, Kollak I. Yoga for Patients with Early Breast Cancer and its Impact on Quality of Life - a Randomized Controlled Trial. *Geburtshilfe und Frauenheilkunde*. 2013;73: 311-317.

82. Cramer H, Rabsilber S, Lauche R, Kummel S, Dobos G. Yoga and meditation for menopausal symptoms in breast cancer survivors - A randomized controlled trial. *Cancer*. 2015;121: 2175-2184.
83. Montgomery GH, David D, Kangas M, et al. Randomized controlled trial of a cognitive-behavioral therapy plus hypnosis intervention to control fatigue in patients undergoing radiotherapy for breast cancer. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2014;32: 557-563.
84. Montgomery GH, Kangas M, David D, et al. Fatigue during breast cancer radiotherapy: an initial randomized study of cognitive-behavioral therapy plus hypnosis. *Health Psychol*. 2009;28: 317-322.
85. National Cancer Institute. Feelings and Cancer. Available from URL: <http://www.cancer.gov/cancertopics/coping/feelings - stress> [accessed 6/18/15].
86. Brintzenhofe-Szoc KM, Levin TT, Li YL, Kissane DW, Zabora JR. Mixed Anxiety/Depression Symptoms in a Large Cancer Cohort: Prevalence by Cancer Type. *Psychosomatics*. 2009;50: 383-391.
87. Eskelinen M, Ollonen P. Assessment of General Anxiety in Patients with Breast Disease and Breast Cancer Using the Spielberger STAI Self Evaluation Test: A Prospective Case-Control Study in Finland. *Anticancer Research*. 2011;31: 1801-1806.
88. Ollonen P, Lehtonen J, Eskelinen M. Anxiety, depression, and the history of psychiatric symptoms in patients with breast disease: A prospective case-control study in Kuopio, Finland. *Anticancer Research*. 2005;25: 2527-2533.
89. Institute of Medicine. Cancer care for the whole patient: meeting psychosocial health needs. . Washington, DC: The National Academic Press, 2007.
90. Valentine A. Mood disorders. MD Anderson manual of psychosocial oncology. New York, NY: McGraw-Hill, Inc., 2011:271-288.
91. Rashid A. Anxiety in cancer patients. MD Anderson manual of psychosocial oncology. New York, NY: McGraw-Hill, Inc., 2011:271-288.
92. Goyal M, Singh S, Sibinga EM, et al. Meditation programs for psychological stress and well-being: a systematic review and meta-analysis. *JAMA Intern Med*. 2014;174: 357-368.

93. Miller JJ, Fletcher K, Kabat-Zinn J. Three-year follow-up and clinical implications of a mindfulness meditation-based stress reduction intervention in the treatment of anxiety disorders. *General hospital psychiatry*. 1995;17: 192-200.
94. Kabat-Zinn J, Massion AO, Kristeller J, et al. Effectiveness of a meditation-based stress reduction program in the treatment of anxiety disorders. *Am J Psychiatry*. 1992;149: 936-943.
95. Hoge EA, Bui E, Marques L, et al. Randomized controlled trial of mindfulness meditation for generalized anxiety disorder: effects on anxiety and stress reactivity. *The Journal of clinical psychiatry*. 2013;74: 786.
96. Speca M, Carlson LE, Goodey E, Angen M. A randomized, wait-list controlled clinical trial: the effect of a mindfulness meditation-based stress reduction program on mood and symptoms of stress in cancer outpatients. *Psychosomatic medicine*. 2000;62: 613-622.
97. Carlson LE, Ursuliak Z, Goodey E, Angen M, Speca M. The effects of a mindfulness meditation-based stress reduction program on mood and symptoms of stress in cancer outpatients: 6-month follow-up. *Supportive care in Cancer*. 2001;9: 112-123.
98. Piet J, Würtzen H, Zachariae R. The effect of mindfulness-based therapy on symptoms of anxiety and depression in adult cancer patients and survivors: A systematic review and meta-analysis. *Journal of consulting and clinical psychology*. 2012;80: 1007.
99. Zainal NZ, Booth S, Huppert FA. The efficacy of mindfulness-based stress reduction on mental health of breast cancer patients: a meta-analysis. *Psycho-Oncology*. 2013;22: 1457-1465.
100. Musial F, Büssing A, Heusser P, Choi K-E, Ostermann T. Mindfulness-based stress reduction for integrative cancer care: a summary of evidence 2011.
101. Rocha T. The Dark Knight of the Soul. *The Atlantic*. 2014;25.
102. Lomas T, Cartwright T, Edginton T, Ridge D. A qualitative analysis of experiential challenges associated with meditation practice. *Mindfulness*. 2014: 1-13.

103. Lengacher CA, Kip KE, Barta M, et al. A pilot study evaluating the effect of mindfulness-based stress reduction on psychological status, physical status, salivary cortisol, and interleukin-6 among advanced-stage cancer patients and their caregivers. *Journal of Holistic Nursing*. 2012;30: 170-185.
104. Tamagawa R, Speca M, Stephen J, Pickering B, Lawlor-Savage L, Carlson LE. Predictors and Effects of Class Attendance and Home Practice of Yoga and Meditation Among Breast Cancer Survivors in a Mindfulness-Based Cancer Recovery (MBCR) Program. *Mindfulness*. 2015: 1-10.
105. Zernicke KA, Campbell TS, Speca M, McCabe-Ruff K, Flowers S, Carlson LE. A Randomized Wait-List Controlled Trial of Feasibility and Efficacy of an Online Mindfulness-Based Cancer Recovery Program: The eTherapy for Cancer Applying Mindfulness Trial. *Psychosomatic medicine*. 2014;76: 257-267.
106. Binns-Turner PG, Wilson LL, Pryor ER, Boyd GL, Prickett CA. Perioperative music and its effects on anxiety, hemodynamics, and pain in women undergoing mastectomy. *AANA Journal*. 2011;79: S21-27.
107. Bulfone T, Quattrin R, Zanotti R, Regattin L, Brusaferrero S. Effectiveness of music therapy for anxiety reduction in women with breast cancer in chemotherapy treatment. *Holist Nurs Pract*. 2009;23: 238-242.
108. Hanser SB, Bauer-Wu S, Kubicek L, et al. Effects of a music therapy intervention on quality of life and distress in women with metastatic breast cancer. *J Soc Integr Oncol*. 2006;4: 116-124.
109. Li XM, Zhou KN, Yan H, Wang DL, Zhang YP. Effects of music therapy on anxiety of patients with breast cancer after radical mastectomy: a randomized clinical trial. *J Adv Nurs*. 2012;68: 1145-1155.
110. Zhou K, Li X, Li J, et al. A clinical randomized controlled trial of music therapy and progressive muscle relaxation training in female breast cancer patients after radical mastectomy: results on depression, anxiety and length of hospital stay. *European journal of oncology nursing : the official journal of European Oncology Nursing Society*. 2015;19: 54-59.

111. Boehm K, Cramer H, Staroszyński T, Ostermann T. Arts therapies for anxiety, depression, and quality of life in breast cancer patients: a systematic review and meta-analysis. *Evid Based Complement Alternat Med.* 2014;2014: 103297.
112. Thyme KE, Sundin EC, Wiberg B, Oster I, Astrom S, Lindh J. Individual brief art therapy can be helpful for women with breast cancer: a randomized controlled clinical study. *Palliat Support Care.* 2009;7: 87-95.
113. Zavotsky KE, Banavage A, James P, Easter K, Pontieri-Lewis V, Lutwin L. The effects of music on pain and anxiety during screening mammography. *Clin J Oncol Nurs.* 2014;18: E45-49.
114. Garssen B, Boomsma MF, Meezenbroek Ede J, et al. Stress management training for breast cancer surgery patients. *Psychooncology.* 2013;22: 572-580.
115. Jacobsen PB, Phillips KM, Jim HS, et al. Effects of self-directed stress management training and home-based exercise on quality of life in cancer patients receiving chemotherapy: a randomized controlled trial. *Psychooncology.* 2013;22: 1229-1235.
116. Aguado Loi CX, Taylor TR, McMillan S, et al. Use and helpfulness of self-administered stress management therapy in patients undergoing cancer chemotherapy in community clinical settings. *J Psychosoc Oncol.* 2012;30: 57-80.
117. Phillips KM, Antoni MH, Lechner SC, et al. Stress management intervention reduces serum cortisol and increases relaxation during treatment for nonmetastatic breast cancer. *Psychosom Med.* 2008;70: 1044-1049.
118. Trijsburg RW, van Knippenberg FC, Rijpma SE. Effects of psychological treatment on cancer patients: a critical review. *Psychosom Med.* 1992;54: 489-517.
119. Banerjee B, Vadiraj H, Ram A, et al. Effects of an integrated yoga program in modulating psychological stress and radiation-induced genotoxic stress in breast cancer patients undergoing radiotherapy. *Integrative Cancer Therapies.* 2007;6: 242-250.

120. Vadiraja H, Raghavendra RM, Nagarathna R, et al. Effects of a yoga program on cortisol rhythm and mood states in early breast cancer patients undergoing adjuvant radiotherapy: a randomized controlled trial. *Integrative Cancer Therapies*. 2009.
121. Raghavendra R, Nagarathna R, Nagendra H, et al. Effects of an integrated yoga programme on chemotherapy-induced nausea and emesis in breast cancer patients. *European Journal of Cancer Care*. 2007;16: 462-474.
122. Rao MR, Raghuram N, Nagendra H, et al. Anxiolytic effects of a yoga program in early breast cancer patients undergoing conventional treatment: a randomized controlled trial. *Complementary therapies in medicine*. 2009;17: 1-8.
123. Bower JE, Garet D, Sternlieb B, et al. Yoga for persistent fatigue in breast cancer survivors: a randomized controlled trial. *Cancer*. 2012;118: 3766-3775.
124. Dhruva A, Miaskowski C, Abrams D, et al. Yoga breathing for cancer chemotherapy-associated symptoms and quality of life: results of a pilot randomized controlled trial. *J Altern Complement Med*. 2012;18: 473-479.
125. Pruthi S, Stan DL, Jenkins SM, et al. A Randomized controlled pilot study assessing feasibility and impact of yoga practice on quality of life, mood, and perceived stress in women with newly diagnosed breast cancer. *Global Advances In Health and Medicine*. 2012;1: 30-35.
126. Chandwani KD, Thornton B, Perkins GH, et al. Yoga improves quality of life and benefit finding in women undergoing radiotherapy for breast cancer. *J Soc Integr Oncol*. 2010;8: 43-55.
127. Smith KB, Pukall CF. An evidence-based review of yoga as a complementary intervention for patients with cancer. *Psycho-Oncology*. 2009;18: 465-475.
128. Culos-Reed SN, Mackenzie MJ, Sohl SJ, Jesse MT, Zahavich ANR, Danhauer SC. Yoga & cancer interventions: a review of the clinical significance of patient reported outcomes for cancer survivors. *Evidence-based Complementary and Alternative Medicine*. 2012;2012.

129. Buffart LM, Van Uffelen JG, Riphagen II, et al. Physical and psychosocial benefits of yoga in cancer patients and survivors, a systematic review and meta-analysis of randomized controlled trials. *BMC cancer*. 2012;12: 559.
130. Sadjia J, Mills PJ. Effects of yoga interventions on fatigue in cancer patients and survivors: a systematic review of randomized controlled trials. *EXPLORE: The Journal of Science and Healing*. 2013;9: 232-243.
131. Lin K-Y, Hu Y-T, Chang K-J, Lin H-F, Tsao J-Y. Effects of yoga on psychological health, quality of life, and physical health of patients with cancer: a meta-analysis. *Evidence-Based Complementary and Alternative Medicine*. 2011;2011.
132. Cramer H, Lange S, Klose P, Paul A, Dobos G. Yoga for breast cancer patients and survivors: a systematic review and meta-analysis. *BMC cancer*. 2012;12: 412.
133. Kirkwood G, Rampes H, Tuffrey V, Richardson J, Pilkington K. Yoga for anxiety: a systematic review of the research evidence. *British Journal of Sports Medicine*. 2005;39: 884-891.
134. Woolery A, Myers H, Sternlieb B, Zeltzer L. A yoga intervention for young adults with elevated symptoms of depression. *Alternative therapies in health and medicine*. 2004;10: 60-63.
135. Cramer H, Krucoff C, Dobos G. Adverse events associated with yoga: a systematic review of published case reports and case series. *PloS one*. 2013;8: e75515.
136. Bao T, Cai L, Snyder C, et al. Patient-reported outcomes in women with breast cancer enrolled in a dual-center, double-blind, randomized controlled trial assessing the effect of acupuncture in reducing aromatase inhibitor-induced musculoskeletal symptoms. *Cancer*. 2014;120: 381-389.
137. Mao JJ, Farrar JT, Bruner D, et al. Electroacupuncture for fatigue, sleep, and psychological distress in breast cancer patients with aromatase inhibitor-related arthralgia: A randomized trial. *Cancer*. 2014;120: 3744-3751.
138. Molassiotis A, Bardy J, Finnegan-John J, et al. Acupuncture for cancer-related fatigue in patients with breast cancer: a pragmatic randomized controlled trial. *J Clin Oncol*. 2012;30: 4470-4476.

139. Billhult A, Bergbom I, Stener-Victorin E. Massage relieves nausea in women with breast cancer who are undergoing chemotherapy. *J Altern Complement Med.* 2007;13: 53-57.
140. Hernandez-Reif M, Ironson G, Field T, et al. Breast cancer patients have improved immune and neuroendocrine functions following massage therapy. *J Psychosom Res.* 2004;57: 45-52.
141. Listing M, Krohn M, Liezmann C, et al. The efficacy of classical massage on stress perception and cortisol following primary treatment of breast cancer. *Arch Womens Ment Health.* 2010;13: 165-173.
142. Wilkinson SM, Love SB, Westcombe AM, et al. Effectiveness of aromatherapy massage in the management of anxiety and depression in patients with cancer: a multicenter randomized controlled trial. *J Clin Oncol.* 2007;25: 532-539.
143. Hilderley M, Holt M. A pilot randomized trial assessing the effects of autogenic training in early stage cancer patients in relation to psychological status and immune system responses. *Eur J Oncol Nurs.* 2004;8: 61-65.
144. Kovacic T, Kovacic M. Impact of relaxation training according to Yoga In Daily Life(R) system on perceived stress after breast cancer surgery. *Integr Cancer Ther.* 2011;10: 16-26.
145. Kovacic T, Zagoricnik M, Kovacic M. Impact of relaxation training according to the Yoga In Daily Life(R) system on anxiety after breast cancer surgery. *J Complement Integr Med.* 2013;10: 153-164.
146. Molassiotis A, Yung HP, Yam BM, Chan FY, Mok TS. The effectiveness of progressive muscle relaxation training in managing chemotherapy-induced nausea and vomiting in Chinese breast cancer patients: a randomised controlled trial. *Support Care Cancer.* 2002;10: 237-246.
147. Nunes DF, Rodriguez AL, da Silva Hoffmann F, et al. Relaxation and guided imagery program in patients with breast cancer undergoing radiotherapy is not associated with neuroimmunomodulatory effects. *J Psychosom Res.* 2007;63: 647-655.
148. Milbury K, Chaoul A, Biegler K, et al. Tibetan sound meditation for cognitive dysfunction: results of a randomized controlled pilot trial. *Psycho-Oncology.* 2013;22: 2354-2363.

149. Hoffman CJ, Ersser SJ, Hopkinson JB, Nicholls PG, Harrington JE, Thomas PW. Effectiveness of mindfulness-based stress reduction in mood, breast- and endocrine-related quality of life, and well-being in stage 0 to III breast cancer: a randomized, controlled trial. *J Clin Oncol*. 2012;30: 1335-1342.
150. Nidich SI, Fields JZ, Rainforth MV, et al. A randomized controlled trial of the effects of transcendental meditation on quality of life in older breast cancer patients. *Integr Cancer Ther*. 2009;8: 228-234.
151. Henderson VP, Massion AO, Clemow L, Hurley TG, Druker S, Hebert JR. A randomized controlled trial of mindfulness-based stress reduction for women with early-stage breast cancer receiving radiotherapy. *Integr Cancer Ther*. 2013;12: 404-413.
152. Dodds SE, Pace TWW, Bell ML, et al. Feasibility of Cognitively-Based Compassion Training (CBCT) for breast cancer survivors: a randomized, wait list controlled pilot study. *Supportive Care in Cancer*. 2015;23: 3599-3608.
153. Teasdale JD, Segal ZV, Williams JMG, Ridgeway VA, Soulsby JM, Lau MA. Prevention of relapse/recurrence in major depression by mindfulness-based cognitive therapy. *Journal of consulting and clinical psychology*. 2000;68: 615.
154. Segal ZV, Teasdale JD, Williams JM, Gemar MC. The mindfulness-based cognitive therapy adherence scale: inter-rater reliability, adherence to protocol and treatment distinctiveness. *Clinical Psychology & Psychotherapy*. 2002;9: 131-138.
155. Hofmann SG, Sawyer AT, Witt AA, Oh D. The effect of mindfulness-based therapy on anxiety and depression: A meta-analytic review. *Journal of consulting and clinical psychology*. 2010;78: 169.
156. Yoo HJ, Ahn SH, Kim SB, Kim WK, Han OS. Efficacy of progressive muscle relaxation training and guided imagery in reducing chemotherapy side effects in patients with breast cancer and in improving their quality of life. *Support Care Cancer*. 2005;13: 826-833.
157. Walker LG, Walker MB, Ogston K, et al. Psychological, clinical and pathological effects of relaxation training and guided imagery during primary chemotherapy. *Br J Cancer*. 1999;80: 262-268.

158. Gudenkauf LM, Antoni MH, Stagl JM, et al. Brief cognitive-behavioral and relaxation training interventions for breast cancer: A randomized controlled trial. *Journal of consulting and clinical psychology*. 2015;83: 677-688.
159. Roffe L, Schmidt K, Ernst E. A systematic review of guided imagery as an adjuvant cancer therapy. *Psychooncology*. 2005;14: 607-617.
160. Demiralp M, Oflaz F, Komurcu S. Effects of relaxation training on sleep quality and fatigue in patients with breast cancer undergoing adjuvant chemotherapy. *J Clin Nurs*. 2010;19: 1073-1083.
161. Goerling U, Jaeger C, Walz A, Stickel A, Mangler M, van der Meer E. The efficacy of short-term psycho-oncological interventions for women with gynaecological cancer: a randomized study. *Oncology*. 2014;87: 114-124.
162. Park ER, Traeger L, Willett J, et al. A relaxation response training for women undergoing breast biopsy: exploring integrated care. *Breast*. 2013;22: 799-805.
163. Song QH, Xu RM, Zhang QH, Ma M, Zhao XP. Relaxation training during chemotherapy for breast cancer improves mental health and lessens adverse events. *Int J Clin Exp Med*. 2013;6: 979-984.
164. Yilmaz SG, Arslan S. Effects of progressive relaxation exercises on anxiety and comfort of Turkish breast cancer patients receiving chemotherapy. *Asian Pac J Cancer Prev*. 2015;16: 217-220.
165. Tanyi RA, Berk LS, Lee JW, Boyd K, Arechiga A. The effects of a psychoneuroimmunology (PNI) based lifestyle intervention in modifying the progression of depression in clinically depressed adults. *Int J Psychiatry Med*. 2011;42: 151-166.
166. Klainin-Yobas P, Oo WN, Suzanne Yew PY, Lau Y. Effects of relaxation interventions on depression and anxiety among older adults: a systematic review. *Aging Ment Health*. 2015: 1-13.
167. Younge JO, Gotink RA, Baena CP, Roos-Hesselink JW, Hunink MM. Mind-body practices for patients with cardiac disease: a systematic review and meta-analysis. *Eur J Prev Cardiol*. 2014.
168. Culos-Reed SN, Carlson LE, Daroux LM, Hatley-Aldous S. A pilot study of yoga for breast cancer survivors: physical and psychological benefits. *Psychooncology*. 2006;15: 891-897.

169. Moadel AB, Shah C, Wylie-Rosett J, et al. Randomized controlled trial of yoga among a multiethnic sample of breast cancer patients: effects on quality of life. *J Clin Oncol*. 2007;25: 4387-4395.
170. Danhauer SC, Mihalko SL, Russell GB, et al. Restorative yoga for women with breast cancer: findings from a randomized pilot study. *Psychooncology*. 2009;18: 360-368.
171. Vadiraja H, Rao MR, Nagarathna R, et al. Effects of yoga program on quality of life and affect in early breast cancer patients undergoing adjuvant radiotherapy: a randomized controlled trial. *Complementary therapies in medicine*. 2009;17: 274-280.
172. Pilkington K, Kirkwood G, Rampes H, Richardson J. Yoga for depression: the research evidence. *Journal of affective disorders*. 2005;89: 13-24.
173. Shapiro D, Cook IA, Davydov DM, Ottaviani C, Leuchter AF, Abrams M. Yoga as a complementary treatment of depression: effects of traits and moods on treatment outcome. *Evidence-based complementary and alternative medicine*. 2007;4: 493-502.
174. Uebelacker LA, Epstein-Lubow G, Gaudiano BA, Tremont G, Battle CL, Miller IW. Hatha yoga for depression: critical review of the evidence for efficacy, plausible mechanisms of action, and directions for future research. *Journal of Psychiatric Practice®*. 2010;16: 22-33.
175. Krohn M, Listing M, Tjahjono G, et al. Depression, mood, stress, and Th1/Th2 immune balance in primary breast cancer patients undergoing classical massage therapy. *Support Care Cancer*. 2011;19: 1303-1311.
176. Listing M, Reissauer A, Krohn M, et al. Massage therapy reduces physical discomfort and improves mood disturbances in women with breast cancer. *Psychooncology*. 2009;18: 1290-1299.
177. Fernandez-Lao C, Cantarero-Villanueva I, Diaz-Rodriguez L, Cuesta-Vargas AI, Fernandez-Delas-Penas C, Arroyo-Morales M. Attitudes towards massage modify effects of manual therapy in breast cancer survivors: a randomised clinical trial with crossover design. *European Journal of Cancer Care*. 2012;21: 233-241.
178. Pan YQ, Yang KH, Wang YL, Zhang LP, Liang HQ. Massage interventions and treatment-related side effects of breast cancer: a systematic review and meta-analysis. *Int J Clin Oncol*. 2014;19: 829-841.

179. Lee MS, Lee EN, Ernst E. Massage therapy for breast cancer patients: a systematic review. *Ann Oncol.* 2011;22: 1459-1461.
180. Kutner JS, Smith MC, Corbin L, et al. Massage therapy versus simple touch to improve pain and mood in patients with advanced cancer: a randomized trial. *Ann Intern Med.* 2008;149: 369-379.
181. Ernst E. The safety of massage therapy. *Rheumatology (Oxford).* 2003;42: 1101-1106.
182. Wilkinson S, Barnes K, Storey L. Massage for symptom relief in patients with cancer: systematic review. *J Adv Nurs.* 2008;63: 430-439.
183. Coelho HF, Boddy K, Ernst E. Massage therapy for the treatment of depression: a systematic review. *Int J Clin Pract.* 2008;62: 325-333.
184. Hou WH, Chiang PT, Hsu TY, Chiu SY, Yen YC. Treatment effects of massage therapy in depressed people: a meta-analysis. *J Clin Psychiatry.* 2010;71: 894-901.
185. Burns DS. The effect of the bonny method of guided imagery and music on the mood and life quality of cancer patients. *J Music Ther.* 2001;38: 51-65.
186. Zhou KN, Li XM, Yan H, Dang SN, Wang DL. Effects of music therapy on depression and duration of hospital stay of breast cancer patients after radical mastectomy. *Chin Med J (Engl).* 2011;124: 2321-2327.
187. Maratos AS, Gold C, Wang X, Crawford MJ. Music therapy for depression. *Cochrane Database Syst Rev.* 2008: CD004517.
188. Nedstrand E, Wijma K, Wyon Y, Hammar M. Vasomotor symptoms decrease in women with breast cancer randomized to treatment with applied relaxation or electro-acupuncture: A preliminary study. *Climacteric.* 2005;8: 243-250.
189. Walker EM, Rodriguez AI, Kohn B, et al. Acupuncture versus venlafaxine for the management of vasomotor symptoms in patients with hormone receptor-positive breast cancer: a randomized controlled trial. *J Clin Oncol.* 2010;28: 634-640.

190. Lechner SC, Whitehead NE, Vargas S, et al. Does a community-based stress management intervention affect psychological adaptation among underserved black breast cancer survivors? *Journal of the National Cancer Institute. Monographs.* 2014;2014: 315-322.
191. Stagl JM, Bouchard LC, Lechner SC, et al. Long-term psychological benefits of cognitive-behavioral stress management for women with breast cancer: 11-year follow-up of a randomized controlled trial. *Cancer.* 2015;121: 1873-1881.
192. National Cancer Institute. Fatigue (PDQ). Available from URL: <http://www.cancer.gov/cancertopics/pdq/supportivecare/fatigue/HealthProfessional> [accessed 6/18/15].
193. Minton O, Stone P. How common is fatigue in disease-free breast cancer survivors? A systematic review of the literature. *Breast Cancer Research and Treatment.* 2008;112: 5-13.
194. Bower JE, Ganz PA, Desmond KA, et al. Fatigue in long-term breast carcinoma survivors - A longitudinal investigation. *Cancer.* 2006;106: 751-758.
195. Stone P, Richardson A, Ream E, et al. Cancer-related fatigue: Inevitable, unimportant and untreatable? Results of a multi-centre patient survey. *Annals of Oncology.* 2000;11: 971-975.
196. Barton DL, Liu H, Dakhil SR, et al. Wisconsin Ginseng (*Panax quinquefolius*) to improve cancer-related fatigue: a randomized, double-blind trial, N07C2. *J Natl Cancer Inst.* 2013;105: 1230-1238.
197. Barton DL, Soori GS, Bauer BA, et al. Pilot study of *Panax quinquefolius* (American ginseng) to improve cancer-related fatigue: a randomized, double-blind, dose-finding evaluation: NCCTG trial N03CA. *Support Care Cancer.* 2010;18: 179-187.
198. Hershman DL, Unger JM, Crew KD, et al. Randomized double-blind placebo-controlled trial of acetyl-L-carnitine for the prevention of taxane-induced neuropathy in women undergoing adjuvant breast cancer therapy. *J Clin Oncol.* 2013;31: 2627-2633.
199. da Costa Miranda V, Trufelli DC, Santos J, et al. Effectiveness of guarana (*Paullinia cupana*) for postradiation fatigue and depression: Results of a pilot double-blind randomized study. *J Altern Complement Med.* 2009;15: 431-433.

200. de Oliveira Campos MP, Riechelmann R, Martins LC, Hassan BJ, Casa FBA, Giglio AD. Guarana (Paullinia cupana) improves fatigue in breast cancer patients undergoing systemic chemotherapy. *J Altern Complement Med.* 2011;17: 505-512.
201. Deng G, Chan Y, Sjoberg D, et al. Acupuncture for the treatment of post-chemotherapy chronic fatigue: a randomized, blinded, sham-controlled trial. *Support Care Cancer.* 2013;21: 1735-1741.
202. Molassiotis A, Bardy J, Finnegan-John J, et al. A randomized, controlled trial of acupuncture self-needling as maintenance therapy for cancer-related fatigue after therapist-delivered acupuncture. *Ann Oncol.* 2013;24: 1645-1652.
203. Smith C, Carmady B, Thornton C, Perz J, Ussher JM. The effect of acupuncture on post-cancer fatigue and well-being for women recovering from breast cancer: a pilot randomised controlled trial. *Acupunct Med.* 2013;31: 9-15.
204. Banasik J, Williams H, Haberman M, Blank SE, Bendel R. Effect of Iyengar yoga practice on fatigue and diurnal salivary cortisol concentration in breast cancer survivors. *Journal of the American Academy of Nurse Practitioners.* 2011;23: 135-142.
205. Littman AJ, Bertram LC, Ceballos R, et al. Randomized controlled pilot trial of yoga in overweight and obese breast cancer survivors: effects on quality of life and anthropometric measures. *Support Care Cancer.* 2012;20: 267-277.
206. Victorson D CD, Wagner L, Kramer L, Smith ML. Measuring quality of life in cancer survivors. . In: Feuerstein M e, editor. *Handbook of cancer survivorship.* New York: Springer, 2007:79-110.
207. Ganz PA, Rowland JH, Desmond K, Meyerowitz BE, Wyatt GE. Life after breast cancer: understanding women's health-related quality of life and sexual functioning. *J Clin Oncol.* 1998;16: 501-514.
208. National Cancer Institute. *Managing Physical Effects.* Available from URL: <http://www.cancer.gov/cancertopics/coping/physicaleffects> [accessed 6/18/15].

209. Ganz PA, Guadagnoli E, Landrum MB, Lash TL, Rakowski W, Silliman RA. Breast cancer in older women: Quality of life and psychosocial adjustment in the 15 months after diagnosis. *Journal of Clinical Oncology*. 2003;21: 4027-4033.
210. Weitzner MA, Meyers CA, Stuebing KK, Saleeba AK. Relationship between quality of life and mood in long-term survivors of breast cancer treated with mastectomy. *Supportive Care in Cancer*. 1997;5: 241-248.
211. Montazeri A, Vahdaninia M, Harirchi I, Ebrahimi M, Khaleghi F, Jarvandi S. Quality of life in patients with breast cancer before and after diagnosis: an eighteen months follow-up study. *Bmc Cancer*. 2008;8.
212. Henderson VP, Clemow L, Massion AO, Hurley TG, Druker S, Hebert JR. The effects of mindfulness-based stress reduction on psychosocial outcomes and quality of life in early-stage breast cancer patients: a randomized trial. *Breast Cancer Res Treat*. 2012;131: 99-109.
213. Foley E, Baillie A, Huxter M, Price M, Sinclair E. Mindfulness-based cognitive therapy for individuals whose lives have been affected by cancer: a randomized controlled trial. *Journal of consulting and clinical psychology*. 2010;78: 72.
214. Kieviet-Stijnen A, Visser A, Garssen B, Hudig W. Mindfulness-based stress reduction training for oncology patients: Patients' appraisal and changes in well-being. *Patient Education and Counseling*. 2008;72: 436-442.
215. Shapiro SL, Bootzin RR, Figueredo AJ, Lopez AM, Schwartz GE. The efficacy of mindfulness-based stress reduction in the treatment of sleep disturbance in women with breast cancer: an exploratory study. *J Psychosom Res*. 2003;54: 85-91.
216. Pruthi S. A Randomized Controlled Pilot Study Assessing Feasibility and Impact of Yoga Practice on Quality of Life, Mood, and Perceived Stress in Women With Newly Diagnosed Breast Cancer. *Global Advances in Health and Medicine*. 2012;1: 30-35.

217. Frisk J, Kallstrom AC, Wall N, Fredrikson M, Hammar M. Acupuncture improves health-related quality-of-life (HRQoL) and sleep in women with breast cancer and hot flushes. *Support Care Cancer*. 2012;20: 715-724.
218. Rostock M, Jaroslowski K, Guethlin C, Ludtke R, Schroder S, Bartsch HH. Chemotherapy-induced peripheral neuropathy in cancer patients: a four-arm randomized trial on the effectiveness of electroacupuncture. *Evid Based Complement Alternat Med*. 2013;2013: 1-9.
219. Semiglasov VF, Stepula VV, Dudov A, Lehmacher W, Mengs U. The Standardised Mistletoe Extract PS76A2 Improves QoL in Patients with Breast Cancer Receiving Adjuvant CMF Chemotherapy: A Randomised, Placebo-controlled, Double-blind, Multicentre Clinical Trial. *Anticancer Research*. 2004;24: 1293-1302.
220. Semiglasov VF, Stepula VV, Dudov A, Schnitker J, Mengs U. Quality of life is improved in breast cancer patients by standardised mistletoe extract PS76A2 during chemotherapy and follow-up: A randomised, placebo-controlled, double-blind, multicentre clinical trial. *Anticancer Research*. 2006;26: 1519-1529.
221. Troger W, Jezdic S, Zdrale Z, Tisma N, Hamre HJ, Matijasevic M. Quality of life and neutropenia in patients with early stage breast cancer: A randomized pilot study comparing additional treatment with mistletoe extract to chemotherapy alone. *Breast Cancer: Basic and Clinical Research*. 2009;3: 35-45.
222. Tröger W, Zdrale Z, Tišma N, Matijašević M. Additional Therapy with a Mistletoe Product during Adjuvant Chemotherapy of Breast Cancer Patients Improves Quality of Life: An Open Randomized Clinical Pilot Trial. *Evidence-based complementary and alternative medicine : eCAM*. 2014;2014: 430518.
223. Chen Z, Meng Z, Milbury K, et al. Qigong improves quality of life in women undergoing radiotherapy for breast cancer: results of a randomized controlled trial. *Cancer*. 2013;119: 1690-1698.
224. Oh B, Butow PN, Mullan BA, et al. Effect of medical Qigong on cognitive function, quality of life, and a biomarker of inflammation in cancer patients: a randomized controlled trial. *Support Care Cancer*. 2012;20: 1235-1242.

225. Dyer J, Thomas K, Sandsund C, Shaw C. Is reflexology as effective as aromatherapy massage for symptom relief in an adult outpatient oncology population? *Complement Ther Clin Pract*. 2013;19: 139-146.
226. Sharp DM, Walker MB, Chaturvedi A, et al. A randomised, controlled trial of the psychological effects of reflexology in early breast cancer. *Eur J Cancer*. 2010;46: 312-322.
227. Wyatt G, Sikorskii A, Rahbar MH, Victorson D, You M. Health-related quality-of-life outcomes: a reflexology trial with patients with advanced-stage breast cancer. *Oncol Nurs Forum*. 2012;39: 568-577.
228. Lerman R, Jarski R, Rea H, Gellish R, Vicini F. Improving symptoms and quality of life of female cancer survivors: a randomized controlled study. *Ann Surg Oncol*. 2012;19: 373-378.
229. Ostermann T, Raak C, Bussing A. Survival of cancer patients treated with mistletoe extract (Iscador): a systematic literature review. *BMC Cancer*. 2009;9: 451.
230. National Cancer Institute. Nausea and Vomiting (PDQ). Available from URL: <http://www.cancer.gov/cancertopics/pdq/supportivecare/nausea/HealthProfessional> [accessed 6/22/15].
231. Roila F, Ruggeri B, Ballatori E, Del Favero A, Tonato M. Aprepitant versus dexamethasone for preventing chemotherapy-induced delayed emesis in patients with breast cancer: a randomized double-blind study. *J Clin Oncol*. 2014;32: 101-106.
232. Rojas C, Raje M, Tsukamoto T, Slusher BS. Molecular mechanisms of 5-HT(3) and NK(1) receptor antagonists in prevention of emesis. *Eur J Pharmacol*. 2014;722: 26-37.
233. Booth CM, Clemons M, Dranitsaris G, et al. Chemotherapy-induced nausea and vomiting in breast cancer patients: a prospective observational study. *J Support Oncol*. 2007;5: 374-380.
234. Lindley CM, Hirsch JD. Nausea and Vomiting and Cancer-Patients Quality-of-Life - Discussion. *British Journal of Cancer*. 1992;66: S26-S29.
235. Fabi A, Barduagni M, Lauro S, et al. Is delayed chemotherapy-induced emesis well managed in oncological clinical practice? An observational study. *Supportive Care in Cancer*. 2003;11: 156-161.

236. Carlotto A, Hogsett VL, Maiorini EM, Razulis JG, Sonis ST. The economic burden of toxicities associated with cancer treatment: review of the literature and analysis of nausea and vomiting, diarrhoea, oral mucositis and fatigue. *Pharmacoeconomics*. 2013;31: 753-766.
237. Dibble SL, Chapman J, Mack KA, Shih AS. Acupressure for nausea: results of a pilot study. *Oncol Nurs Forum*. 2000;27: 41-47.
238. Dibble SL, Luce J, Cooper BA, et al. Acupressure for chemotherapy-induced nausea and vomiting: a randomized clinical trial. *Oncol Nurs Forum*. 2007;34: 813-820.
239. Molassiotis A, Helin AM, Dabbour R, Hummerston S. The effects of P6 acupressure in the prophylaxis of chemotherapy-related nausea and vomiting in breast cancer patients. *Complement Ther Med*. 2007;15: 3-12.
240. Lee J, Dodd M, Dibble S, Abrams D. Review of acupressure studies for chemotherapy-induced nausea and vomiting control. *J Pain Symptom Manage*. 2008;36: 529-544.
241. Kwon JH, Shin Y, Juon HS. Effects of Nei-Guan (P6) Acupressure Wristband: On Nausea, Vomiting, and Retching in Women After Thyroidectomy. *Cancer Nurs*. 2015.
242. Noroozinia H, Mahoori A, Hasani E, Gerami-Fahim M, Sepehrvand N. The effect of acupressure on nausea and vomiting after cesarean section under spinal anesthesia. *Acta Med Iran*. 2013;51: 163-167.
243. Collins KB, Thomas DJ. Acupuncture and acupressure for the management of chemotherapy-induced nausea and vomiting. *J Am Acad Nurse Pract*. 2004;16: 76-80.
244. Lee J, Dibble S, Dodd M, Abrams D, Burns B. The relationship of chemotherapy-induced nausea to the frequency of pericardium 6 digital acupressure. *Oncol Nurs Forum*. 2010;37: E419-425.
245. Molassiotis A, Russell W, Hughes J, et al. The effectiveness and cost-effectiveness of acupressure for the control and management of chemotherapy-related acute and delayed nausea: Assessment of Nausea in Chemotherapy Research (ANCHoR), a randomised controlled trial. *Health Technol Assess*. 2013;17: 1-114.

246. Beith JM, Oh B, Chatfield MD, Davis E, Venkateswaran R. Electroacupuncture for nausea, vomiting, and myelosuppression in women receiving adjuvant chemotherapy for early breast cancer: a randomized controlled pilot trial. *Medical Acupuncture*. 2012;24: 241-248.
247. Shen J, Wenger N, Glaspy J, et al. Electroacupuncture for control of myeloablative chemotherapy-induced emesis: a randomized controlled trial. *JAMA* 2000;284: 2755-2761.
248. NIH Consensus Conference. Acupuncture. *JAMA*. 1998;280: 1518-1524.
249. Yang Y, Zhang Y, Jing NC, et al. [Electroacupuncture at Zusanli (ST 36) for treatment of nausea and vomiting caused by the chemotherapy of the malignant tumor: a multicentral randomized controlled trial]. *Zhongguo Zhen Jiu*. 2009;29: 955-958.
250. Dundee JW, Ghaly RG, Fitzpatrick KT, Abram WP, Lynch GA. Acupuncture prophylaxis of cancer chemotherapy-induced sickness. *J R Soc Med*. 1989;82: 268-271.
251. Dundee JW, Ghaly RG, Fitzpatrick KT, Lynch GA, Abram WP. Acupuncture to prevent cisplatin-associated vomiting. *Lancet*. 1987;1: 1083.
252. Ezzo J, Vickers A, Richardson MA, et al. Acupuncture-point stimulation for chemotherapy-induced nausea and vomiting. *J Clin Oncol*. 2005;23: 7188-7198.
253. Gan TJ, Jiao KR, Zenn M, Georgiade G. A randomized controlled comparison of electro-acupoint stimulation or ondansetron versus placebo for the prevention of postoperative nausea and vomiting. *Anesth Analg*. 2004;99: 1070-1075.
254. Wang XQ, Yu JL, Du ZY, Xu R, Jiang CC, Gao X. Electroacupoint stimulation for postoperative nausea and vomiting in patients undergoing supratentorial craniotomy. *J Neurosurg Anesthesiol*. 2010;22: 128-131.
255. El-Deeb AM, Ahmady MS. Effect of acupuncture on nausea and/or vomiting during and after cesarean section in comparison with ondansetron. *J Anesth*. 2011;25: 698-703.
256. Rusy LM, Hoffman GM, Weisman SJ. Electroacupuncture prophylaxis of postoperative nausea and vomiting following pediatric tonsillectomy with or without adenoidectomy. *Anesthesiology*. 2002;96: 300-305.

257. Lee S, Lee MS, Choi DH, Lee SK. Electroacupuncture on PC6 prevents opioid-induced nausea and vomiting after laparoscopic surgery. *Chin J Integr Med.* 2013;19: 277-281.
258. Zhang H, Wang L, Zhang M, et al. [Effects of electroacupuncture on postoperative functional recovery in patients with gynaecological laparoscopic surgery]. *Zhongguo Zhen Jiu.* 2014;34: 273-278.
259. White PF, Issioui T, Hu J, et al. Comparative efficacy of acustimulation (ReliefBand) versus ondansetron (Zofran) in combination with droperidol for preventing nausea and vomiting. *Anesthesiology.* 2002;97: 1075-1081.
260. Genc F, Tan M. The effect of acupressure application on chemotherapy-induced nausea, vomiting, and anxiety in patients with breast cancer. *Palliat Support Care.* 2014: 1-10.
261. Arslan M. Oral Intake of Ginger for Chemotherapy-Induced Nausea and Vomiting Among Women With Breast Cancer. Pittsburgh, Pennsylvania: Oncology Nursing Society, 2015:E92-E97.
262. Panahi Y, Saadat A, Sahebkar A, Hashemian F, Taghikhani M, Abolhasani E. Effect of ginger on acute and delayed chemotherapy-induced nausea and vomiting: a pilot, randomized, open-label clinical trial. *Integr Cancer Ther.* 2012;11: 204-211.
263. Ryan JL, Heckler CE, Roscoe JA, et al. Ginger (*Zingiber officinale*) reduces acute chemotherapy-induced nausea: A URCC CCOP study of 576 patients. *Support Care Cancer.* 2012;20: 1479-1489.
264. Bozzetti F, Biganzoli L, Gavazzi C, et al. Glutamine supplementation in cancer patients receiving chemotherapy: a double-blind randomized study. *Nutrition.* 1997;13: 748-751.
265. Peterson DE, Jones JB, Petit Li RG. Randomized, placebo-controlled trial of Saforis for prevention and treatment of oral mucositis in breast cancer patients receiving anthracycline-based chemotherapy. *Cancer.* 2007;109: 322-331.
266. National Cancer Institute. Radiation Therapy. Available from URL: <http://www.cancer.gov/about-cancer/treatment/types/radiation-therapy - RTCCSE> [accessed 6/27/16].
267. National Cancer Institute. Skin and Nail Changes. Available from URL: <http://www.cancer.gov/about-cancer/treatment/side-effects/skin-nail-changes> [accessed 6/27/16].

268. Heggie S, Bryant GP, Tripcony L, et al. A Phase III study on the efficacy of topical aloe vera gel on irradiated breast tissue. *Cancer Nurs.* 2002;25: 442-451.
269. Williams MS, Burk M, Loprinzi CL, et al. Phase III double-blind evaluation of an aloe vera gel as a prophylactic agent for radiation-induced skin toxicity. *Int J Radiat Oncol Biol Phys.* 1996;36: 345-349.
270. Kirova YM, Fromantin I, De Rycke Y, et al. Can we decrease the skin reaction in breast cancer patients using hyaluronic acid during radiation therapy? Results of phase III randomised trial. *Radiother Oncol.* 2011;100: 205-209.
271. Pinnix C, Perkins GH, Strom EA, et al. Topical hyaluronic acid vs. standard of care for the prevention of radiation dermatitis after adjuvant radiotherapy for breast cancer: single-blind randomized phase III clinical trial. *Int J Radiat Oncol Biol Phys.* 2012;83: 1089-1094.
272. National Cancer Institute. Hot flashes and night sweats (PDQ). Available from URL: <https://www.cancer.gov/about-cancer/treatment/side-effects/sexuality-fertility-women/hot-flashes-hp-pdq> [accessed 12/19/16].
273. Vincent A. Management of menopause in women with breast cancer. *Climacteric.* 2014;8: 690-701.
274. Bokmand S, Flyger H. Acupuncture relieves menopausal discomfort in breast cancer patients: a prospective, double blinded, randomized study. *Breast.* 2013;22: 320-323.
275. Deng G, Vickers A, Yeung S, et al. Randomized, controlled trial of acupuncture for the treatment of hot flashes in breast cancer patients. *J Clin Oncol.* 2007;25: 5584-5590.
276. Frisk J, Carlhall S, Kallstrom AC, Lindh-Astrand L, Malmstrom A, Hammar M. Long-term follow-up of acupuncture and hormone therapy on hot flushes in women with breast cancer: a prospective, randomized, controlled multicenter trial. *Climacteric.* 2008;11: 166-174.
277. Hervik J, Mjaland O. Quality of life of breast cancer patients medicated with anti-estrogens, 2 years after acupuncture treatment: a qualitative study. *Int J Womens Health.* 2010;2: 319-325.

278. Liljegren A, Gunnarsson P, Landgren BM, Robeus N, Johansson H, Rotstein S. Reducing vasomotor symptoms with acupuncture in breast cancer patients treated with adjuvant tamoxifen: a randomized controlled trial. *Breast Cancer Res Treat.* 2012;135: 791-798.
279. Mao J, Bowman M, Xie S, Bruner D, De Michele A, Farrar J. Electroacupuncture versus gabapentin for hot flashes among breast cancer survivors: A randomized placebo-controlled trial. *Journal of Clinical Oncology.* 2015;33: 3615-3620.
280. MacGregor CA, Canney PA, Patterson G, McDonald R, Paul J. A randomised double-blind controlled trial of oral soy supplements versus placebo for treatment of menopausal symptoms in patients with early breast cancer. *Eur J Cancer.* 2005;41: 708-714.
281. Quella SK, Loprinzi CL, Barton DL, et al. Evaluation of soy phytoestrogens for the treatment of hot flashes in breast cancer survivors: A North Central Cancer Treatment Group trial. *J Clin Oncol.* 2000;18: 1068-1074.
282. Van Patten CL, Olivotto IA, Chambers GK, et al. Effect of soy phytoestrogens on hot flashes in postmenopausal women with breast cancer: a randomized, controlled clinical trial. *J Clin Oncol.* 2002;20: 1449-1455.
283. Paskett ED, Naughton MJ, McCoy TP, Case LD, Abbott JM. The epidemiology of arm and hand swelling in premenopausal breast cancer survivors. *Cancer Epidemiol Biomarkers Prev.* 2007;16: 775-782.
284. Ridner SH. Quality of life and a symptom cluster associated with breast cancer treatment-related lymphedema. *Support Care Cancer.* 2005;13: 904-911.
285. Pyszel A, Malyszczak K, Pyszel K, Andrzejak R, Szuba A. Disability, psychological distress and quality of life in breast cancer survivors with arm lymphedema. *Lymphology.* 2006;39: 185-192.
286. Ahmed Omar MT, Abd-El-Gayed Ebid A, El Morsy AM. Treatment of post-mastectomy lymphedema with laser therapy: double blind placebo control randomized study. *J Surg Res.* 2011;165: 82-90.

287. Ridner SH, Poage-Hooper E, Kanar C, Doersam JK, Bond SM, Dietrich MS. A pilot randomized trial evaluating low-level laser therapy as an alternative treatment to manual lymphatic drainage for breast cancer-related lymphedema. *Oncol Nurs Forum*. 2013;40: 383-393.
288. Andersen L, Hojris I, Erlandsen M, Andersen J. Treatment of breast-cancer-related lymphedema with or without manual lymphatic drainage: a randomized study. *Acta Oncol*. 2000;39: 399-405.
289. Devoogdt N, Christiaens MR, Geraerts I, et al. Effect of manual lymph drainage in addition to guidelines and exercise therapy on arm lymphoedema related to breast cancer: randomised controlled trial. *BMJ*. 2011;343:d5326.
290. Gurdal SO, Kostanoglu A, Cavdar I, et al. Comparison of intermittent pneumatic compression with manual lymphatic drainage for treatment of breast cancer-related lymphedema. *Lymphat Res Biol*. 2012;10: 129-135.
291. Maher J, Refshauge K, Ward L, Paterson R, Kilbreath S. Change in extracellular fluid and arm volumes as a consequence of a single session of lymphatic massage followed by rest with or without compression. *Support Care Cancer*. 2012;20: 3079-3086.
292. McNeely ML, Magee DJ, Lees AW, Bagnall KM, Haykowsky M, Hanson J. The addition of manual lymph drainage to compression therapy for breast cancer related lymphedema: a randomized controlled trial. *Breast Cancer Res Treat*. 2004;86: 95-106.
293. Williams AF, Vadgama A, Franks PJ, Mortimer PS. A randomized controlled crossover study of manual lymphatic drainage therapy in women with breast cancer-related lymphoedema. *Eur J Cancer Care (Engl)*. 2002;11: 254-261.
294. Dayes IS, Whelan TJ, Julian JA, et al. Randomized trial of decongestive lymphatic therapy for the treatment of lymphedema in women with breast cancer. *J Clin Oncol*. 2013;31: 3758-3763.
295. National Cancer Institute. Nerve Problems. Available from URL: <http://www.cancer.gov/about-cancer/treatment/side-effects/nerve-problems> [accessed 11/27/16].

296. International Association for the Study of Pain. Epidemiology of Cancer Pain. Available from URL: <http://www.iasppain.org/AM/Template.cfm?Section=Home&Template=/CM/ContentDisplay.cfm&ContentID=7395> [accessed 6/6/16].
297. National Cancer Institute. Cancer Pain (PDQ). Available from URL: <http://www.cancer.gov/about-cancer/treatment/side-effects/pain/pain-hp-pdq> [accessed 6/27/16].
298. Montgomery GH, Bovbjerg DH, Schnur JB, et al. A randomized clinical trial of a brief hypnosis intervention to control side effects in breast surgery patients. *J Natl Cancer Inst.* 2007;99: 1304-1312.
299. Montgomery GH, Wertz CR, Seltz M, Bovbjerg DH. Brief presurgery hypnosis reduces distress and pain in excisional breast biopsy patients. *Int J Clin Exp Hypn.* 2002;50: 17-32.
300. Bao T, Cai L, Giles JT, et al. A dual-center randomized controlled double blind trial assessing the effect of acupuncture in reducing musculoskeletal symptoms in breast cancer patients taking aromatase inhibitors. *Breast Cancer Res Treat.* 2013;138: 167-174.
301. Crew KD, Capodice JL, Greenlee H, et al. Pilot study of acupuncture for the treatment of joint symptoms related to adjuvant aromatase inhibitor therapy in postmenopausal breast cancer patients. *J Cancer Surviv.* 2007;1: 283-291.
302. Crew KD, Capodice JL, Greenlee H, et al. Randomized, blinded, sham-controlled trial of acupuncture for the management of aromatase inhibitor-associated joint symptoms in women with early-stage breast cancer. *J Clin Oncol.* 2010;28: 1154-1160.
303. Oh B, Kimble B, Costa DSJ, et al. Acupuncture for treatment of arthralgia secondary to aromatase inhibitor therapy in women with early breast cancer: pilot study. *Acupunct Med.* 2013;31: 264-271.
304. Mao JJ, Xie SX, Farrar JT, et al. A randomised trial of electro-acupuncture for arthralgia related to aromatase inhibitor use. *European journal of cancer (Oxford, England : 1990).* 2014;50: 267-276.
305. Savard J, Morin CM. Insomnia in the context of cancer: a review of a neglected problem. *J Clin Oncol.* 2001;19: 895-908.

306. National Cancer Institute. Sleep Disorders (PDQ). Available from URL: <http://www.cancer.gov/about-cancer/treatment/side-effects/sleep-disorders-hp-pdq> [accessed 6/27/16].
307. Greenlee H, Shi Z, Sardo Molmenti CL, Rundle A, Tsai WY. Trends in Obesity Prevalence in Adults With a History of Cancer: Results From the US National Health Interview Survey, 1997 to 2014. *J Clin Oncol*. 2016;34: 3133-3140.
308. Bardwell WA, Profant J, Casden DR, et al. The relative importance of specific risk factors for insomnia in women treated for early-stage breast cancer. *Psychooncology*. 2008;17: 9-18.
309. Mustian K, Sprod L, Janelins M, et al. Multicenter, randomized controlled trial of yoga for sleep quality among cancer survivors. *J Clin Oncol*. 2013;31: 3233-3241.
310. Witt CM, Cardoso MJ. Complementary and integrative medicine for breast cancer patients - Evidence based practical recommendations. *Breast*. 2016;28: 37-44.
311. Gabriella M. John DLH, Laura Falci, Zaixing Shi, Wei-Yann Tsai, Heather Greenlee. Out-of-Pocket Costs of Vitamin/Mineral and Complementary and Alternative Medicine Use among Cancer Survivors in the United States: Results from the 2012 National Health Interview Survey. *Journal of Cancer Survivorship*. In review.
312. Society for Oncology Massage. Society for Oncology Massage. Available from URL: <http://www.s4om.org/> [accessed 6/18/15].
313. OncANP: Oncology Association of Naturopathic Physicians. Available from URL: <https://oncnp.org> [accessed 12/14/16].
314. Kluetz PG, Chingos DT, Basch EM, Mitchell SA. Patient-Reported Outcomes in Cancer Clinical Trials: Measuring Symptomatic Adverse Events With the National Cancer Institute's Patient-Reported Outcomes Version of the Common Terminology Criteria for Adverse Events (PRO-CTCAE). *Am Soc Clin Oncol Educ Book*. 2016;35: 67-73.

315. National Cancer Institute. CTCAE Files. Available from URL: <https://evs.nci.nih.gov/ftp1/CTCAE/About.html> [accessed 10/26/16].
316. John GM, Hershman DL, Falci L, Shi Z, Tsai WY, Greenlee H. Complementary and alternative medicine use among US cancer survivors. *J Cancer Surviv.* 2016;10: 850-864.
317. Zick SM, Alrawi S, Merel G, et al. Relaxation acupuncture reduces persistent cancer-related fatigue. *Evid Based Complement Alternat Med.* 2011;2011: 1-10.
318. Targ EF, Levine EG. The efficacy of a mind-body-spirit group for women with breast cancer: a randomized controlled trial. *Gen Hosp Psychiatry.* 2002;24: 238-248.
319. Yaal-Hahoshen N, Maimon Y, Siegelmann-Danieli N, et al. A prospective, controlled study of the botanical compound mixture LCS101 for chemotherapy-induced hematological complications in breast cancer. *Oncologist.* 2011;16: 1197-1202.
320. Zhuang SR, Chiu HF, Chen SL, et al. Effects of a Chinese medical herbs complex on cellular immunity and toxicity-related conditions of breast cancer patients. *Br J Nutr.* 2012;107: 712-718.
321. Yang B, Li H, Qi Y, Lu M, Bo Y. Clinical study on treatment of mammary cancer by Shenqi Fuzheng Injection in cooperation with chemotherapy. *Chin J Integr Med.* 2007;13: 37-40.
322. Monti DA, Kash KM, Kunkel EJ, et al. Changes in cerebral blood flow and anxiety associated with an 8-week mindfulness programme in women with breast cancer. *Stress Health.* 2012;28: 397-407.
323. Gaston-Johansson F, Fall-Dickson JM, Nanda J, et al. The effectiveness of the comprehensive coping strategy program on clinical outcomes in breast cancer autologous bone marrow transplantation. *Cancer Nurs.* 2000;23: 277-285.
324. Robb KA, Newham DJ, Williams JE. Transcutaneous electrical nerve stimulation vs. transcutaneous spinal electroanalgesia for chronic pain associated with breast cancer treatments. *J Pain Symptom Manage.* 2007;33: 410-419.
325. Frank LS, Frank JL, March D, Makari-Judson G, Barham RB, Mertens WC. Does therapeutic touch ease the discomfort or distress of patients undergoing stereotactic core breast biopsy? A randomized clinical trial. *Pain Med.* 2007;8: 419-424.

326. Schnur JB, Bovbjerg DH, David D, et al. Hypnosis decreases presurgical distress in excisional breast biopsy patients. *Anesth Analg*. 2008;106: 440-444.
327. Marshall-Mckenna R, Paul L, McFadyen AK, et al. Myofascial release for women undergoing radiotherapy for breast cancer: A pilot study. *European Journal of Physiotherapy*. 2014;16: 58-64.
328. Baker BS, Harrington JE, Choi BS, Kropf P, Muller I, Hoffman CJ. A randomised controlled pilot feasibility study of the physical and psychological effects of an integrated support programme in breast cancer. *Complement Ther Clin Pract*. 2012;18: 182-189.
329. Liu CJ, Hsiung PC, Chang KJ, et al. A study on the efficacy of body-mind-spirit group therapy for patients with breast cancer. *J Clin Nurs*. 2008;17: 2539-2549.
330. Stephenson NL, Weinrich SP, Tavakoli AS. The effects of foot reflexology on anxiety and pain in patients with breast and lung cancer. *Oncol Nurs Forum*. 2000;27: 67-72.
331. Potter PJ. Breast biopsy and distress: feasibility of testing a reiki intervention. *J Holist Nurs*. 2007;25: 238-251.
332. Robins JLW, McCain NL, Elswick RK, Walter JM, Gray DP, Tuck I. Psychoneuroimmunology-based stress management during adjuvant chemotherapy for early breast cancer. *Evid Based Complement Alternat Med*. 2013;2013: 1-7.
333. Jo SH, Kim LS, Kim SA, et al. Evaluation of short-term use of N-acetylcysteine as a strategy for prevention of anthracycline-induced cardiomyopathy: EPOCH trial - A prospective randomized study. *Korean Circ J*. 2013;43: 174-181.
334. Roscoe JA, Matteson SE, Morrow GR, et al. Acustimulation wrist bands are not effective for the control of chemotherapy-induced nausea in women with breast cancer. *J Pain Symptom Manage*. 2005;29: 376-384.
335. Lua PL, Salihah N, Mazlan N. Effects of inhaled ginger aromatherapy on chemotherapy-induced nausea and vomiting and health-related quality of life in women with breast cancer. *Complementary therapies in medicine*. 2015;23: 396-404.

336. Valadares F, Novaes MR, Canete R. Effect of *Agaricus sylvaticus* supplementation on nutritional status and adverse events of chemotherapy of breast cancer: a randomized, placebo-controlled, double-blind clinical trial. *Indian J Pharmacol.* 2013;45: 217-222.
337. Perol D, Provencal J, Hardy-Bessard AC, et al. Can treatment with Cocculine improve the control of chemotherapy-induced emesis in early breast cancer patients? A randomized, multi-centered, double-blind, placebo-controlled Phase III trial. *BMC Cancer.* 2012;12: 603.
338. Moradian S, Walshe C, Shahidsales S, Ghavam Nasiri MR, Pilling M, Molassiotis A. Nevasic audio program for the prevention of chemotherapy induced nausea and vomiting: A feasibility study using a randomized controlled trial design. *European journal of oncology nursing : the official journal of European Oncology Nursing Society.* 2015;19: 282-291.
339. Cimprich B, Ronis DL. An environmental intervention to restore attention in women with newly diagnosed breast cancer. *Cancer Nurs.* 2003;26: 284-292.
340. Derry HM, Bennett JM, Jaremka LM, et al. Yoga and self-reported cognitive problems: A randomized controlled trial for breast cancer survivors. *2014;76: A-3.*
341. Hanai A, Ishiguro H, Sozu T, et al. Effects of a self-management program on antiemetic-induced constipation during chemotherapy among breast cancer patients: a randomized controlled clinical trial. *Breast Cancer Res Treat.* 2016;155: 99-107.
342. Jain S, Pavlik D, Distefan J, et al. Complementary medicine for fatigue and cortisol variability in breast cancer survivors. *Cancer.* 2012;118: 777-787.
343. Lesser GJ, Case D, Stark N, et al. A randomized, double-blind, placebo-controlled study of oral coenzyme Q10 to relieve self-reported treatment-related fatigue in newly diagnosed patients with breast cancer. *J Support Oncol.* 2013;11: 31-42.
344. Zhao H, Zhang Q, Zhao L, Huang X, Wang J, Kang X. Spore powder of *Ganoderma lucidum* improves cancer-related fatigue in breast cancer patients undergoing endocrine therapy: a pilot clinical trial. *Evid Based Complement Alternat Med.* 2012;2012: 1-8.

345. Schnur JB, David D, Kangas M, Green S, Bovbjerg DH, Montgomery GH. A randomized trial of a cognitive-behavioral therapy and hypnosis intervention on positive and negative affect during breast cancer radiotherapy. *J Clin Psychol.* 2009;65: 443-455.
346. Andersen BL, Farrar WB, Golden-Kreutz DM, et al. Psychological, behavioral, and immune changes after a psychological intervention: a clinical trial. *J Clin Oncol.* 2004;22: 3570-3580.
347. Spahn G, Choi KE, Kennemann C, et al. Can a multimodal mind-body program enhance the treatment effects of physical activity in breast cancer survivors with chronic tumor-associated fatigue? A randomized controlled trial. *Integr Cancer Ther.* 2013;12: 291-300.
348. Balk J, Day R, Rosenzweig M, Beriwal S. Pilot, randomized, modified, double-blind, placebo-controlled trial of acupuncture for cancer-related fatigue. *J Soc Integr Oncol.* 2009;7: 4-11.
349. Johnston MF, Hays RD, Subramanian SK, et al. Patient education integrated with acupuncture for relief of cancer-related fatigue randomized controlled feasibility study. *BMC Complement Altern Med.* 2011;11: 49.
350. Ancoli-Israel S, Rissling M, Neikrug A, et al. Light treatment prevents fatigue in women undergoing chemotherapy for breast cancer. *Support Care Cancer.* 2012;20: 1211-1219.
351. van der Lee ML, Garssen B. Mindfulness-based cognitive therapy reduces chronic cancer-related fatigue: a treatment study. *Psychooncology.* 2012;21: 264-272.
352. Reis D, Walsh ME, Young-McCaughan S, Jones T. Effects of Nia exercise in women receiving radiation therapy for breast cancer. *Oncology Nursing Forum.* 2013;40: E374-381.
353. Adamsen L, Quist M, Andersen C, et al. Effect of a multimodal high intensity exercise intervention in cancer patients undergoing chemotherapy: randomised controlled trial. *BMJ.* 2009;339: 895-898.
354. Bjornekleit HG, Lindemalm C, Rosenblad A, et al. A randomised controlled trial of support group intervention after breast cancer treatment: results on anxiety and depression. *Acta Oncol.* 2012;51: 198-207.
355. de Souza Fede AB, Bensi CG, Trufelli DC, et al. Multivitamins do not improve radiation therapy-related fatigue: results of a double-blind randomized crossover trial. *Am J Clin Oncol.* 2007;30: 432-436.

356. Roscoe J, Matteson S, Mustian K, Padmanaban D, Morrow G. Treatment of radiotherapy-induced fatigue through a nonpharmacological approach. *Integr Cancer Ther.* 2005;4: 8-13.
357. Mustian KM, Roscoe JA, Palesh OG, et al. Polarity Therapy for cancer-related fatigue in patients with breast cancer receiving radiation therapy: a randomized controlled pilot study. *Integr Cancer Ther.* 2011;10: 27-37.
358. Rissanen R, Arving C, Ahlgren J, Nordin K. Group versus individual stress management intervention in breast cancer patients for fatigue and emotional reactivity: A randomised intervention study. *Acta Oncologica.* 2014;53: 1221-1229.
359. Steindorf K, Schmidt ME, Klassen O, et al. Randomized, controlled trial of resistance training in breast cancer patients receiving adjuvant radiotherapy: results on cancer-related fatigue and quality of life. *Annals of Oncology.* 2014;25: 2237-2243.
360. Schmidt ME, Wiskemann J, Armbrust P, Schneeweiss A, Ulrich CM, Steindorf K. Effects of resistance exercise on fatigue and quality of life in breast cancer patients undergoing adjuvant chemotherapy: A randomized controlled trial. *International journal of cancer. Journal international du cancer.* 2015;137: 471-480.
361. Cluzan RV, Alliot F, Ghabboun S, Pascot M. Treatment of secondary lymphedema of the upper limb with CYCLO 3 FORT. *Lymphology.* 1996;29: 29-35.
362. Belmonte R, Tejero M, Ferrer M, et al. Efficacy of low-frequency low-intensity electrotherapy in the treatment of breast cancer-related lymphoedema: a cross-over randomized trial. *Clin Rehabil.* 2012;26: 607-618.
363. Cluzan RV, Pecking AP, Mathiex-Fortunet H, Leger Picherit E. Efficacy of BN165 (Ginkor Fort) in breast cancer related upper limb lymphedema: a preliminary study. *Lymphology.* 2004;37: 47-52.
364. Gothard L, Cornes P, Earl J, et al. Double-blind placebo-controlled randomised trial of vitamin E and pentoxifylline in patients with chronic arm lymphoedema and fibrosis after surgery and radiotherapy for breast cancer. *Radiother Oncol.* 2004;73: 133-139.

365. Magnusson M, Hoglund P, Johansson K, et al. Pentoxifylline and vitamin E treatment for prevention of radiation-induced side-effects in women with breast cancer: a phase two, double-blind, placebo-controlled randomised clinical trial (Ptx-5). *Eur J Cancer*. 2009;45: 2488-2495.
366. Loudon A, Barnett T, Piller N, Immink MA, Williams AD. Yoga management of breast cancer-related lymphoedema: a randomised controlled pilot-trial. *BMC complementary and alternative medicine*. 2014;14: 214.
367. Ghoreishi Z, Esfahani A, Djazayeri A, et al. Omega-3 fatty acids are protective against paclitaxel-induced peripheral neuropathy: a randomized double-blind placebo controlled trial. *BMC Cancer*. 2012;12: 355.
368. Argyriou AA, Chroni E, Koutras A, et al. Preventing paclitaxel-induced peripheral neuropathy: a phase II trial of vitamin E supplementation. *J Pain Symptom Manage*. 2006;32: 237-244.
369. Pace A, Giannarelli D, Galie E, et al. Vitamin E neuroprotection for cisplatin neuropathy: a randomized, placebo-controlled trial. *Neurology*. 2010;74: 762-766.
370. Santos Araujo MDC, Farias IL, Gutierrez J, et al. *Uncaria tomentosa* - adjuvant treatment for breast cancer: clinical trial. *Evid Based Complement Alternat Med*. 2012;2012: 1-8.
371. Rastelli AL, Taylor ME, Gao F, et al. Vitamin D and aromatase inhibitor-induced musculoskeletal symptoms (AIMSS): a phase II, double-blind, placebo-controlled, randomized trial. *Breast Cancer Res Treat*. 2011;129: 107-116.
372. Arathuzik D. Effects of cognitive-behavioral strategies on pain in cancer patients. *Cancer Nurs*. 1994;17: 207-214.
373. Butler LD, Koopman C, Neri E, et al. Effects of supportive-expressive group therapy on pain in women with metastatic breast cancer. *Health Psychol*. 2009;28: 579-587.
374. Sharp L, Finnila K, Johansson H, Abrahamsson M, Hatschek T, Bergenmar M. No differences between *Calendula* cream and aqueous cream in the prevention of acute radiation skin reactions--results from a randomised blinded trial. *Eur J Oncol Nurs*. 2013;17: 429-435.

375. Brisbois TD, de Kock IH, Watanabe SM, et al. Delta-9-tetrahydrocannabinol may palliate altered chemosensory perception in cancer patients: results of a randomized, double-blind, placebo-controlled pilot trial. *Ann Oncol.* 2011;22: 2086-2093.
376. Noguchi N, Maruyama I, Yamada A. The influence of chlorella and its hot water extract supplementation on quality of life in patients with breast cancer. *Evidence-based complementary and alternative medicine : eCAM.* 2014;2014: 704619.
377. Panahi Y, Saadat A, Beiraghdar F, Sahebkar A. Adjuvant therapy with bioavailability-boosted curcuminoids suppresses systemic inflammation and improves quality of life in patients with solid tumors: a randomized double-blind placebo-controlled trial. *Phytother Res.* 2014;28: 1461-1467.
378. Pruthi S, Qin R, Terstreip SA, et al. A phase III, randomized, placebo-controlled, double-blind trial of flaxseed for the treatment of hot flashes: North Central Cancer Treatment Group N08C7. *Menopause.* 2012;19: 48-53.
379. Richardson M, Post-White J, Grimm E, Moye L, Singletary S, Justice B. Coping, life attitudes, and immune responses to imagery and group support after breast cancer treatment. *Alternative therapies in health and medicine.* 1997;3: 62-70.
380. Frass M, Friehs H, Thallinger C, et al. Influence of adjunctive classical homeopathy on global health status and subjective wellbeing in cancer patients - A pragmatic randomized controlled trial. *Complementary therapies in medicine.* 2015;23: 309-317.
381. Jacobs J, Herman P, Heron K, Olsen S, Vaughters L. Homeopathy for menopausal symptoms in breast cancer survivors: A preliminary randomized controlled trial. *J Altern Complement Med.* 2005;11: 21-27.
382. Thompson EA, Oxon BA, Montgomery A, Douglas D, Reilly D. A pilot, randomized, double-blinded, placebo-controlled trial of individualized homeopathy for symptoms of estrogen withdrawal in breast-cancer survivors. *J Altern Complement Med.* 2005;11: 13-20.
383. Sandel SL, Judge JO, Landry N, Faria L, Ouellette R, Majczak M. Dance and movement program improves quality-of-life measures in breast cancer survivors. *Cancer Nurs.* 2005;28: 301-309.

384. Witt CM, Ausserer O, Baier S, et al. Effectiveness of an additional individualized multi-component complementary medicine treatment on health-related quality of life in breast cancer patients: a pragmatic randomized trial. *Breast Cancer Research and Treatment*. 2015;149: 449-460.
385. Loprinzi CL, Levitt R, Barton DL, et al. Evaluation of shark cartilage in patients with advanced cancer: a North Central Cancer Treatment Group trial. *Cancer*. 2005;104: 176-182.
386. Kissane DW, Grabsch B, Clarke DM, et al. Supportive-expressive group therapy for women with metastatic breast cancer: survival and psychosocial outcome from a randomized controlled trial. *Psycho-Oncology*. 2007;16: 277-286.
387. Campo RA, O'Connor K, Light KC, et al. Feasibility and acceptability of a Tai Chi Chih randomized controlled trial in senior female cancer survivors. *Integr Cancer Ther*. 2013;12: 464-474.
388. Mustian KM, Katula JA, Gill DL, Roscoe JA, Lang D, Murphy K. Tai Chi Chuan, health-related quality of life and self-esteem: a randomized trial with breast cancer survivors. *Support Care Cancer*. 2004;12: 871-876.
389. Sprod LK, Janelins MC, Palesh OG, et al. Health-related quality of life and biomarkers in breast cancer survivors participating in tai chi chuan. *J Cancer Surviv*. 2012;6: 146-154.
390. Harder H, Langridge C, Solis-Trapala I, et al. Post-operative exercises after breast cancer surgery: Results of a RCT evaluating standard care versus standard care plus additional yoga exercise. *European Journal of Integrative Medicine*. 2015;7: 202-210.
391. Huang C, Hou M, Kan J, et al. Prophylactic Treatment with Adlay Bran Extract Reduces the Risk of Severe Acute Radiation Dermatitis: A Prospective, Randomized, Double-Blind Study. *Evidence-based Complementary and Alternative Medicine*. 2015;2015.
392. Ansari M, Dehsara F, Mosalaei A, Omidvari S, Ahmadloo N, Mohammadianpanah M. Efficacy of topical Alpha ointment (containing natural henna) compared to topical hydrocortisone (1%) in the healing of radiation-induced dermatitis in patients with breast cancer: a randomized controlled clinical trial. *Iran J Med Sci*. 2013;38: 293-300.

393. Gosselin TK, Schneider SM, Plambeck MA, Rowe K. A prospective randomized, placebo-controlled skin care study in women diagnosed with breast cancer undergoing radiation therapy. *Oncol Nurs Forum*. 2010;37: 619-626.
394. Togni S, Maramaldi G, Bonetta A, Giacomelli L, Di Pierro F. Clinical evaluation of safety and efficacy of Boswellia-based cream for prevention of adjuvant radiotherapy skin damage in mammary carcinoma: a randomized placebo controlled trial. *Eur Rev Med Pharmacol Sci*. 2015;19: 1338-1344.
395. Pommier P, Gomez F, Sunyach MP, D'Hombres A, Carrie C, Montbarbon X. Phase III randomized trial of Calendula Officinalis compared with trolamine for the prevention of acute dermatitis during irradiation for breast cancer. *J Clin Oncol*. 2004;22: 1447-1453.
396. Maïche AG, Grohn P, Maki-Hokkonen H. Effect of chamomile cream and almond ointment on acute radiation skin reaction. *Acta Oncol*. 1991;30: 395-396.
397. Ryan JL, Heckler CE, Ling M, et al. Curcumin for radiation dermatitis: a randomized, double-blind, placebo-controlled clinical trial of thirty breast cancer patients. *Radiat Res*. 2013;180: 34-43.
398. Rubio I, Suva LJ, Todorova V, et al. Oral glutamine reduces radiation morbidity in breast conservation surgery. *J Parenter Enteral Nutr*. 2013;37: 623-630.
399. Balzarini A, Felisi E, Martini A, De Conno F. Efficacy of homeopathic treatment of skin reactions during radiotherapy for breast cancer: a randomised, double-blind clinical trial. *Br Homeopath J*. 2000;89: 8-12.
400. Moolenaar M, Poorter RL, van der Toorn PP, Lenderink AW, Poortmans P, Egberts AC. The effect of honey compared to conventional treatment on healing of radiotherapy-induced skin toxicity in breast cancer patients. *Acta Oncol*. 2006;45: 623-624.
401. Shoma A, Eldars W, Noman N, et al. Pentoxifylline and local honey for radiation-induced burn following breast conservative surgery. *Curr Clin Pharmacol*. 2010;5: 251-256.
402. Kunos CAMDP, Abdallah RRLA, Lyons JAMD. Hydration During Breast Radiotherapy May Lower Skin Toxicity. *Breast Journal* November/December. 2014;20: 679-681.

403. Bourgeois JF, Gourgou S, Kramar A, Lagarde JM, Guillot B. A randomized, prospective study using the LPG technique in treating radiation-induced skin fibrosis: clinical and profilometric analysis. *Skin Res Technol*. 2008;14: 71-76.
404. Jensen JM, Gau T, Schultze J, et al. Treatment of acute radiodermatitis with an oil-in-water emulsion following radiation therapy for breast cancer: a controlled, randomized trial. *Strahlenther Onkol*. 2011;187: 378-384.
405. Enomoto TM, Johnson T, Peterson N, et al. Combination glutathione and anthocyanins as an alternative for skin care during external-beam radiation. *Am J Surgery*. 2005;189: 627-631.
406. Wheat J, Currie G, Coulter K. Wheatgrass extract as a topical skin agent for acute radiation skin toxicity in breast radiation therapy: a randomised controlled trial. *Journal of the Australian Traditional-Medicine Society*. 2006;12: 135-137.
407. Jacobson G, Bhatia S, Smith BJ, Button AM, Bodeker K, Buatti J. Randomized trial of pentoxifylline and vitamin E vs standard follow-up after breast irradiation to prevent breast fibrosis, evaluated by tissue compliance meter. *Int J Radiat Oncol Biol Phys*. 2013;85: 604-608.
408. Delanian S, Porcher R, Balla-Mekias S, Lefaix JL. Randomized, placebo-controlled trial of combined pentoxifylline and tocopherol for regression of superficial radiation-induced fibrosis. *J Clin Oncol*. 2003;21: 2545-2550.
409. Andersen SR, Wurtzen H, Steding-Jessen M, et al. Effect of mindfulness-based stress reduction on sleep quality: results of a randomized trial among Danish breast cancer patients. *Acta Oncol*. 2013;52: 336-344.
410. Lengacher CA, Reich RR, Paterson CL, et al. The effects of mindfulness-based stress reduction on objective and subjective sleep parameters in women with breast cancer: a randomized controlled trial. *Psycho-oncology*. 2015;24: 424-432.
411. Hernandez Munoz G, Pluchino S. *Cimicifuga racemosa* for the treatment of hot flashes in women surviving breast cancer. *Maturitas*. 2003;44: S59-S65.

412. Jacobson JS, Troxel AB, Evans J, et al. Randomized trial of black cohosh for the treatment of hot flashes among women with a history of breast cancer. *J Clin Oncol*. 2001;19: 2739-2745.
413. Elkins G, Marcus J, Stearns V, et al. Randomized trial of a hypnosis intervention for treatment of hot flashes among breast cancer survivors. *J Clin Oncol*. 2008;26: 5022-5026.
414. Carpenter JS, Wells N, Lambert B, et al. A pilot study of magnetic therapy for hot flashes after breast cancer. *Cancer Nurs*. 2002;25: 104-109.
415. Dyer J, Ashley S, Shaw C. A study to look at the effects of a hydrolat spray on hot flushes in women being treated for breast cancer. *Complement Ther Clin Pract*. 2008;14: 273-279.
416. Barton DL, Loprinzi CL, Quella SK, et al. Prospective evaluation of vitamin E for hot flashes in breast cancer survivors. *J Clin Oncol*. 1998;16: 495-500.
417. Carson JW, Carson KM, Porter LS, Keefe FJ, Seewaldt VL. Yoga of Awareness program for menopausal symptoms in breast cancer survivors: results from a randomized trial. *Support Care Cancer*. 2009;17: 1301-1309.

Accepted Article