

Nutritional status of children with Wilms' tumour on admission to a South African hospital and its influence on outcome

Lauren F. Lifson¹ | G. P. Hadley² | Nicola L. Wiles¹ | Kirthee Pillay¹

¹Dietetics and Human Nutrition, University of KwaZulu-Natal, Pietermaritzburg, South Africa

²Department of Paediatric Surgery, Nelson R Mandela School of Medicine, University of KwaZulu-Natal, Durban, South Africa

Correspondence

Nicola L. Wiles, Dietetics and Human Nutrition, University of KwaZulu-Natal, Pietermaritzburg 3201, South Africa.

Email: wilesn@ukzn.ac.za

Abstract

Background: In developing countries up to 77% of children with cancer have been shown to be malnourished on admission. High rates of malnutrition occur due to factors such as poverty and advanced disease. Weight can be an inaccurate parameter for nutritional assessment of children with solid tumours as it is influenced by tumour mass. This study aimed to assess the prevalence of malnutrition amongst children with Wilms tumour (WT), the level of nutritional support received on admission and the influence of nutritional status on outcome.

Methods: Seventy-six children diagnosed with WT and admitted to Inkosi Albert Luthuli Central Hospital between 2004 and 2012 were studied prospectively. Nutritional assessment was conducted using weight, height, mid-upper arm circumference (MUAC) and triceps skinfold thickness (TSFT) prior to initiating treatment. Outcome was determined 2 years after admission. Time until commencement of nutritional resuscitation and nature, thereof, were recorded.

Results: Stunting and wasting was evident in 12% and 15% of patients, respectively. The prevalence of malnutrition was 66% when MUAC, TSFT and albumin were used. Malnutrition was not a predictor of poor outcome and did not predict advanced disease. The majority of patients (84%) received nutritional resuscitation within 2 weeks of admission.

Conclusions: When classifying nutritional status in children with WT, the utilisation of weight and height in isolation can lead to an underestimation of the prevalence of malnutrition. Nutritional assessment of children with WT should also include MUAC and TSFT. Early aggressive nutritional resuscitation is recommended.

KEYWORDS

arm anthropometry, cancer, children, malnutrition, nutrition, Wilms tumour

1 | INTRODUCTION

Wilms' tumour (WT), also known as nephroblastoma, contributes to approximately 5% of all childhood cancer cases in developed countries^{1,2} and more than 10% of childhood cancers in many African countries.³ In South Africa, WT is the fourth most common childhood cancer according to hospital-based registries.⁴

There is a paucity of information regarding the incidence of cancer amongst children in developing countries. This is due to

under-diagnosis of paediatric cancers in these countries and a lack of simple measures for recording and capturing patient-related data, such as cancer registries.⁵ As in most developing countries, South African children with WT face many challenges that can affect prognosis. These include delayed presentation, high rates of treatment abandonment and poverty.⁶⁻⁸ The strongest prognostic indicators amongst this population are considered to be disease stage and histology.⁹

Studies have shown that 45-77% of children with cancer in developing countries are malnourished on admission to hospital.¹⁰⁻¹² Malnutrition amongst this population can lead to higher mortality and treatment abandonment rates,¹⁰ and increased complication risks.¹³ Some risk factors for a poor outcome amongst children with cancer, such as age, are not modifiable. Yet, malnutrition is a risk factor that can be modified if appropriate steps are taken.¹⁴

Abbreviations: AHOPCA, Asociacion de Hemato-Oncología Pediátrica de Centro America; BMI, body mass index; HFA, height for age; IALCH, Inkosi Albert Luthuli Central Hospital; KZN, KwaZulu-Natal; MUAC, mid-upper arm circumference; NG, nasogastric; TSFT, triceps skinfold thickness; WFA, weight for age; WFH, weight for height; WFL, weight for length; WT, Wilms tumour

The assessment of nutritional status of children with cancer needs to be standardised amongst centres treating these patients.¹⁵ When weight and height are used in isolation, malnutrition in children with WT can be masked^{12,16,17} due to tumour mass and fluid shifts.¹⁸ In addition, utilisation of serum albumin in isolation as an indicator of nutritional status can lead to underestimation of malnutrition, as albumin levels can be affected by several disease-related factors.¹⁷ Many studies have used weight, height and body mass index (BMI) or albumin levels in isolation to classify nutritional status.^{17,19} It can, therefore, be posited that malnutrition rates amongst WT patients could have been under-reported, leading to a significant underestimation of the actual prevalence of malnutrition amongst this population to date.

A growing body of research has emphasised the importance of including arm anthropometry measurements, specifically mid-upper arm circumference (MUAC) and triceps skinfold thickness (TSFT), as part of the classification criteria^{10,12,17,20} as these measurements are independent of tumour size.^{21,22} Utilisation of biochemistry (namely albumin) and anthropometry measurements *in combination* has been shown to improve the sensitivity of classification of nutritional status.¹⁰

When assessing the influence of nutritional status at the time of admission on outcome, there is a paucity of data relating to children with cancer, particularly WT. Although some studies have found a significant correlation between malnutrition on admission and poorer outcome,^{10,23,24} others have failed to show any significant link.^{25–27} Of concern regarding previous studies assessing this relationship is that many children may have been incorrectly classified as being well-nourished when in fact they were malnourished.^{12,16,17}

The purpose of this study was to determine the prevalence of malnutrition amongst children with WT on admission to Inkosi Albert Luthuli Central Hospital (IALCH). Furthermore, it aimed to determine the relationship between nutritional status on admission and outcome, and the level of nutritional intervention that the subjects received within the first 2 weeks of admission.

2 | METHODS

This prospective observational study utilised a cohort study design amongst children with WT at IALCH between 2004 and 2012. The study design was approved by the Biomedical Research Committee of the University of KwaZulu-Natal. IALCH is classified as a National Central Hospital and services a large catchment area spanning two of South Africa's nine provinces (KwaZulu-Natal [KZN] and Eastern Cape). It is a referral hospital for all tertiary and district hospitals in the KZN province²⁸ and has a catchment area of over 10 million people.²⁹

The study population included 88 children aged 6 months–13 years diagnosed with WT and admitted to the paediatric surgical ward. The files of 12 children had to be excluded (anthropometric data of two children were collected after the initiation of nutritional intervention, while the medical files of 10 children could not be located, leading to incomplete data), therefore 76 children made up the final sample for analysis.

2.1 | Anthropometry and biochemistry

Between 2004 and 2012, subjects were assessed by the ward dietitian after admission. Anthropometric measurements were taken before any interventions in the form of nutritional supplementation or chemo-/radiotherapy.

Nutritional status was assessed using weight, height, MUAC and TSFT measurements. Body weight and height were measured while subjects wore light clothing and no shoes using a Nagata electronic weight and height measurement scale (model BW-1122H). Weight was recorded to the nearest 100 g and height and length to the nearest 1 cm. Height measurements were taken while subjects stood straight on the electronic scale with their back against the height meter, looking forward. Infants were weighed on a Nagata electronic baby scale (model BW-20) and their length measured with a measuring tape while supine. These scales are calibrated annually by the manufacturer in order to maintain accuracy. Excised tumour weight was recorded by the paediatric surgeon post-operatively and the corrected weight was utilised for statistical analysis.

In order to measure MUAC, the midpoint between the acromion process of the scapula and olecranon process of the ulna was determined while the left forearm was bent at a right angle. A non-stretchable measuring tape was used to measure the circumference at the midpoint while the arm hung straight down to the nearest 0.1 cm. TSFT was measured on the same midpoint using a Harpenden skinfold calliper and recorded to the nearest 0.1 cm. To ensure accuracy, measurements were performed in triplicate and the average recorded.

Weight and height measurements were utilised to determine weight for age (WFA), length for age/height for age (HFA), weight for length/weight for height (WFH) and BMI depending on the subject's age. These were classified according to age- and gender-matched norms utilising the STAT GrowthCharts app,³⁰ which classifies subjects according to Z-scores using the WHO growth charts. MUAC and TSFT measurements were compared with age- and gender-matched norms and interpreted using the Frisancho percentile charts.³¹ Albumin levels were routinely assessed on admission to hospital.

2.2 | Classification of nutritional status

A combination of arm anthropometry and albumin were used to classify subjects. Subjects who met at least one of the following criteria were classified as malnourished:

- (1) MUAC and TSFT between the 5th and 10th percentiles.
- (2) MUAC or TSFT below the 5th percentile.
- (3) Albumin less than 35 g/l.

Those with a MUAC and/or TSFT below the 5th percentile, or albumin levels below 32 g/l, were classified as being severely malnourished. This combination of criteria, which have been modified and utilised in previous studies,^{17,32} was adapted from the guidelines developed at the Asociacion de Hemato-Oncologia Pediatrica de Centro America (AHOPCA) congress in 2004.³³

TABLE 1 Age and anthropometric characteristics of the study population

	Mean	Median	IQR	Min	Max	SD
Age (months)						
Combined (n = 76)	56	47	37	11	149	35
Males (n = 38)	57	47	32	11	149	35
Females (n = 38)	55	46	49	14	147	35
Weight (kg)						
Combined (n = 76)	16.8	15.3	5.7	7.5	43.2	7.0
Males (n = 38)	16.8	15.5	4.6	8.5	34.6	6.0
Females (n = 38)	17.0	14.6	7.9	7.5	43.2	7.9
Height (m)						
Combined (n = 75)	1.0	1.0	0.2	0.6	1.6	0.2
Males (n = 37 ^a)	1.0	1.0	0.2	0.6	1.5	0.2
Females (n = 38)	1.0	1.0	0.3	0.7	1.6	0.2
BMI (kg/m²)						
Combined (n = 75)	15.7	15.9	3.1	10.3	23.9	2.3
Males (n = 37 ^a)	15.3	15.3	3.6	12.7	20.1	1.8
Females (n = 38)	16.2	16.7	2.5	10.3	23.9	2.7

^aHeight was not available for one of the subjects; SD = Standard deviation.

2.3 | Level of nutritional support

At the time of intervention, the dietitian recorded any nutritional support that subjects received in their electronic files. Early intervention was considered as dietary intervention that took place within 2 weeks of admission. Data relating to both the assessment and nutritional intervention type (oral, enteral or parenteral) were documented.

2.4 | Data analysis

To ensure confidentiality, anthropometric measurements were only accessible to the Dietetics Department at IALCH. Outcome data were obtained using electronic patient records recorded by the doctor. The data were analysed using the Statistical Package for Social Sciences version 21. Descriptive tests as well as the χ^2 test of independence were conducted. A *P*-value <0.05 was considered significant.

3 | RESULTS

The male and female distribution amongst the sample of 76 children with WT was exactly half (50%; n = 38). Most consisted of black African children (97.3%; n = 74), while 2.6% (n = 2) comprised white and coloured children. The mean age of the subjects was 4 years, 8 months (SD \pm 2.89) and the median age was 3 years, 10 months (Interquartile ranges (IQR) = 37.32 months). Anthropometric characteristics are presented in Table 1. Median height for males and females was 0.98 m (IQR = 0.23), while on average females weighed slightly more than males (16.99 kg vs. 16.56 kg). Consequently, the mean female BMI [16.17 kg/m² (SD \pm 2.69)] was higher than the males [15.30 kg/m² (SD \pm 1.83)]. The mean albumin level was 38.61 g/l (SD \pm 5.87).

Table 2 shows malnutrition prevalence according to various anthropometric measurements. According to WFA, HFA and BMI results,

most of the sample was classified as well nourished. However, according to arm anthropometry measurements, the malnutrition prevalence was much higher.

Figure 1 shows a comparison between different combinations of measurements when classifying nutritional status. When albumin was utilised to classify nutritional status in isolation, 70% (n = 53) of subjects were well nourished. In this study, a combination of MUAC, TSFT and albumin was used to classify overall nutritional status. The results showed that two-thirds (66.67%; n = 48) of the sample (n = 72) were malnourished on admission to hospital.

3.1 | Stage of disease and nutritional status

Neoadjuvant chemotherapy was offered to all patients. Patients without metastatic disease received 4 weeks of actinomycin and vincristine, while those with metastases received additional epi-adriamycin over 6 weeks. Post-operative chemotherapy was determined by the operative staging and histology using Societe Internationale d'Oncologie Pediatrique (International Society of Pediatric Oncology) guidelines, with radiotherapy offered to patients with stages III and IV disease. While most of the patients presented in stages III–V (71%; n = 54), 29% presented in stage I or II (n = 22). The influence of stage of disease on nutritional status was analysed using the χ^2 test of independence. This analysis showed a trend towards increased malnutrition prevalence amongst those in stages II–IV (Table 3). Those in stage V had a lower malnutrition prevalence compared to stages II–IV. Overall, the influence of stage of disease on nutritional status was not significant.

3.2 | Nutritional status and outcome

The influence of nutritional status upon admission and outcome after 2 years is presented in Tables 4 and 5. Overall, nutritional status could

TABLE 2 Anthropometric classification of study population

	Female		Male		Combined		Total %
	n	%	n	%	n	%	
WFA (n = 71 ^a)							
Normal	25	54.35	21	45.65	46	64.79	NW
Underweight, mild (<-1 SD)	9	56.25	7	43.75	16	22.53	64.79
Underweight, moderate (<-2 SD)	3	37.50	5	62.50	8	11.27	UW
Underweight, severe (<-3 SD)	0	0.00	1	100.00	1	1.41	35.21
HFA (n = 75 ^b)							
Normal	28	60.87	18	39.13	46	61.33	NW
Underweight, mild (<-1 SD)	9	45.00	11	55.00	20	26.67	61.33
Underweight, moderate (<-2 SD)	1	16.67	5	83.33	6	8.00	UW
Underweight, severe (<-3 SD)	0	0.00	3	100.00	3	4.00	38.67
BMI (n = 75 ^b)							
Normal	25	47.17	28	52.83	53	70.67	NW
Underweight, mild (<-1 SD)	10	76.92	3	23.08	13	17.33	70.67
Underweight, moderate (<-2 SD)	2	40.00	3	60.00	5	6.67	UW
Underweight, severe (<-3 SD)	1	25.00	3	75.00	4	5.33	29.33
MUAC (n = 74 ^c)							
Normal	10	66.7	5	33.3	15	21.1	NW
Underweight, mild (<-1 SD)	8	44.4	10	55.5	18	24.3	21.2
Underweight, moderate (<-2 SD)	5	55.5	4	44.4	9	12.2	UW
Underweight, severe (<-3 SD)	15	46.9	17	53.1	32	43.2	79.7
TSFT (n = 71 ^d)							
Normal	12	60.0	8	40.0	20	28.2	NW
Underweight, mild (<-1 SD)	4	28.6	10	71.4	14	19.7	28.2
Underweight, moderate (<-2 SD)	8	57.1	6	42.9	14	19.7	UW
Underweight, severe (<-3 SD)	12	52.2	11	47.8	23	32.4	71.8

^aFive subjects could not be classified due to insufficient age specific growth charts - WHO WFA growth charts not available for children over the age of ten years.

^bHeight information not available for one subject.

^cMUAC not available for two subjects.

^dTSFT not available for five subjects.

NW = Normal Weight; UW = Underweight.

not be classified for five subjects as their TSFT measurement was not recorded. Outcome was defined as the subject surviving or dying. It should be noted that the outcome of 14 of the subjects could not be described as they were lost to follow up or did not return for scheduled follow-up visits 2 years post-admission. The rate of treatment abandonment was 2.6%.

Two-thirds (n = 41) of the 62 subjects were classified as being malnourished according to their MUAC, TSFT and albumin results. Seventeen (27.42%) of the 62 subjects died during the 2-year follow-up period. Of those who died, 13 (76.47%) were malnourished on admission. Of the 45 subjects who survived, 28 (62.22%) were malnourished on admission. There was no significant relationship between nutritional status on admission and outcome after 2 years.

No significant relationship was found between malnutrition severity and outcome. Table 5 shows that on admission both well-nourished and severely malnourished subjects were equally prevalent in the group who survived.

Binary logistic regression analysis was conducted to account for confounding variables known to have a significant influence of prognosis (stage and histology). No significant relationship between nutritional status and outcome was found, despite correcting for these variables ($P = 0.711$).

3.3 | Level of nutritional support received

Most of the sample (84.2%; n = 64) received nutritional supplementation within 2 weeks of admission. Just over 10% (n = 8) received supplements more than 2 weeks after admission and were not included. Only 5.3% (n = 4) of the sample did not receive any nutritional supplementation. Of those who received supplementation, 87.5% (n = 56) received oral supplementation, which refers to a nutritional supplement, a nutritious snack (e.g. yoghurt or a sandwich) or a combination of the two. Of the remaining subjects, 9.4% (n = 6) received nasogastric (NG) feeds, whereas 3.1% (n = 2) received a combination of oral and




FIGURE 1 Prevalence of malnutrition using biochemical and anthropometric measurements. Y axis: number of subjects; X axis: well nourished (first three bars) and malnourished (second three bars). Key: Blue bar represents TSFT + MUAC + albumin ($n = 72$); yellow bar represents TSFT + MUAC ($n = 72$); green bar represents albumin ($n = 76$).

NG supplementation. There was no significant relationship between malnutrition severity and whether or not the subjects received supplements ($P = 0.768$). Furthermore, there was no significant relationship between malnutrition severity and type of nutritional support received ($P = 0.620$).

4 | DISCUSSION

To date, there is no published research assessing the prevalence of malnutrition amongst South African children with WT utilising arm anthropometry measurements.

This study found the prevalence of malnutrition to be just over 66%. Previous studies in developing countries have found malnutrition prevalence amongst children with cancer ranging from 45% to 77%.^{10–12,20} Sala et al. conducted a large study that included

1,787 children with cancer from seven developing countries in Central America. The authors used similar categories to classify nutritional status to those in this study. Malnutrition was found to be present in 63.8% of the sample while 72.4% of those with solid tumours were malnourished.

A comprehensive search of the literature found four studies assessing the prevalence of malnutrition amongst South African children with WT.^{26,34–36} Wessels et al. found that 35% of their sample was malnourished, whereas Davidson et al. and Visser et al. found the prevalence of malnutrition to be 20.7% and 10.3%, respectively. Holzinger et al., who conducted their study at IALCH, found a prevalence of 45%. This study was performed from 2002 to 2005 and did not include the same patient population as the present study. Unfortunately, these studies were limited by the fact that they used weight and height in isolation and did not include arm anthropometry in their classification. Wessels et al. did, however, try to compensate for this limitation by classifying any subject with a WFH less than 90% of the expected value as poorly nourished. A final limitation of the studies by Wessels et al. and Holzinger et al. (2007) was that they utilised relatively small sample sizes (59 and 37, respectively).

Assessment of the nutritional status of children with solid tumours is problematic due to the effect of tumour weight on body weight measurements.¹⁸ Although corrected weight was used in this sample (admission weight minus excised tumour weight), tumour weight would have shrunk from the time of admission due to neoadjuvant treatment. When comparing the prevalence of malnutrition according to WFA, HFA and BMI to MUAC and TSFT measurements in this study, the lack of correlation between these different types of anthropometric measurements was striking. It is concerning that the prevalence of severe malnutrition remained below 6% when arm anthropometry was not utilised, even after correcting for excised tumour weight, and increased to over 43% and 32% when MUAC and TSFT were assessed, respectively. The results of this study are in agreement with several other studies that showed that utilising body weight measurements for classification of nutritional status amongst children with cancer,

TABLE 3 The association between stage of disease and nutritional status

	Stage 1 ($n = 5$)	Stage 2 ($n = 13$)	Stage 3 ($n = 19$)	Stage 4 ($n = 29$)	Stage 5 ($n = 6$)	Total ($n = 72$) ^a	<i>p</i> -value
Yes, <i>n</i> , (%)	1 (20)	9 (69)	13 (68)	22 (76)	3 (50)	48 (67)	$p = 0.152$
No, <i>n</i> (%)	4 (80)	4 (31)	6 (32)	7 (24)	3 (50)	24 (33)	

^aFour subjects did not have TSFT therefore their overall nutritional status could not be classified.

TABLE 4 The association between malnutrition on admission and outcome (2 years)^a

	Outcome				Total	
	Dead		Alive			
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
Malnourished						
Yes	13	32	28	68	41	100
No	4	19	17	81	21	100
Total	17	27	45	73	62	100

^a χ^2 test, $p = 0.443$.

TABLE 5 The association between severity of malnutrition and outcome^a

Outcome	Degree of malnutrition						Total	
	Normal		Malnourished		Severely malnourished			
	n	%	n	%	n	%	n	%
Dead	4	25	6	37.5	6	37.5	16	100.00
Alive	17	37	12	26	17	37	46	100.00
Total	21	34	18	29	23	37	62	100.00

^a χ^2 test, $p = 0.163$.

without including arm anthropometry measurements, often leads to an underestimation of the prevalence of malnutrition.^{12,17,20}

The AHOPCA algorithm incorporates arm anthropometry, albumin, weight loss and percentage of ideal body weight measurements. Initially, the algorithm stated that both MUAC and TSFT had to be below a certain value for a subject to be classified as malnourished. However, to increase the sensitivity of the algorithm and to encourage interventions amongst a larger proportion of subjects these criteria have since changed, stipulating that all subjects with either MUAC or TSFT below a certain level be classified as malnourished.¹⁰ Because many studies in the past assessed the prevalence of malnutrition amongst children with solid tumours using weight and height measurements in isolation,^{19,25,26} it is possible that the prevalence of malnutrition amongst this population has been grossly underestimated.

The results from this study showed that, in isolation, albumin considerably underestimated the prevalence of malnutrition (30%). This is in agreement with previous research by Tazi et al., which showed that malnutrition prevalence according to albumin was 28%. However, when the authors classified nutritional status according to arm anthropometry, the prevalence of malnutrition increased to 39% and 50% using MUAC and TSFT, respectively. When including albumin in combination with MUAC and TSFT in the current study, the prevalence of malnutrition increased by almost 10%. Because of its low specificity, albumin is considered to be a poor indicator of nutritional status when assessed in isolation.³⁷ However, previous research has shown the inclusion of albumin as one of a combination of criteria when classifying nutritional status does add considerable value to the accuracy and sensitivity of the classification methodology.¹⁰

Nutritional status was not shown to be significantly related to outcome, although there was a trend towards poorer outcome amongst malnourished children. The correlation between nutritional status on admission and outcome has not been conclusively established in previous studies.

Most research has focused on analysing this relationship amongst children with haematological malignancies, with few studies focusing on children with solid tumours. A large study by Hoffmeister et al.³⁸ analysed 733 American children with haematological malignancies who had undergone a haematopoietic stem cell transplant. They found those with an arm muscle area below the 5th percentile had significantly worse event-free survival at 100 days and 3 years post-transplant. Hoffmeister's study was strong as it had a large sample size and utilised arm anthropometry when classifying nutritional status. In

contrast, a United Kingdom study with a large sample of 1,025 children with haematological malignancies found no evidence that nutritional status on admission was a prognostic indicator of outcome. While this sample size was substantial, arm anthropometry was not used when classifying nutritional status.²⁷

Results from previous studies amongst children with solid tumours are similarly inconclusive. The strengths of the aforementioned study conducted by Sala et al. included its large sample size, accurate classification of nutritional status and multiple locations from which children were sampled. The authors found a significant relationship between malnutrition and higher mortality rates amongst children with solid tumours, but not haematological malignancies.¹⁰ Similar studies found no significant relationship between these variables; however, none of these studies took arm anthropometry into account.^{19,25,26}

Late presentation and advanced disease stage on admission can have extensive consequences, as children with larger tumours on admission have been shown to be more severely malnourished than those with smaller tumours.¹¹ This, in turn, can negatively affect their outcome¹⁰ and can contribute to poorer prognosis.⁹

In this study, nutritional interventions within 2 weeks of admission were implemented for more than 80% of the study sample. This included an initial assessment of nutritional status and implementation of oral and/or NG supplements.

Children with cancer often struggle to meet their nutritional requirements,³⁹ making nutritional intervention extremely important. Nutritional intervention has been shown to lead to effective weight gain amongst children with cancer.^{32,35,39–41} Wessels et al. emphasised the importance of utilising clinical judgement in addition to anthropometric assessment in order to determine which patients are in need of nutritional intervention. When treating these patients, the challenge faced by the dietitian is to prevent further deterioration of nutritional status and to attempt to reverse any deterioration in nutritional status that may have already occurred. Children with solid tumours in particular are often in need of nutritional interventions as they practice food restrictions more often and may experience significantly worse eating problems including nausea, vomiting and loss of appetite.²¹

An international study that assessed the nutrition-related practices amongst 125 institutions found inconsistent practices related to nutritional interventions.⁴² Similar results were demonstrated by a survey of several South African hospitals.⁴³ It is unfortunate that there are no standardised guidelines related to nutritional intervention that can be implemented throughout centres that treat children with cancer, as the

efficacy of nutritional intervention is well documented.^{32,39–41} The fact that there was no significant relationship between severity of malnutrition and type of support received suggests that both oral nutritional supplements as well as NG feeds have a role in the nutritional management of malnourished children with cancer.

The greatest weight changes amongst children with cancer have been shown to occur within 3 months of diagnosis,⁴⁴ thus emphasising the importance of early and aggressive nutritional intervention. A study by Antillon et al. looking at children with leukaemia showed that multidisciplinary management, in the form of a combination of nutritional intervention as well as chemotherapy treatment, contributed to improved nutritional status after 6 months. Furthermore, of those who were classified as being severely depleted on admission, the nutritional status of almost two-thirds of the subjects improved. For the subjects who were severely malnourished but for whom an improvement in nutritional status was seen by 6 months, the overall survival probability at 5 years was similar to those whose nutritional status was adequate on admission.³²

A similar study by Orgel et al. found that children with leukaemia who were underweight for more than 50% of their treatment period had significantly worse outcomes. Those who were initially underweight but who gained weight and maintained their normal weight for more than 50% of treatment had a similar risk for recurrence as those who maintained a normal weight throughout. Therefore, nutritional monitoring is imperative amongst this population, and nutritional intervention should be implemented as early as possible. By reducing the time that children are classified as being malnourished during their treatment, the potential negative effects of poor nutritional status on admission can be significantly reduced.¹⁴

4.1 | Study limitations

A limitation of this study was the sample size, which was relatively small compared to other international studies. This study may be subject to selection bias due to the fact that it was conducted at a government hospital. However, IALCH serves a catchment area spanning two provinces in South Africa and is therefore representative of a wide spectrum of the South African population. The use of the Frisancho charts to classify nutritional status in this study may have introduced some bias as these charts are based on a population made up of only Caucasian subjects.

5 | CONCLUSION

Nutritional status should be assessed as early as possible in order to improve nutritional status or prevent malnutrition in children with WT. Utilisation of a combination of arm anthropometry and albumin provides accurate criteria for assessing nutritional status amongst children with WT.

Malnutrition on admission did not contribute to significantly poorer outcomes. Aggressive early nutritional assessment, intervention and management of these children may help to reduce the negative impact that malnutrition on admission may have on outcome.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

REFERENCES

1. Ward E, DeSantis C, Robbins A, Kohler B, Jemal A. Childhood and Adolescent Cancer Statistics, 2014. *Cancer J Clin*. 2014;64:83–103.
2. Stack M, Walsh PM, Comber H, Ryan CA, O'Lorcain P. Childhood cancer in Ireland: A population-based study. *Arch Dis Child*. 2007;92:890–897.
3. Stefan DC. Patterns of distribution of childhood cancer in Africa. *J Trop Pediatr*. 2015;61(3):165–173
4. Stefan DC, Stones DK. The South African Paediatric Tumour Registry—25 Years of activity. *South Afr Med J*. 2012;102(7):605–606.
5. Stefan DC, Baadjies B, Kruger M. Incidence of childhood cancer in Namibia: The need for registries in Africa. *Pan Afr Med J*. 2014;17:191.
6. Paintsil V, David H, Kambugu J, et al. The collaborative Wilms tumour Africa Project; baseline evaluation of Wilms tumour treatment and outcome in eight institutes in sub-Saharan Africa. *Eur J Cancer*. 2015;51(1):84–91.
7. Wilde JCH, Lameris W, van Hasselt EH, et al. Challenges and outcome of Wilms' tumour management in a resource-constrained setting. *Afr J Paediatr Surg*. 2010;7(3):159–162.
8. Abuidris DO, Elimam ME, Nugud FM, Elgaili EM, Ahmed ME, Arora RS. Wilms tumour in Sudan. *Pediatr Blood Cancer*. 2008;50:1135–1137.
9. Kaste SC, Dome JS, Babyn PS, et al. Wilms tumour: Prognostic factors, staging, therapy and late effects. *Paediatr Radiol*. 2008;38:2–17.
10. Sala A, Rossi E, Antillon F, et al. Nutritional status at diagnosis is related to clinical outcomes in children and adolescents with cancer: A perspective from Central America. *Eur J Cancer*. 2012;48:243–252.
11. Israels T, Borgstein E, Jamali M, de Kraker J, Caron HN, Molyneux EM. Acute malnutrition is common in Malawian patients with a Wilms tumour: A role for peanut butter. *Pediatr Blood Cancer*. 2009;53:1221–1226.
12. Israels T, Chirambo C, Caron HN, Molyneux EM. Nutritional status at admission of children with cancer in Malawi. *Pediatr Blood Cancer*. 2008;51:626–628.
13. Murry DJ, Riva L, Poplack DG. Impact of nutrition on pharmacokinetics of anti-neoplastic agents. *Int J Cancer*. 1998;11:48–51.
14. Orgel E, Sposto R, Malvar J, et al. Impact on survival and toxicity by duration of weight extremes during treatment for pediatric acute lymphoblastic leukemia: A report from the Children's Oncology Group. *J Clin Oncol*. 2014;32(13):1331–1337.
15. Rogers PC, Melnick SJ, Ladas EJ, Halton J, Baillargeon J, Sacks N. Children's Oncology Group (COG) nutrition committee. *Pediatr Blood Cancer*. 2008;50:447–450.
16. Lemos Pdos SM, de Oliveira FLC, Caran EMM. Nutritional status of children and adolescents at diagnosis of hematological and solid malignancies. *Rev Bras Hematol Hemoter*. 2014;36(6):420–423.
17. Tazi I, Hidane Z, Zafad S, Harif M, Benchekroun S, Ribeiro R. Nutritional status at diagnosis of children with malignancies in Casablanca. *Pediatr Blood Cancer*. 2008;51:495–498.
18. Rogers PCJ. Nutritional status as a prognostic indicator for pediatric malignancies. *J Clin Oncol*. 2014;32(13):1293–1294.
19. Burke ME, Lyden ER, Meza JL, et al. Does body mass index at diagnosis or weight change during therapy predict toxicity or survival in intermediate risk rhabdomyosarcoma? A Report From the Children's Oncology Group Soft Tissue Sarcoma Committee. *Pediatr Blood Cancer*. 2013;60:748–753.

20. Garofolo A, Lopez FA, Petrilli AS. High prevalence of malnutrition among patients with solid non-hematological tumors as found by using skinfold and circumference measurements. *Sao Paulo Med J*. 2005;123(6):277–281.
21. Tah PC, Nik Shanita S, Poh BK. Nutritional status among pediatric cancer patients: A comparison between hematological malignancies and solid tumors. *J Spec Pediatr Nurs*. 2012;17:301–311.
22. Brennan BMD, Gill M, Pennells L, Eden OB, Thomas AG, Clayton PE. Insulin-like growth factor I, IGF binding protein 3, and IGFBP protease activity: relation to anthropometric indices in solid tumours or leukaemia. *Arch Dis Child*. 1999;80:226–230.
23. Mejia-Arangure JM, Fajardo-Gutierrez A, Reyes-Ruiz NI, et al. Malnutrition in childhood lymphoblastic leukemia: A predictor of early mortality during the induction-to-remission phase of the treatment. *Arch Med Res*. 1999;30:150–159.
24. Viana MB, Murao M, Ramos G, et al. Malnutrition as a prognostic factor in lymphoblastic leukaemia: A multivariate analysis. *Arch Dis Child*. 1994;71:304–310.
25. Pedrosa F, Bonilla M, Liu A, et al. Effect of malnutrition at the time of diagnosis on the survival of children treated for cancer in El Salvador and Northern Brazil. *J Pediatr Hematol/Oncol*. 2000;22(6):502–505.
26. Wessels G, Hesseling PB, Van Ommeren KH, Boonstra V. Nutrition, morbidity, and survival in South African children with Wilms' tumor. *Pediatr Hematol Oncol*. 1999;16:321–327.
27. Weir J, Reilly JJ, McColl JH, Gibson BES. No evidence for an effect of nutritional status at diagnosis on prognosis in children with acute lymphoblastic leukemia. *J Pediatr Hematol/Oncol*. 1998;20(6):534–538.
28. Department of Health. The hospital Dietetic Department: A guide to the management of the Dietetic Department and Dietetic Patient Care. Circular Minute Number G8/2014. Pietermaritzburg: KZN Health; 2014.
29. Statistics South Africa. Census 2011. Statistical release (Revised) P0301.4. <http://www.statssa.gov.za/publications/P03014/P03014-2011.pdf>. Accessed June 30, 2015.
30. Austin Physician Productivity, LLC. STAT GrowthCharts™ WHO Lite. (iPhone application). Vers. 1.3. 2012. <https://itunes.apple.com/us/app/stat-growthcharts-who-lite/id384332193?mt=8>. Accessed on May 20, 2014.
31. Frisancho AR. New norms of upper limb fat and muscle areas for assessment of nutritional status. *Am J Clin Nutr*. 1981;34:2540–2545.
32. Antillon F, Rossi E, Molina AL, et al. Nutritional status of children during treatment for acute lymphoblastic leukemia in Guatemala. *Pediatr Blood Cancer*. 2013;60:911–915.
33. Sala A, Antillon F, Pencharz P, Barr R, AHOPCA Consortium. Nutritional status in children with cancer: A report from the AHOPCA workshop held in Guatemala City, August 31-September 5, 2004. *Pediatr Blood Cancer*. 2005;45(2):230–236.
34. Visser YT, Uys R, van Zyl A, Stefan DC. Nephroblastoma—A 25-year review of a South African unit. *J Med Life*. 2014;7(3):445–449.
35. Holzinger TT, Shaik AS, Hadley GP. The role of nutritional intervention in children with nephroblastoma. *South Afr J Clin Nutr*. 2007;20(3):96–99.
36. Davidson A, Hartley P, Desai F, Daubenton J, Rode H, Millar A. Wilms tumour experience in a South African centre. *Pediatr Blood Cancer*. 2006;46:465–471.
37. Fuhrman MP. The albumin-nutrition connection: Separating myth from fact. *Nutrition*. 2002;18:199–200.
38. Hoffmeister PA, Storer BE, Macris PC, Carpenter PA, Baker KS. Relationship of body mass index and arm anthropometry to outcomes after pediatric allogeneic hematopoietic cell transplantation for hematologic malignancies. *Biol Blood Marrow Transplant*. 2013;19:1081–1086.
39. Barbosa JM, Pedrosa F, Cabral PC. Nutritional status and adequacy of enteral nutrition in pediatric cancer patients at a reference center in northeastern Brazil. *Nutr Hosp*. 2012;27(4):1099–1105.
40. De Waele E, Mattens S, Honore P, Spapen H, De Greve J, Pen JJ. Nutrition therapy in cachectic cancer patients. The Tight Caloric Control (TiCaCo) pilot trial. *Appetite*. 2015;91:298–301.
41. Sacks N, Hwang WT, Lange BJ, et al. Proactive enteral tube feeding in pediatric patients undergoing chemotherapy. *Pediatr Blood Cancer*. 2014;61:281–285.
42. Ladas EJ, Sacks N, Brophy P, Rogers PC. Standards of nutritional care in pediatric oncology: Results from a nationwide survey on the standards of practice in pediatric oncology. A Children's Oncology Group Study. *Pediatr Blood Cancer*. 2006;46:339–344.
43. Schoeman J. Dietetic survey: Paediatric oncology unit [Unpublished survey]. 2011.
44. Brinksma A, Roodbol PF, Sulkers E, et al. Changes in nutritional status in childhood cancer patients: A prospective cohort study. *Clin Nutr*. 2015;34:66–73.

How to cite this article: Lifson LF, Hadley GP, Wiles NL, Pillay K. Nutritional status of children with Wilms' Tumour on admission to a South African hospital and its influence on outcome. *Pediatr Blood Cancer*. 2017;64:e26382. DOI: 10.1002/pbc.26382