

Author Manuscript

Title: Heteroatom Tuning of Bimolecular Criegee Reactions and Its Implications

Authors: Joseph S. Francisco; Manoj Kumar

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record.

To be cited as: 10.1002/anie.201604848

Link to VoR: <http://dx.doi.org/10.1002/anie.201604848>

Heteroatom Tuning of Bimolecular Criegee Reactions and Its Implications

Manoj Kumar^[a] and Joseph S. Francisco^{*[a]}

Abstract: High-level quantum chemical calculations have been performed to understand the key reactivity determinants of bimolecular Criegee + H₂X (X = O, S, Se, and Te) reactions. Criegee intermediates are implicated as key intermediates in atmospheric, synthetic organic and enzymatic chemistry. Generally, it is believed that the nature and location of substituent at the Criegee carbon plays a key role in determining the reactivity. However, the present work suggests that it is not only the Criegee substitution, but the nature of heteroatom in H₂X that also plays a crucial role in determining the reactivity of the Criegee-H₂X interaction. The barriers for the Criegee-H₂X reactions satisfy an inverse correlation with the bond strength of X-H in H₂X, and a direct correlation with the first pK_a of H₂X. This heteroatom tuning causes a substantial barrier lowering of 8-11 kcal/mol in the Criegee reaction barrier in going from H₂O to H₂Te. The reactions of sulfur dioxide and sulfur trioxide with H₂X also satisfy similar correlations, indicating that the heteroatom-induced barrier tuning could be a ubiquitous reactivity determinant in addition reactions. An important implication of these results is that the Criegee-H₂S reaction could be a source of thioaldehydes, which are important in planetary atmospheres and synthetic organic chemistry. By performing the Criegee-H₂S reaction under water or acid catalysis, thioaldehydes could be detected in a hydrogen-bonded complexed state, which is significantly more stable than their uncomplexed form. As a result, simpler aliphatic thioaldehydes could be selectively synthesized in laboratory, which, otherwise, has been a significant synthetic challenge because of their ability to oligomerize.

Understanding Criegee chemistry has become a hot topic of research in the recent past. Criegee intermediates are carbonyl oxides that are principally produced in the ozone-olefin cycloadditions.¹ The unimolecular and bimolecular Criegee reactions play a crucial role in the tropospheric budgets of hydroxyl radicals, organic acids, hydroperoxides, nitrates, sulfates and secondary organic aerosols.²⁻⁹ At the same time, Criegee chemistry is a key component of ozonolysis-based syntheses that offer safe and scalable routes for synthesizing pharmaceutical intermediates and other useful commodities.¹⁰⁻¹² Criegee intermediates are also implicated in the reaction cycles of flavin-dependent Baeyer-Villiger monooxygenases¹³ that provide an environmentally benign route for forming enantiopure drugs.¹⁴

Because of its broad profile, factors influencing Criegee reactivity has been extensively investigated by means of experimental and theoretical means. The nature and location of the substituents on the Criegee intermediate significantly tunes its reactivity.¹⁵ Criegee intermediates having a methyl substituent in the *anti* position react faster than Criegee intermediates having a methyl substituent in *syn* position. Criegee intermediates with electron withdrawing substituents and zwitterionic character react faster than those having electron releasing substituents. Substitution in the Criegee intermediate can cause up to ten orders of

magnitude differences in its reactivity. The unimolecular and bimolecular reactions of the Criegee intermediate CH₃CHOO represent one of the well-studied conformer-dependent Criegee reactivity profile.

Criegee intermediates participate in various bimolecular reactions. Though the effect of Criegee substitution in bimolecular reactions has been well explored, the effect of substituent in the non-Criegee reacting partner is less so. For example, the most probable bimolecular Criegee reaction in troposphere is the reaction with water. However, it is not clear whether the nature of heteroatom X in H₂X (X = O, S, Se, and Te) would alter the energetics of the Criegee-H₂X reaction i.e., does there exist any correlation between the Criegee reaction barrier and the heteroatom nature? If yes, what are its implications? If not, what other factors influence the Criegee-H₂X reaction? A detailed knowledge of these determinants may provide general guidelines for synthesizing useful organic compounds, which are otherwise difficult to synthesize, and in addition, may also reveal new source of atmospherically important thioaldehydes, thus improving molecular details of existing atmospheric models.

Here it is shown using high-level quantum chemical calculations that such a tuning of the barrier for the Criegee-H₂X reaction is indeed possible and more importantly, this heteroatom tuning is also found to be a general trend among other atmospherically important reactions. Firstly, examining the gas-phase reaction between the simplest Criegee intermediate, CH₂OO and H₂X; the reaction energetics are computed by performing single point calculations using the coupled cluster single and double substitution method with a perturbative treatment of triple excitations (CCSD(T)) and the augmented correlation-consistent triplet zeta basis set, aug-cc-pVTZ at the M06-2X/aug-cc-pVTZ optimized geometries. (See Supporting Information for details). The CH₂OO-H₂X reaction is a multi-step reaction that converts the Criegee moiety, -COO into -C=X functional group. The reaction consists of two main steps namely chalcogen hydride (H₂X) addition and hydrogen peroxide (H₂O₂) elimination with the latter step being its rate-limiting step (**Figure 1** and **Table S1**). Overall, the reaction is at least 24.1 kcal/mol exoergic. For the CH₂OO-H₂O reaction, the addition step has a barrier of 9.2 kcal/mol relative to **Int**₁ whereas the elimination step has a barrier of 49.1 kcal/mol relative to **Int**₃, H₂C(OH)(OOH). These barriers are significantly reduced upon changing the heteroatom X in H₂X. For the CH₂OO-H₂S reaction, the




Figure 1. CCSD(T)/aug-cc-pVTZ/M06-2X/aug-cc-pVTZ calculated zero-point-corrected reaction profiles for the reaction between the simplest Criegee intermediate, CH₂OO and various chalcogen hydrides, H₂X (X = O, S, Se and Te). The energies are given in kcal/mol units.

[a] Dr. M. Kumar, Prof. J. S. Francisco
Department of Chemistry
University of Nebraska-Lincoln
Lincoln, NE 68588 (USA)
E-mail: jfrancisco3@unl.edu

Supporting information for this article is given via a link at the end of the document.




Figure 2. CCSD(T)/aug-cc-pVTZ//M06-2X/aug-cc-pVTZ calculated zero-point-corrected barriers (*top panels*: addition reactions and *bottom panels*: elimination reactions) for the reactions between various Criegee intermediates and chalcogen hydrides. The barriers are given in kcal/mol units. The correlations between the calculated barriers for the reactions of the simplest Criegee intermediate, CH₂OO with chalcogen hydrides and X-H (X = O, S, Se, and Te) bond distances are given in right side panels.

addition barrier is just 2.5 kcal/mol and the H₂O₂ elimination barrier is 44.1 kcal/mol. For the H₂Se and H₂Te reactions, the addition steps involve barriers < 1.0 kcal/mol whereas the barriers for the elimination step are lowered to 41.1 and 38.4 kcal/mol, respectively. It is important to mention here that the hydroperoxide (*Int*₂ or *Int*₃) in the addition step is formed with an excess energy of at least 42.2 kcal/mol that significantly compensates the high barrier for the elimination step. These results clearly suggest that there is a correlation between the Criegee reaction barrier and the nature of X in H₂X. This correlation can be explained in terms of the H-X bond length or atomic radius of X (**Figure 2**). Moving down the chalcogen group, the atomic radius increases and the H-X bond becomes weaker. Since the H₂X addition to Criegee intermediate involves breakage of one of the H-X bonds, a weaker H-X bond in H₂X would promote the H₂X addition across the Criegee moiety. Oxygen atom has the shortest atomic radius of 66 pm, and the O-H bond (96 pm) with an enthalpy of 463 kJ/mol is the strongest X-H bond whereas Te atom has the largest radius (138 pm), and the Te-H bond (169 pm) with an enthalpy of 267 kJ/mol is the weakest X-H bond.¹⁶ This explains why the H₂O reaction has the largest barriers whereas the H₂Te reaction has the lowest ones. The barriers for the H₂X reactions also directly correlate with first pK_a of H₂X (**Figure S1**). H₂O has the largest pK_a of 15.7 where H₂Te has the lowest pK_a of 3.0, which is consistent with the barrier trends.¹⁷ This implies that chalcogen hydrides with heteroatoms lying in the lower end of periodic table would react faster with Criegee intermediate.

We then examined the reactions of the next larger Criegee intermediate, CH₃CHOO with H₂X. There are two conformers possible for CH₃CHOO; (i) *anti*-CH₃CHOO, and (ii) *syn*-CH₃CHOO. We studied reactions involving both of these conformers. The calculations suggest that the barriers for the *anti*-CH₃CHOO reactions are relatively lower than those for the *syn*-CH₃CHOO reactions (**Figure 2**), which is consistent with previous studies^{8,15} suggesting that *anti*-CH₃CHOO is more reactive than *syn*-CH₃CHOO towards bimolecular reactions. For the H₂O or H₂S reactions, the addition barrier for the *anti*-CH₃CHOO case is ~5.0 kcal/mol lower than that for the *syn*-CH₃CHOO case. More importantly, the calculated barriers for the addition and elimination steps

of the *anti*- and *syn*-CH₃CHOO-H₂X reactions correlate well with the bond length of X-H bond of H₂X (**Figures S2 and S3**) or their first pK_a values. In going from H₂O → H₂Te, the addition barrier for the *anti*-CH₃CHOO case is lowered by 7.8 kcal/mol and the H₂X addition occurs in a near-barrierless manner for X = Se or Te. For the *syn*-CH₃CHOO case, the larger barrier lowering of 11.0 kcal/mol is observed in moving from H₂O → H₂Te. Since the H₂X addition to *anti* or *syn*-CH₃CHOO should lead to the same hydroperoxide, (HX)(CH₃)(H)COOH, we only examined H₂O₂ elimination from the *anti*-CH₃CHOO + H₂X reaction. The barrier for the H₂O₂ elimination from the (HX)(CH₃)(H)COOH is lowered from 45.3 kcal/mol to 37.3 kcal/mol in going from H₂O → H₂Te, which again inversely correlates with the X-H bond strength or directly correlates with first pK_a of H₂X.

The calculated addition and elimination barriers for the reactions of (CH₃)₂COO with H₂X also satisfy a correlation with the X-H bond length (**Figures 2, S2 and S3**) or first pK_a of H₂X, which is indicative of the fact that the correlation between the Criegee reaction barrier and the nature of X in H₂X is a common reactivity trend, and is independent of the structure of the Criegee intermediate involved in the reaction. The comparative analysis suggests that the calculated elimination barriers for the (CH₃)₂COO reactions are not only lower than those for the *syn*-CH₃CHOO, but are the lowest ones. For example, the elimination barriers for H₂O and H₂Te reactions are lowered from 49.1 and 38.4 kcal/mol to 43.9 and 36.2 kcal/mol, respectively, in going from CH₂OO to (CH₃)₂COO. This is due to the fact that the substitution at the hydroperoxyl carbon of (HX)(R₁)(R₂)COOH promotes the H₂O₂ elimination, *i.e.*, greater the extent of substitution, lower would be the elimination barrier.




Figure 3. CCSD(T)/aug-cc-pVTZ//M06-2X/aug-cc-pVTZ calculated zero-point-corrected reaction profiles for the reactions between various Criegee intermediates and hydrogen sulfide with (bottom panel) and without water catalyst (top panel). The energetics are given in kcal/mol units.

The correlation between the Criegee reaction barrier and the nature of heteroatom in H₂X may have important implications for atmospheric and synthetic organic chemistry. For example, thioformaldehyde (HCHS) has been detected in dark clouds, the interstellar clouds¹⁸⁻²⁰ and circumstellar envelope around an asymptotic giant branch star²¹. Few lines of HCHS have also been identified in the Orion KL nebula²²

and in the atmosphere of the comet Hale-Bopp²³. The present calculations suggest that HCHS in the atmosphere could be formed in a near-barrierless manner from the reaction between CH₂OO and H₂S. This reaction could be feasible in the atmosphere because the estimated H₂S concentration in the lower troposphere lies in the 1-1000 ppt range. Another important implication of our results is for the laboratory synthesis of thioaldehydes. In synthetic organic chemistry, thioaldehydes are used as key building blocks. The Diels-Alder additions of thioaldehydes and thioketones offer useful routes for the incorporation of sulfur heteroatoms into the synthesis of heterocycles such as substituted pyridines, furans, and dihydrothiopyrans or natural products.²⁴⁻²⁶ However, experimental detection of simpler aliphatic thioaldehydes such as HCHS has been a significant challenge because it is unstable at room temperature and readily trimerizes to s-trithiane.²⁷ The findings from this work suggest that if the Criegee-H₂S reaction could be performed under water or acid catalysis (Figure 3 and Table S2), thioaldehydes would be barrierlessly formed in an extensively hydrogen-bonded complexed state, which is significantly more stable than their uncomplexed one. Thus, the catalyzed Criegee-H₂S reaction could selectively produce thioaldehyde, avoiding any oligomerization. This could prove useful synthetic guidelines for the laboratory synthesis of simpler thioaldehydes that does not require harsh experimental conditions.

In short, high level quantum chemical calculations reveal an inverse correlation between the Criegee-H₂X reaction barrier and the bond length of the X-H bond in H₂X or a direct correlation with the first pK_a of H₂X. This heteroatom-induced tuning causes a 8-12 kcal/mol lowering in the reaction barrier, thus significantly impacting the overall energetics of the Criegee-H₂X reaction. Considering that bimolecular chemistries are abundant in atmosphere, the heteroatom tuning of the reaction barrier could be a common reactivity determinant of atmospheric addition reactions.

Keywords: Criegee Intermediate • organocatalysis • water catalysis • CCSD(T) theory • atmospheric chemistry • chalcogen hydride • Thioaldehydes

- [1] D. Johnson, G. Marston, *Chem. Soc. Rev.* **2008**, *37*, 699-716.
 [2] K. E. Leather, M. R. McGillen, M. C. Cooke, S. R. Utembe, A. T. Archibald, M. E. Jenkin, R. G. Derwent, D. E. Shallcross, and C. J. Percival, *Atmos. Chem. Phys.* **2012**, *12*, 469.
 [3] R. L. Mauldin, T. Berndt, M. Sipil., P. Paasonen, T. Petäjä, S. Kim, T. Kurtén, F. Stratmann, V.-M. Kerminen, M. Kulmala, *Nature* **2012**, *488*, 193.
 [4] K. J. Heaton, R. L. Sleigher, P. G. Hatcher, W. A. Hall, M. V. Johnston, *Environ. Sci. Technol.* **2009**, *43*, 7797.
 [5] Y. Ma, R. A. Porter, D. Chappell, A. T. Russell, G. Marston, *Phys. Chem. Chem. Phys.* **2009**, *11*, 4184.
 [6] C. A. Taatjes, O. Welz, A. J. Eskola, J. D. Savee, D. L. Osborn, E. P. F. Lee, J. M. Dyke, D. W. K. Mok, D. E. Shallcross, C. J. Percival, *Phys. Chem. Chem. Phys.* **2012**, *14*, 10391.
 [7] O. Welz, J. D. Savee, D. L. Osborn, S. S. Vasu, C. J. Percival, D. E. Shallcross, and C. A. Taatjes, *Science* **2012**, *335*, 204.
 [8] C. A. Taatjes, O. Welz, A. J. Eskola, J. D. Savee, A. M. Scheer, D. E. Shallcross, B. Rotavera, E. P. F. Lee, J. M. Dyke, D. K. W. Mok, D. L. Osborn, and C. J. Percival, *Science* **2013**, *340*, 177.
 [9] L. Vereecken, *Science* **2013**, *340*, 154-155.
 [10] J. A. Ragan, D. J. am Ende, S. J. Brenek, S. A. Eisenbreis, R. A. Singer, D. L. Tickner, J. J. Teixeira, B. C. Vanderplas, N. Weston, *Org. Process Res. Dev.* **2003**, *7*, 155-160.
 [11] R. H. Callighan and M. H. Wilt, *J. Org. Chem.* **1961**, *26*, 4912-4914.
 [12] S. G. Van Omum, R. M. Champeau and R. Pariza, *Chem. Rev.* **2006**, *106*, 2990-3001.
 [13] H. Leisch, K. Morley and P. Lau, *Chem. Rev.* **2011**, *111*, 4165-4222.
 [14] A. Baeyer, V. Villiger, *Ber. Dtsch. Chem. Ges.* **1899**, *32*, 3625-3633.
 [15] J. M. Anglada, J. Gonzalez, M. Torrent-Sucarrat, *Phys. Chem. Chem. Phys.* **2011**, *13*, 13034.
 [16] D. A. Johnson, *Some Thermodynamic Aspects of Inorganic Chemistry*, 2nd Ed., Part of Cambridge Texts in Chemistry and Biochemistry, **1982**, p. 207.
 [17] R. P. Bell, "The proton in Chemistry," 2nd Ed., Cornell University Press, Ithaca, NY, **1973**.
 [18] N. J. Evans, C. H. Townes, H. F. Weaver, D. R. Williams, *Science*


1970, *169*, 680-681.

- [19] M. W. Sinclair, N. Fourikis, J. C. Ribes, B. J. Robinson, R. D. Brown, P. D. Godfrey, *Aust. J. Phys.* **1973**, *26*, 85-91.
 [20] Y. C. Minh, W. M. Irvine, M. K. Brewer, *Astron. Astrophys.* **1991**, *244*, 181-9.
 [21] M. Agundez, J. P. Fonfria, J. Cernicharo, J. R. Pardo, M. Guelin, *Astron. Astrophys.* **2008**, *479*, 493-501.
 [22] C. Comito, P. Schilke, T. G. Phillips, D. C. Lis, F. Motte, D. Mehringer, *Astrophys. J. Suppl. Ser.* **2005**, *156*, 127-167.
 [23] L. M. Woodney, M. F. A'Hearn, J. McMullin, N. Samarasingha, *Earth Moon Planets* **1997**, *78*, 69-70.
 [24] E. Vedejs, T. H. Eberlein, D. L. Varie, *J. Am. Chem. Soc.* **1982**, *104*, 1445-1447.
 [25] R. Okazaki, In *Organosulfur Chemistry*; Page, P. D., Ed.; Academic Press: London, **1995**; p 226.
 [26] F. Duus, 'Comprehensive Organic Chemistry', Vol 3, ed D. H. R. Barton, W. D. Ollis, **1979**, Pergamon, Oxford, p 373.
 [27] R. H. Judge, G. W. King, *Can. J. Phys.* **1975**, *53*, 1927-1928.

Entry for the Table of Contents

COMMUNICATION

The calculated barriers for the bimolecular Criegee-H₂X reactions inversely correlate with the bond strength of X-H bond of H₂X or directly correlate with first pK_a of H₂X. Thus, it is not only the Criegee substitution, but the nature of heteroatom in H₂X that also plays a crucial role in determining the reactivity of the Criegee-H₂X interaction.



Manoj Kumar and Joseph S. Francisco*

Heteroatom Tuning of Bimolecular Criegee Reactions and Its Implications.