

Inflammasome assembly in the chorioamniotic membranes during spontaneous labor at term

Nardhy Gomez-Lopez^{1,2,3} | Roberto Romero^{2,4,5,6}  | Yi Xu^{1,2} | Valeria Garcia-Flores^{1,2} | Yaozhu Leng^{1,2} | Bogdan Panaitescu^{1,2} | Derek Miller^{1,2} | Vikki M. Abrahams⁷ | Sonia S. Hassan^{1,2}

¹Department of Obstetrics and Gynecology, Wayne State University School of Medicine, Detroit, MI, USA

²Perinatology Research Branch, Program for Perinatal Research and Obstetrics, Division of Intramural Research, Eunice Kennedy Shriver National Institute of Child Health and Human Development, NICHD/NIH/DHHS, Bethesda, MD and Detroit, MI, USA

³Department of Immunology and Microbiology, Wayne State University School of Medicine, Detroit, MI, USA

⁴Department of Obstetrics and Gynecology, University of Michigan, Ann Arbor, MI, USA

⁵Department of Epidemiology and Biostatistics, Michigan State University, East Lansing, MI, USA

⁶Center for Molecular Medicine and Genetics, Wayne State University, Detroit, MI, USA

⁷Department of Obstetrics, Gynecology and Reproductive Sciences, Yale University School of Medicine, New Haven, CT, USA

Correspondence

Nardhy Gomez-Lopez, Department of Obstetrics and Gynecology, Wayne State University School of Medicine, Perinatal Research Initiative, Detroit, MI, USA.
Email: ngomezlo@med.wayne.edu and

Roberto Romero, Perinatology Research Branch, NICHD/NIH/DHHS, Hutzel Women's Hospital, Detroit, MI, USA.
Email: prbchiefstaff@med.wayne.edu

Funding information

Eunice Kennedy Shriver National Institute of Child Health and Human Development, Grant/Award Number: HHSN275201300006C; National Institutes of Health; Department of Health and Human Services; Wayne State University Perinatal Initiative in Maternal, Perinatal and Child Health.

Problem: Inflammasome activation requires two steps: priming and assembly of the multimeric complex. The second step includes assembly of the sensor molecule and adaptor protein ASC (an apoptosis-associated speck-like protein containing a CARD), which results in ASC speck formation and the recruitment of caspase (CASP)-1. Herein, we investigated whether there is inflammasome assembly in the chorioamniotic membranes and chorionic leukocytes from women who underwent spontaneous labor at term.

Method of Study: Using in situ proximity ligation assays, ASC/CASP-1 complexes were determined in the chorioamniotic membranes from women who delivered at term without labor or underwent spontaneous labor at term with or without acute histologic chorioamnionitis (n=10-11 each). Also, ASC speck formation was determined by flow cytometry in the chorionic leukocytes isolated from women who delivered at term with or without spontaneous labor (n=9-12 each).

Results: (i) ASC/CASP-1 complexes were detected in the chorioamniotic membranes; (ii) ASC/CASP-1 complexes were greater in the chorioamniotic membranes from women who underwent spontaneous labor at term than in those without labor; (iii) ASC/CASP-1 complexes were even more abundant in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis than in those without this placental lesion; (iv) ASC speck formation was detected in the chorionic leukocytes; and (v) ASC speck formation was greater in the chorionic leukocytes isolated from women who underwent spontaneous labor at term than in those without labor.

Conclusion: There is inflammasome assembly in the chorioamniotic membranes and chorionic leukocytes during spontaneous labor at term.

KEYWORDS

acute chorioamnionitis, apoptosis-associated speck-like protein containing a CARD (ASC), ASC speck, caspase-1, interleukin-1 beta, intra-amniotic infection/inflammation

1 | INTRODUCTION

Inflammasomes are multimeric protein complexes located in the cytoplasm capable of detecting microorganisms, damage-derived signals, and other cellular stressors.^{1–14} The inflammasome structure includes: (i) a sensor molecule or pattern recognition receptor (PRR), (ii) pro-caspase 1 (pro-CASP-1), and (iii) the adaptor protein ASC (an apoptosis-associated speck-like protein containing a CARD).^{1–14} The ASC protein is encoded by *PYCARD* and includes two death-fold domains: one pyrin domain and one caspase activation and recruitment domain (CARD).^{15,16} The pyrin domain interacts with the sensor molecule of the inflammasome, triggering the assembly of the ASC protein into a large complex termed “speck” that consists of multimers of ASC dimers.^{17,18} The CARD domain of the ASC protein recruits monomers of pro-CASP-1 (ASC/CASP-1 complexes), initiating the self-cleavage of this enzyme and the formation of its active subunits (p10 and p20).¹⁹ These subunits can then assemble to form active CASP-1 hetero-tetramers, which are able to convert pro-interleukin (IL)-1 β and pro-IL-18 into their mature and secreted forms.^{20–27} The secretion of large amounts of pro-inflammatory cytokines along with the activation of CASP-1 induces a programmed cell death pathway termed pyroptosis.²⁸

Most of the inflammasomes contain sensor molecules from the NOD-like receptor (NLR) family, namely NLRP1,¹ NLRP3 (also known as cryopyrin or NALP3),² NLRP6,^{29,30} NLRP7,^{31,32} and NLRP12 or NLRC4 (NLR family CARD domain-containing protein 4 or IPAF).³³ Yet, there are inflammasomes that do not contain an NLR (eg, interferon gamma-inducible protein 16 (IFI16),³⁴ absent in melanoma 2 (AIM2),^{35–39} and pyrin^{40–42}). The best-characterized inflammasome is the NLRP3, and its activation includes a two-step process.⁴³ The first is a priming step that includes the activation of the nuclear factor kappa B (NF- κ B) pathway, which induces the upregulation of pro-IL-1 β and the NLR protein to a functional level.^{43,44} The second step includes the assembly and activation of the inflammasome complex, which culminates in the activation of CASP-1 and the consequent processing of pro-IL-1 β into its mature form.⁴³ Priming and assembly of the NLRP3 inflammasome can be triggered by both exogenous and endogenous molecules; therefore, such signaling platforms are implicated in the mechanisms that lead to sterile inflammation and microbial-derived inflammation.⁴³

Recently, we provided evidence supporting a role for the NLRP3 inflammasome in the mechanisms responsible for sterile and microbial-associated inflammation in the chorioamniotic membranes (eg, acute histologic chorioamnionitis).^{45–47} Specifically, we demonstrated that there is priming of the inflammasome in the chorioamniotic membranes during spontaneous labor at term with⁴⁶ or without⁴⁵ acute histologic chorioamnionitis. This concept is supported by the fact that the chorioamniotic membranes from women who underwent spontaneous labor at term express increased mRNA and protein levels of IL-1 β and NLRP3,⁴⁵ all of which are even greater in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis.⁴⁶ However, whether there is inflammasome assembly in the chorioamniotic membranes from

women who underwent spontaneous labor at term with or without acute histologic chorioamnionitis is unknown.

Inflammasome assembly can be assessed by identifying ASC/CASP-1 complexes using in situ proximity ligation assays⁴⁷ or by evaluating ASC speck formation using flow cytometry.⁴⁸ Herein, we determined ASC/CASP-1 complexes in the chorioamniotic membranes and ASC speck formation in the choriodecidual leukocytes as a readout for inflammasome assembly during spontaneous labor at term.

2 | MATERIALS AND METHODS

2.1 | Human subjects, clinical specimens, and definitions

Chorioamniotic membrane samples were obtained from the Bank of Biological Specimens of the Detroit Medical Center, Wayne State University, and the Perinatology Research Branch (Detroit, MI, USA), an intramural program of the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, National Institutes of Health, U. S. Department of Health and Human Services (NICHD/NIH/DHHS). The collection and utilization of biological materials for research purposes were approved by the Institutional Review Boards of Wayne State University and NICHD. All participating women provided written informed consent. The following three study groups were included: (i) women who delivered at term without labor; (ii) women who underwent spontaneous labor at term without acute histologic chorioamnionitis; and (iii) women who underwent spontaneous labor at term with acute histologic chorioamnionitis. Table 1 contains the demographic and clinical characteristics of the three study populations utilized for in situ proximity ligation assays. Table 2 contains the demographic and clinical characteristics of the two study populations utilized for the determination of ASC speck formation by flow cytometry. Multiparous women or women with neonates having congenital or chromosomal abnormalities were excluded. Labor at term was defined by the presence of regular uterine contractions occurring at a frequency of at least two contractions every 10 minutes with cervical changes resulting in delivery.

2.2 | Placental histopathological examinations

Five-micrometer-thick sections of formalin-fixed, paraffin-embedded tissue specimens were cut and mounted on SuperFrost™ Plus microscope slides (Erie Scientific LLC, Portsmouth, NH, USA). In each case, several tissue sections of the chorioamniotic membranes, umbilical cord, and placental disk were examined. After deparaffinization, slides were rehydrated, stained with hematoxylin-eosin, and evaluated by pathologists who were blinded to the clinical outcome, according to the published criteria.^{49,50} The diagnosis of acute histologic chorioamnionitis was made when the infiltration of neutrophils into the chorionic trophoblast layer or chorioamniotic connective tissue was observed.^{49,50}

TABLE 1 Demographic and clinical characteristics of the study populations utilized for in situ proximity ligation assays

	Term no labor (n=10)	Term in labor (n=10)	Term in labor with acute histologic chorioamnionitis (n=11)	P value
Age (y; median [IQR]) ^a	29.0 (25.5-30.8)	23.5 (19.3-31.0)	23.0 (21.0-26.5)	NS
Body mass index (kg/m ² ; median [IQR]) ^a	32.3 (29.4-36.6)	24.1 (22.5-31.0)	28.2 (23.9-33.6)	.028
Gestational age at delivery (wk; median [IQR]) ^a	39.3 (39.0-39.4)	39.3 (39.0-39.9)	39.1 (38.8-39.9)	NS
Birth weight (g; median [IQR]) ^a	3705.0 (3272.5-3842.5)	3220.0 (3011.3-3304.0)	3270.0 (3072.5-3585.0)	NS
Race (n[%]) ^b				
African-American	7 (70%)	10 (100%)	11 (100%)	NS
Caucasian	1 (10%)	0 (0%)	0 (0%)	
Other	2 (20%)	0 (0%)	0 (0%)	
Primiparity (n[%]) ^b	0 (0%)	4 (40%)	1 (9.1%)	NS
C-section (n[%]) ^b	10 (100%)	1 (10%)	2 (18.2%)	NS
Acute fetal inflammatory response (n[%]) ^b				
No fetal inflammatory response	10 (100%)	10 (100%)	0 (0%)	.001
Stage 1 (acute phlebitis/chorionic vasculitis)	0 (0%)	0 (0%)	3 (27.3%)	NS
Stage 2 (acute arteritis)	0 (0%)	0 (0%)	8 (72.7%)	NS
Stage 3 (necrotizing funisitis)	0 (0%)	0 (0%)	0 (0%)	NS

^aKruskal-Wallis test.^bFisher's exact test.

IQR=interquartile range.

TABLE 2 Demographic and clinical characteristics of the study populations utilized for the determination of ASC speck formation by flow cytometry

	Term no labor (n=12)	Term in labor (n=9)	P value
Age (y; median [IQR]) ^a	28.5 (27.0-35.0)	24.0 (22.0-27.0)	.032
Body mass index (kg/m ² ; median [IQR]) ^a	32.4 (29.3-45.4)	32.3 (26.6-37.0)	NS
Gestational age at delivery (wk; median [IQR]) ^a	39.0 (38.8-39.9)	38.0 (38.0-39.0)	NS
Birth weight (g; median [IQR]) ^a	3287.5 (2820.0-3438.8)	3140.0 (2950.0-3275.0)	NS
Race (n[%]) ^b			
African-American	10 (83.3%)	7 (77.8%)	NS
Caucasian	2 (16.7%)	1 (11.1%)	
Hispanic	0 (0%)	1 (11.1%)	
Primiparity (n[%]) ^b	0 (0%)	1 (11.1%)	NS
C-section (n[%]) ^b	12 (100%)	0 (0%)	.001

^aKruskal-Wallis test.^bFisher's exact test.

IQR, interquartile range.

None of the samples showed signs of a fetal inflammatory response.

2.3 | In Situ proximity ligation assay

ASC/CASP-1 complexes were detected by co-localizing ASC and CASP-1 using a Duolink[®] in situ proximity ligation assay kit (Olink Bioscience, Uppsala, Sweden), following the manufacturer's instructions. Briefly, chorioamniotic membrane tissues were frozen in Tissue-Plus O.C.T. compound (Fisher HealthCare, Houston, TX, USA) immediately after collection. Cryogenic sections were cut to 10 μm and placed onto glass microscope slides (Fisherbrand Superfrost Plus

slides; Thermo Scientific, Waltham, MA, USA). The sections were fixed using 4% paraformaldehyde (Electron Microscopy Sciences, Hatfield, PA, USA) for 20 minutes at room temperature, rinsed with 1X PBS, and permeabilized using 0.25% Triton X-100 (Promega, Madison, WI, USA) for 5 minutes at room temperature. Prior to staining, non-specific antibody interactions were blocked using serum-free protein blocker (catalog number X09090; DAKO North America, Carpinteria, CA, USA) for 30 minutes at room temperature. The sections were then stained with the following antibodies at 4°C overnight: rabbit antihuman ASC

(catalog number AG-25B-0006-C100; Adipogen, San Diego, CA, USA) and mouse antihuman CASP-1 (catalog number MAB6251, clone 661228; R&D Systems, Minneapolis, MN, USA). Rabbit IgG and mouse IgG2A were used as negative controls, respectively. Following staining, slides were briefly washed with 1X Wash Buffer A from the Duolink® kit and incubated with the provided proximity ligation assay probes for 1 hour at 37°C, followed by a second wash with 1X Wash Buffer A. The slides were then incubated with the Duolink® ligase solution for 30 minutes at 37°C, washed with 1X Wash Buffer A, and incubated for 100 minutes with the Duolink® polymerase solution. Finally, the slides were washed with 1X Wash Buffer B and mounted with Duolink® mounting media with DAPI. Immunofluorescence was visualized using an Olympus BX60 fluorescence microscope (Olympus, Tokyo, Japan) at 400X magnification. Images were acquired using an Olympus DP71 camera and DP Controller software (Olympus). Semi-quantification was performed using the Duolink® image analysis software. Images and videos were acquired using a Zeiss LSM 800 laser scanning confocal microscope (Carl Zeiss Microscopy, Jena, Germany) at the Microscopy, Imaging, and Cytometry Resources Core at the Wayne State University School of Medicine (<http://micr.med.wayne.edu/>). ASC/CASP-1 complexes were calculated by dividing the number of signals over the area of the tissue. The area of the tissue is expressed in pixels.

2.4 | ASC speck formation

ASC speck formation in inflammasome-competent cells can be assessed by flow cytometry.⁴⁸ Choriondecidual leukocytes were isolated, as previously described⁵¹ (purity >85% CD45+ cells within the forward scatter [FSC] vs size scatter [SSC] gate for leukocytes). Isolated choriondecidual leukocytes were incubated with 20 µL of human FcR blocking reagent (Miltenyi Biotec, San Diego, CA, USA) in 80 µL of stain buffer (catalog number 554656; BD Biosciences, San Jose, CA, USA) for 10 minutes at 4°C. Choriondecidual leukocytes were then washed with 1 mL of stain buffer, re-suspended in 250 µL of BD Cytofix/Cytoperm Fixation and Permeabilization Solution (BD Biosciences), and incubated for 30 minutes at 4°C in the dark. After fixation, choriondecidual leukocytes were washed with 1 mL of 1X BD Perm/Wash Buffer (BD Biosciences) and re-suspended in 50 µL of the 1X BD Perm/Wash Buffer. Choriondecidual leukocytes were then stained with anti-ASC-PE (clone HASC-71; catalog number 653903; BioLegend, San Diego, CA, USA) or the IgG isotype control (catalog number 400139; BioLegend) for 30 minutes at 4°C in the dark. Finally, the stained cells were washed with 1 mL of 1X BD Perm/Wash Buffer, re-suspended in 0.5 mL of stain buffer, and acquired using the BD LSR Fortessa Flow Cytometer and BD FACSDiva 6.0 software. The analysis and figures were performed using FlowJo software version 10 (FlowJo, Ashland, OR, USA).

As a positive control for ASC speck formation, THP-1 cells (monocyte cell line; American Type Culture Collection; ATCC; Manassas, VA; 2.5×10⁶ cells/mL) were incubated with 1 µg/mL of lipopolysaccharide (LPS; Escherichia coli 0111:B4; Sigma-Aldrich, 2 hours) and 20 µM of nigericin (catalog number N7142, Sigma-Aldrich, St. Louis, MO, 1 hour). Non-treated THP-1 cells were used as a negative control.

2.5 | Statistical analyses

The SPSS v.19.0 software (IBM Corp., Armonk, NY, USA) was used to analyze the demographic and clinical characteristics of the study populations. Normality of the data was tested using the Shapiro-Wilk test. Comparisons among groups were performed using the Kruskal-Wallis test and/or the Mann-Whitney *U* test. Comparisons of proportions were made using the Fisher's exact test. A *P* value of ≤.05 was used to determine statistical significance.

3 | RESULTS

3.1 | ASC/CASP-1 complex formation is increased in the chorioamniotic membranes from women who underwent spontaneous labor at term

As a readout for inflammasome assembly, we first performed *in situ* proximity ligation assays to identify ASC/CASP-1 complex formation.⁴⁷ Figure 1 shows representative images of ASC/CASP-1 complexes in the chorioamniotic membranes. ASC/CASP-1 complexes were rarely found in the chorioamniotic membranes from women who delivered at term without labor (Figure 1A). A 3-D reconstruction shows that ASC/CASP-1 complexes were scarce in the chorioamniotic membranes from women who delivered at term without labor (Video S1). ASC/CASP-1 complex formation was evident in the chorioamniotic membranes from women who underwent spontaneous labor at term (Figure 1B), and a 3-D reconstruction shows that such complexes are mostly localized in the compact layer of the amnion (Video S2). Yet, ASC/CASP-1 complexes were more abundant in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis than in those without this placental lesion (Figure 1C vs 1B). A 3-D reconstruction shows that ASC/CASP-1 complexes were abundant in both the amnion and chorion from women who underwent spontaneous labor at term with acute histologic chorioamnionitis (Video S3). The isotype controls did not show any non-specific background signal (Figure 1D-F). Semi-quantification of the fluorescence signal revealed that, indeed, ASC/CASP-1 complexes were greater in the chorioamniotic membranes from women who underwent spontaneous labor at term than in those who delivered at term without labor (Figure 2). ASC/CASP-1 complexes were even greater in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis than in those without this placental lesion (Figure 2). Collectively, these results provide evidence that there is inflammasome assembly in the chorioamniotic membranes during spontaneous labor at term and that such an event is enriched in cases with acute histologic chorioamnionitis.

3.2 | ASC speck formation is increased in the choriondecidual leukocytes isolated from women who underwent spontaneous labor at term

The ASC adaptor protein is diffusely located throughout the nucleus and cytoplasm; yet, upon inflammasome activation, it forms a large




FIGURE 1 ASC/CASP-1 complex formation in the chorioamniotic membranes. In situ proximity ligation assays for ASC and CASP-1 in the chorioamniotic membranes from women who delivered at term without labor or underwent spontaneous labor at term with or without acute histologic chorioamnionitis. Representative images of ASC/CASP-1 complexes (red signal) in each group (A-C). Representative images of isotype controls in each group (D-F). The blue signal is DAPI (nuclei). 400 \times magnification. CASP-1, caspase 1; ASC, apoptosis-associated speck-like protein containing a CARD




FIGURE 2 Semi-quantification of ASC/CASP-1 complexes in the chorioamniotic membranes. ASC/CASP-1 complexes were calculated by dividing the number of signals over the area of the tissue. The area of the tissue is expressed in pixels (N=10-11 each)

perinuclear agglutination termed "speck."^{15,18} Therefore, the presence of ASC specks represents inflammasome activation.¹⁸ Recently, a novel flow cytometry method was developed to determine ASC speck

formation in inflammasome-competent cells.⁴⁸ Using this method, we evaluated whether there is ASC speck formation, as a readout of inflammasome activation, in the choriodecidual leukocytes isolated from women who underwent spontaneous labor at term. The gating strategy used to evaluate ASC speck formation is shown in Figure 3A. Briefly, choriodecidual leukocytes were first gated within the FSC-area vs SSC-area gate to exclude cellular debris and stromal cells. Next, doublets were excluded using FSC-area vs FSC-width profile characteristics. Choriodecidual leukocytes stained for ASC were then gated, and the detection of ASC specks was analyzed within the ASC-area vs ASC-width gate. We used untreated (Figure 3B) or treated (LPS+nigericin; Figure 3C) THP-1 cells as negative and positive controls, respectively. Isotype controls for THP-1 cells are shown in Figure 3E, and 3F. Detection of ASC specks in the choriodecidual leukocytes isolated from women who underwent spontaneous labor at term is shown in Figure 3D along with its isotype control (Figure 3G). Flow cytometric quantification revealed that ASC speck formation is greater in the choriodecidual leukocytes isolated from women who underwent spontaneous labor at term than in those who delivered without labor (Figure 4). These results indicate that there is inflammasome activation in the choriodecidual leukocytes isolated from women who underwent spontaneous labor at term.




FIGURE 3 ASC speck formation in the choriodecidual leukocytes. (A) Flow cytometry gating strategy used to determine ASC speck formation in the choriodecidual leukocytes. THP-1 cells were untreated (negative control, B) or treated with lipopolysaccharide (LPS) plus nigericin (positive control, C). A representative image of ASC speck formation in the choriodecidual leukocytes isolated from women who underwent spontaneous labor at term (D). All experiments included isotype controls




FIGURE 4 Flow cytometric quantification of ASC speck formation in the choriodecidual leukocytes. ASC speck formation in the choriodecidual leukocytes isolated from women who delivered at term without labor or underwent spontaneous labor at term (N=9-12 each)

4 | DISCUSSION

4.1 | Principal findings of the study

(i) ASC/CASP-1 complexes were detected in the chorioamniotic membranes; (ii) ASC/CASP-1 complexes were greater in the chorioamniotic membranes from women who underwent spontaneous labor at term than in those without labor; (iii) ASC/CASP-1 complexes were even more abundant in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis than in those without this placental lesion; (iv) ASC speck formation was detected in the choriodecidual leukocytes; and (v) ASC speck formation was greater in the choriodecidual leukocytes isolated from women who underwent spontaneous labor at term than in those without labor. Collectively, these results provide evidence that there

is assembly of the inflammasome in the chorioamniotic membranes and chorionic leukocytes during spontaneous labor at term.

Spontaneous labor at term is a state of physiological inflammation⁵²⁻⁵⁷ that, in most women, is considered sterile because it occurs in the absence of intra-amniotic infection.^{58,59} This concept is supported by evidence demonstrating that there is an increased bioavailability of pro-inflammatory mediators, such as cytokines, in the amniotic fluid of women who underwent spontaneous labor at term.⁶⁰⁻⁷⁹ Of all these cytokines, IL-1 β seems to be a central mediator in the process of parturition because its concentrations are increased in the amniotic fluid^{61,62} and chorioamniotic membranes^{45,80,81} from women who underwent spontaneous labor at term. Indeed, the systemic⁸² or intra-amniotic^{83,84} administration of IL-1 β induces preterm labor and birth. The function of this cytokine during the process of labor includes, but is not limited to, the induction of (i) the biosynthesis of prostaglandin E2 by the human amnion⁸⁵ and myometrial cells;^{86,87} (ii) the expression of cyclooxygenase-2 in human myometrial cells;⁸⁸ and (iii) the expression of matrix-metabolizing enzymes (MMP-1, MMP-3, MMP-9, and cathepsin S) in human cervical smooth muscle cells⁸⁹. However, the mechanisms that lead to the processing of the bioactive and mature form of IL-1 β during labor are poorly understood. Recently, we provided evidence that there is priming of the inflammasome (ie, upregulation of pro-IL-1 β and the NLRP3 protein) in the chorioamniotic membranes of women who underwent spontaneous labor at term, which may be implicated in the processing of mature IL-1 β .⁴⁵ In the study herein, we found that there is assembly of the inflammasome (ie, the ASC protein forms complexes with CASP-1) in the chorioamniotic membranes during spontaneous labor at term. These findings along with the fact that there is activation of CASP-1 and processing of mature IL-1 β in the chorioamniotic membranes⁴⁵ support the concept that the inflammasome is involved in the processing of IL-1 β by the chorioamniotic membranes during spontaneous labor at term.^{45,90,91}

In the current study, we also observed that there is increased assembly of the inflammasome in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis. This finding is in line with our previous observations demonstrating that the chorioamniotic membranes from women who underwent spontaneous labor at term without acute histologic chorioamnionitis, when compared to those with this placental lesion: (i) express higher mRNA and protein levels of IL-1 β and NLRP3 (ie, priming of the inflammasome); (ii) contain increased quantities of the active forms of CASP-1; and (iii) release greater concentrations of mature IL-1 β .⁴⁶ Together, these data suggest that, in the pathological process of acute histologic chorioamnionitis, there is an exacerbated activation of the inflammasome in the chorioamniotic membranes.

Chorionic leukocytes express pro-inflammatory mediators implicated in the processes of term and preterm labor.^{57,81,92-99} Indeed, the chorionic leukocytes isolated from women who underwent spontaneous labor at term exhibit an increased expression of IL-1 β ,⁸¹ which likely contributes to the amount of this cytokine released by the chorioamniotic membranes⁴⁵ and decidual cells.¹⁰⁰ These findings prompted us to investigate whether the chorionic leukocytes could express inflammasome components. Herein, we observed that the chorionic leukocytes display increased ASC speck formation

(ie, inflammasome assembly/activation) during spontaneous labor at term. Although most of the isolated cells were leukocytes (purity >85%), we do not overlook the possibility that, in some decidual cells, there is inflammasome activation during parturition. Together, these data suggest that there is inflammasome assembly and activation in the chorionic leukocytes during spontaneous labor at term.

In summary, the study herein provides further evidence that there is inflammasome activation (assembly of the ASC protein with CASP-1 and ASC speck formation) in the chorioamniotic membranes and chorionic leukocytes from women who underwent spontaneous labor at term. These data also support the hypothesis stating that the inflammasome mediates the activation of CASP-1 and the consequent release of mature IL-1 β during physiological and pathological inflammation of the chorioamniotic membranes in spontaneous labor at term.^{45,46,90,91}

ACKNOWLEDGMENTS

This research was supported, in part, by the Perinatology Research Branch, Division of Intramural Research, Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institutes of Health, U. S. Department of Health and Human Services (NICHD/NIH/DHHS), and, in part, with federal funds from the NICHD/NIH/DHHS under Contract No. HHSN275201300006C. This research was also supported by the Wayne State University Perinatal Initiative in Maternal, Perinatal and Child Health. We thank the physicians and nurses from the Center for Advanced Obstetrical Care and Research, the Intrapartum Unit, and the research assistants from the PRB Clinical Laboratory, for their help in collecting human samples. We also thank staff members of the PRB Histology and Pathology Units for their examination of the pathological sections, and Tara Mial and Maureen McGerty, M.A., for their critical readings of the manuscript.

CONFLICT OF INTEREST

The authors disclose no conflicts of interest.

REFERENCES

1. Martinon F, Burns K, Tschopp J. The inflammasome: a molecular platform triggering activation of inflammatory caspases and processing of proIL-beta. *Mol Cell*. 2002;10:417-426.
2. Agostini L, Martinon F, Burns K, McDermott MF, Hawkins PN, Tschopp J. NALP3 forms an IL-1beta-processing inflammasome with increased activity in Muckle-Wells autoinflammatory disorder. *Immunity*. 2004;20:319-325.
3. Petrilli V, Papin S, Tschopp J. The inflammasome. *Curr Biol*. 2005;15:R581.
4. Ogura Y, Sutterwala FS, Flavell RA. The inflammasome: first line of the immune response to cell stress. *Cell*. 2006;126:659-662.
5. Stutz A, Golenbock DT, Latz E. Inflammasomes: too big to miss. *J Clin Invest*. 2009;119:3502-3511.
6. Franchi L, Eigenbrod T, Munoz-Planillo R, Nunez G. The inflammasome: a caspase-1-activation platform that regulates immune responses and disease pathogenesis. *Nat Immunol*. 2009;10:241-247.

7. Jha S, Ting JP. Inflammasome-associated nucleotide-binding domain, leucine-rich repeat proteins and inflammatory diseases. *J Immunol.* 2009;183:7623–7629.
8. Pedra JH, Cassel SL, Sutterwala FS. Sensing pathogens and danger signals by the inflammasome. *Curr Opin Immunol.* 2009;21:10–16.
9. Lamkanfi M, Dixit VM. Modulation of inflammasome pathways by bacterial and viral pathogens. *J Immunol.* 2011;187:597–602.
10. van de Veerdonk FL, Netea MG, Dinarello CA, Joosten LA. Inflammasome activation and IL-1beta and IL-18 processing during infection. *Trends Immunol.* 2011;32:110–116.
11. Franchi L, Munoz-Planillo R, Nunez G. Sensing and reacting to microbes through the inflammasomes. *Nat Immunol.* 2012;13:325–332.
12. Latz E, Xiao TS, Stutz A. Activation and regulation of the inflammasomes. *Nat Rev Immunol.* 2013;13:397–411.
13. Vanaja SK, Rathinam VA, Fitzgerald KA. Mechanisms of inflammasome activation: recent advances and novel insights. *Trends Cell Biol.* 2015;25:308–315.
14. Guo H, Callaway JB, Ting JP. Inflammasomes: mechanism of action, role in disease, and therapeutics. *Nat Med.* 2015;21:677–687.
15. Masumoto J, Taniguchi S, Ayukawa K, et al. ASC, a novel 22-kDa protein, aggregates during apoptosis of human promyelocytic leukemia HL-60 cells. *J Biol Chem.* 1999;274:33835–33838.
16. Masumoto J, Taniguchi S, Sagara J. Pyrin N-terminal homology domain- and caspase recruitment domain-dependent oligomerization of ASC. *Biochem Biophys Res Commun.* 2001;280:652–655.
17. Vajjhala PR, Mirams RE, Hill JM. Multiple binding sites on the pyrin domain of ASC protein allow self-association and interaction with NLRP3 protein. *J Biol Chem.* 2012;287:41732–41743.
18. Fernandes-Alnemri T, Wu J, Yu JW, et al. The pyroptosome: a supramolecular assembly of ASC dimers mediating inflammatory cell death via caspase-1 activation. *Cell Death Differ.* 2007;14:1590–1604.
19. Srinivasula SM, Poyet JL, Razmara M, Datta P, Zhang Z, Alnemri ES. The PYRIN-CARD protein ASC is an activating adaptor for caspase-1. *J Biol Chem.* 2002;277:21119–21122.
20. Black RA, Kronheim SR, Merriam JE, March CJ, Hopp TP. A pre-aspartate-specific protease from human leukocytes that cleaves pro-interleukin-1 beta. *J Biol Chem.* 1989;264:5323–5326.
21. Kostura MJ, Tocci MJ, Limjuco G, et al. Identification of a monocyte specific pre-interleukin 1 beta convertase activity. *Proc Natl Acad Sci U S A.* 1989;86:5227–5231.
22. Thornberry NA, Bull HG, Calaycay JR, et al. A novel heterodimeric cysteine protease is required for interleukin-1 beta processing in monocytes. *Nature.* 1992;356:768–774.
23. Wilson KP, Black JA, Thomson JA, et al. Structure and mechanism of interleukin-1 beta converting enzyme. *Nature.* 1994;370:270–275.
24. Gu Y, Kuida K, Tsutsui H, et al. Activation of interferon-gamma inducing factor mediated by interleukin-1beta converting enzyme. *Science.* 1997;275:206–209.
25. Ghayur T, Banerjee S, Hugunin M, et al. Caspase-1 processes IFN-gamma-inducing factor and regulates LPS-induced IFN-gamma production. *Nature.* 1997;386:619–623.
26. Dinarello CA. Interleukin-1 beta, interleukin-18, and the interleukin-1 beta converting enzyme. *Ann N Y Acad Sci.* 1998;856:1–11.
27. Fantuzzi G, Dinarello CA. Interleukin-18 and interleukin-1 beta: two cytokine substrates for ICE (caspase-1). *J Clin Immunol.* 1999;19:1–11.
28. Bergsbaken T, Fink SL, Cookson BT. Pyroptosis: host cell death and inflammation. *Nat Rev Microbiol.* 2009;7:99–109.
29. Grenier JM, Wang L, Manji GA, et al. Functional screening of five PYPAF family members identifies PYPAF5 as a novel regulator of NF-kappaB and caspase-1. *FEBS Lett.* 2002;530:73–78.
30. Chen GY, Liu M, Wang F, Bertin J, Nunez G. A functional role for Nlrp6 in intestinal inflammation and tumorigenesis. *J Immunol.* 2011;186:7187–7194.
31. Kinoshita T, Wang Y, Hasegawa M, Imamura R, Suda T. PYPAF3, a PYRIN-containing APAF-1-like protein, is a feedback regulator of caspase-1-dependent interleukin-1beta secretion. *J Biol Chem.* 2005;280:21720–21725.
32. Khare S, Dorfleutner A, Bryan NB, et al. An NLRP7-containing inflammasome mediates recognition of microbial lipopeptides in human macrophages. *Immunity.* 2012;36:464–476.
33. Sutterwala FS, Flavell RA. NLRC4/IPAF: a CARD carrying member of the NLR family. *Clin Immunol.* 2009;130:2–6.
34. Kerur N, Veettil MV, Sharma-Walia N, et al. IFI16 acts as a nuclear pathogen sensor to induce the inflammasome in response to Kaposi Sarcoma-associated herpesvirus infection. *Cell Host Microbe.* 2011;9:363–375.
35. Burckstummer T, Baumann C, Bluml S, et al. An orthogonal proteomic-genomic screen identifies AIM2 as a cytoplasmic DNA sensor for the inflammasome. *Nat Immunol.* 2009;10:266–272.
36. Fernandes-Alnemri T, Yu JW, Datta P, Wu J, Alnemri ES. AIM2 activates the inflammasome and cell death in response to cytoplasmic DNA. *Nature.* 2009;458:509–513.
37. Roberts TL, Idris A, Dunn JA, et al. HIN-200 proteins regulate caspase activation in response to foreign cytoplasmic DNA. *Science.* 2009;323:1057–1060.
38. Hornung V, Ablasser A, Charrel-Dennis M, et al. AIM2 recognizes cytosolic dsDNA and forms a caspase-1-activating inflammasome with ASC. *Nature.* 2009;458:514–518.
39. Franchi L, Nunez G. AIM2 joins the gang of microbial sensors. *Cell Host Microbe.* 2010;7:340–341.
40. Park YH, Wood G, Kastner DL, Chae JJ. Pyrin inflammasome activation and RhoA signaling in the autoinflammatory diseases FMF and HIDS. *Nat Immunol.* 2016;17:914–921.
41. Akula MK, Shi M, Jiang Z, et al. Control of the innate immune response by the mevalonate pathway. *Nat Immunol.* 2016;17:922–929.
42. Dorfleutner A, Stehlik C. A dRAsTic RHOA block of Pyrin inflammasome activation. *Nat Immunol.* 2016;17:900–902.
43. Sutterwala FS, Haasken S, Cassel SL. Mechanism of NLRP3 inflammasome activation. *Ann N Y Acad Sci.* 2014;1319:82–95.
44. Bauernfeind FG, Horvath G, Stutz A, et al. Cutting edge: NF-kappaB activating pattern recognition and cytokine receptors license NLRP3 inflammasome activation by regulating NLRP3 expression. *J Immunol.* 2009;183:787–791.
45. Romero R, Xu Y, Plazyo O, et al. A role for the inflammasome in spontaneous labor at term. *Am J Reprod Immunol.* 2016; doi: 10.1111/aji.12440. [Epub ahead of print].
46. Gomez-Lopez N, Romero R, Xu Y, et al. A role for the inflammasome in spontaneous labor at term with acute histologic chorioamnionitis. *Reprod Sci.* 2016; doi: 10.1177/1933719116675058. [Epub ahead of print].
47. Gomez-Lopez N, Romero R, Xu Y, et al. A role for the inflammasome in spontaneous preterm labor with acute histologic chorioamnionitis. *Reprod Sci.* 2017; doi: 10.1177/193371911668765. [Epub ahead of print].
48. Sester DP, Thygesen SJ, Sagulenko V, et al. A novel flow cytometric method to assess inflammasome formation. *J Immunol.* 2015;194:455–462.
49. Kim CJ, Romero R, Chaemsaitong P, Chaiyasit N, Yoon BH, Kim YM. Acute chorioamnionitis and funisitis: definition, pathologic features, and clinical significance. *Am J Obstet Gynecol.* 2015;213:S29–S52.
50. Redline RW. Classification of placental lesions. *Am J Obstet Gynecol.* 2015;213:S21–S28.
51. Xu Y, Plazyo O, Romero R, Hassan SS, Gomez-Lopez N. Isolation of leukocytes from the human maternal-fetal interface. *J Vis Exp.* 2015;e52863.
52. Romero R, Espinoza J, Goncalves LF, Kusanovic JP, Friel LA, Nien JK. Inflammation in preterm and term labour and delivery. *Semin Fetal Neonatal Med.* 2006;11:317–326.

53. Haddad R, Tromp G, Kuivaniemi H, et al. Human spontaneous labor without histologic chorioamnionitis is characterized by an acute inflammation gene expression signature. *Am J Obstet Gynecol.* 2006;195:394. e391-324.
54. Hassan SS, Romero R, Haddad R, et al. The transcriptome of the uterine cervix before and after spontaneous term parturition. *Am J Obstet Gynecol.* 2006;195:778-786.
55. Romero R, Gotsch F, Pineles B, Kusanovic JP. Inflammation in pregnancy: its roles in reproductive physiology, obstetrical complications, and fetal injury. *Nutr Rev.* 2007;65:S194-S202.
56. Mittal P, Romero R, Tarca AL, et al. Characterization of the myometrial transcriptome and biological pathways of spontaneous human labor at term. *J Perinat Med.* 2010;38:617-643.
57. Gomez-Lopez N, StLouis D, Lehr MA, Sanchez-Rodriguez EN, Arenas-Hernandez M. Immune cells in term and preterm labor. *Cell Mol Immunol.* 2014;11:571-581.
58. Romero R, Nores J, Mazor M, et al. Microbial invasion of the amniotic cavity during term labor. Prevalence and clinical significance. *J Reprod Med.* 1993;38:543-548.
59. Seong HS, Lee SE, Kang JH, Romero R, Yoon BH. The frequency of microbial invasion of the amniotic cavity and histologic chorioamnionitis in women at term with intact membranes in the presence or absence of labor. *Am J Obstet Gynecol.* 2008;199:375. e371-375.
60. Romero R, Brody DT, Oyarzun E, et al. Infection and labor. III. Interleukin-1: a signal for the onset of parturition. *Am J Obstet Gynecol.* 1989;160:1117-1123.
61. Romero R, Parvizi ST, Oyarzun E, et al. Amniotic fluid interleukin-1 in spontaneous labor at term. *J Reprod Med.* 1990;35:235-238.
62. Romero R, Mazor M, Brandt F, et al. Interleukin-1 alpha and interleukin-1 beta in preterm and term human parturition. *Am J Reprod Immunol.* 1992;27:117-123.
63. Romero R, Sepulveda W, Mazor M, et al. The natural interleukin-1 receptor antagonist in term and preterm parturition. *Am J Obstet Gynecol.* 1992;167:863-872.
64. Romero R, Gomez R, Galasso M, et al. The natural interleukin-1 receptor antagonist in the fetal, maternal, and amniotic fluid compartments: the effect of gestational age, fetal gender, and intrauterine infection. *Am J Obstet Gynecol.* 1994;171:912-921.
65. Romero R, Mazor M, Sepulveda W, Avila C, Copeland D, Williams J. Tumor necrosis factor in preterm and term labor. *Am J Obstet Gynecol.* 1992;166:1576-1587.
66. Opsjln SL, Wathen NC, Tingulstad S, et al. Tumor necrosis factor, interleukin-1, and interleukin-6 in normal human pregnancy. *Am J Obstet Gynecol.* 1993;169:397-404.
67. Maymon E, Ghezzi F, Edwin SS, et al. The tumor necrosis factor alpha and its soluble receptor profile in term and preterm parturition. *Am J Obstet Gynecol.* 1999;181:1142-1148.
68. Saito S, Kasahara T, Kato Y, Ishihara Y, Ichijo M. Elevation of amniotic fluid interleukin 6 (IL-6), IL-8 and granulocyte colony stimulating factor (G-CSF) in term and preterm parturition. *Cytokine.* 1993;5:81-88.
69. Andrews WW, Hauth JC, Goldenberg RL, Gomez R, Romero R, Cassell GH. Amniotic fluid interleukin-6: correlation with upper genital tract microbial colonization and gestational age in women delivered after spontaneous labor versus indicated delivery. *Am J Obstet Gynecol.* 1995;173:606-612.
70. Romero R, Ceska M, Avila C, Mazor M, Behnke E, Lindley I. Neutrophil attractant/activating peptide-1/interleukin-8 in term and preterm parturition. *Am J Obstet Gynecol.* 1991;165:813-820.
71. Olah KS, Vince GS, Neilson JP, Deniz G, Johnson PM. Interleukin-6, interferon-gamma, interleukin-8, and granulocyte-macrophage colony stimulating factor levels in human amniotic fluid at term. *J Reprod Immunol.* 1996;32:89-98.
72. Romero R, Gomez R, Galasso M, et al. Macrophage inflammatory protein-1 alpha in term and preterm parturition: effect of microbial invasion of the amniotic cavity. *Am J Reprod Immunol.* 1994;32:108-113.
73. Dudley DJ, Hunter C, Mitchell MD, Varner MW. Elevations of amniotic fluid macrophage inflammatory protein-1 alpha concentrations in women during term and preterm labor. *Obstet Gynecol.* 1996;87:94-98.
74. Esplin MS, Romero R, Chaiworapongsa T, et al. Amniotic fluid levels of immunoreactive monocyte chemoattractant protein-1 increase during term parturition. *J Matern Fetal Neonatal Med.* 2003;14:51-56.
75. Athayde N, Romero R, Maymon E, et al. A role for the novel cytokine RANTES in pregnancy and parturition. *Am J Obstet Gynecol.* 1999;181:989-994.
76. Gotsch F, Romero R, Kusanovic JP, et al. The anti-inflammatory limb of the immune response in preterm labor, intra-amniotic infection/inflammation, and spontaneous parturition at term: a role for interleukin-10. *J Matern Fetal Neonatal Med.* 2008;21:529-547.
77. Hamill N, Romero R, Gotsch F, et al. Exodus-1 (CCL20): evidence for the participation of this chemokine in spontaneous labor at term, preterm labor, and intrauterine infection. *J Perinat Med.* 2008;36:217-227.
78. Keelan JA, Blumenstein M, Helliwell RJ, Sato TA, Marvin KW, Mitchell MD. Cytokines, prostaglandins and parturition—a review. *Placenta* 2003;24(Suppl A):S33-S46.
79. Martinez-Varea A, Romero R, Xu Y, et al. Clinical chorioamnionitis at term VII: the amniotic fluid cellular immune response. *J Perinat Med.* 2016; doi: 10.1515/jpm-2016-0225. [Epub ahead of print].
80. Keelan JA, Marvin KW, Sato TA, Coleman M, McCowan LM, Mitchell MD. Cytokine abundance in placental tissues: evidence of inflammatory activation in gestational membranes with term and preterm parturition. *Am J Obstet Gynecol.* 1999;181:1530-1536.
81. Gomez-Lopez N, Vega-Sanchez R, Castillo-Castrejon M, Romero R, Cubeiro-Arreola K, Vadillo-Ortega F. Evidence for a role for the adaptive immune response in human term parturition. *Am J Reprod Immunol.* 2013;69:212-230.
82. Romero R, Mazor M, Tartakovsky B. Systemic administration of interleukin-1 induces preterm parturition in mice. *Am J Obstet Gynecol.* 1991;165:969-971.
83. Gravett MG, Witkin SS, Haluska GJ, Edwards JL, Cook MJ, Novy MJ. An experimental model for intraamniotic infection and preterm labor in rhesus monkeys. *Am J Obstet Gynecol.* 1994;171:1660-1667.
84. Witkin SS, Gravett MG, Haluska GJ, Novy MJ. Induction of interleukin-1 receptor antagonist in rhesus monkeys after intraamniotic infection with group B streptococci or interleukin-1 infusion. *Am J Obstet Gynecol.* 1994;171:1668-1672.
85. Romero R, Durum S, Dinarello CA, Oyarzun E, Hobbins JC, Mitchell MD. Interleukin-1 stimulates prostaglandin biosynthesis by human amnion. *Prostaglandins.* 1989;37:13-22.
86. Hertelendy F, Romero R, Molnar M, Todd H, Baldassare JJ. Cytokine-initiated signal transduction in human myometrial cells. *Am J Reprod Immunol.* 1993;30:49-57.
87. Hertelendy F, Rastogi P, Molnar M, Romero R. Interleukin-1beta-induced prostaglandin E2 production in human myometrial cells: role of a pertussis toxin-sensitive component. *Am J Reprod Immunol.* 2001;45:142-147.
88. Belt AR, Baldassare JJ, Molnar M, Romero R, Hertelendy F. The nuclear transcription factor NF-kappaB mediates interleukin-1beta-induced expression of cyclooxygenase-2 in human myometrial cells. *Am J Obstet Gynecol.* 1999;181:359-366.
89. Watari M, Watari H, DiSanto ME, Chacko S, Shi GP, Strauss JF 3rd. Pro-inflammatory cytokines induce expression of matrix-metabolizing enzymes in human cervical smooth muscle cells. *Am J Pathol.* 1999;154:1755-1762.
90. Pineles BL, Romero R, Montenegro D, et al. "The inflammasome" in human parturition. *Reprod Sci.* 2007;14:59A.
91. Gotsch F, Romero R, Chaiworapongsa T, et al. Evidence of the involvement of caspase-1 under physiologic and pathologic cellular

- stress during human pregnancy: a link between the inflammasome and parturition. *J Matern Fetal Neonatal Med.* 2008;21:605–616.
92. Young A, Thomson AJ, Ledingham M, Jordan F, Greer IA, Norman JE. Immunolocalization of proinflammatory cytokines in myometrium, cervix, and fetal membranes during human parturition at term. *Biol Reprod.* 2002;66:445–449.
93. Osman I, Young A, Ledingham MA, et al. Leukocyte density and proinflammatory cytokine expression in human fetal membranes, decidua, cervix and myometrium before and during labour at term. *Mol Hum Reprod.* 2003;9:41–45.
94. Hamilton SA, Tower CL, Jones RL. Identification of chemokines associated with the recruitment of decidual leukocytes in human labour: potential novel targets for preterm labour. *PLoS ONE.* 2013;8:e56946.
95. Stephen GL, Lui S, Hamilton SA, et al. Transcriptomic profiling of human choriodecidua during term labor: inflammation as a key driver of labor. *Am J Reprod Immunol.* 2015;73:36–55.
96. Gomez-Lopez N, Tong WC, Arenas-Hernandez M, et al. Chemotactic activity of gestational tissues through late pregnancy, term labor, and RU486-induced preterm labor in Guinea pigs. *Am J Reprod Immunol.* 2015;73:341–352.
97. Svensson-Arvelund J, Ernerudh J. The Role of Macrophages in Promoting and Maintaining Homeostasis at the Fetal-Maternal Interface. *Am J Reprod Immunol.* 2015;74:100–109.
98. Xu Y, Romero R, Miller D, et al. An M1-like Macrophage Polarization in Decidual Tissue during Spontaneous Preterm Labor That Is Attenuated by Rosiglitazone Treatment. *J Immunol.* 2016;196:2476–2491.
99. Ning F, Liu H, Lash GE. The Role of Decidual Macrophages During Normal and Pathological Pregnancy. *Am J Reprod Immunol.* 2016;75:298–309.
100. Romero R, Wu YK, Brody DT, Oyarzun E, Duff GW, Durum SK. Human decidua: a source of interleukin-1. *Obstet Gynecol.* 1989;73:31–34.

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

How to cite this article: Gomez-Lopez N, Romero R, Xu Y, et al. Inflammasome assembly in the chorioamniotic membranes during spontaneous labor at term. *Am J Reprod Immunol.* 2017;77:e12648. <https://doi.org/10.1111/aji.12648>