

REGULAR ARTICLE

Biventricular function on early echocardiograms in neonatal hypoxic–ischaemic encephalopathy

Sanjeev Aggarwal, Girija Natarajan (gnatara@med.wayne.edu)

Department of Pediatrics, Wayne State University, Children's Hospital of Michigan, Detroit, MI, USA

Keywords

Echocardiogram, Hypoxic–ischaemic encephalopathy, Pulmonary hypertension

Correspondence

Girija Natarajan MD, Division of Neonatal-Perinatal Medicine, Department of Pediatrics, Children's Hospital of Michigan, 3901 Beaubien Blvd, Detroit, MI, USA.

Tel: 313 745 5638 |

Fax: 313-745-5867 |

Email: gnatara@med.wayne.edu

Received

12 October 2016; revised 30 January 2017; accepted 30 March 2017.

DOI:10.1111/apa.13866

ABSTRACT**Aim:** To compare early (<24 hours) echocardiograms (ECHOs) in infants with perinatal hypoxic–ischaemic encephalopathy (HIE) undergoing (i) therapeutic hypothermia (TH), (ii) normothermia and (iii) normal controls.**Methods:** This was a single-centre retrospective review of clinical early ECHOs of term infants with moderate or severe HIE and controls (with a normal ECHO <72 hours of age). Right (RVO) and left ventricular output (LVO), RV and LV myocardial performance index (MPI), systolic to diastolic duration ratio (S/D) and eccentricity indices (EI) in systole and diastole were compared using ANOVA.**Results:** Among infants with HIE (n = 56, 38 in the TH and 18 in normothermia groups), 14 (25%) infants died and 42 survived. Significantly elevated biventricular MPI, lower RVO and LVO and pulmonary hypertension (abnormal EI, higher RV S/D and bidirectional or right-to-left ductal shunt) were found in groups with HIE, compared to controls (n = 35). LV MPI was lower in HIE-TH, compared to the HIE-normothermia group. Infants with HIE who died (n = 14) had a significantly lower Eld [0.77 (0.09) vs. 0.83 (0.08), p = 0.021] compared to survivors (n = 42).**Conclusion:** Infants with perinatal HIE have ventricular dysfunction; those who died had significantly lower Eld than survivors; this association needs to be further validated.**INTRODUCTION**

The incidence of neonatal hypoxic–ischaemic encephalopathy (HIE) due to asphyxia is 1–4/1000 live births in the developed world (1). HIE is the fifth commonest cause of death in children below five years and accounts for 23% of all neonatal deaths worldwide each year (2). The Cochrane meta-analysis of randomised controlled trials on therapeutic hypothermia (TH) in HIE demonstrates a beneficial effect, with a reduction in mortality or major neurodevelopmental disability at 18 months of age (RR: 0.76; 95% CI: 0.65–0.89) (3). Despite this, mortality remains significant at 26% (4). Cardiac dysfunction may be a primary cause of death in this population.

The major randomised controlled trials of TH in HIE have reported that the majority of infants have hypotension during TH (5–7). In the National Institute of Child Health and Human Development Neonatal Research Network TH trial, 59% of infants required pressor support at any of four

time points until 72 hours and 19% were on pressor support throughout the 72 hours of TH (5). The reported rates of hypotension and severe hypotension were 55% and 3% in cooled infants in the CoolCap study (6). In the Total Body Hypothermia for Neonatal Encephalopathy Trial, persistent hypotension, defined as mean BP < 40 mm Hg, was noted in 77% of cooled infants (7). In the Infant Cooling Evaluation trial of 221 infants, hypotension requiring inotropes was noted in 46% of infants (8). In small studies, up to 60%–80% of neonates with HIE have cardiac dysfunction on echocardiograms (ECHOs), compared to controls (9–13). The few studies which have evaluated

Abbreviations

ECHO, Echocardiogram; ECMO, Extracorporeal membrane oxygenator; Eld, Eccentricity index in diastole; EIs, Eccentricity index in systole; LVO, Left ventricular output; MPI, Myocardial performance index; PPHN, Persistent pulmonary hypertension of the newborn; RVO, Right ventricular output; S/D, Ratio of systolic to diastolic durations.

Key notes

- Limited data suggest decreased cardiac output in infants with perinatal hypoxic–ischaemic encephalopathy during therapeutic hypothermia.
- We compared early (<24 hours) echocardiogram measures in groups of infants with hypoxic–ischaemic encephalopathy undergoing (i) therapeutic hypothermia (TH), (ii) normothermia and (iii) normal controls and evaluated their association with death.
- Infants with perinatal HIE have profound cardiac dysfunction, compared to controls. Those who died had significantly lower eccentricity index in diastole.

ECHOs of infants with HIE undergoing TH have demonstrated a decrease in cardiac output (CO), heart rate (HR) and stroke volume during TH with recovery following rewarming (14–17). In this study, we sought to compare multiple early (within 24 hours of age) ECHO measures of biventricular function in groups of infants with HIE undergoing (i) TH or (ii) normothermia and (iii) normal controls and to evaluate their association, if any, with death.

PATIENTS AND METHODS

This was a retrospective review of ECHOs performed for clinical indications at <24 hours of age in term or late preterm (≥ 36 weeks gestational age) infants with moderate or severe HIE and admitted to the neonatal intensive care units (NICUs) at Children's Hospital of Michigan or Hutzel Women's Hospital between July 2002 and December 2012. Study infants were identified from the NICU database using 'HIE (diagnosis) and ECHO (Procedure)' as search words. The study was approved by the Institutional Review Board of Wayne State University as was waiver of parental consent. Moderate or severe HIE was defined using NICHD screening criteria along with moderate or severe encephalopathy on neurologic examination (modified Sarnat staging) or seizures (5). At our centre, infants with moderate or severe HIE have received whole body TH to 33.5°C for 72 hours with the Cincinnati Subzero Hypothermia system initiated within six hours of age since 2006. Rewarming is performed at 0.5°C per hour until the oesophageal temperature is ≥ 36.5 °C for four hours. Healthy controls were identified from the ECHO database as having normal ECHOs within 72 hours of age for evaluation of a murmur. Infants with major congenital or chromosomal abnormalities, severe growth restriction (≤ 1800 g birthweight), moribund infants in whom further aggressive treatment was limited and congenital heart disease other than a ventricular septal defect, patent foramen ovale or patent ductus arteriosus (PDA) were excluded.

ECHOs were performed using neonatal transducers on the Philips machine and reviewed by a single paediatric cardiologist (SA), who was blinded to clinical details. Each ECHO included M-mode, two-dimensional (subcostal, apical, left parasternal and suprasternal notch views), Doppler and tissue Doppler imaging. The following ECHO functional measurements were performed offline:

- Systolic function

- 1 *LV and RV output (LVO and RVO)*: Aortic and pulmonary valve annulus diameters and velocity time integral (VTI) were measured from parasternal long-axis and apical five-chamber views, respectively, on Doppler ultrasound (18). LVO and RVO were calculated using standard formulae. HR was taken as the average of three readings during the ECHO.

- 2 *Fractional Shortening (FS)* was obtained on M-mode imaging as the ratio of the difference between the LV end-diastolic and end-systolic diameters to the LV end-diastolic diameter (18). FS has limitations in neonates because of high right-sided pressures and reduced septal motion (19,20).

- RV and LV Global myocardial function

- 1 *LV and RV MPI* were assessed by dividing the sum of isovolumic contraction and relaxation times by the ejection time and standard formula (21). MPI is inversely related to myocardial function, and an increase in MPI indicates impaired global myocardial function. It is independent of HR and blood pressure, relatively load-independent and is unaffected by angle insonation or PDA.
- 2 *RV S/D ratio* was measured offline in triplicate from the best Doppler signal of tricuspid regurgitation (TR) on either apical four-chamber view or parasternal long-axis view. The systolic duration was calculated from the onset to the termination of TR and diastolic duration as the time between two jets of TR (22,23).

- PDA: A high parasternal ductal view was utilised to assess presence, size and direction of ductal shunt.
- LV mass: LV internal dimension, posterior wall thickness, interventricular septum thickness and LV mass were measured in systole and end-diastole from M-mode ECHOs according to American Society of Echocardiography recommendations (24). Left ventricular mass index (LVMI) was indexed to height in cm^2 .⁷
- Eccentricity index (EI): The EI in diastole (EId) and systole (EIs) was calculated as the ratio of perpendicular diameters in parasternal short axis at the level of papillary muscle, using the method of Ryan et al. (25).

Data collection

Demographic data (gestational age, gender) and clinical characteristics (birthweight, severity of HIE, cord pH, mode of delivery, Apgar scores at five minutes, perinatal events) need for inotropes, high-frequency ventilation, nitric oxide and length of stay were collected.

Data analyses

Calculated means of three cardiac cycles were used for ECHO parameters. ECHO data were described as mean (SD), median (IQR) or 95% CI as appropriate. Comparisons of ECHO measures between groups were made using t-tests and analysis of variance (ANOVA) with Bonferroni correction for post hoc comparisons for continuous variables. Chi-square test was used for comparisons of categorical data. Statistical analysis was performed using SPSS version 19 (SPSS, Chicago, IL, USA) and a p value <0.05 taken as significant.

RESULTS

Our study cohort included 56 infants with moderate or severe HIE with a mean (SD) GA of 38.7 (1.6) weeks and birthweight of 3.385 (0.719) kg. Males comprised 50% of the cohort; 38 infants underwent TH and 18 did not, either because they ($n = 12$) were born before the TH era or admitted after 6 hours of age ($n = 6$). ECHO was performed at a mean (SD) age of 10.4 (4.8) hours of age. C-section was the mode of delivery in the majority [38 (67.9%)] of cases. The documented perinatal events included uterine rupture [5 (8.9%)], foetal decelerations [21 (37.5%)], placental abruption [3 (5.4%)] and meconium stained amniotic fluid [2 (3.6%)] among others. The median (IQR) cord pH was 6.93 (6.82–7.06) and 5-minute Apgar score was 4 (1–6). Pressors were required in 48 (85.7%) infants with 34 (60.7%) infants requiring two or more pressors. High-frequency ventilation was administered in 25 (44.6%) infants and inhaled nitric oxide in 30 (53.6%) infants. Fourteen (25%) infants died, seven of whom had severe HIE, three moderate HIE, and four had missing information. Among the 42 infants who survived, the mean (SD) length of hospital stay was 24.0 (16.2) days and the length of ventilation was 15.1 (11.7) days.

There were no significant differences in clinical characteristics between infants with HIE who underwent TH or normothermia except for a higher requirement of inhaled nitric oxide in the normothermia group (Table 1). The normal control infants ($n = 35$), 17 (48.6%) of whom were males, had a mean (SD) birthweight of 3.103 (0.638) kg. A comparison of ECHO measures in the two groups of infants

with HIE who underwent TH and normothermia and normal controls is shown (Table 2). There was no significant difference in mean (SD) birthweight or GA between the three groups. Significantly elevated biventricular MPI, lower biventricular VO and EI in infants with HIE, compared to normal controls, confirmed cardiac dysfunction in this population. The mean HR among infants with HIE who underwent normothermia was significantly higher than in those who underwent TH. The only ECHO measure that was significantly different in the two groups of infants with HIE was LV MPI. Among the controls, three (8.6%) infants had mild tricuspid regurgitation (TR); in infants with HIE who underwent TH and were normothermic, 11 (29.0%) and four (22.2%) had mild and eight (21.1%) and four (22.2%) had moderate TR (combined $p = 0.002$). One infant with HIE in the normothermia had severe TR. LVO below 150 ml/kg/min and RV S/D ratio above 1.3 were also analysed in view of previous data establishing these cut-offs (18, 23). In infants with HIE, none of the ECHO parameters were associated with clinical severity measures such as need for high-frequency ventilation, nitric oxide, pressors or severity of HIE. Table 3 shows a comparison of ECHO measures between the groups of infants with HIE who died ($n = 14$) and survived ($n = 42$). The only significant difference was in the EID.

DISCUSSION

This was a retrospective analysis of ECHOs performed within 24 hours of age for clinical indications in a cohort of infants with moderate or severe HIE. The majority (54%) of our study cohort received inhaled nitric oxide therapy for pulmonary hypertension and 86% of infants were on pressor therapy. About half the infants in whom data were available had severe HIE and 25% died.

On a detailed early ECHO, infants with HIE had evidence of decreased biventricular outputs, global systolic and diastolic dysfunction reflected by elevated biventricular MPI and pulmonary hypertension (abnormal EI, higher RV S/D and bidirectional or right-to-left shunt across the PDA), compared to normal controls. The FS values were comparable to those of control infants. Previous studies in small numbers of asphyxiated infants have reported ventricular dysfunction as reflected in a reduced FS, tricuspid insufficiency or elevated LV and RV MPI, compared to controls, although results of specific measures have varied (9,11,12). Other ECHO measures such as longitudinal peak systolic strain and peak systolic strain rate by tissue Doppler in 18 segments of the heart on day 1 of life in 20 asphyxiated neonates were significantly lower than in the 48 healthy term controls. FS was similar in the two groups [29.2% (26.8, 31.5) vs. 29.0% (27.9, 30.1); $p = 0.874$] (10).

A few previous studies have evaluated ECHO measures of ventricular function, coronary and superior venacaval flow in infants with HIE who underwent TH (14–17). As the most notable changes in hemodynamics following perinatal asphyxia occur in the first days of life (26), we limited our study to ECHOs within the initial 24 hours of

Table 1 A comparison of clinical characteristics between infants with HIE who underwent TH or normothermia

Mean (SD) or n (%)	HIE-TH (n = 38)	HIE-Normothermia (n = 18)	p value
Birthweight (g)	3365 (689)	3429 (803)	0.766
GA (weeks)	38.7 (1.5)	38.7 (2.0)	0.949
Male gender	18 (47.4%)	10 (58.8%)	0.562
C-section	29 (76.3%)	9 (50%)	0.116
Perinatal event			0.733
Uterine rupture	3 (7.9%)	2 (11.2%)	
Foetal decelerations	14 (36.8%)	7 (38.9%)	
Placental abruption	2 (5.2%)	1 (5.6%)	
Cord pH	6.89 (0.18)	6.94 (0.18)	0.432
5-min Apgar <5	27 (71.1%)	9 (50%)	0.229
Severe HIE	12/24 (50%)	3/7 (42.8%)	0.693
Pressors in initial 72 hours			0.087
None	6 (15.8%)	0 (0%)	
1 pressor	11 (29%)	3 (16.7%)	
2 or more pressors	21 (55.3%)	15 (83.3%)	
Age at ECHO (hours)	10.4 (5.1)	12.5 (1.4)	0.579
Nitric oxide	17 (44.7%)	13 (72.2%)	0.041
High-frequency oscillation	15 (39.5%)	10 (55.6%)	0.245
Days of ventilation	13.0 (12.7)	12.4 (8.6)	0.868
Days of hospital stay	19.6 (17.2)	19.7 (13.9)	0.988
Death	8 (21.1%)	6 (33.3%)	0.322

Table 2 A comparison of ECHO measures in the two groups of infants with HIE who underwent TH and normothermia and normal controls (n = 35)

Mean (SD)	Normal controls (n = 35)	HIE-TH (n = 38)	HIE-Normothermia (n = 18)	Combined p value
HR (bpm)	137 (17)	124 (32)*	161 (23)**	0.0001
RVO (mL/kg)	343 (123)***	195 (73)	216 (83)**	0.0001
RVO <150 mL/kg	0	11 (29.0%)	4 (11.1%)	0.012
LVO (mL/kg)	393 (107)***	269 (87)	292 (101)**	0.0001
LVO <150 mL/kg	0	3 (7.9%)	2 (11.1%)	0.169
FS (%)	0.38 (0.07)	0.35 (0.07)	0.34 (0.09)	0.074
RV MPI	0.25 (0.08)***	0.58 (0.21)	0.66 (0.23)**	0.0001
LV MPI	0.29 (0.06)***	0.47 (0.16)*	0.60 (0.13)**	0.0001
RV S/D	1.10 (0.14)***	1.58 (0.44)	1.73 (0.26)**	0.0001
RV S/D > 1.3	2 (5.7%)	24 (63.2%)	15 (83.3%)	0.0001
LV mass	7.25 (2.14)	8.05 (2.12)	7.41 (2.89)	0.304
LVMI (g/m 2.7)	46.65 (12.40)	51.49 (12.72)	46.45 (16.93)	0.322
Els	0.87 (0.16)***	0.75 (0.11)	0.76 (0.13)**#	0.001
Eld	0.90 (0.09)***	0.81 (0.09)	0.81 (0.07)**	0.0001
PDA	12 (34.3%)	33 (86.8%)	13 (72.2%)	0.0001
PDA shunt				0.0001
Right-left	0	0	2 (11.1%)	
Bidirectional	0	21 (55.3%)	5 (27.8%)	
PDA size (cm)	0.19 (0.06)***	0.33 (0.15)	0.35 (0.13)**	0.002

*Post hoc analyses p < 0.01 HIE-TH compared to HIE-normothermia.

**p < 0.01 HIE-normothermia compared to controls.

***p < 0.01 HIE-TH compared to controls.

**#p = 0.021 HIE-normothermia compared to controls.

Table 3 A comparison of ECHO measures in the two groups of infants with HIE who survived (n = 42) and died (n = 14)

Mean (SD) or n (%)	Survivors (n = 42)	Infants who died (n = 14)
HR (bpm)	133 (34)	139 (32)
RVO (mL/kg)	200.4 (73.5)	198.9 (87.0)
LVO (mL/kg)	273.1 (93.1)	279.2 (87.0)
FS (%)	0.35 (0.07)	0.34 (0.09)
RV MPI	0.59 (0.20)	0.63 (0.26)
LV MPI	0.49 (0.15)	0.56 (0.18)
RV S/D	1.59 (0.40)	1.72 (0.42)
RV S/D >1.3	25 (59.5%)	13 (92.9%)
LV mass	7.92 (2.16)	7.45 (3.02)
LVMI (g/m 2.7)	50.66 (11.67)	47.78 (20.26)
Els	0.77 (0.11)	0.72 (0.11)
Eld	0.83 (0.08)	0.77 (0.09)*

*p = 0.021.

life and for those who underwent TH, within a few hours of induction of TH. We found that infants who underwent TH had decreased biventricular outputs, elevated MPIs and abnormal EIs, compared to controls. The only significant differences between infants who underwent TH and normothermia were the significantly lower HRs and LV MPI in the TH group. Whether the improvement in LV MPI, a sensitive marker of function, represents some degree of cardioprotection remains to be determined. Consistent with our data, the few previous studies carried out during the TH era have shown lower CO, HR, superior venacaval and

coronary flows and Doppler deformation indices of myocardial function and impaired peak systolic strain rate and LV MPI, compared to controls (14–17,26). Other studies have also shown that the decrease in ventricular function in infants with HIE is transient during TH, with subsequent recovery (14,16).

We also evaluated measures of pulmonary hypertension in infants with HIE receiving TH in the current study. We found the EIs and RV S/D ratio to be significantly deranged, compared to controls. The directionality of the shunt across PDA confirmed the pulmonary hypertension in infants with HIE. The association between persistent pulmonary hypertension of the newborn and perinatal asphyxia either because of direct effects of hypoxia and acidosis on pulmonary vascular resistance or due to indirect effects of coexisting morbidities such as meconium aspiration syndrome and sepsis/pneumonia has been previously recognised (27). Pulmonary hypertension, in turn, is associated with ventricular dysfunction. In one previous study, measures of pulmonary hypertension such as pulmonary arterial diastolic pressure, pulmonary arterial resistance and pulmonary arterial resistance/ systemic resistance ratio were all significantly elevated in 40 term infants with HIE, compared to 40 healthy controls on day 1 of life (28).

When we compared infants with HIE who died and survivors, the mean Eld was significantly different between groups. A few previous studies have examined ECHO measures as biomarkers of mortality with largely negative results. In one study of 34 term newborns with mild to severe HIE, only one of nine infants who died had mild

decrease in ejection fraction in the initial 24–48 hours of life (29). In another study of 25 asphyxiated and 20 nonasphyxiated term infants, FS and Doppler tissue imaging measures performed during the initial 72 hours of age did not show any significant predictive value for mortality (9). We have previously shown RV S/D ratio, an index of RV function, was significantly higher in infants with persistent pulmonary hypertension or congenital diaphragmatic hernia who died or required ECMO, compared to survivors without ECMO (22,30). RV S/D ratio >1.3 had a sensitivity of 93% for prediction of death (22). In the current data set, while the proportion of infants with RV S/D > 1.3 was greater in those who died, the association did not reach statistical significance. The correlation of EID with mortality may be a reflection of pulmonary hypertension in infants with severe, rather than moderate HIE.

We acknowledge the limitations of our study. As we evaluated ECHOs performed for clinical indications, our cohort included a group of 'sick' infants. While such children have not been previously studied in any detail, this was a selection bias. As the numbers of infants who underwent normothermia were relatively small, the comparative effects of TH on ventricular function and hemodynamics could be studied in a limited way. Whether the cardiac dysfunction was related to intrapartum birth events resulting in HIE, effects of HIE itself or TH could not be ascertained. The difference in ages between the groups with HIE and normal controls (<72 hours of age) was a source of potential bias. Nonetheless, the current observational study adds to our understanding of the cardiac function in infants with HIE undergoing TH through a comprehensive ECHO within the initial day of life. Specifically, novel ECHO measures of pulmonary hypertension such as EID appear to correlate with survival outcomes in this population. Further studies are needed to validate this association in larger numbers of infants.

CONFLICT OF INTEREST

The authors have no conflict of interests to declare.

FUNDING

None of the authors have any relevant financial disclosures. The first draft of the article was written by Girija Natarajan, who did not receive any honorarium or grant for the purpose.

References

- Zanelli SA. Hypoxic-ischemic encephalopathy—Diseases and conditions. 2015. <http://emedicine.medscape.com/article/973501>.
- Bryce J, Boschi-Pinto C, Shibuya K, Black RE. WHO estimates of the causes of death in children. *Lancet* 2005; 365: 1147–52.
- Jacobs S, Hunt R, Tarnow-Mordi W, Inder T, Davis P. Cooling for newborns with hypoxic ischaemic encephalopathy. *Cochrane Database Syst Rev* 2013; 1: CD003311.
- Tagin MA, Woolcott CG, Vincer MJ, Whyte RK, Stinson DA. Hypothermia for neonatal hypoxic ischemic encephalopathy. An updated systematic review and meta-analysis. *Arch Pediatr Adolesc Med* 2012; 166: 558–66.
- Shankaran S, Pappas A, Laptook AR, McDonald SA, Ehrenkranz RA, Tyson JE, et al. NICHD Neonatal Research Network. Outcomes of safety and effectiveness in a multicenter randomized, controlled trial of whole-body hypothermia for neonatal hypoxic-ischemic encephalopathy. *Pediatrics* 2008; 122: e791–8.
- Battin MR, Thoresen M, Robinson E, Polin RA, Edwards AD, Gunn AJ. Does head cooling with mild systemic hypothermia affect requirement for blood pressure support? *Pediatrics* 2009; 123: 1031–6.
- Azzopardi DV, Strohm B, Edwards AD, Dyet L, Halliday HL, Juszczak E. Moderate hypothermia to treat perinatal asphyxial encephalopathy. *N Engl J Med* 2009; 361: 1349–58.
- Jacobs SE, Morley CJ, Inder TE, Stewart MJ, Smith KR, McNamara PJ, et al. Whole body hypothermia for term and near-term newborns with hypoxic ischemic encephalopathy. *Arch Pediatr Adolesc Med* 2011; 165: 692–700.
- Matter M, Abdel-Hady H, Attia G, Hafez M, Seliem W, Al-Arman M. Myocardial performance in asphyxiated full-term infants assessed by Doppler tissue imaging. *Pediatr Cardiol* 2010; 31: 634–42.
- Nestaas E, Støylen A, Brunvand L, Fugelseth D. Longitudinal strain and strain rate by tissue Doppler are more sensitive indices than fractional shortening for assessing the reduced myocardial function in asphyxiated neonates. *Cardiol Young* 2011; 21: 1–7.
- Barberi I, Calabro MP, Cordaro S, Gitto E, Sottile A, Prudente D, et al. Myocardial ischemia in neonates with perinatal asphyxia. *Eur J Pediatr* 1999; 158: 742–7.
- Szymankiewicz M, Matuszczak-Wleklak M, Hodgman JE, Gadzinowski J. Usefulness of cardiac troponin T and echocardiography in the diagnosis of hypoxic myocardial injury of full-term neonates. *Biol Neonate* 2005; 88: 19–23.
- Turner Gomes SA, Izukawa T, Rowe RD. Persistence of atrioventricular valve regurgitation and electrocardiographic abnormalities following transient myocardial ischemia of the newborn. *Pediatr Cardiol* 1989; 10: 191–4.
- Gebauer CM, Knuepfer M, Robel-Tillig E, Pulzer F, Vogtmann C. Hemodynamics among neonates with hypoxic-ischemic encephalopathy during whole-body hypothermia and passive rewarming. *Pediatrics* 2006; 117: 843–50.
- Czernik C, Rhode S, Helfer S, Schmalisch G, Beuhrer C. Left ventricular longitudinal strain and strain rate measured by 2-D speckle tracking echocardiography in neonates during whole body hypothermia. *Ultrasound Med Biol* 2013; 39: 1343–9.
- Hochwald O, Jabr M, Osiovič H, Miller SP, McNamara PJ, Lavoie PM. Preferential cephalic redistribution of left ventricular cardiac output during therapeutic hypothermia for perinatal hypoxic-ischemic encephalopathy. *J Pediatr* 2014; 164: 999–1004.
- Sehgal A, Wong F, Mehta S. Reduced cardiac output and its correlation with coronary blood flow and troponin in asphyxiated infants treated with therapeutic hypothermia. *Eur J Pediatr* 2012; 171: 1511–7.
- Mertens L, Seri I, Marek J, Arlettaz R, Barker P, McNamara P, et al. Targeted neonatal echocardiography in the neonatal intensive care unit: practice guidelines and recommendations for training. *J Am Soc Echocardiogr* 2011; 24: 1057–78.
- Lee LA, Kimball TR, Daniels SR, Khoury P, Meyer RA. Left ventricular mechanics in the preterm infant and their effect on the measurement of cardiac performance. *J Pediatr* 1992; 120: 114–9.

20. Kluckow M. Functional echocardiogram in assessment of the cardiovascular system in asphyxiated neonates. *J Pediatr* 2011; 158(2 Suppl): e13–8.
21. Ichihashi K, Yada Y, Takahashi N, Honma Y, Momoi M. Utility of a Doppler-derived index combining systolic and diastolic performance (Tei index) for detecting hypoxic cardiac damage in newborns. *J Perinat Med* 2005; 33: 549–52.
22. Aggarwal S, Stockman PT, Klein MD, Natarajan G. The right ventricular systolic to diastolic duration ratio: a simple prognostic marker in congenital diaphragmatic hernia? *Acta Paediatr* 2011; 100: 1315–8.
23. Friedberg MK, Silverman NH. The systolic to diastolic duration ratio in children with hypoplastic left heart syndrome: a novel Doppler index of right ventricular function. *J Am Soc Echocardiogr* 2007; 20: 749–55.
24. Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, et al. Recommendations for chamber quantification: a report from the American Society of Echocardiography's Guidelines and Standards Committee and the Chamber Quantification Writing Group, developed in conjunction with the European Association of Echocardiography, a branch of the European Society of Cardiology. *J Am Soc Echocardiogr* 2005; 18: 1440–63.
25. Ryan T, Petrovic O, Dillon JC, Feigenbaum H, Conley MJ, Armstrong WF. An echocardiographic index for separation of right ventricular volume and pressure overload. *J Am Coll Cardiol* 1985; 5: 918–27.
26. Nestaas E, Skranes JH, Støylen A, Brunvand L, Fugelseth D. The myocardial function during and after whole-body therapeutic hypothermia for hypoxic-ischemic encephalopathy, a cohort study. *Early Hum Dev* 2014; 90: 247–52.
27. Lapointe A, Barrington KJ. Pulmonary hypertension and the asphyxiated newborn. *J Pediatr* 2011; 158: e19–24.
28. Liu J, Feng ZC. Changes in pulmonary arterial pressure in term-infants with hypoxic-ischemic encephalopathy. *Pediatr Int* 2009; 51: 786–9.
29. Kanik E, Ozer EA, Bakiler AR, Aydinlioglu H, Dorak C, Dogrusoz B, et al. Assessment of myocardial dysfunction in neonates with hypoxic-ischemic encephalopathy: Is it a significant predictor of mortality? *J Matern Fetal Neonatal Med* 2009; 22: 239–42.
30. Aggarwal S, Natarajan G. Echocardiographic correlates of persistent pulmonary hypertension of the newborn. *Early Hum Dev* 2015; 91: 285–9.