

A 37-Year-Old Man With Primary Antiphospholipid Syndrome Presenting With Respiratory Distress and Worsening Toe Ischemia

NAYEF M. KAZZAZ, ALLECIA M. WILSON, RUBA KADO, GEOFFREY D. BARNES, AND JASON S. KNIGHT

CASE PRESENTATION

History of the present illness

This is a 37-year-old man with a history of primary antiphospholipid syndrome (APS), with clinical manifestations including recurrent venous thrombosis, lower-extremity ulceration, and toe ischemia. He also had history of thrombocytopenia and seizure disorder. Laboratory findings were consistent with APS, including prolonged dilute Russell's viper venom time (dRVVT), as well as an abnormal hexagonal phase phospholipid neutralization assay. He also had high levels of both IgG anticardiolipin and IgG anti- β_2 -glycoprotein I (anti- β_2 GPI; both consistently >40 units). All of these laboratory findings were durably positive over many years.

The patient presented with a 3-day history of dyspnea on exertion, now occurring at rest on the day of presentation. He denied productive cough, hemoptysis, chest pain, nausea, vomiting, abdominal pain, change in bowel habits, or dysuria. He also denied arthralgia, photosensitivity, Raynaud's phenomenon, ocular symptoms, mucosal ulceration, alopecia, pleurisy, rash, weakness, or paresthesia. He did report worsening bilateral foot pain for about 1 month prior to presentation, with the pain especially severe on the right. He denied typical symptoms of claudication. For 2

days prior to admission, the pain had been especially severe: present at rest, dull in nature, and nonradiating.

Past medical history and medications

The APS diagnosis was made at age 22 years when he presented with an extensive left lower-extremity deep venous thrombosis (DVT) event involving the common femoral, popliteal, and peroneal veins. He had 2 recurrences in the same extremity within 2 years, both apparently in the setting of subtherapeutic anticoagulation with warfarin.

Around age 30 years, he developed intermittent discoloration of his feet, more notable on the right, as well as hypocomplementemia and thrombocytopenia: C3 level of 74 mg/dl (normal range 83–240), C4 of 11 mg/dl (normal range 13–60), and platelets of 90,000/ μ l. Aspirin 81 mg daily and hydroxychloroquine 400 mg daily were added to his warfarin therapy. Given the hypocomplementemia, the hydroxychloroquine was added by his rheumatologist, although there were no symptoms beyond these laboratory abnormalities to suggest a diagnosis of lupus. At this time, his hematologist also began monitoring and adjusting his warfarin dosing based on chromogenic factor X assay, rather than the international normalized ratio (INR), given concern that the lupus anticoagulant was interfering with INR monitoring. The clinical and laboratory abnormalities all normalized with these various medication adjustments. Also around this time, computed tomography (CT) angiography of the lower extremities demonstrated peripheral arterial disease, moderate to severe in the distal right common iliac artery, and extending to the proximal right external iliac artery. Mild narrowing of the distal left common iliac artery was also present. Regarding risk factors for atherosclerosis, there was no evidence of dyslipidemia or human immunodeficiency virus on multiple laboratory tests. The patient had also never been exposed to regular corticosteroids. Transthoracic echocardiography at that time showed mild mitral regurgitation with no evidence of vegetation on good quality imaging.

At age 31 years, he developed tonic-clonic seizures, which were subsequently well controlled with levetiracetam. Migraine headaches ensued and persisted. His doctors felt

Dr. Kazzaz's work was supported by Security Forces Hospital Program, Ministry of Interior, Riyadh, Saudi Arabia. Dr. Barnes's work was supported by the NIH (grant T32-HL-007853). Dr. Knight's work was supported by career development awards from the NIH (grant K08-AR-066569), the Burroughs Wellcome Fund, the Arthritis National Research Foundation, and the Rheumatology Research Foundation.

Nayef M. Kazzaz, MD, Allecia M. Wilson, MD, Ruba Kado, MD, Geoffrey D. Barnes, MD, Jason S. Knight, MD, PhD: University of Michigan Medical School, Ann Arbor.

Address correspondence to Jason S. Knight, MD, PhD, 5520A MSRB1, 1150 W Medical Center Drive, SPC 5680, Ann Arbor, MI 48109-5680. E-mail: jsknight@umich.edu.

Submitted for publication August 24, 2016; accepted in revised form December 6, 2016.

that both the seizures and migraine headaches could be attributable to APS, but there was no definitive proof.

At age 35 years, he developed a difficult-to-heal left lateral malleolar ulceration, first emerging after minor trauma. Given concerns about fluctuating INR testing and an inability to obtain chromogenic factor X levels locally, he was switched by his local physicians to the factor Xa inhibitor, rivaroxaban. It appears that rivaroxaban was chosen over low molecular weight heparin primarily because of patient preference. The ulceration was an on-and-off issue for the next 2 years, and was managed primarily by the wound clinic. Although the location was a bit unusual, the prevailing opinion was that the ulcer was based in venous stasis more so than arterial insufficiency. His course was otherwise stable. There were no new thrombotic events or notable lab abnormalities (such as hypocomplementemia or thrombocytopenia). His seizure disorder was stable. He continued to work.

Although the patient lacked a symptom history to suggest a concomitant diagnosis of lupus (no arthritis, serositis, glomerulonephritis, or lupus-specific rash), "lupus labs" were obtained from time to time over the years. These revealed borderline positive test results for antinuclear antibody (ANA) by HEp-2 assay and anti-double-stranded DNA (anti-dsDNA) by radioimmunoassay (see below for more details). Extractable nuclear antigen testing was consistently negative for anti-SSA, anti-SSB, anti-RNP, anti-Sm, anti-SmRNP, antichromatin, anti-RiboP, anticentromere, anti-Scl70, and anti-Jo-1 antibodies. Complement levels tended to be normal with the exception mentioned above.

Social and family history

There was no family history of systemic autoimmunity, inflammatory arthritis, coagulation disorders, cardiovascular disease, or malignancy. He was a university graduate, employed, and married with 2 children. He denied smoking or the use of illicit drugs, and reported drinking alcohol occasionally.

Physical examination

Examination on presentation to the hospital was notable for tachycardia (129 beats per minute), tachypnea (36 breaths per minute), hypoxemia (83% on room air), and fever (38.5°C). Pulmonary examination demonstrated decreased breath sounds and coarse crackles at the bilateral bases. On cardiovascular examination, normal first and second heart sounds were noted with no appreciable rubs, gallops, or murmurs. Dorsalis pedis, posterior tibialis, and popliteal artery pulses were absent bilaterally. Femoral pulses were faint. He was noted to have a 2-cm × 3-cm left lateral malleolar ulceration that was stable compared to examination 1 month prior. Additionally, the right first, second, and third toes demonstrated evidence of necrosis, compared to pinkish discoloration on recent outpatient examinations.

Laboratory testing and radiology

His historical laboratory findings were notable for a prolonged dRVVT at 146.6 seconds (normal <42.9 seconds), dRVVT ratio 2.1 (essentially an assessment of phospholipid quenching, normal <1.19), and hexagonal phospholipid neutralization assay prolonged at 144 seconds (normal <7.9 seconds). While he was on warfarin at the time of this

testing, the hexagonal assay involves mixing 1:1 with normal plasma, and prolongation due to factor deficiency is therefore unlikely. IgG anticardiolipin was >100 IgG phospholipid units (normal <22), and IgG anti- β_2 GPI was >100 (normal <20 standard IgG anti- β_2 GPI units). Additionally, ANA was positive at a titer of 1:80 by HEp-2 immunofluorescence assay, with a speckled pattern. Over the years, transient hypocomplementemia was once present (as above). He had also had slight elevations in anti-dsDNA antibodies in the 30s range (normal range <7.0 IU/ml). Urinalysis was typically bland through the years, albeit with a mildly elevated urine protein:creatinine ratio of 0.26 (normal range <0.18).

In the acute setting, laboratory results were notable for a total white blood cell count of 15,000/ μ l with neutrophilia of 12,900/ μ l, new normochromic normocytic anemia (hemoglobin 9.5 gm/dl, decreased from a baseline of 13 gm/dl), and new thrombocytopenia (platelets 68,000/ μ l, decreased from a recently normal baseline). Peripheral blood smear did not show evidence of microangiopathy. Haptoglobin, ferritin, fibrinogen, and D-dimer findings were all normal. A comprehensive metabolic panel was notable for newly elevated creatinine of 1.64 mg/dl, blood urea nitrogen of 40 mg/dl, and a slight increase in aspartate aminotransferase from 32 to 57 IU/liter. Lactate was normal on presentation. Urinalysis was bland with no dysmorphic red blood cells. Urine protein-creatinine ratio showed a slight increase from 0.26 to 0.46 (normal <0.18). The erythrocyte sedimentation rate was 53 mm/hour, and the C-reactive protein level was 15.2 mg/dl (normal <0.6). Both of these tests had been recently normal. Complements C3 and C4 were low at 59 mg/dl and 5 mg/dl, respectively. Anti-dsDNA was positive at a level of 65 IU/ml (the hospital laboratory had recently changed from a radioimmunoassay to an enzyme-linked immunosorbent-based assay, with 30–75 on the new assay considered "borderline positive"). Troponin was elevated at 12.77 ng/ml (normal range <0.29), trending down to 9.86 ng/ml over the next 48 hours. Blood culture findings (both bacterial and fungal) were negative.

Portable chest radiograph demonstrated bilateral patchy opacities. Chest CT angiography (Figure 1) showed extensive multifocal patchy and ground-glass airspace opacities most pronounced at the lung bases where there was partial consolidation, left greater than right. There was no evidence of pulmonary embolism. Electrocardiogram showed sinus




Figure 1. Multifocal patchy and ground-glass airspace opacities. These are most pronounced at the lung bases where there is partial consolidation, left greater than right.

tachycardia, possible left atrial enlargement, and RSR' or QR pattern in V1 suggestive of right ventricular conduction delay. Transthoracic echocardiogram showed a new (as compared to a study 6 years prior) and severe decrease in left and right ventricular function, a dilated inferior vena cava, and worsening mitral regurgitation (previously mild, now moderate). Right ventricular systolic pressure was estimated at 33 mm Hg with right atrial pressure of 10 mm Hg; there was no evidence of tricuspid valve abnormality. Abdominal and retroperitoneal ultrasounds with Doppler were normal. Bilateral ankle-brachial indices were normal; however, an abnormality was noted on toe-brachial indices.

CASE SUMMARY

This is a 37-year-old man who carries a diagnosis of primary APS with presumed associations including recurrent DVT complicated by post-thrombotic syndrome and ulceration, lower-extremity atherosclerotic disease predisposing to toe ischemia, episodic thrombocytopenia, and seizure disorder. At the time of presentation, he was being treated with rivaroxaban, aspirin, and hydroxychloroquine. He presented with dyspnea, fever, hypoxemia with bilateral pulmonary ground glass opacities, worsening toe ischemia, systolic cardiac failure, renal failure, normochromic normocytic anemia, and thrombocytopenia. There was no evidence of hemolysis or disseminated intravascular coagulation, as demonstrated by normal haptoglobin and lack of schistocytes on peripheral blood smear.

DIFFERENTIAL DIAGNOSIS

Catastrophic antiphospholipid syndrome (CAPS)

APS is an autoimmune disease characterized by thrombosis and pregnancy loss (1). The catastrophic variety of APS (CAPS) is diagnosed in the setting of multiorgan involvement, typically with microvascular thromboses on biopsy findings (2). A definitive diagnosis is often only made in hindsight, as histologic confirmation of microvascular thrombosis is typically lacking at the time of presentation (3). In this case, there is indication of multiple compromised organs, including the hypoxemic respiratory failure, rising creatinine, rising troponin with reduction in cardiac function, and toe ischemia. Pointing away from CAPS is the absence of definitive microthromboses (with the caveat that such specimens are often difficult to obtain in critically ill patients). Further, some patients with CAPS will present with hemolysis and even microangiopathy, which this patient does not have.

APS-associated diffuse alveolar hemorrhage

The development of acute hypoxemia in an APS patient, hemoglobin drop, and new infiltrate on imaging should raise suspicion for diffuse alveolar hemorrhage. This manifestation has been described in both primary APS and CAPS (4–6). Interestingly, the pathophysiology of diffuse alveolar hemorrhage in APS is typically of inflammatory vasculitis/capillaritis, rather than thrombosis, even in patients with primary APS (5,6). Hemoptysis was absent here, although that will be the case for many cases of diffuse alveolar

hemorrhage. Diffuse alveolar hemorrhage is, of course, a rare entity and there may be other explanations for the hemoglobin drop and lung infiltrate in this patient.

Lupus flare

Despite a historic diagnosis that had been labeled as “primary APS,” it is warranted to assess for overlapping systemic autoimmunity, with lupus being the most common culprit. Thrombocytopenia and worsening hypocomplementemia both raise suspicion for lupus; however, these can also be seen at modest levels in primary APS (7). Diffuse alveolar hemorrhage can be seen in lupus, with secondary APS and thrombocytopenia as recognized risk factors for that complication (8). Renal disease could be based in lupus glomerulonephritis, although this is unlikely in the absence of active renal sediment and with a negligible amount of proteinuria. Further, his physicians had never diagnosed him with lupus, given the absence of clinical signs and symptoms (as discussed in detail above).

Infectious or Libman-Sacks endocarditis

The worsening of mitral regurgitation raises suspicion for endocarditis. Further, primary APS is regularly associated with valvular vegetations, similar to those seen in patients with lupus (7,9). Valvular vegetations can precipitate embolic phenomena (10), including ischemic events (11). The worsening of renal function could be the result of either frank embolic disease or endocarditis/immune-complex-mediated glomerulonephritis. The foot pain and toe necrosis could be secondary to embolism. Negative blood cultures point away (although do not rule out) infectious endocarditis. There were also no stigmata of peripheral embolic disease, such as Janeway lesions or Osler nodes. While transthoracic echocardiogram findings were negative, a transesophageal echocardiogram (which would have been more definitive) was not performed.

Thrombotic thrombocytopenic purpura

With new anemia, thrombocytopenia, and renal failure, thrombotic thrombocytopenic purpura can be considered. However, with no evidence of microangiopathy on blood smear, or hemolysis by biochemical parameters, this becomes less likely (4,12).

Sepsis/community acquired pneumonia

Infectious pulmonary infiltrates culminating in sepsis could explain the presentation. The worsening toe ischemia could potentially stem from poor perfusion in the setting of systolic heart failure. Infection is also well known to trigger CAPS (13). Anemia and thrombocytopenia could result from disseminated intravascular coagulation in the setting of sepsis, although the normal D-dimer and fibrinogen make this less likely.

HOSPITAL COURSE

The patient's hypoxemic respiratory failure worsened, requiring intubation. He developed hypotension that was initially reversible with intravenous fluid. He received




Figure 2. Left ventricle, microscopic view. Broad areas of dense collagenous scar (left) and subacute changes (in response to ischemia) with granulation tissue and early collagen deposition (right).

piperacillin/tazobactam, vancomycin, and azithromycin as treatment for suspected pneumonia and sepsis. From a rheumatologic perspective, he was managed as a possible case of CAPS with pulse-dose corticosteroids (methylprednisolone 1,000 mg \times 2), heparin infusion, and plasma exchange (3). Cultures obtained by mini bronchoalveolar lavage were negative. The mini bronchoalveolar was not sufficient to prove or disprove diffuse alveolar hemorrhage (i.e., classical assessment by serial lavage); that assessment was not performed for fear of decompensating the patient. On the third day of admission, he was improving with minimal ventilator requirements (fraction of inspired oxygen 35%, positive

end-expiratory pressure 5). He then developed a sudden drop in systolic blood pressure requiring norepinephrine infusion. He developed ectopic beats, and amlodipine was added. Cardiac arrest occurred within 1 hour of the initial blood pressure drop. He could not be resuscitated.

AUTOPSY

The heart was enlarged in size at 630 grams (normal mean 350 grams), with broad areas of fibrosis seen both grossly and microscopically (Figure 2). Although no acute ischemia was




Figure 3. Left anterior descending artery, microscopic view. Focal complete luminal occlusion with dense intramural inflammatory response, especially prominent at the media-adventitia interface (A). There is a predominance of CD4+ lymphocytes (B), with less CD8+ lymphocytes (C), and natural killer cell (D) infiltration.




Figure 4. Pulmonary capillaritis, microscopic view.

noted grossly, there were multiple areas of microscopic ischemia (Figure 2). Coronaries were significantly and chronically stenosed with complete occlusion of the proximal left anterior descending artery and 90% of the left circumflex proximally. Microscopically, there were diffuse atheromatous plaques with chronic inflammatory changes including lymphocytes infiltrating the adventitia of all 3 major coronary arteries (Figure 3). The infiltrates were predominantly comprised of CD4+ lymphocytes. Two firm orange-tan vegetations were present on the mitral valve (0.5 cm × 0.4 cm and 0.3 cm × 0.2 cm), along with 1 friable pink-tan vegetation on the aortic valve (1.5 cm × 1 cm). Microscopically, the valves demonstrated sterile fibrinous vegetations with chronic inflammation (lymphocytes more so than neutrophils), which would be consistent with Libman-Sacks endocarditis (14), rather than infectious endocarditis (15). Determining the precise etiology of Libman-Sacks endocarditis (lupus versus APS) is probably not feasible, as noted in the literature (16). The distal abdominal aorta and bilateral common iliac arteries were remarkable for moderate calcific plaques.

The pulmonary parenchyma was markedly hemorrhagic and congested. This was unlikely to be explained by post-resuscitation trauma given the widespread nature of the bleeding noted macroscopically (17). Pulmonary arteries were free from thrombi. Microscopically, the lungs had extensive acute intraalveolar hemorrhage and intraalveolar macrophages (no definitive hemosiderin-laden macrophages were detected). Some capillaries showed surrounding acute inflammation (Figure 4). The alveolar septae had patchy fibrosis. Focal areas of intraalveolar acute inflammation were seen. Kidneys were notable for numerous sclerotic glomeruli, chronic interstitial inflammation, and tubular calcification. The renal pathology was hard to delineate further given post-mortem changes. With that caveat, there were no features to suggest active glomerulonephritis (as in lupus) or microangiopathy/intimal hyperplasia (as in APS). The interstitial inflammation was likely nonspecific and related to either antibiotics (piperillin-tazobactam is notorious) or even atherosclerosis (18). Although there was no evidence of thrombi, atherosclerosis, or vasculitis in the vessels at the base of the brain, there was brown-yellow mottled discoloration near the left occipital horn of the lateral ventricle indicative of old ischemia. The liver and spleen weighed 2,300

grams (normal mean 1,500 grams) and 320 grams (normal mean 150 grams), respectively, with no evidence of thrombi in either.

DISCUSSION

APS is an autoimmune disease mediated by pathogenic autoantibodies (especially to the phospholipid-binding plasma protein β_2 GPI), which promote thromboinflammatory outcomes by engaging cellular membranes (19). As mentioned above, the hallmarks of the disease are obstetric complications and vascular thromboses, with both arteries and veins at risk (1,20). There are other associated (sometimes called “non-criteria”) manifestations of APS that cannot be explained by thrombotic events (21), such as thrombocytopenia, heart valve lesions, and seizures. Further, patients with APS have an elevated risk of cardiovascular/coronary disease (22,23), and accelerated atherosclerosis (24). APS has also occasionally been linked to a traditional small-vessel vasculitis, especially in the lung capillaries (5). Other “named” vasculitides that have rarely been connected to antiphospholipid antibodies in the literature include polyarteritis nodosa and Takayasu’s arteritis (25–29).

Speaking more broadly, the intersection of inflammation and coagulation pathways in APS is well-recognized, with the two going hand-in-hand to promote thrombotic events. Several cell types may be activated by antiphospholipid antibodies, including endothelial cells, platelets, monocytes, and neutrophils (19,30,31). Mechanistically, antiphospholipid antibodies promote activation of both the endothelium (for example, upregulation of vascular adhesion molecules) and the complement cascade. The result is recruitment of neutrophils and monocytes (which may themselves be primed by antiphospholipid antibodies) to the vascular lumen, where they activate coagulation pathways and trigger thrombosis (32).

In this patient, there is coronary disease that is well beyond what would be expected based on his age and known traditional risk factors. The adventitial layer of the coronary arteries is strikingly infiltrated by lymphocytes. Traditionally, the histopathologic description of atherosclerosis is for lipid deposition, followed by stages of blood vessel wall thickening and finally fibrosis (33). While inflammatory cells may infiltrate the adventitia in atherosclerosis, they especially surround the lipid-rich regions of the vessel wall (the intima) (34,35). Here, the histopathology and concentration of CD4+ lymphocytes in the adventitia is quite compelling (and reminiscent of what might be seen in conditions like giant cell arteritis). We speculate that some element of frank coronary vasculitis contributed to this patient’s vessel wall derangements, resulting in significant, progressive stenosis of both the left anterior descending and left circumflex arteries. To our knowledge, coronary vasculitis has not been previously described in APS. Of course, an important comment is that even in the absence of definite clinical or histopathologic features of lupus in this patient, we cannot completely rule out overlapping or emerging lupus. For example, recent work by our group and others has shown a type I interferon signature in patients with primary APS (36,37), similar to that seen in lupus, which could have implications for the emergence of lupus features over time. Having said that, his

cardiovascular findings and rapid development of multiorgan failure are well out of proportion to any solid clinical evidence for lupus here, and more likely relate to his well-established diagnosis of severe APS.

Regarding other key pathologic findings, the valvular disease in this patient is characterized by fibrinous inflammation and calcification, which is typical of what has been described in APS (i.e., Libman-Sacks endocarditis) (38). The diffuse alveolar hemorrhage here seems to be driven by a capillaritis, which has also been reported in APS (5,6,39).

In summary, the patient had severe coronary disease, with no clear risk factors beyond the primary APS, and with pathology supporting a potential role for coronary vasculitis. The APS-related diffuse alveolar hemorrhage was the final stressor that set in motion the events that culminated in his death.

FINAL DIAGNOSIS

Diffuse alveolar hemorrhage in a patient with limited cardiovascular reserve attributable to severe coronary disease, with pathology supporting a potential role for coronary vasculitis.

AUTHOR CONTRIBUTIONS

All authors were involved in drafting the article or revising it critically for important intellectual content, and all authors approved the final version to be submitted for publication. Dr. Knight had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study conception and design. Kazzaz, Knight.

Acquisition of data. Kazzaz, Wilson, Knight.

Analysis and interpretation of data. Kazzaz, Wilson, Kado, Barnes, Knight.

REFERENCES

- Miyakis S, Lockshin MD, Atsumi T, Branch DW, Brey RL, Cervera R, et al. International consensus statement on an update of the classification criteria for definite antiphospholipid syndrome (APS). *J Thromb Haemost* 2006;4:295–306.
- Cervera R, Font J, Gomez-Puerta JA, Espinosa G, Cucho M, Bucciarelli S, et al. Validation of the preliminary criteria for the classification of catastrophic antiphospholipid syndrome. *Ann Rheum Dis* 2005;64:1205–9.
- Kazzaz NM, McCune WJ, Knight JS. Treatment of catastrophic antiphospholipid syndrome. *Curr Opin Rheumatol* 2016;28:218–27.
- Rodriguez-Pinto I, Espinosa G, Cervera R. Catastrophic APS in the context of other thrombotic microangiopathies. *Curr Rheumatol Rep* 2015;17:482.
- Deane KD, West SG. Antiphospholipid antibodies as a cause of pulmonary capillaritis and diffuse alveolar hemorrhage: a case series and literature review. *Sem Arthritis Rheum* 2005;35:154–65.
- Cartin-Ceba R, Peikert T, Ashrani A, Keogh K, Wylam ME, Ytterberg S, et al. Primary antiphospholipid syndrome-associated diffuse alveolar hemorrhage. *Arthritis Care Res (Hoboken)* 2014;66:301–10.
- Cervera R, Piette JC, Font J, Khamashta MA, Shoenfeld Y, Camps MT, et al. Antiphospholipid syndrome: clinical and immunologic manifestations and patterns of disease expression in a cohort of 1,000 patients. *Arthritis Rheum* 2002;46:1019–27.
- Kazzaz NM, Coit P, Lewis EE, McCune WJ, Sawalha AH, Knight JS. Systemic lupus erythematosus complicated by diffuse alveolar haemorrhage: risk factors, therapy and survival. *Lupus Sci Med* 2015;2:e000117.
- Gleason CB, Stoddard MF, Wagner SG, Longaker RA, Pierangeli S, Harris EN. A comparison of cardiac valvular involvement in the primary antiphospholipid syndrome versus anticardiolipin-negative systemic lupus erythematosus. *Am Heart J* 1993;125:1123–9.
- Asopa S, Patel A, Khan OA, Sharma R, Ohri SK. Non-bacterial thrombotic endocarditis. *Eur J Cardiothorac Surg* 2007;32:696–701.
- Turiel M, Muzzupappa S, Gottardi B, Crema C, Sarzi-Puttini P, Rossi E. Evaluation of cardiac abnormalities and embolic sources in primary antiphospholipid syndrome by transesophageal echocardiography. *Lupus* 2000;9:406–12.
- Montecucco C, Di Lauro M, Bobbio-Pallavicini E, Longhi M, Caporali R, De Gennaro F, et al. Anti-phospholipid antibodies and thrombotic thrombocytopenic purpura. *Clin Exp Rheumatol* 1987;5:355–8.
- Cervera R, Rodriguez-Pinto I, Colafrancesco S, Conti F, Valesini G, Rosario C, et al. 14th international congress on antiphospholipid antibodies task force report on catastrophic antiphospholipid syndrome. *Autoimmun Rev* 2014;13:699–707.
- Bulkley BH, Roberts WC. The heart in systemic lupus erythematosus and the changes induced in it by corticosteroid therapy: a study of 36 necropsy patients. *Am J Med* 1975;58:243–64.
- Lee JL, Naguwa SM, Cheema GS, Gershwin ME. Revisiting Libman-Sacks endocarditis: a historical review and update. *Clin Rev Allergy Immunol* 2009;36:126–30.
- Eiken PW, Edwards WD, Tazelaar HD, McBane RD, Zehr KJ. Surgical pathology of nonbacterial thrombotic endocarditis in 30 patients, 1985–2000. *Mayo Clin Proc* 2001;76:1204–12.
- Ornato JP, Ryschon TW, Gonzalez ER, Bredthauer JL. Rapid change in pulmonary vascular hemodynamics with pulmonary-edema during cardiopulmonary resuscitation. *Am J Emerg Med* 1985;3:137–42.
- Chade AR, Rodriguez-Porcel M, Grande JP, Krier JD, Lerman A, Romero JC, et al. Distinct renal injury in early atherosclerosis and renovascular disease. *Circulation* 2002;106:1165–71.
- Giannakopoulos B, Krilis SA. The pathogenesis of the antiphospholipid syndrome. *New Engl J Med* 2013;368:1033–44.
- Davies JO, Hunt BJ. Myocardial infarction in young patients without coronary atherosclerosis: assume primary antiphospholipid syndrome until proved otherwise. *Int J Clin Pract* 2007;61:379–84.
- Abreu MM, Danowski A, Wahl DG, Amigo MC, Tektonidou M, Pacheco MS, et al. The relevance of “non-criteria” clinical manifestations of antiphospholipid syndrome: 14th international congress on antiphospholipid antibodies technical task force report on antiphospholipid syndrome clinical features. *Autoimmun Rev* 2015;14:401–14.
- Soltesz P, Szekanecz Z, Kiss E, Shoenfeld Y. Cardiac manifestations in antiphospholipid syndrome. *Autoimmun Rev* 2007;6:379–86.
- Cervera R. Coronary and valvular syndromes and antiphospholipid antibodies. *Thromb Res* 2004;114:501–7.
- Djokovic A, Stojanovich L, Stanisavljevic N, Bisenic V, Radovanovic S, Soldatovic I, et al. Does the presence of secondary antiphospholipid syndrome in patients with systemic lupus erythematosus accelerate carotid arteries intima-media thickness changes? *Rheumatol Int* 2014;34:321–7.
- Han BK, Inaganti K, Fahmi S, Reimold A. Polyarteritis nodosa complicated by catastrophic antiphospholipid syndrome. *J Clin Rheumatol* 2004;10:210–3.
- Schoonjans R, van Vlem B, Weyers S, De Keyser F, Praet M, Versieck J, et al. Polyarteritis nodosa and the antiphospholipid syndrome. *Clin Rheumatol* 1996;15:410–3.
- Norden DK, Ostrov BE, Shafritz AB, von Feldt JM. Vasculitis associated with antiphospholipid syndrome. *Semin Arthritis Rheum* 1995;24:273–81.

28. Yokoi K, Hosoi E, Akaike M, Shigeakiyo T, Saito S. Takayasu's arteritis associated with antiphospholipid antibodies: report of two cases. *Angiology* 1996;47:315–9.
29. Dongola NA, Foord KD. Angiographic features associated with antiphospholipid syndrome. *Brit J Radiol* 2000;73:1215–8.
30. Yalavarthi S, Gould TJ, Rao AN, Mazza LF, Morris AE, Nunez-Alvarez C, et al. Release of neutrophil extracellular traps by neutrophils stimulated with antiphospholipid antibodies: a newly identified mechanism of thrombosis in the antiphospholipid syndrome. *Arthritis Rheumatol* 2015;67:2990–3003.
31. Meng H, Yalavarthi S, Kanthi Y, Mazza LF, Elflin MA, Luke CE, et al. In vivo role of neutrophil extracellular traps in antiphospholipid antibody-mediated venous thrombosis. *Arthritis Rheumatol* 2017;69:655–67.
32. Lally L, Sammaritano LR. Vasculitis in antiphospholipid syndrome. *Rheum Dis Clin North Am* 2015;41:109–23.
33. Stary HC, Chandler AB, Dinsmore RE, Fuster V, Glagov S, Insull W Jr, et al. A definition of advanced types of atherosclerotic lesions and a histological classification of atherosclerosis: a report from the Committee on Vascular Lesions of the Council on Arteriosclerosis, American Heart Association. *Arterioscler Thromb Vasc Biol* 1995;15:1512–31.
34. Van der Wal AC. Coronary artery pathology. *Heart* 2007;93:1484–9.
35. Van der Wal AC, Becker AE. Atherosclerotic plaque rupture: pathologic basis of plaque stability and instability. *Cardiovasc Res* 1999;41:334–44.
36. Grenn RC, Yalavarthi S, Gandhi AA, Kazzaz NM, Nunez-Alvarez C, Hernandez-Ramirez D, et al. Endothelial progenitor dysfunction associates with a type I interferon signature in primary antiphospholipid syndrome. *Ann Rheum Dis* 2016.
37. Van den Hoogen LL, Fritsch-Stork RD, Versnel MA, Derksen RH, van Roon JA, Radstake TR. Monocyte type I interferon signature in antiphospholipid syndrome is related to proinflammatory monocyte subsets, hydroxychloroquine and statin use. *Ann Rheum Dis* 2016;75:e81.
38. Garcia-Torres R, Amigo MC, de la Rosa A, Moron A, Reyes PA. Valvular heart disease in primary antiphospholipid syndrome (PAPS): clinical and morphological findings. *Lupus* 1996;5:56–61.
39. Asherson RA, Greenblatt MA. Recurrent alveolar hemorrhage and pulmonary capillaritis in the “primary” antiphospholipid syndrome. *J Clin Rheumatol* 2001;7:30–3.