

REVIEWS

Antiviral Therapy for Chronic Hepatitis B Viral Infection in Adults: A Systematic Review and Meta-Analysis

Anna S.F. Lok,¹ Brian J. McMahon,² Robert S. Brown, Jr.,³ John B. Wong,⁴ Ahmed T. Ahmed,^{5,6} Wigdan Farah,^{5,6} Jehad Almasri,^{5,6} Fares Alahdab,^{5,6} Khalid Benkhadra,^{5,6} Mohamed A. Mouchli,⁷ Siddharth Singh,⁸ Essa A. Mohamed,⁸ Abd Moain Abu Dabrh,⁶ Larry J. Prokop,⁹ Zhen Wang,^{5,6} Mohammad Hassan Murad,^{5,6,10} and Khaled Mohammed^{5,6,10}

Chronic hepatitis B viral (HBV) infection remains a significant global health problem. Evidence-based guidelines are needed to help providers determine when treatment should be initiated, which medication is most appropriate, and when treatment can safely be stopped. The American Association for the Study of Liver Diseases HBV guideline methodology and writing committees developed a protocol *a priori* for this systematic review. We searched multiple databases for randomized controlled trials and controlled observational studies that enrolled adults ≥ 18 years old diagnosed with chronic HBV infection who received antiviral therapy. Data extraction was done by pairs of independent reviewers. We included 73 studies, of which 59 (15 randomized controlled trials and 44 observational studies) reported clinical outcomes. Moderate-quality evidence supported the effectiveness of antiviral therapy in patients with immune active chronic HBV infection in reducing the risk of cirrhosis, decompensated liver disease, and hepatocellular carcinoma. In immune tolerant patients, moderate-quality evidence supports improved intermediate outcomes with antiviral therapy. Only very low-quality evidence informed the questions about discontinuing versus continuing antiviral therapy in hepatitis B e antigen-positive patients who seroconverted from hepatitis B e antigen to hepatitis B e antibody and about the safety of entecavir versus tenofovir. Noncomparative and indirect evidence was available for questions about stopping versus continuing antiviral therapy in hepatitis B e antigen-negative patients, monotherapy versus adding a second agent in patients with persistent viremia during treatment, and the effectiveness of antivirals in compensated cirrhosis with low-level viremia. **Conclusion:** Most of the current literature focuses on the immune active phases of chronic HBV infection; decision-making in other commonly encountered and challenging clinical settings depends on indirect evidence. (HEPATOLOGY 2016;63:284-306)

Chronic hepatitis B viral (HBV) infection remains a significant global health problem. Despite the availability of HBV vaccines for three decades, the global prevalence of chronic HBV infection has only declined slightly, from 4.2% in 1990 to 3.7% in 2005.¹ Worldwide, however, the absolute

number of persons chronically infected has increased from 223 million in 1990 to 240 million in 2005. In the United States, based on 1999-2006 data from the National Health and Nutrition Examination Survey, the prevalence of chronic HBV infection was estimated to be 0.27%.² However, the National Health and

Abbreviations: AASLD, American Association for the Study of Liver Diseases; ALT, alanine aminotransferase; CI, confidence interval; HBeAg, hepatitis B e antigen; HBsAg, hepatitis B surface antigen; HBV, hepatitis B virus; HCC, hepatocellular carcinoma; IFN, interferon; RCT, randomized controlled trial; RR, risk ratio.

From the ¹Division of Gastroenterology and Hepatology, University of Michigan, Ann Arbor, MI; ²Liver Diseases and Hepatitis Program, Alaska Native Tribal Health Consortium, Anchorage, AK; ³Division of Gastroenterology and Hepatology, Weill Cornell Medical College, New York, NY; ⁴Division of Clinical Decision Making, Tufts Medical Center, Boston, MA; ⁵Evidence-Based Practice Research Program, Mayo Clinic, Rochester, MN; ⁶Center for the Science of Health Care Delivery, Mayo Clinic, Rochester, MN; ⁷Division of Hospital Internal Medicine, Mayo Clinic, Rochester, MN; ⁸Division of Gastroenterology and Hepatology, Mayo Clinic, Rochester, MN; ⁹Library Public Services, Mayo Clinic, Rochester, MN; ¹⁰Division of Preventive, Occupational and Aerospace Medicine, Mayo Clinic, Rochester, MN

This Review published with an accompanying Practice Guideline and Reviews by Jonas et al. and Brown et al.

Nutrition Examination Survey undersampled high-prevalence groups, so when accounting for immigration from endemic countries, as many as 2.2 million US residents (instead of 730,000) may have chronic HBV infection.³

The natural course of chronic HBV infection consists of four characteristic phases: immune tolerant, hepatitis B e antigen (HBeAg)-positive immune active, inactive, and HBeAg-negative immune active phases.⁴ The immune tolerant phase is characterized by the presence of HBeAg, normal alanine aminotransferase (ALT) levels, and high levels of HBV DNA, usually well over 20,000 IU/mL. The immune active phases, also called HBeAg-positive or HBeAg-negative chronic hepatitis, are characterized by intermittently or persistently elevated ALT with active hepatic inflammation and HBV DNA generally above 2000 IU/mL. The inactive phase is characterized by absence of HBeAg and presence of hepatitis B e antibody, normal ALT in the absence of other concomitant liver diseases, and undetectable or low levels of HBV DNA, generally below 2000 IU/mL. Although not all patients go through each phase and immune responses to HBV during each phase have not been fully characterized, this classification schema provides a useful framework when developing a management approach for chronic HBV infection.

Currently, seven medications are approved for treatment of chronic HBV infection: two formulations of interferon (IFN), standard and pegylated, and five nucleos(t)ide analogues: lamivudine, telbivudine, entecavir, adefovir, and tenofovir. These medications suppress HBV replication and ameliorate hepatic inflammation but do not eradicate HBV. While IFN is given for a finite duration, nucleos(t)ide analogues are administered for many years and often for life. Long durations of treatment are associated with risks of adverse reactions, drug resistance, nonadherence, and increased cost. Therefore, there is a need to have evidence-based guidelines to help providers determine when treatment should be initiated, which medication is most appropriate, and when treatment can safely be stopped.

Materials and Methods

The American Association for the Study of Liver Diseases (AASLD) HBV guideline methodology and writing committees developed a protocol *a priori* for this systematic review. The reporting of this review follows the standards set in the Preferred Reporting Items for Systematic Reviews and Meta-analysis statement.⁵ The committee identified and developed a protocol for seven key Population Intervention Comparison Outcome questions (Supporting Table S1). The outcomes of interest were clinical outcomes (cirrhosis, liver decompensation, hepatocellular carcinoma [HCC], and all-cause mortality); however, when such outcome data were unavailable, surrogate (intermediate) outcomes were sought, specifically durability of HBeAg seroconversion, loss of hepatitis B surface (HBsAg), long-term suppression of HBV DNA, and normalization of ALT.

Eligibility Criteria. We included randomized controlled trials (RCTs) and controlled observational studies that enrolled adults ≥ 18 years old diagnosed with chronic HBV infection who received antiviral therapy. We excluded studies that included patients with acute HBV infection; patients who were pregnant; patients coinfecting with hepatitis C or D or human immunodeficiency virus; patients receiving corticosteroids, chemotherapy, or immunosuppressive therapy; transplant recipients; and hemodialysis patients, as well as studies without control or comparison groups. Supporting Table S1 summarizes the inclusion and exclusion criteria for each key question.

Search Strategy. An experienced Mayo Clinic librarian conducted a comprehensive search of Medline In-Process & Other Non-Indexed Citations, MEDLINE, EMBASE, Cochrane Central Register of Controlled Trials, Cochrane Database of Systematic Reviews, and Scopus from early 1988 to September 16, 2014. Controlled vocabulary supplemented with keywords was used to search for comparative studies of antivirals for chronic hepatitis B. No language restrictions were used. Members from the AASLD HBV guideline methodology and writing committees helped identify

Additional Supporting Information may be found at <http://onlinelibrary.wiley.com/doi/10.1002/hep.28280/supinfo>.

Received August 18, 2015; accepted September 23, 2015.

Address reprint requests to: Anna S. Lok, M.D., University of Michigan Health System, 1500 E. Medical Center Drive, 3912 Taubman Center, SPC 5362, Ann Arbor, MI 48109. E-mail: asllok@umich.edu; tel: +1-734-936-7511; fax: +1-734-936-7392.

Copyright © 2015 by the American Association for the Study of Liver Diseases.

View this article online at wileyonlinelibrary.com.

DOI 10.1002/hep.28280

Potential conflict of interest: Dr. Lok advises and received grants from Gilead. She consults for GlaxoSmithKline and Merck and received grants from Bristol-Myers Squibb. Dr. Brown consults for Gilead and Bristol-Myers Squibb.

additional studies. [Supporting Table S2](#) specifies the detailed search strategy.

Study Selections. Two reviewers independently screened titles and abstracts for potential eligibility using an online reference management system (DistillerSR; Evidence Partners, Inc.). Full texts of the included abstracts were retrieved and screened in duplicate. Disagreements were resolved by seeking consensus or arbitration by a third reviewer. Interreviewer agreement (kappa) was calculated during each screening level to assess agreement between reviewers. For Population Intervention Comparison Outcome questions where no studies meeting the predefined criteria were found, the AASLD HBV guideline methodology committee performed manual searches for uncontrolled observational studies. Data from these studies were summarized narratively and in general consistent with low-quality evidence.

Data Extraction. Data extraction was done using a standardized, piloted form. We extracted data on study characteristics, patient characteristics, intervention details, and outcomes of interest.

Methodological Quality and Risk of Bias Assessment. We used the Cochrane Risk of Bias assessment tool and modified Newcastle-Ottawa Scale to assess the risk of bias in RCTs and observational studies, respectively. Quality of evidence (i.e., certainty in the estimates) was evaluated using the Grading of Recommendations Assessment, Development, and Evaluation approach. Criteria used to evaluate quality of evidence were risk of bias, indirectness (surrogate outcomes), imprecision (wide confidence intervals), inconsistency (heterogeneity), and publication bias.⁶

Statistical Analysis. For dichotomized outcomes, we calculated risk ratios (RRs) and 95% confidence intervals (95% CI) using binomial distribution. We then pooled the log-transformed RRs using the DerSimonian and Laird random-effects models and estimated heterogeneity using the Mantel-Haenszel model. To measure the overall heterogeneity across the included studies, we calculated the I^2 statistic, where $I^2 > 50\%$ suggests a high degree of heterogeneity. All statistical analyses were conducted using STATA, version 13 (StataCorp LP, College Station, TX). To explore heterogeneity, we conducted subgroup analysis for studies enrolling patients with more advanced liver disease; we performed stratified analysis for the following groups: compensated cirrhosis, decompensated cirrhosis, acute on chronic liver failure, and severe acute exacerbations of chronic hepatitis B. We explored the impact of publication bias using the Egger regression asymmetry test and constructing funnel plots if a sufficient number of studies

(>20) per outcome was available and heterogeneity was low.⁷

Results


A total of 73 studies were included. Figure 1 describes the details of the selection process. The average weighted kappa for study selection was 0.78. Controlled studies that reported the outcomes of interest were only available for questions 1, 2, 3, and 5. Uncontrolled studies that are relevant to questions 4, 6, and 7 are summarized in [Supporting Information](#). [Supporting Table S4](#) provides the Grading of Recommendations Assessment, Development, and Evaluation summary of the evidence.

Question 1: Effectiveness of Antiviral Therapy in Patients With Immune Active Chronic HBV Infection

We included 59 studies (15 RCTs and 44 observational studies) that evaluated antiviral therapy and reported clinical outcomes. Forty-two studies compared antiviral therapy versus control, and 18 studies compared one antiviral agent versus another.

Effectiveness of Antiviral Therapy Compared to Control in Patients With Chronic Hepatitis B Infection. Among 42 studies comparing antiviral therapy versus control in 62,731 patients, 16 studies⁸⁻²³ compared IFN versus no treatment, 16 studies²⁴⁻³⁹ compared lamivudine versus no treatment, seven studies^{28,40-45} compared entecavir versus no treatment, one study each compared telbivudine⁴⁴ and tenofovir⁴⁶ versus placebo, and three studies⁴⁷⁻⁴⁹ compared a variety of oral antiviral versus no treatment. Eleven studies enrolled only patients with compensated cirrhosis, five studies enrolled only patients with acute on chronic liver failure, two studies enrolled only patients with decompensated liver disease, three studies enrolled only patients with severe acute exacerbations of chronic hepatitis B, and 21 studies enrolled patients with stable chronic hepatitis B. Study characteristics are illustrated in Table 1. Risk of bias assessment for RCTs was low to moderate as two of the included RCTs reported the randomization method, two reported use of allocation concealment, and six reported the blinding method used. Most of the observational studies were at high risk of bias due to lack of clear description of the selection process of the population and inadequate exposure and outcome ascertainment. Risk of bias is described in Tables 2 and 3.

In seven RCTs^{8,23-25,29,33,46} involving 3463 subjects with a mean follow-up of 28 months, antiviral therapy versus control (Fig. 2) significantly decreased the overall



*Articles may be included in more than one question.

Fig. 1. Flow diagram showing selection process for studies to include. Abbreviations: HCV, hepatitis C virus; HIV, human immunodeficiency virus.

risk of decompensated liver disease (one RCT, RR = 0.4, 95% CI 0.3-0.7) and cirrhosis (one RCT, RR = 0.4, 95% CI 0.2-0.8). No significant differences were found in all-cause mortality (four RCTs, RR = 0.5, 95% CI 0.2-1.3, I² = 72.9%) or HCC incidence (three RCTs, RR = 0.6, 95% CI 0.3-1.1, I² = 0%). The quality of the evidence was low to moderate. One RCT²⁹ examined adverse events including death and decompensation as outcomes, but no events were observed in either the intervention or the control group.

In 35 observational studies involving 59,201 patients with a mean follow-up of 60 months, meta-analysis showed that antiviral therapy versus control decreased the risk of HCC (23 studies, RR = 0.5, 95% CI 0.4-0.7, I² = 87.4%), all-cause mortality (23 studies, RR = 0.6, 95% CI 0.5-0.8, I² = 92.3%), and cirrhosis (four studies, RR = 0.6, 95% CI 0.4-0.8, I² = 0%) but did not significantly reduce the risk of decompensated liver disease (six studies, RR = 0.7, 95% CI 0.3-1.9, I² = 96.5%) when compared to untreated controls (Figs. 3-5). The quality of this evidence overall was low; however, these studies included large numbers of patients with long duration of follow-up, yielding precise and narrow 95% CIs.

Effectiveness of antiviral therapy compared to control in the subgroup with stable chronic hepatitis B. Of the 21 studies that enrolled patients with stable chronic hepatitis B, 0%-91% of the 54,719 patients included had compensated cirrhosis. Reduction in risk of decompensated cirrhosis was shown in only one RCT and reduction in HCC in 11 observational studies. No studies demonstrated reduction in all-cause mortality.

Effectiveness of Antiviral Therapy Compared to Control in Patients With Chronic HBV Infection and Compensated Cirrhosis. In one RCT²⁵ enrolling 222 patients with cirrhosis and a follow-up of 53 months, lamivudine versus control reduced all-cause mortality (RR = 0.1, 95% CI 0.1-0.3, moderate-quality evidence).

In 10 observational studies (Fig. 3) involving patients with compensated cirrhosis (mean follow-up 60 months), antiviral therapy decreased the risk of HCC (10 studies, RR = 0.6, 95% CI 0.4-0.8, I² = 36.3%), decompensated liver disease (two studies, RR = 0.5, 95% CI 0.2-0.9, I² = 67.2%), and all-cause mortality (three studies, RR = 0.5, 95% CI 0.4-0.6, I² = 0%).

Table 1. Characteristics of the Included Studies

Authors, Year	Country	Patients (N)	Interventions	Age (Years)	HBeAg Positive (N)	Baseline ALT (U/L)	Baseline HBV DNA (log ₁₀ IU/mL)*	Follow-up duration (months)	Baseline cirrhosis (%)	Study design
Question 1: Effectiveness of antiviral therapy in patients with immune active chronic HBV infection (antiviral versus control)										
Anderson et al., 1987 ⁸	England	14	IFN- α	36	14	77% elevated ALT	NR	12	20	RCT
		16	Control	35	16	77% elevated ALT	NR	12	20	
IHCSG, 1998 ⁹	Italy and Argentina	49	IFN- α	54	NR	NR	NR	69.6	100	Case-control
		97	Control	54	NR	NR	NR	82.2	100	
Lin et al., 2007 ¹⁰	Taiwan	233	IFN- α	32 \pm 7	233	175 \pm 112	40% > 7.7	81.6 \pm 38.4	8.1	Cohort
		233	Control	31 \pm 8	233	187 \pm 109	40% > 7.7	73.2 \pm 36	10.7	
Truong et al., 2005 ¹¹	Japan	27	IFN- α	33.2 \pm 10.4	17	238.6 \pm 250.1	NR	84 \pm 30	3	Case-control
		35	Control	36.6 \pm 10.9	20	142.3 \pm 152.1	NR	74.4 \pm 34.8	14.3	
Tangkijvanich et al., 2001 ¹²	Thailand	67	IFN- α	36.9 \pm 10.5	67	180.7 \pm 137.9	NR	59.4 \pm 30.9	17.9	Case-control
		72	Control	39.9 \pm 13.7	72	93.3 \pm 114.4	NR	60.1 \pm 35.3	22.2	
Papatheodoridis et al., 2001 ¹³	Greece	209	IFN- α	46.8 \pm 11.3	0	112 (13-1905)	5.4	72 \pm 32.4	27.3	Cohort
		195	Control	48.8 \pm 13.7	0	68 (20-1335)	5.4	73.2 \pm 46.8	34.9	
Niederer et al., 1996 ¹⁴	Germany	103	IFN- α	NR	103	NR	NR	50.0 \pm 19.8	27	Cohort
		53	Control	NR	53	NR	NR	38.5 \pm 18.2	16	
Lin et al., 2004 ¹⁵	Taiwan	109	IFN- α	31 \pm 9	NR	132 \pm 86	NR	84.5	90	Cohort
		34	Control	32 \pm 6	NR	256 \pm 232	NR	92	85	
Benvegno et al., 1998 ¹⁶	Italy	13	IFN- α	57	NR	NR	NR	72	100	Cohort
		24	Control	60	NR	NR	NR	72	100	
Tong et al., 2006 ¹⁷	USA	22	IFN- α	48	49%	NR	NR	84	35	Cohort
		378	Control	48	NR	NR	NR	84	35	
Di Marco et al., 1999 ¹⁸	Italy	109	IFN- α	33	NR	NR	NR	93.6	29	Cohort
		193	Control	35	NR	NR	NR	93.6	29	
Brunetto et al., 2002 ¹⁹	Italy	103	IFN- α	40	0	NR	NR	72	38	Cohort
		61	Control	40	0	NR	NR	72	38	
Mahmood et al., 2005 ²⁰	Japan	23	IFN- α	49	NR	NR	NR	84	100	Case-control
		68	Control	49	NR	NR	NR	84	100	
Ikeda et al., 1998 ²¹	Japan	94	IFN- α	41	NR	NR	NR	81.6	100	Case-control
		219	Control	44	NR	NR	NR	84	100	
Fattovich et al., 1997 ²²	Italy	40	IFN- α	47 \pm 1.8	40	5.3 (0.61 \times ULN)	NR	74.4	100	Cohort
		50	Control	45 \pm 2.2	50	5.3 (0.61 \times ULN)	NR	74.4	100	
Krogsgaard et al., 1998 ²³	Europe	210	IFN- α	36	210	100% elevated ALT	NR	15.6	19	RCT
		98	Control	36	98	NR	NR	15.6	19	
Chan et al., 2007 ²⁴	China	89	Lamivudine	39 \pm 11	6	2.1 \pm 1.7 (\times ULN)	5 \pm 0.09	120	31	RCT
		47	Placebo	39 \pm 11	4	2.6 \pm 2.3 (\times ULN)	4.9 \pm 0.8	120	21	
Eun et al., 2007 ²⁵	Korea	111	Lamivudine	NR	NR	NR	NR	52.8	100	RCT
		111	Placebo	NR	NR	NR	NR	52.8	100	
Tong et al., 2009 ²⁶	USA	27	Lamivudine	40	NR	NR	NR	63.6	100	Cohort

Table 1. Continued

Authors, Year	Country	Patients (N)	Interventions	Age (Years)	HBeAg Positive (N)	Baseline ALT (U/L)	Baseline HBV DNA (log ₁₀ IU/mL)*	Follow-up duration (months)	Baseline cirrhosis (%)	Study design
Das et al., 2010 ²⁷	India	101	Control	46	NR	NR	NR	63.6	100	Case-control
		151	Lamivudine and adefovir	42	45%	NR	NR	48	100	
Cui et al., 2010 ²⁸	China	102	Control	46	NR	NR	NR	45.6	100	Cohort
		33	Entecavir	38.4 ± 10.8	10	364 (47-2861)	5.2 ± 0.8	0.2-41.5	NR	
		34	Lamivudine	39.4 ± 10.6	13	226.5 (22-2314)	5.1 ± 0.6	0.2-41.5	NR	
		37	Control	41.0 ± 11.5	11	287 (17-2535)	5 ± 0.9	0.2-41.5	NR	
Dienstag et al., 1999 ²⁹	USA	66	Lamivudine	40 (18-73)	66	125 (46-401)	6.7 (4.6-7.9)	12	6	RCT
		71	Placebo	38 (20-67)	71	135 (33-592)	6.5 (4.6-7.6)	12	14	
Chan et al., 2002 ³⁰	Hong Kong	28	Lamivudine	42.7 ± 13.5	16	1416.6 ± 577.7	NR	12	NR	Cohort
		18	Control	47.2 ± 14	2	1659.5 ± 1928.4	NR	12	NR	
Lok et al., 2003 ³¹	Multinational	998	Lamivudine	32.0 (15-73)	998	1.6 (0.2-23.4) (U/LN)	6.7 (4.7-8.1)	48	10	Cohort
		200	Placebo	34.5 (15-67)	200	2.3 (0.4-4.14) (U/LN)	6.6 (4.7-7.8)	12	13	
Manolakopoulos et al., 2004 ³²	UK	30	Lamivudine	63.1 ± 1.7	30	77 (26-280)	4.9 (3.2-7)	18 (3-36)	100	Case-control
		30	Control	62.8 ± 1.4	30	80 (30-199)	NR	22 (2-55)	100	
Liaw et al., 2004 ³³	Multinational	436	Lamivudine	43 (17-74)	252	70 (14-959)	6.4 (<5.1-10.3)	32 (0-42)	31	RCT
		215	Placebo	44 (22-71)	124	68 (7-821)	6.6 (<5.1-8.9)	32 (0-42)	39	
Matsumoto et al., 2005 ³⁴	Japan	657	Lamivudine	40.9 ± 11.0	355	183.4 ± 211.1	NR	58.8 ± 52.8	14.9	Case-control
		2138	Control	37.3 ± 12.4	1272	163.5 ± 234.3	NR	74.4 ± 66	15.5	
Ma et al., 2007 ³⁵	China	51	Lamivudine	NR	12	NR	NR	35	100	Cohort
		166	Control	NR	39	NR	NR	35	100	
Yuen et al., 2007 ³⁶	Hong Kong	142	Lamivudine	33.9 (20.2-54.4)	142	125 (47-514)	8 (3.5-11)	89.9 (26.5-128.3)	0	Cohort
		124	Control	33.4 (20.8-59)	124	125 (47-514)	6.1 (0.8-8.9)	107.8 (30.9-127.3)	0	
Sun et al., 2009 ³⁷	China	130	Lamivudine	44.3 ± 3.5	90	474.1 ± 83.4	>4.3	3	10	Cohort
		130	Control	45.2 ± 3.6	95	492.3 ± 82.6	>4.3	3	10	
Kim et al., 2012 ³⁸	Korea	240	Lamivudine	49.6 ± 10.9	145	159 ± 265.4	6.2 ± 0.6	46.4 (1-124)	100	Cohort
		481	Control	46.4 ± 10.3	280	90.2 ± 136.3	NR	51.4 (2-94)	100	
Eun et al., 2010 ³⁹	Korea	872	Lamivudine	40.1 ± 12.2	694	161 ± 183.8	7.1 ± 0.4	56.4 ± 28.8	47.4	Cohort
		699	Control	35.5 ± 12.9	637	141.3 ± 199.1	6.7 ± 0.3	68.4 ± 50.4	37.2	
Wong et al., 2013 ⁴⁰	Hong Kong	1466	Entecavir	51 ± 12	443	145 ± 319	5	36 ± 13	100	Cohort
		424	Control	41 ± 13	155	84 ± 113	5	114 ± 31	100	
Hosaka et al., 2013 ⁴¹	Japan	472	Entecavir	42 ± 12.4	219	70 (42-163)	6 (4.6-7.3)	38.4 (25.2-51.6)	25	Cohort
		1143	Control	39 ± 13.1	398	33 (20-68)	5.1 (3.3-6.8)	114 (52.8-193.2)	17	
Lin et al., 2013 ⁴²	China	53	Entecavir	38 (32-49)	16	360 (181-704)	5.8 ± 0.8	12	32.1	Cohort
		55	Control	40 (34-47)	20	467 (107-1192)	5.3 ± 0.7	12	27.3	
Xiao et al., 2009 ⁴³	China	39	Entecavir	NR	NR	NR	NR	NR	NR	Cohort
		39	Control	NR	NR	NR	NR	NR	NR	

Table 1. Continued

Authors, Year	Country	Patients (N)	Interventions	Age (Years)	HBeAg Positive (N)	Baseline ALT (U/L)	Baseline HBV DNA (log ₁₀ IU/mL)*	Follow-up duration (months)	Baseline cirrhosis (%)	Study design
Xu et al., 2009 ⁴⁴	China	133	Telbivudine, entecavir, or lamivudine	40.6 ± 11.4	NR	534 ± 712.8	4.3	NR	NR	Cohort
Chen et al., 2009 ⁴⁵	China	215	Control	40.6 ± 10.5	NR	526.1 ± 688.5	3.8	NR	NR	Cohort
Garg et al., 2011 ⁴⁶	India	55	Entecavir	43.6 ± 10.9	14	357 ± 405.2	5 ± 0.65	3	NR	Cohort
Wu et al., 2014 ⁴⁷	Taiwan	74	Control	40.3 ± 11.7	25	451.9 ± 464.6	4.4 ± 0.1.1	3	NR	RCT
Gordon et al., 2014 ⁴⁸	USA	14	Tenofovir	47.5 (16-62)	13	226 (188-1185)	5.2	3	NR	RCT
		13	Placebo	45 (16-67)	12	206 (186-2000)	5.5	3	NR	Cohort
		21595	Variety of oral antivirals	43.5 ± 13.4	26	179	5.3 ± 0.3	40 (16.8-66)	13.2	Cohort
		21595	Control	43.6 ± 13.6	12	185	5.3 ± 1.3	78 (42.5-84)	14	Cohort
		820	IFN and variety of oral antiviral	NR	820	NR	NR	62.4 (36-108)	32.9	Cohort
Kumada et al., 2013 ⁴⁹	Japan	1851	Control	NR	1851	NR	NR	62.4 (36-108)	14.6	Cohort
		148	Variety of oral antiviral	53 (26-81)	76	65 (7-1088)	6.3 (1.9-8.9)	153.6 (37.2-235.2)	62	Cohort
		637	Control	48 (4-85)	151	26 (5-3410)	3.1 (1.6-9.2)	164.4 (37.2-240)	91	Cohort
Question 1. Head-to-head studies comparing individual antiviral agents										
Cui et al., 2010 ²⁸	China	33	Entecavir	38.4 ± 10.8	10	364 (47-2861)	5.2 ± 0.8	0.2-41.5	NR	Cohort
		34	Lamivudine	39.4 ± 10.6	13	226.5 (22-2314)	5.1 ± 0.6	0.2-41.5	NR	Cohort
		37	Control	41.03 ± 11.5	11	287 (17-2535)	5 ± 0.9	0.2-41.5	NR	Cohort
Chan et al., 2012 ⁵⁰	China	114	Telbivudine	49.6 ± 10.9	61	75.1 ± 54.4	6.9 ± 1.2	24	100	RCT
		114	Lamivudine	51.9 ± 10	55	84 ± 87.8	6.9 ± 1.2	24	100	RCT
Chang et al., 2006 ⁵¹	Multinational	354	Entecavir	35 ± 13	348	140.5 ± 114.3	8.9 ± 1.3	12	8	RCT
		355	Lamivudine	35 ± 13	351	146.3 ± 132.3	9 ± 1.3	12	8	RCT
Lai et al., 2006 ⁵²	Multinational	325	Entecavir	44 ± 11	3	141 ± 114.7	6.9 ± 1.1	12	5	RCT
		313	Lamivudine	44 ± 11	4	143 ± 119.4	6.9 ± 1	12	10	RCT
Lau et al., 2005 ⁵³	Multinational	271	Peg-IFN plus placebo	32.5 ± 9.6	271	114.6 ± 114.3	9.2 ± 1.4	18	18	RCT
		271	Peg-IFN plus lamivudine	31.7 ± 10.3	271	114.9 ± 94.1	9.4 ± 1.2	18	15	RCT
		272	Lamivudine	31.6 ± 9.7	272	102.3 ± 78.4	9.4 ± 1.3	18	17	RCT
Marcellin et al., 2004 ⁵⁴	Multinational	177	Peg-IFN plus placebo	40 ± 11.7	0	94.4 ± 85.9	6.4 ± 1.1	18	31	RCT
		179	Peg-IFN plus lamivudine	41 ± 10.8	0	90.8 ± 76.2	6.5 ± 1.1	18	22	RCT
		181	Lamivudine	40 ± 11.1	0	105.7 ± 128.2	6.5 ± 1.1	18	29	RCT
Wang et al., 2013 ⁵⁵	China	102	Adefovir	44 ± 9.5	NR	72.76 ± 61.8	6.2 ± 1.2	24	100	RCT
		104	Lamivudine	44.9 ± 10.03	NR	72.6 ± 46.4	6.1 ± 1.1	24	100	RCT
Yang et al., 2009 ⁵⁶	China	32	Adefovir	31-62	NR	NR	NR	NR	100	RCT
		30	Lamivudine	25-69	NR	NR	NR	NR	100	RCT
Liaw et al., 2011 ⁵⁷	Taiwan	100	Entecavir	51 ± 1.2	54	99.2 ± 11.1	6.8 ± 0.01	24	100	RCT
		91	Adefovir	53 ± 1.1	50	100 ± 8.6	7.5 ± 0.01	24	100	RCT
Lim et al., 2014 ⁵⁸	Korea	2000	Entecavir	47 ± 11	1168	101 (53-190)	7.1 ± 1.6	37.2 (26.4-51.6)	53.6	Cohort
		3374	Lamivudine	43 ± 11	2421	128 (68-244)	7.5 ± 1.2	104.4 (78-138)	48	Cohort
Hsu et al., 2012 ⁵⁹	Taiwan	53	Entecavir	48 (40-56)	18	467 (78-879)	6.1	12	45.3	Cohort
		73	Lamivudine	46 (37-58)	17	391 (68-1530)	6.3	12	48	Cohort
Wong et al., 2011 ⁶⁰	Hong Kong	36	Entecavir	51 ± 13	13	1151 ± 724	6.6 ± 1.4	18 ± 12	14	Cohort

Table 1. Continued

Authors, Year	Country	Patients (N)	Interventions	Age (Years)	HBeAg Positive (N)	Baseline ALT (U/L)	Baseline HBV DNA (log ₁₀ IU/mL)*	Follow-up duration (months)	Baseline cirrhosis (%)	Study design
Liang et al., 2009 ⁶¹	China	117	Lamivudine	44 ± 14	55	1499 ± 841	6.8 ± 0.9	79 ± 6	21	Cohort
		40	Telbivudine	51.8 ± 10.7	20	NR	5.8 ± 0.6	12	100	Cohort
		40	Lamivudine	52.4 ± 8.5	18	NR	5.7 ± 0.6	12	100	Cohort
Chen et al., 2014 ⁶²	Taiwan	215	Lamivudine	49.5 ± 14.4	60	1239.4 ± 941.7	5.8 ± 1	20 (6.5-71.3)	42.8	Cohort
		107	Entecavir	48.6 ± 14.1	35	1045.3 ± 782.8	5.8 ± 1.2	20 (6.5-71.3)	49.5	Cohort
Zhang et al., 2014 ⁶³	China	65	Entecavir	42.8 ± 13.1	21	352.5 ± 77.2	6.3 ± 0.7	12	NR	Cohort
		54	Lamivudine	45.6 ± 11.4	23	345.2 ± 89.5	6.5 ± 0.9	12	NR	Cohort
Tsai et al., 2014 ⁶⁴	Taiwan	53	Entecavir	49 ± 13	15	1287 ± 788	8.2 ± 6.8	4	NR	Cohort
		114	Lamivudine	43 ± 15	47	1629 ± 1011	7.5 ± 6.9	4	NR	Cohort
Tsai et al., 2014 ⁶⁵	Taiwan	88	Telbivudine	55.7 ± 11.4	20	102.5 ± 137.5	5.1 ± 0.5	27.6	100	Cohort
		88	Entecavir	56.1 ± 9.8	17	125.8 ± 179	5.3 ± 0.4	53.1	100	Cohort
Koklu et al., 2013 ⁶⁶	Turkey	72	Tenofovir	54.2 ± 10.5	9	115.2 ± 217.1	4.9 ± 1.2	12	100	Cohort
		76	Entecavir	54.2 ± 11.2	17	86.2 ± 115.6	5 ± 1.2	12	100	Cohort
		74	Lamivudine	56.8 ± 11.4	10	53.2 ± 44.5	4 ± 1.3	12	100	Cohort
Question 2. Effectiveness of antiviral therapy in patients with immune tolerant chronic HBV infection										
Chan et al., 2014 ⁶⁷	Multinational	64	Tenofovir and placebo	33 ± 9.5	63	26.9 ± 14.05	8.4 ± 0.4	48	NR	RCT
		62	Tenofovir and entricitabine	33 ± 11.2	62	26.2 ± 9.88	8.4 ± 0.4	48	NR	RCT
Lu et al., 2015 ⁶⁸	China	30	Peg-IFN and adefovir	26.8 ± 3.1	30	<40	>5	6	NR	Cohort
		38	Control	26.8 ± 3.1	30			6	NR	Cohort
Question 3: Discontinuing versus continuing antiviral therapy in HBeAg-positive patients who seroconverted from HBeAg to anti-HBe										
Chaung et al., 2012 ⁶⁹	USA	49	Variety of oral antiviral alone or in combination	39 ± 12	NR	87 (16-1281)	7 ± 1.3	12	NR	Cohort
Fung et al., 2009 ⁷⁰	Hong Kong	39	Discontinued therapy	34 ± 10	NR	139 (37-576)	7 ± 1.2	12	NR	Cohort
		79	Lamivudine, continued therapy	32 (21-55)	NR	158 (21-2069)	7.9 (3-10.3)	45	NR	Cohort
		22	Discontinued therapy			176 (46-1670)	8.7 (6.4-10.2)	45	NR	Cohort
Question 5. Safety of entecavir compared to tenofovir										
Koklu et al., 2013 ⁶⁶	Turkey	54	Tenofovir	54.2 ± 10.2	9	115.2 ± 217.1	4.9 ± 1.2	21.4 ± 9.7	100	Cohort
		60	Entecavir	52.4 ± 11.2	17	86.2 ± 115.6	5 ± 1.2	24.0 ± 13.3	100	Cohort
Liaw et al., 2011 ⁷¹	Multinational	45	Tenofovir	52 (48-57)	14	48 (31-73)	5 (4.2-5.9)	12	NR	RCT
		45	Tenofovir and Entricitabine	50 (42-58)	18	54 (34-98)	5.6 (3.8-6.6)	12	NR	RCT
Dogan et al., 2012 ⁷²	Turkey	22	Entecavir	54 (47-58)	7	52 (41-66)	5.2 (3.5-6.7)	12	NR	Cohort
		65	Tenofovir	NR	29	114 ± 181	7 ± 6.9	12	NR	Cohort
		29	Entecavir	NR	10	84 ± 69	7.2 ± 7.6	12	NR	Cohort
Batirel et al., 2014 ⁷³	Turkey	90	Tenofovir	43.3 ± 12.9	29	116.7 ± 92.6	7.6 ± 4.6	30.2 ± 15.7	NR	Cohort

Table 1. Continued

Authors, Year	Country	Patients (N)	Interventions	Age (Years)	HBeAg Positive (N)	Baseline ALT (U/L)	Baseline HBV DNA (log ₁₀ IU/mL)*	Follow-up duration (months)	Baseline cirrhosis (%)	Study design
Cholongitas et al., 2015 ⁷⁴	Greece	105	Entecavir	42.0 ± 11.2	36	120 ± 96.6	7.6 ± 4.3	30.2 ± 15.7	NR	Cohort
		31	Tenofovir	60 ± 10	NR	57 ± 40	3.8 (>0-5.6)	25 (6-66)	100	Cohort
		21	Entecavir	58 ± 9	NR	75 ± 34	4.6 (>0-7.4)	18 (7-68)	100	Cohort
Huang et al., 2015 ⁷⁵	China	33	Tenofovir	35 (26-61)	NR	194.1 ± 128.5	6.50 ± 0.69	13.4 (6.2-28.0)	NR	Cohort
		65	Entecavir	39 (20-67)	NR	157.6 ± 216.8	6.15 ± 1.36	16 (6.0-27.0)	NR	Cohort
Hung et al., 2015 ⁷⁶	Taiwan	41	Tenofovir	49.8 ± 13.1	NR	1104 ± 918	6.3 ± 1.2	6	20	Cohort
		148	Entecavir	50.6 ± 14.7	NR	1084 ± 830	5.8 ± 1.2	6	34	Cohort
Mallet et al., 2014 ⁷⁷	France	70	Tenofovir	47 (37.8-56)	NR	52 (32-107)	4.4 (2.9-6.6)	22	NR	Cohort
		61	Entecavir	47 (37.8-56)	NR	52 (32-107)	4.4 (2.9-6.6)	22	NR	Cohort
Mauss et al., 2011 ⁷⁸	Germany	37	Tenofovir	43 (19-75)	11	73 (21-528)	5.58 (2.41- >8.04)	12 (6-36)	NR	Cohort
		32	Entecavir	43 (20-73)	16	72 (18-2230)	6.38 (3.49- >8.04)	24 (6-48)	NR	Cohort
Tien et al., 2014 ⁷⁹	USA	42	Tenofovir	49 ± 12	11	NR	NR	26 ± 13	20	Cohort
		44	Entecavir	51 ± 9	8	NR	NR	32 ± 24	10	Cohort
Gish et al., 2012 ⁸⁰	USA	80	Tenofovir	54.5 ± 13	NR	NR	6.99 (0-8.8)	20 (2-45)	NR	Retrospective cohort
		80	Entecavir	55.1 ± 12	NR	NR	7.36 (0-8.7)	29 (1-55)	NR	Retrospective cohort

*Baseline HBV DNA in studies that used different units were converted using the formulas 1 copy = 0.2 IU and 1 pg = 283,000 copies or 56,000 IU.

Abbreviations: anti-HBe, hepatitis B e antibody; NR, not reported; Peg, pegylated; ULN, upper limit of normal.

Table 2. Risk of Bias Assessment in the Included RCTs

Author, Year	Sequence Generation	Allocation Concealment	Blinding			Baseline Imbalance	Attrition Bias or Lost to Follow-Up
			Participants	Providers	Outcome Assessors		
Question 1: Effectiveness of antiviral therapy compared to control in patients with immune active chronic HBV infection (antiviral versus control)							
Anderson et al., 1987 ⁸	NR	NR	Yes	Yes	Yes	NR	NR
Krogsgaard et al., 1998 ²³	NR	NR	Yes	Yes	Yes	NR	NR
Chan et al., 2007 ²⁴	Randomized; randomization was centralized and stratified according to geographical region	NR	Yes	Yes	Yes	No	>15%
Eun et al., 2007 ²⁵	Randomized	NR	NR	NR	NR	NR	NR
Dienstag et al., 1999 ²⁹	Randomized	Yes	Yes	Yes	NR	No	10%-15%
Liaw et al., 2004 ³³	Randomized	NR	Yes	NR	Yes	NR	NR
Garg et al., 2011 ⁴⁶	Randomized; randomization was done with a random number table	Yes	Yes	Yes	NR	No	<10%
Question 1. Head-to-head studies comparing individual antiviral agents							
Chan et al., 2012 ⁵⁰	Randomized; centralized, stratifying based on screening CTP score and ALT level	Yes	Yes	Yes	Yes	No	<10%
Chang et al., 2006 ⁵¹	Randomized	NR	Yes	Yes	Yes	NR	NR
Lai et al., 2006 ⁵²	Randomized	NR	Yes	NR	Yes	NR	NR
Lau et al., 2005 ⁵³	Randomized; centralized and stratified according to geographic region and ALT levels	NR	NR	NR	NR	NR	NR
Marcellin et al., 2004 ⁵⁴	Randomized; centralized and stratified according to geographic region and ALT levels	NR	Yes	Yes	Yes	NR	NR
Wang et al., 2013 ⁵⁵	Randomized	NR	NR	NR	NR	No	NR
Yang et al., 2009 ⁵⁶	Randomized	NR	NR	NR	NR	NR	<10%
Liaw et al., 2011 ⁵⁷	Randomized; randomization was not blocked or stratified	NR	No	No	No	No	<10%
Question 2. Effectiveness of antiviral therapy in patients with immune tolerant chronic HBV infection							
Chan et al., 2014 ⁶⁷	Randomization	NR	Yes	Yes	NR	None	<10%
Question 5. Safety of entecavir compared to tenofovir							
Liaw et al., 2011 ⁷¹	Randomization	NR	Yes	Yes	NR	None	<10%

Abbreviations: CTP, Child-Turcotte-Pugh; NR, not reported.

In five observational studies^{25,26,35,38,41} (Fig. 4) with a mean follow-up of 84 months, IFN- α compared to no treatment significantly decreased the risk of HCC (five studies, RR = 0.6, 95% CI 0.4-0.9, $I^2 = 0\%$) but not of all-cause mortality (one study, RR = 0.7, 95% CI 0.5-2.4, $I^2 = 56.9\%$) or decompensated liver disease (one study, RR = 0.7, 95% CI 0.3-1.5).

In four observational studies^{26,35,38,41} (Fig. 5) with a mean follow-up of 45 months, lamivudine versus no treatment significantly reduced the risk of HCC (four studies, RR = 0.6, 95% CI 0.4-0.96, $I^2 = 49.9\%$), all-cause mortality (one study, RR = 0.4, 95% CI 0.3-0.6), and decompensated liver disease (one study, RR = 0.3, 95% CI 0.3-0.5). In one cohort study⁴⁰ of 1980 patients with cirrhosis followed for a mean of 52 months, entecavir versus control reduced the risk of HCC (RR = 0.3, 95% CI 0.1-0.5) and death (RR = 0.6, 95% CI 0.3-0.98).

Effectiveness of Antiviral Therapy Compared to Control in Patients With Chronic HBV Infection and Decompensated Cirrhosis. In two observational studies with follow-up of 29 months,^{27,32} lamivudine versus control reduced all-cause mortality (two studies, RR = 0.5, 95% CI 0.3-0.8, $I^2 = 0\%$).

Effectiveness of Antiviral Therapy Compared to Control in Patients With Chronic HBV Infection Experiencing Acute on Chronic Liver Failure. In one RCT⁴⁶ involving 26 patients followed for 1 year, tenofovir reduced all-cause mortality (RR = 0.5, 95% CI 0.3-0.99, moderate-quality evidence). In four observational studies^{28,37,42,44} with a mean follow-up of 26 months, antiviral therapy versus no therapy reduced all-cause mortality (RR = 0.7, 95% CI 0.6-0.8, $I^2 = 5.4\%$). Similarly, reduced mortality was also found in studies evaluating individual therapies including lamivudine (RR = 0.8, 95% CI 0.7-0.9, $I^2 = 50.2\%$),^{28,37,44}

Table 3. Risk of Bias Assessment for the Included Nonrandomized Studies

Author, Year	Selection of Cohort/Patients		Ascertainment of Exposure	Assessment and Clear Ascertainment of Outcome	Adequacy of Follow-Up	Funding Sources
	Exposed Cohort	Nonexposed Cohort/Control				
Question 1: Effectiveness of antiviral therapy compared to control in patients with immune active chronic HBV infection (antiviral versus control)						
IHCSCG, 1998 ⁹	Selected group of users	No description of the derivation of the nonexposed cohort	No description	No description	NR	NR
Lin et al., 2007 ¹⁰	Selected group of users	No description of the derivation of the nonexposed cohort	No description	No description	NR	NR
Truong et al., 2005 ¹¹	Somewhat representative of the community or population	Drawn from the same community as the exposed cohort	Secure records	Record linkage	NA	NR
Tangkijvanich et al., 2001 ¹²	Selected group of users	No description of the derivation of the nonexposed cohort	No description	No description	NR	NR
Papatheodoridis et al., 2001 ¹³	No description	No description of the derivation of the nonexposed cohort	Secure records	Record linkage	Complete follow-up	NR
Niederlau et al., 1996 ¹⁴	Selected group of users	No description of the derivation of the nonexposed cohort	No description	No description	NR	NR
Lin et al., 2004 ¹⁵	Somewhat representative of the community or population	Drawn from a different community or population from the exposed cohort	Secure records	Record linkage	Complete follow-up	Reported
Benvegna et al., 1998 ¹⁶	No description	No description	No description	No description	NR	NR
Tong et al., 2006 ¹⁷	No description	No description	No description	No description	NR	NR
Di Marco et al., 1999 ¹⁸	No description	No description	No description	No description	NR	NR
Brunetto et al., 2002 ¹⁹	No description	No description	No description	No description	NR	NR
Mahmood et al., 2005 ²⁰	Selected group of users	No description of the derivation of the nonexposed cohort	No description	No description	NR	NR
Ikeda et al., 1998 ²¹	Selected group of users	No description of the derivation of the nonexposed cohort	No description	No description	NR	NR
Fattovich et al., 1997 ²²	Selected group of users	No description of the derivation of the nonexposed cohort	Secure records	Record linkage	NR	NR
Tong et al., 2009 ²⁶	Selected group of users	No description of the derivation of the nonexposed cohort	No description	No description	NR	NR
Das et al., 2010 ²⁷	Selected group of users	No description of the derivation of the nonexposed cohort	No description	No description	NR	NR
Cui et al., 2010 ²⁸	Truly representative of the community or population	Drawn from the same community as the exposed cohort	Secure records	Record linkage	Complete follow-up	NR
Chan et al., 2002 ³⁰	Selected group of users	Drawn from a different community or population from the exposed cohort	Secure record	Record linkage	NR	NR
Lok et al., 2003 ³¹	Somewhat representative of the community or population	Drawn from the same community as the exposed cohort	Secure records	Record linkage	Follow-up rate <90% and no description of the reasons for loss to follow-up	NR
Manolakopoulos et al., 2004 ³²	Selected group of users	No description of the derivation of the nonexposed cohort	No description	No description	NR	NR
Matsumoto et al., 2005 ³⁴	Truly representative of the community or population	Drawn from the same community as the exposed cohort	Secure records	Record linkage	NA	Reported

Table 3. Continued

Author, Year	Selection of Cohort/Patients		Ascertainment of Exposure	Assessment and Clear Ascertainment of Outcome	Adequacy of Follow-Up	Funding Sources
	Exposed Cohort	Nonexposed Cohort/Control				
Ma et al., 2007 ³⁵	No description	Drawn from the same community as the exposed cohort	No description	No description	NR	NR
Yuen et al., 2007 ³⁶	Truly representative of the community or population	Drawn from a different community or population from the exposed cohort	Secure records	Record linkage	Follow-up rate <90% and no description of the reasons for loss to follow-up	Reported
Sun et al., 2010 ³⁷	Truly representative of the community or population	Drawn from the same community as the exposed cohort	Secure records	Record linkage	Complete follow-up	NR
Kim et al., 2012 ³⁸	Truly representative of the community or population	Drawn from the same community as the exposed cohort	Secure records	Record linkage	Complete follow-up	Reported
Eun et al., 2010 ³⁹	Truly representative of the community or population	Drawn from the same community as the exposed cohort	Secure records	Record linkage	Complete follow-up	Reported
Wong et al., 2013 ⁴⁰	Somewhat representative of the community or population	Drawn from the same community as the exposed cohort	No description	No description	NR	NR
Hosaka et al., 2013 ⁴¹	Somewhat representative of the community or population	Drawn from the same community as the exposed cohort	No description	No description	NR	NR
Lin et al., 2013 ⁴²	Truly representative of the community or population	Drawn from the same community as the exposed cohort	Secure records	Record linkage	Subjects lost to follow-up unlikely to introduce bias, small number lost to follow-up	Reported
Xiao et al., 2009 ⁴³	No description of the derivation of the cohort	Drawn from the same community as the exposed cohort	No description	No description	NR	NR
Xu et al., 2009 ⁴⁴	Truly representative of the community or population	No description of the derivation of the nonexposed cohort	No description	No description	NR	NR
Chen et al., 2009 ⁴⁵	Somewhat representative of the community or population	Drawn from the same community as the exposed cohort	No description	Record linkage	Complete follow-up, all subjects accounted for	Reported
Wu et al., 2014 ⁴⁷	Truly representative of the community or population	Drawn from the same community as the exposed cohort	Secure records	Record linkage	NR	NR
Gordon et al., 2014 ⁴⁸	Truly representative of the community or population	Drawn from the same community as the exposed cohort	Secure records	Record linkage	NR	Reported
Kumada et al., 2013 ⁴⁹	Truly representative of the community or population	Drawn from the same community as the exposed cohort	Secure records	Record linkage	NR	Reported
Question 1. Head-to-head studies comparing individual antiviral agents						
Cui et al., 2010 ²⁸	Truly representative of the community or population	Drawn from the same community as the exposed cohort	Secure records	Record linkage	Complete follow-up	NR
Lim et al., 2014 ⁵⁸	Selected group of users	Drawn from a different community or population from the exposed cohort	Secure records	Record linkage	Complete follow-up	Reported
Hsu et al., 2012 ⁵⁹	Somewhat representative of the community or population	Drawn from the same community as the exposed cohort	Secure records	No description	NR	Reported
Wong et al., 2011 ⁶⁰	Truly representative of the community or population	Drawn from the same community as the exposed cohort	Secure records	Independent blind assessment	Follow-up rate <90% and no description	Reported

Table 3. Continued

Author, Year	Selection of Cohort/Patients		Ascertainment of Exposure	Assessment and Clear Ascertainment of Outcome	Adequacy of Follow-Up	Funding Sources
	Exposed Cohort	Nonexposed Cohort/Control				
Liang et al., 2009 ⁶¹	No description	Drawn from the same community as the exposed cohort	Secure records	No description	Not reported of the reasons for loss to follow-up	NR
Chen et al., 2014 ⁶²	Somewhat representative of the community or population	Drawn from the same community as the exposed cohort	Secure records	Record linkage	NR	Reported
Zhang et al., 2014 ⁶³	No description of the derivation of the cohort	No description of the derivation of the non-exposed cohort	Secure records	Record linkage	Follow-up rate <90% and no description of the reasons for loss to follow-up	NR
Tsai et al., 2014 ⁶⁴	Selected group of users	Drawn from a different community or population from the exposed cohort	Secure records	Independent blind assessment	NR	NR
Tsai et al., 2014 ⁶⁵	Truly representative of the community or population	Drawn from the same community as the exposed cohort	Secure records	Record linkage	Follow-up rate <90% and no description of the reasons for loss to follow-up	Reported
Koklu et al., 2013 ⁶⁶	Truly representative of the community or population	Drawn from the same community as the exposed cohort	Secure records	Record linkage	Complete follow-up	NR
Question 2. Effectiveness of antiviral therapy in patients with immune tolerant chronic HBV infection						
Lu et al., 2015 ⁶⁸	Selected group of users	Drawn from the same community as the exposed cohort	Secure records	Record linkage	NR	NR
Question 3: Discontinuing versus continuing antiviral therapy in HBeAg-positive patients who seroconverted from HBeAg to anti-HBe						
Chaung et al., 2012 ⁶⁹	Selected group of users	Drawn from the same community as the exposed cohort	Secure records	Record linkage	NR	NR
Fung et al., 2009 ⁷⁰	Selected group of users	Drawn from the same community as the exposed cohort	Secure records	Record linkage	NR	NR
Question 5. Safety of entecavir compared to tenofovir						
Koklu et al., 2013 ⁶⁶	Selected group of users	Drawn from the same community as the exposed cohort	Secure records	Record linkage	NR	NR
Dogan et al., 2012 ⁷²	Selected group of users	Drawn from the same community as the exposed cohort	Secure records	Record linkage	NR	NR
Batirel et al., 2014 ⁷³	Selected group of users	Drawn from the same community as the exposed cohort	Secure records	Record linkage	NR	NR
Cholongitas et al., 2015 ⁷⁴	Selected group of users	Drawn from the same community as the exposed cohort	Secure records	Record linkage	NR	NR
Huang et al., 2015 ⁷⁵	Selected group of users	Drawn from the same community as the exposed cohort	Secure records	Record linkage	NR	NR
Hung et al., 2015 ⁷⁶	Selected group of users	Drawn from the same community as the exposed cohort	Secure records	Record linkage	NR	NR
Mallet et al., 2014 ⁷⁷	Selected group of users	Drawn from the same community as the exposed cohort	Secure records	Record linkage	NR	NR
Mauss et al., 2011 ⁷⁸	Selected group of users	Drawn from the same community as the exposed cohort	Secure records	Record linkage	NR	NR

Table 3. Continued

Author, Year	Selection of Cohort/Patients		Ascertainment of Exposure	Assessment and Clear Ascertainment of Outcome	Adequacy of Follow-Up	Funding Sources
	Exposed Cohort	Nonexposed Cohort/Control				
Tien et al., 2014 ⁷⁹	Selected group of users	Drawn from the same community as the exposed cohort	Secure records	Record linkage	NR	NR
Gish et al., 2012 ⁸⁰	Selected group of users	Drawn from the same community as the exposed cohort	Secure records	Record linkage	NR	NR

Abbreviations: anti-HBe, hepatitis B e antibody; NR, not reported.

entecavir (RR = 0.7, 95% CI 0.6-0.8, $I^2 = 0\%$),^{28,42,44} and telbivudine (RR = 0.4, 95% CI 0.2-0.9).⁴⁴

Effectiveness of Antiviral Therapy Compared to Control in Patients With Chronic HBV Infection With Severe Acute Exacerbations. In three observational studies^{30,43,45} with more than 12-month mean follow-up, meta-analysis of antiviral therapy versus control showed no statistically significant reduction in all-cause mortality (RR = 0.9, 95% CI 0.5-1.5, $I^2 = 54.5\%$), which was consistent with studies evaluating the effect of individual agents: lamivudine (RR = 0.5, 95% CI 0.2-1.7)³⁰ and entecavir (RR = 0.9, 95% CI 0.5-1.9, $I^2 = 71.3\%$).^{43,45}

Head-to-Head Studies Comparing Individual Antiviral Agents. We included eight RCTs⁵⁰⁻⁵⁷ enrolling 2318 patients and 10 observational studies^{28,58-66} enrolling 6737 patients that compared one antiviral agent with another. We considered most of these RCTs^{52,55-57} to have high risk of bias due to unclear randomization methods, allocation concealment, blinding, and loss to follow-up. The observational studies were also limited by the unclear description of the characteristics for cohort selection, ascertainment of the outcomes, and inadequate follow-up. Tables 1 and 2 describe the details of the included studies and risk of bias.

Among five studies enrolling 3300 patients with chronic HBV infection and compensated cirrhosis (mean follow-up 22 months), one RCT⁵⁵ compared adefovir versus lamivudine and four observational studies compared entecavir versus lamivudine,⁵⁸ entecavir versus telbivudine,⁶⁵ lamivudine versus tenofovir,⁶⁶ and telbivudine versus lamivudine, respectively.⁶¹ Only 1 study⁵⁸ showed a significant difference in outcome with reduction in all-cause mortality in patients who received entecavir versus lamivudine (one study, RR = 0.4, 95% CI 0.3-0.6, very low-quality evidence).

Four studies enrolled 607 patients with chronic HBV infection and decompensated cirrhosis (mean follow-up 28 months). Three RCTs compared entecavir versus ade-

fovir,⁵⁷ adefovir versus lamivudine,⁵⁶ and telbivudine versus lamivudine, respectively⁵⁰; and one cohort study⁵⁹ compared entecavir versus lamivudine. Reduction in risk of HCC was observed in the RCT⁵⁷ comparing entecavir versus adefovir (RR = 0.4, 95% CI 0.2-0.8), and reduction in all-cause mortality was observed in the cohort study comparing entecavir versus lamivudine (RR = 0.4, 95% CI 0.3-0.7) in patients who received entecavir.

Three cohort studies^{28,62,63} that enrolled 508 patients with acute on chronic liver failure and compared entecavir to lamivudine (mean follow-up 32 months) showed no significant effect on all-cause mortality.

Two cohort studies^{60,64} that compared entecavir versus lamivudine in 320 patients with severe acute exacerbation of chronic hepatitis B (mean follow-up 32 months) showed no significant effect on mortality.

Question 2. Effectiveness of Antiviral Therapy in Patients With Immune-Tolerant Chronic HBV Infection

Two studies^{67,68} evaluated antiviral therapy in HBeAg-positive patients with normal ALT levels. Detailed study characteristics and risk of bias are described in Tables 1 and 2.

One RCT⁶⁷ compared tenofovir (64 patients) to a combination of tenofovir and emtricitabine (62 patients) for 192 weeks. Although no long-term clinical outcomes were reported, tenofovir and emtricitabine versus tenofovir showed a statistically significant increase in viral suppression (RR = 1.4, 95% CI 1.1-1.8, moderate-quality evidence) but no statistically significant increase in HBeAg loss (RR = 0.3, 95% CI 0.03-2.2), HBeAg seroconversion (RR = 0.1, 95% CI 0.01-2.8), or HBsAg clearance (RR = 1.0, 95% CI 0.3-3.9). The quality of evidence was low due to indirectness and imprecision.

In a cohort study⁶⁸ of 68 HBeAg-positive postpartum women, pegylated IFN and adefovir versus untreated control significantly improved rates of HBeAg




Fig. 2. Forest plot of clinical outcomes for randomized controlled trials comparing any antiviral vs. no treatment. I-square and P values for study heterogeneity cannot be computed for outcomes with only one study.

seroconversion (RR = 41.8, 95% CI 2.6-666.9) and HBeAg loss (RR = 20.3, 95% CI 1.2-337.7). The quality of evidence was very low, down-rated due to the observational nature of the study, risk of bias, and imprecision.

Question 3: Discontinuing Compared to Continuing Antiviral Therapy in HBeAg-Positive Patients Who Seroconverted From HBeAg to Hepatitis B e Antibody

Two observational studies^{69,70} compared patients with chronic hepatitis B who stopped therapy (61 patients) after HBeAg seroconversion to those who continued (128 patients) to receive antiviral therapy. For both studies, the median (range) duration of therapy leading to HBeAg seroconversion was 21 (1-120) months, median follow-up after stopping therapy was

40 (range 2-120) months, and median duration of consolidation treatment after HBeAg seroconversion was 12 (range 1-55) months. Characteristics and risk of bias for both studies are illustrated in Tables 1 and 3.

Compared to continued antiviral therapy, very low-quality evidence suggests increased risk of relapse of viremia in patients who stopped antiviral therapy (RR = 94.4, 95% CI 13.3-670.7, I² = 0%) with no effect on ALT flares. The rate of HBeAg seroreversion was 8% after a median of 6 months in 1 study,⁶⁹ with a cumulative incidence of 9% at 5 years in another study.⁷⁰ No clinical outcomes were reported. The quality of evidence was very low due to increased risk of bias, indirectness, and imprecision. Additional noncomparative and indirect evidence is summarized in the [Supporting Information](#).




Fig. 3. Forest plot of clinical outcomes for observational studies comparing antiviral therapy vs. no treatment in patients with chronic HBV infection and compensated cirrhosis.

Question 4. Stopping Compared to Continuing Antiviral Therapy In HBeAg-Negative Adults With Immune Active Chronic HBV Infection

We were unable to find comparative studies for this question. The Supporting Information summarizes uncontrolled studies and indirect evidence that may address this question. Data from these studies indicate a high rate of viral relapse when treatment was stopped, but rates of clinical relapse were lower.

Question 5. Safety of Entecavir Compared to Tenofovir

Eleven studies (one RCT⁷¹ and 10 observational studies^{66,72-80}) compared entecavir versus tenofovir in 1300 patients with a mean follow-up of 18.6 months.

Characteristics of the included studies and risk of bias are described in Tables 1 and 2.

Meta-analysis of the studies included showed no statistically significant difference between entecavir and tenofovir in renal safety profiles or hypophosphatemia, but duration of observation was short. No studies reported on bone density. Table 4 describes the detailed outcomes reported for each study.

Question 6. Adding a Second Antiviral Agent Compared to Continuing Monotherapy (Entecavir or Tenofovir) in Patients With Chronic HBV Infection and Persistent Viremia

We were unable to identify comparative studies for this question. Uncontrolled studies and indirect




Fig. 4. Forest plot of clinical outcomes for observational studies comparing IFN- α vs. no treatment in patients with chronic HBV infection and compensated cirrhosis. I-squared and P values for study heterogeneity cannot be computed for outcomes with only one study.

evidence (Supporting Information) showed little to no benefit in adding a second antiviral agent compared to continuing monotherapy with entecavir or tenofovir.

Question 7. Antiviral Therapy in Patients With Chronic HBV Infection and Compensated Cirrhosis and Low-Level Viremia (HBV DNA <2000 IU/mL)

We were unable to identify comparative studies on outcomes of these patients with or without antiviral therapy. The Supporting Information summarizes uncontrolled studies and indirect evidence that address this question. In patients with compensated cirrhosis and low-level viremia, one study specifically examined the benefit of antiviral therapy and found a decrease in incidence of HCC, but the results could be confounded by differences in the characteristics of treated versus untreated patients.⁸¹

Publication Bias. We were unable to evaluate publication bias due to high heterogeneity and the small number of studies for each outcome.

Discussion

The members of the AASLD methodology and writing committees for the HBV Practice Guideline developed seven key clinical questions that challenge clinicians and patients in daily practice. The methodologists performed an extensive literature search, selected studies that included a comparison group and data on clinical outcomes, and then rated the quality of the evidence. Sufficient comparative evidence was found for four of the key questions, but evidence was sparse or absent for the remaining three questions: when to stop therapy in persons with immune active chronic HBV infection who are HBeAg-negative, the benefit of adding




Fig. 5. Forest plot of clinical outcomes for observational studies comparing lamivudine vs. no treatment in patients with chronic HBV infection and compensated cirrhosis. I-square and P values for study heterogeneity cannot be computed for outcomes with only one study.

either entecavir or tenofovir in persons who fail to suppress HBV DNA to undetectable levels with either of these drugs alone, and whether antiviral therapy should be used in patients with compensated cirrhosis and HBV DNA levels below 2000 IU/mL. For these three questions, the committee identified indirect and non-comparative evidence (Supporting Information).

Antiviral therapy in patients with immune active chronic HBV infection had 59 published studies available for review and evaluation. Moderate-quality to low-quality evidence supported the benefit of therapy in reducing adverse outcomes of chronic HBV infection including progression to cirrhosis, liver decompensation, and all-cause mortality. Because the observational studies had more patients (59,201 versus 3463) and longer follow-up (60 versus 28 months), data on mortality and HCC from 35 observational studies were sufficiently precise, whereas data from seven RCTs were imprecise. These larger sample sizes and longer follow-up in the

observational studies account for the significant benefit of antiviral treatment on HCC and mortality found in the observational studies but not in the RCTs.

Given the indolent nature of chronic HBV infection, it is not surprising that evidence supporting the benefit of antiviral treatment on clinical outcomes was found only when the analysis was limited to patients with more advanced disease: compensated cirrhosis, decompensated cirrhosis, or acute on chronic liver failure. Indeed, most RCTs of antiviral therapy in chronic HBV infection enrolled only or mostly patients with no cirrhosis, and very few trials that enrolled predominantly patients with no cirrhosis provided data on clinical outcomes. Provision of evidence to support that antiviral therapy improves clinical outcomes in patients with chronic HBV infection and no cirrhosis would require thousands of patients followed for many years and withholding treatment in the control group until the completion of the study. Such a study would be unethical

Table 4. Outcomes Reported for Tenofovir Versus Entecavir in Chronic HBV Infection

Author, Year	Outcomes Reported	Tenofovir	Entecavir	RR (95% CI)
		Events/Total	Events/Total	
Koklu et al., 2013 ⁶⁶	Renal impairment	1/72	0/77	3.21 (0.13-77.44)
	Hypophosphatemia	1/72	0/77	3.21 (0.13-77.44)
	Increase of creatinine kinase	0/72	1/77	0.36 (0.01-8.60)
Liaw et al., 2011 ⁷¹	Increase in creatinine ≥ 0.5 mg/dL from baseline	4/45	1/22	1.96 (0.23-16.47)
	Phosphorus < 2.0 mg/dL	1/45	0/22	1.50 (0.00-35.40)
Batirel et al., 2014 ⁷³	Hypophosphatemia	2/90	0/105	5.82 (0.28-119.75)
Cholongitas et al., 2015 ⁷⁴	eGFR < 50 mL/minute	3/31	2/21	1.02 (0.19-5.57)
	Serum phosphate levels	NR	NR	NA
Hung et al., 2015 ⁷⁶	Baseline serum creatinine 0.5 mg/dL	2/30	2/99	3.30 (0.49-22.44)
	Reduction of eGFR	108 to 87 189 mL/min/1.73 m ²	92 to 84 mL/ min/1.73 m ²	NA
Huang et al., 2015 ⁷⁵	CK levels 2 times over the upper limit of normal	1/33	1/65	1.97 (0.13-30.50)
Mallet et al., 2014 ⁷⁷	Mean eGFR variation	0.6 (-0.8 to 1.94)	-0.1 (-1.5 to 1.3)	NA
Mauss et al., 2011 ⁷⁸	Changes in eGFR (CKD-EPI formula)	-0.92 mL/min	-1.00 mL/min	NA
	Decrease of eGFR > 20 mL/min	1/37	2/32	0.43 (0.04-4.55)
Tien et al., 2015 ⁷⁹	Phosphate threshold for renal tubular reabsorption < 2.8 mg/dL	18/42	10/44	1.89 (0.99-3.60)
	GFR by Cockcroft-Gault < 60 mL/min	1/42	2/44	0.52 (0.05-5.56)
	GFR by MDRD < 60 mL/min	1/42	2/44	0.52 (0.05-5.56)
	Serum phosphate (mg/dL) < 2.8 mg/dL	6/42	2/44	3.14 (0.67-14.71)
	SCr (mg/dL) > 1.5 mg/dL	0/42	0/44	NA
	Serum alkaline phosphatase > 145 U/L	0/42	1/44	0.35 (0.01-8.33)
Gish et al., 2012 ⁸⁰	Confirmed SCr increase 0.5 mg/dL	3/80	11/80	0.27 (0.08-0.94)
	New Cockcroft-Gault eGFR < 60 mL/min	15/80	6/80	2.50 (1.02-6.12)
	Decrease in eGFR 20% (MDRD)	33/80	35/80	0.94 (0.66-1.35)

Abbreviations: CK, creatine kinase; CKD-EPI, Chronic Kidney Disease Epidemiology Collaboration; eGFR, estimated glomerular filtration rate; MDRD, Modification of Diet in Renal Disease; NA, not available; NR, not reported; SCr, serum creatinine.

and likely infeasible. Thus, evidence supporting the benefit of antiviral therapy in patients without cirrhosis has to rely on intermediate outcomes such as HBV DNA suppression, ALT normalization, HBeAg seroconversion, HBsAg loss, and cirrhosis prevention or regression. These intermediate outcomes have been shown to correlate with improvement in clinical outcomes and represent a series of steps toward the ultimate goal of improving clinical outcome. For example, HBV DNA suppression precedes HBeAg seroconversion, which precedes HBsAg loss; and HBsAg loss has been associated with decreased risk of HCC, particularly if it occurs before the development of cirrhosis.

Recent studies showed that high levels of HBV viremia are associated with an increased risk of cirrhosis, HCC, and liver-related mortality.⁸²⁻⁸⁴ Patients in the immune tolerant phase have the highest level of viremia. In the two studies exclusively enrolling patients in the immune tolerant phase, clinical outcomes were not reported but rates of intermediate outcomes were lower than those in patients in the HBeAg-positive immune active phase.

In the two observational studies comparing the risk of viral relapse and HBeAg seroreversion in HBeAg-positive patients who achieved HBeAg seroconversion

during nucleos(t)ide analogue therapy and who stopped versus continued therapy, very low-quality evidence suggests an increased risk of relapse of viremia with stopping. Other observational studies (see [Supporting Information](#)) showed durable HBeAg seroconversion varying from 20% to 90% depending on the duration of consolidation therapy after achieving HBeAg seroconversion, the most consistent predictor of durable response. Studies directly comparing stopping versus continuing therapy in HBeAg-negative patients on nucleos(t)ide analogue therapy were not found; however, observational studies in the literature on the virologic, serologic, and biochemical outcomes of patients who stopped therapy showed that viral relapse is universal but that sustained clinical remission and even HBsAg loss are possible (see [Supporting Information](#)). Because hepatitis flares and hepatic decompensation may occur after stopping treatment, close monitoring after discontinuation of treatment is important, especially for those with cirrhosis at the start of therapy who have the highest risk for decompensation.

Entecavir and tenofovir have been used as first-line nucleos(t)ide analogues because of their potent antiviral activity and low risk of antiviral drug resistance. Tenofovir can cause impairment in renal function, renal tubular

dysfunction including Fanconi anemia, and decreased bone mineral density. Meta-analysis of studies comparing monotherapy with entecavir or tenofovir did not show a significant difference in serum creatinine level, estimated glomerular filtration rate, or serum phosphate level; however, the duration of treatment was short in these studies.

While entecavir and tenofovir have potent antiviral activity, some patients have persistent viremia despite being adherent to medication. This is more common among HBeAg-positive patients with high baseline serum HBV DNA. Studies comparing continuing entecavir or tenofovir monotherapy versus adding a second antiviral agent in patients with persistent viremia were not found. Observational studies of patients who continued entecavir or tenofovir monotherapy showed that most patients ultimately achieved undetectable HBV DNA.

Patients with compensated cirrhosis have a high risk of liver failure and HCC, particularly those with high levels of HBV DNA. The benefit of antiviral therapy in patients with compensated cirrhosis and low levels of HBV DNA has not been established. One retrospective study comparing outcomes of patients with compensated cirrhosis and low levels of HBV DNA (<2000 IU/mL) with or without antiviral therapy suggests a benefit of antiviral therapy in decreasing the incidence of HCC; but patients who received treatment differed substantially from those who did not receive treatment, and in most patients the HBV DNA level was higher than 2000 IU/mL at the time treatment was started.⁸¹

Several questions that had been addressed in the previous AASLD HBV Guidelines were not included in this systematic review: who should be screened for HBV infection, who should be vaccinated against HBV, what clinical and laboratory criteria (levels of HBV DNA and ALT) should be used to initiate antiviral therapy, who should undergo surveillance for HCC, and how frequently patients with chronic HBV infection who are not receiving antiviral therapy should be monitored. Management of special populations, such as those with human immunodeficiency virus or hepatitis C or D viral coinfection and those requiring immunosuppressive therapy, was also not addressed in the current review because data from controlled studies for these patient populations were sparse. Additional recommendations can be found in the previous AASLD HBV Guideline and in the Centers for Disease Control and Prevention and the World Health Organization guidelines.⁸⁵⁻⁸⁸

In conclusion, most of the current literature focuses on the immune active phases of chronic HBV infection.

Decision-making in other commonly encountered and challenging clinical settings depends on indirect evidence. In addition to evidence-based data, management of patients with chronic HBV infection should take into consideration individual patient preference and available resources. Recommendations for management of adults with chronic HBV infection based on this systematic review are provided in the updated AASLD guidelines.⁸⁹

References

- Ott JJ, Stevens GA, Groeger J, Wiersma ST. Global epidemiology of hepatitis B virus infection: new estimates of age-specific HBsAg seroprevalence and endemicity. *Vaccine* 2012;30:2212-2219.
- Wasley A, Kruszon-Moran D, Kuhnert W, Simard EP, Finelli L, McQuillan G, et al. The prevalence of hepatitis B virus infection in the United States in the era of vaccination. *J Infect Dis* 2010;202:192-201.
- Kowdley KV, Wang CC, Welch S, Roberts H, Brosgart CL. Prevalence of chronic hepatitis B among foreign-born persons living in the United States by country of origin. *HEPATOLOGY* 2012;56:422-433.
- Lok AS, McMahon BJ. Chronic hepatitis B. *HEPATOLOGY* 2007;45:507-539.
- Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.
- Murad MH, Montori VM, Ioannidis JP, Jaeschke R, Devereaux PJ, Prasad K, et al. How to read a systematic review and meta-analysis and apply the results to patient care: users' guides to the medical literature. *JAMA* 2014;312:171-179.
- Sterne JA, Sutton AJ, Ioannidis JP, Terrin N, Jones DR, Lau J, et al. Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials. *BMJ* 2011;343:d4002.
- Anderson MG, Harrison TJ, Alexander G, Zuckerman AJ, Murray-Lyon IM. Randomised controlled trial of lymphoblastoid interferon for chronic active hepatitis B. *Gut* 1987;28:619-622.
- International Interferon-alpha Hepatocellular Carcinoma Study Group. Effect of interferon-alpha on progression of cirrhosis to hepatocellular carcinoma: a retrospective cohort study. *Lancet* 1998;351:1535-1539.
- Lin S-M, Yu M-L, Lee C-M, Chien R-N, Sheen IS, Chu C-M, et al. Interferon therapy in HBeAg positive chronic hepatitis reduces progression to cirrhosis and hepatocellular carcinoma. *J Hepatol* 2007;46:45-52.
- Truong BX, Seo Y, Kato M, Hamano K, Ninomiya T, Katayama M, et al. Long-term follow-up of Japanese patients with chronic hepatitis B treated with interferon-alpha. *Int J Mol Med* 2005;16:279-284.
- Tangkijvanich P, Thong-ngam D, Mahachai V, Klachareon N, Suwangool P, Kullavanijaya P. Long-term effect of interferon therapy on incidence of cirrhosis and hepatocellular carcinoma in Thai patients with chronic hepatitis B. *Southeast Asian J Trop Med Public Health* 2001;32:452-458.
- Papatheodoridis GV, Manesis E, Hadziyannis SJ. The long-term outcome of interferon-alpha treated and untreated patients with HBeAg-negative chronic hepatitis B. *J Hepatol* 2001;34:306-313.
- Niederau C, Heintges T, Lange S, Goldmann G, Niederau CM, Mohr L, et al. Long-term follow-up of HBeAg-positive patients treated with interferon alfa for chronic hepatitis B. *N Engl J Med* 1996;334:1422-1427.
- Lin SM, Tai DI, Chien RN, Sheen IS, Chu CM, Liaw YF. Comparison of long-term effects of lymphoblastoid interferon alpha and

- recombinant interferon alpha-2a therapy in patients with chronic hepatitis B. *J Viral Hepat* 2004;11:349-357.
16. Benvegna L, Chemello L, Noventa F, Fattovich G, Pontisso P, Alberti A. Retrospective analysis of the effect of interferon therapy on the clinical outcome of patients with viral cirrhosis. *Cancer* 1998;83:901-909.
 17. Tong MJ, Blatt LM, Tyson KB, Kao VWC. Death from liver disease and development of hepatocellular carcinoma in patients with chronic hepatitis B virus infection: a prospective study. *Gastroenterol Hepatol (Qué)* 2006;2:41-47.
 18. Di Marco V, Lo Iacono O, Camma C, Vaccaro A, Giunta M, Matorana G, et al. The long-term course of chronic hepatitis B. *HEPATOLOGY* 1999;30:257-264.
 19. Brunetto MR, Oliveri F, Coco B, Leandro G, Colombatto P, Gorin JM, et al. Outcome of anti-HBe positive chronic hepatitis B in alpha-interferon treated and untreated patients: a long term cohort study. *J Hepatol* 2002;36:263-270.
 20. Mahmood S, Niiyama G, Kamei A, Izumi A, Nakata K, Ikeda H, et al. Influence of viral load and genotype in the progression of hepatitis B-associated liver cirrhosis to hepatocellular carcinoma. *Liver Int* 2005;25:220-225.
 21. Ikeda K, Saitoh S, Suzuki Y, Kobayashi M, Tsubota A, Fukuda M, et al. Interferon decreases hepatocellular carcinogenesis in patients with cirrhosis caused by the hepatitis B virus: a pilot study. *Cancer* 1998;82:827-835.
 22. Fattovich G, Giustina G, Realdi G, Corrocher R, Schalm SW. Long-term outcome of hepatitis B e antigen-positive patients with compensated cirrhosis treated with interferon alfa. *HEPATOLOGY* 1997;26:1338-1342.
 23. Krogsgaard K, Thomas HC, Farrell G, Cooksley WGE, Moroni M, Perez V, et al. The long-term effect of treatment with interferon-alpha2a in chronic hepatitis B. *J Viral Hepat* 1998;5:389-397.
 24. Chan HLY, Wang H, Niu J, Chim AML, Sung JY. Two-year lamivudine treatment for hepatitis B e antigen-negative chronic hepatitis B: a double-blind, placebo-controlled trial. *Antivir Ther* 2007;12:345-353.
 25. Eun LH Jr, Lee SH, Kim TN, Jang BIK, Choi JW, Park YS, et al. The effect of lamivudine and adefovir dipivoxil on preventing hepatocellular carcinoma in HBV-related liver cirrhosis. *HEPATOLOGY* 2007;46:664A-665A.
 26. Tong MJ, Hsien C, Song JJ, Kao JH, Sun HE, Hsu L, et al. Factors associated with progression to hepatocellular carcinoma and to death from liver complications in patients with HBsAg-positive cirrhosis. *Dig Dis Sci* 2009;54:1337-1346.
 27. Das K, Das K, Datta S, Pal S, Hembram JR, Dhali GK, et al. Course of disease and survival after onset of decompensation in hepatitis B virus-related cirrhosis. *Liver Int* 2010;30:1033-1042.
 28. Cui Y-L, Yan F, Wang Y-B, Song X-Q, Liu L, Lei X-Z, et al. Nucleoside analogue can improve the long-term prognosis of patients with hepatitis B virus infection-associated acute on chronic liver failure. *Dig Dis Sci* 2010;55:2373-2380.
 29. Dienstag JL, Schiff ER, Wright TL, Perrillo RP, Hann HW, Goodman Z, et al. Lamivudine as initial treatment for chronic hepatitis B in the United States. *N Engl J Med* 1999;341:1256-1263.
 30. Chan HLY, Tsang SWC, Hui Y, Leung NWY, Chan FKL, Sung JY. The role of lamivudine and predictors of mortality in severe flare-up of chronic hepatitis B with jaundice. *J Viral Hepat* 2002;9:424-428.
 31. Lok ASF, Lai C-L, Leung N, Yao G-B, Cui Z-Y, Schiff ER, et al. Long-term safety of lamivudine treatment in patients with chronic hepatitis B. *Gastroenterology* 2003;125:1714-1722.
 32. Manolakopoulos S, Karatapanis S, Elefsiniotis J, Mathou N, Vlachogiannakos J, Iliadou E, et al. Clinical course of lamivudine monotherapy in patients with decompensated cirrhosis due to HBeAg negative chronic HBV infection. *Am J Gastroenterol* 2004;99:57-63.
 33. Liaw Y-F, Sung JY, Chow WC, Farrell G, Lee C-Z, Yuen H, et al. Lamivudine for patients with chronic hepatitis B and advanced liver disease. *N Engl J Med* 2004;351:1521-1531.
 34. Matsumoto A, Tanaka E, Rokuhara A, Omata M, Iino S, Tanikawa K, et al. Efficacy of lamivudine for preventing hepatocellular carcinoma in chronic hepatitis B: a multicenter retrospective study of 2795 patients. *Hepatol Res* 2005;32:173-184.
 35. Ma H, Guo F, Wei L, Sun Y, Wang H. The prospective study of the clinical features and outcome of HBeAg-negative and HBeAg-positive cirrhosis in patients with chronic type B hepatitis. [in Chinese] *Zhonghua Yi Xue Za Zhi* 2007;87:1832-1835.
 36. Yuen M-F, Seto W-K, Chow DH-F, Tsui K, Wong DK-H, Ngai VW-S, et al. Long-term lamivudine therapy reduces the risk of long-term complications of chronic hepatitis B infection even in patients without advanced disease. *Antivir Ther* 2007;12:1295-1303.
 37. Sun L-J, Yu J-W, Zhao Y-H, Kang P, Li S-C. Influential factors of prognosis in lamivudine treatment for patients with acute-on-chronic hepatitis B liver failure. *J Gastroenterol Hepatol* 2010;25:583-590.
 38. Kim CH, Um SH, Seo YS, Jung JY, Kim JD, Yim HJ, et al. Prognosis of hepatitis B-related liver cirrhosis in the era of oral nucleos(t)ide analog antiviral agents. *J Gastroenterol Hepatol* 2012;27:1589-1595.
 39. Eun JR, Lee HJ, Kim TN, Lee KS. Risk assessment for the development of hepatocellular carcinoma: according to on-treatment viral response during long-term lamivudine therapy in hepatitis B virus-related liver disease. *J Hepatol* 2010;53:118-125.
 40. Wong GL-H, Chan HL-Y, Mak CW-H, Lee SK-Y, Ip ZM-Y, Lam AT-H, et al. Entecavir treatment reduces hepatic events and deaths in chronic hepatitis B patients with liver cirrhosis. *HEPATOLOGY* 2013;58:1537-1547.
 41. Hosaka T, Suzuki F, Kobayashi M, Seko Y, Kawamura Y, Sezaki H, et al. Long-term entecavir treatment reduces hepatocellular carcinoma incidence in patients with hepatitis B virus infection. *HEPATOLOGY* 2013;58:98-107.
 42. Lin B, Pan CQ, Xie D, Xie J, Xie S, Zhang X, et al. Entecavir improves the outcome of acute-on-chronic liver failure due to the acute exacerbation of chronic hepatitis B. *Hepatol Int* 2013;7:460-467.
 43. Xiao G-M, He K-Y, Jia W-D, Lei C-L, Yang Z. Case-controlled study of entecavir treatment for chronic severe hepatitis B. [in Chinese] *Zhonghua Shi Yan He Lin Chuang Bing Du Xue Za Zhi* 2009;23:56-58.
 44. Xu Q-h, Chen L-b, Xu Z, Shu X, Chen N, Cao H, et al. The short-term efficacy of antiviral treatment in patients with acute-on-chronic hepatitis B liver failure. [in Chinese] *Zhonghua Shi Yan He Lin Chuang Bing Du Xue Za Zhi* 2009;23:467-469.
 45. Chen J, Han JH, Liu C, Yu RH, Li FZ, Li QF, et al. Short-term entecavir therapy of chronic severe hepatitis B. *Hepatobiliary Pancreat Dis Int* 2009;8:261-266.
 46. Garg H, Sarin SK, Kumar M, Garg V, Sharma BC, Kumar A. Tenofovir improves the outcome in patients with spontaneous reactivation of hepatitis B presenting as acute-on-chronic liver failure. *HEPATOLOGY* 2011;53:774-780. Erratum in *HEPATOLOGY* 2011;54:1114.
 47. Wu CY, Lin JT, Ho HJ, Su CW, Lee TY, Wang SY, et al. Association of nucleos(t)ide analogue therapy with reduced risk of hepatocellular carcinoma in patients with chronic hepatitis B-A nationwide cohort study. *Gastroenterology* 2014;147:143-151.
 48. Gordon SC, Lamerato LE, Rupp LB, Li J, Holmberg SD, Moorman AC, et al. Antiviral therapy for chronic hepatitis B virus infection and development of hepatocellular carcinoma in a US population. *Clin Gastroenterol Hepatol* 2014;12:885-893.
 49. Kumada T, Toyoda H, Tada T, Kiriya S, Tanikawa M, Hisanaga Y, et al. Effect of nucleos(t)ide analogue therapy on hepatocarcinogenesis in chronic hepatitis B patients: a propensity score analysis. *J Hepatol* 2013;58:427-433.
 50. Chan HLY, Chen YC, Gane EJ, Sarin SK, Suh DJ, Piratvisuth T, et al. Randomized clinical trial: efficacy and safety of telbivudine and lamivudine in treatment-naive patients with HBV-related decompensated cirrhosis. *J Viral Hepat* 2012;19:732-743.
 51. Chang T-T, Gish RG, de Man R, Gadano A, Sollano J, Chao Y-C, et al. A comparison of entecavir and lamivudine for HBeAg-positive chronic hepatitis B. *N Engl J Med* 2006;354:1001-1010.
 52. Lai C-L, Shouval D, Lok AS, Chang T-T, Cheinquer H, Goodman Z, et al. Entecavir versus lamivudine for patients with HBeAg-negative

- chronic hepatitis B. *N Engl J Med* 2006;354:1011-1020. Erratum in *N Engl J Med* 2006;354:1863.
53. Lau GKK, Piratvisuth T, Kang XL, Marcellin P, Thongsawat S, Cooksley G, et al. Peginterferon alfa-2a, lamivudine, and the combination for HBeAg-positive chronic hepatitis B. *N Engl J Med* 2005;352:2682-2695.
 54. Marcellin P, Lau GKK, Bonino F, Farci P, Hadziyannis S, Jin R, et al. Peginterferon alfa-2a alone, lamivudine alone, and the two in combination in patients with HBeAg-negative chronic hepatitis B. *N Engl J Med* 2004;351:1206-1217.
 55. Wang H, Ji YY, Yao GB, Ma XY, Xie Q, Pang HY, et al. Two years efficiency of lamivudine and adefovir dipivoxil combined therapy in chronic hepatitis B patients. *Eur Rev Med Pharmacol Sci* 2013;17:636-643.
 56. Yang Q, Gong Z-j, Hu D-f. A clinical study of adefovir dipivoxil treatment for chronic hepatitis patients with cirrhosis in their decompensation period. [in Chinese] *Chung Hua Kan Tsang Ping Tsa Chih* 2009;17:515-519.
 57. Liaw Y-F, Raptopoulou-Gigi M, Cheinquer H, Sarin SK, Tanwandee T, Leung N, et al. Efficacy and safety of entecavir versus adefovir in chronic hepatitis B patients with hepatic decompensation: a randomized, open-label study. *HEPATOLOGY* 2011;54:91-100.
 58. Lim Y-S, Han S, Heo N-Y, Shim JH, Lee HC, Suh DJ. Mortality, liver transplantation, and hepatocellular carcinoma among patients with chronic hepatitis B treated with entecavir vs lamivudine. *Gastroenterology* 2014;147:152-161.
 59. Hsu Y-C, Mo L-R, Chang C-Y, Perng D-S, Tseng C-H, Lo G-H, et al. Entecavir versus lamivudine in the treatment of chronic hepatitis B patients with hepatic decompensation. *Antivir Ther* 2012;17:605-612.
 60. Wong VW-S, Wong GL-H, Yiu KK-L, Chim AM-L, Chu SH-T, Chan H-Y, et al. Entecavir treatment in patients with severe acute exacerbation of chronic hepatitis B. *J Hepatol* 2011;54:236-242.
 61. Liang J, Han T, Xiao S-X. Telbivudine treatment on cirrhosis resulting from chronic hepatitis B. [in Chinese] *Chung Hua Kan Tsang Ping Tsa Chih* 2009;17:24-27.
 62. **Chen CH, Lin CL**, Hu TH, Hung CH, Tseng PL, Wang JH, et al. Entecavir versus lamivudine in chronic hepatitis B patients with severe acute exacerbation and hepatic decompensation. *J Hepatol* 2014;60:1127-1134.
 63. Zhang Y, Hu XY, Zhong S, Yang F, Zhou TY, Chen G, et al. Entecavir vs lamivudine therapy for naive patients with spontaneous reactivation of hepatitis B presenting as acute-on-chronic liver failure. *World J Gastroenterol* 2014;20:4745-4752.
 64. Tsai WL, Chiang PH, Chan HH, Lin HS, Lai KH, Cheng JS, et al. Early entecavir treatment for chronic hepatitis B with severe acute exacerbation. *Antimicrob Agents Chemother* 2014;58:1918-1921.
 65. Tsai MC, Yu HC, Hung CH, Lee CM, Chiu KW, Lin MT, et al. Comparing the efficacy and clinical outcome of telbivudine and entecavir naive patients with hepatitis B virus-related compensated cirrhosis. *J Gastroenterol Hepatol* 2014;29:568-575.
 66. Koklu S, Tuna Y, Gulsen MT, Demir M, Koksak AS, Kockar MC, et al. Long-term efficacy and safety of lamivudine, entecavir, and tenofovir for treatment of hepatitis B virus-related cirrhosis. *Clin Gastroenterol Hepatol* 2013;11:88-94.
 67. Chan HL, Chan CK, Hui AJ, Chan S, Poordad F, Chang TT, et al. Effects of tenofovir disoproxil fumarate in hepatitis B e antigen-positive patients with normal levels of alanine aminotransferase and high levels of hepatitis B virus DNA. *Gastroenterology* 2014;146:1240-1248.
 68. Lu J, Zhang S, Liu Y, Du X, Ren S, Zhang H, et al. Effect of peginterferon alpha-2a combined with adefovir in HBV postpartum women with normal levels of ALT and high levels of HBV DNA. *Liver Int* 2015;35:1692-1699.
 69. Chung KT, Ha NB, Trinh HN, Garcia RT, Nguyen HA, Nguyen KK, et al. High frequency of recurrent viremia after hepatitis B e antigen seroconversion and consolidation therapy. *J Clin Gastroenterol* 2012;46:865-870.
 70. Fung J, Lai CL, Tanaka Y, Mizokami M, Yuen J, Wong DKH, et al. The duration of lamivudine therapy for chronic hepatitis B: cessation versus continuation of treatment after HBeAg seroconversion. *Am J Gastroenterol* 2009;104:1940-1947.
 71. Liaw Y-F, Sheen IS, Lee C-M, Akarca US, Papatheodoridis GV, Suet-Hing Wong F, et al. Tenofovir disoproxil fumarate (TDF), emtricitabine/TDF, and entecavir in patients with decompensated chronic hepatitis B liver disease. *HEPATOLOGY* 2011;53:62-72.
 72. Dogan UB, Kara B, Gumurdulu Y, Soyulu A, Akin MS. Comparison of the efficacy of tenofovir and entecavir for the treatment of nucleos(t)ide-naïve patients with chronic hepatitis B. *Turk J Gastroenterol* 2012;23:247-252.
 73. Batirel A, Guclu E, Arslan F, Kocak F, Karabay O, Ozer S, et al. Comparable efficacy of tenofovir versus entecavir and predictors of response in treatment-naïve patients with chronic hepatitis B: a multicenter real-life study. *Int J Infect Dis* 2014;28:153-159.
 74. Cholongitas E, Papatheodoridis GV, Goulis J, Vlachogiannakos J, Karatapanis S, Ketikoglou J, et al. The impact of newer nucleos(t)ide analogues on patients with hepatitis B decompensated cirrhosis. *Ann Gastroenterol* 2015;28:109-117.
 75. Huang M, Jie Y, Shi H, Li X, Wu Y, Lin G, et al. Comparison of the efficacy of tenofovir disoproxil fumarate and entecavir for initial treatment of patient with chronic hepatitis B in China. *Int J Clin Exp Med* 2015;8:666-673.
 76. Hung CH, Hu TH, Lu SN, Lee CM, Chen CH, Kee KM, et al. Tenofovir versus entecavir in the treatment of chronic hepatitis B with severe acute exacerbation. *Antimicrob Agents Chemother* 2015;59:3168-3173.
 77. Mallet V, Schwarzinger M, Vallet-Pichard A, Fontaine H, Corouge M, Sogni P, et al. Effect of nucleoside and nucleotide analogues on renal function in patients with chronic hepatitis B virus mono-infection. *Clin Gastroenterol Hepatol* 2014;13:1181-1188.
 78. Mauss S, Berger F, Filmann N, Hueppe D, Henke J, Hegener P, et al. Effect of HBV polymerase inhibitors on renal function in patients with chronic hepatitis B. *J Hepatol* 2011;55:1235-1240.
 79. Tien C, Xu JJ, Chan LS, Chang M, Lim C, Lee S, et al. Long-term treatment with tenofovir in Asian-American chronic hepatitis B patients is associated with abnormal renal phosphate handling. *Dig Dis Sci* 2015;60:566-572.
 80. Gish RG, Clark MD, Kane SD, Shaw RE, Mangahas MF, Baqai S. Similar risk of renal events among patients treated with tenofovir or entecavir for chronic hepatitis B. *Clin Gastroenterol Hepatol* 2012;10:941-946; e968.
 81. Sinn DH, Lee J, Goo J, Kim K, Gwak GY, Paik YH, et al. Hepatocellular carcinoma risk in chronic hepatitis B virus-infected compensated cirrhosis patients with low viral load. *HEPATOLOGY* 2015;62:694-701.
 82. Chen CJ, Yang HI, Su J, Jen CL, You SL, Lu SN, et al. Risk of hepatocellular carcinoma across a biological gradient of serum hepatitis B virus DNA level. *JAMA* 2006;295:65-73.
 83. Iloeje UH, Yang HI, Jen CL, Su J, Wang LY, You SL, et al. Risk and predictors of mortality associated with chronic hepatitis B infection. *Clin Gastroenterol Hepatol* 2007;5:921-931.
 84. Iloeje UH, Yang HI, Su J, Jen CL, You SL, Chen CJ. Predicting cirrhosis risk based on the level of circulating hepatitis B viral load. *Gastroenterology* 2006;130:678-686.
 85. Lok AS, McMahon BJ. Chronic hepatitis B: update 2009. *HEPATOLOGY* 2009;50:661-662.
 86. Weinbaum CM, Williams I, Mast EE, Wang SA, Finelli L, Wasley A, et al. Recommendations for identification and public health management of persons with chronic hepatitis B virus infection. *MMWR Recomm Rep* 2008;57:1-20.
 87. Mast EE, Weinbaum CM, Fiore AE, Alter MJ, Bell BP, Finelli L, et al. A comprehensive immunization strategy to eliminate transmission of hepatitis B virus infection in the United States: recommendations of the Advisory Committee on Immunization Practices (ACIP). Part II:

- Immunization of adults. *MMWR Recomm Rep* 2006;55:1-33; CE31-34.
88. World Health Organization. Guidelines for the prevention, care and treatment of persons with chronic hepatitis B infection. <http://www.who.int/hepatitis/publications/hepatitis-b-guidelines/en/>. Published March 2015. Accessed on October 23, 2015.
89. Terrault N, Bzowej N, Chang K-M, Hwang J, Jonas M, Murad H. AASLD guidelines for treatment of chronic hepatitis B. *HEPATOLOGY* 2016;63:261-283. <http://onlinelibrary.wiley.com/doi/10.1002/hep.28156/full>.

Author names in bold indicate shared co-first authorship.

Supporting Information

Additional Supporting Information may be found at <http://onlinelibrary.wiley.com/doi/10.1002/hep.28280/supinfo>.