

Article Type: Brief Communication

Subheading: Gynecology

BRIEF COMMUNICATION

Root causes and social consequences of birth injuries in Western Uganda

Jennifer N. Angell ^{1*}, Priscilla Busingye ², Gabriel Y-K. Ganyaglo ³, Cheryl A. Moyer ^{4,5}

¹ University of Michigan Medical School, Ann Arbor, Michigan, USA.

² Holy Family Virika Hospital, Department of Obstetrics and Gynecology, Fort Portal, Uganda.

³ Korle Bu Teaching Hospital, Department of Obstetrics and Gynecology, Accra, Ghana.

⁴ University of Michigan Medical School, Department of Learning Health Sciences, Ann Arbor, Michigan, USA.

⁵ University of Michigan Medical School, Department of Obstetrics and Gynecology, Ann Arbor Michigan, USA.

* Corresponding author: Jennifer N. Angell

335 Catherine St., Ann Arbor, MI, 48105, USA. Tel.: +1 248 882 4536.

E-mail address: Angellj@umich.edu

Keywords: Incontinence; Obstetric complications; Obstetric fistula; Qualitative research; Risk factors; Root cause; Social stigma; Uganda.

Synopsis: A case series of Ugandan women with fistula and other birth injuries suggested that the root causes included social, cultural, and system-based factors.

Incontinence-inducing delivery injuries, such as obstetric fistula and anal-sphincter tears, are devastating sequelae of prolonged and obstructed labor, and can only be

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/ijgo.12257](https://doi.org/10.1002/ijgo.12257)

This article is protected by copyright. All rights reserved

repaired through surgery. Such injuries are most prevalent in settings where delays and limitations in accessing emergency obstetric services are common [1, 2]. The 2011 Demographic and Health Survey of Uganda [3] estimated that 2% of Ugandan women have experienced obstetric fistula and resulting trans-vaginal urine or fecal incontinence.

Whereas the clinical causes of delivery injuries are well-understood, the contribution of social, cultural, and health-system factors have received limited attention. The present exploratory pilot study, approved by the institutional review board of the University of Michigan Medical School, Ann Arbor, Michigan, USA, sought to address this gap by employing in-depth case studies for 10 women in rural Western Uganda.

Virika Hospital, Fort Portal, Uganda conducts quarterly birth-injury camps for up to 50 women with a range of birth injuries. Virika Hospital provides surgical repair, recovery, and assisted community re-entry. All patients who presented to the May 2016 camp with vesicovaginal fistula, rectovaginal fistula, or fourth-degree anal-sphincter tears were invited to participate in an in-depth, semi-structured interview; written informed consent was obtained. The interviews assessed demographic data, obstetric history, rehabilitative services received, and the social impact of injuries. Data from each respondent were treated as individual case studies and the narrative portion of their interviews were used to trace potential root causes of injuries and to identify themes regarding social consequences.

The mean age of participants was 31 years (range 18–48) and the mean parity was 4 (range 1–10). Of the 10 women, four had been unmarried and developed injuries after their first pregnancies and five had been living with their injury for at least 5 years. The root-cause analysis suggested the following upstream causes of birth injuries: preference for traditional providers; lack of autonomy; lack of transportation; lack of clarity on presenting in labor; referral/access challenges; and lack of recognition or explicit discussion of injury (Table 1). There were 8 (80%) patients who reported their birth injuries as having a significant impact on day-to-day activities, including not being

allowed to prepare food, difficulty continuing to work outside the home, social isolation, limited sexual intercourse, and fear of partners leaving.

In Western Uganda, the high prevalence of obstetric fistula is entwined with many social, cultural, and system-based root causes. The present study is but one step toward a broader understanding of the factors contributing to birth injuries; further research is needed to effectively address these complex factors.

Author contributions

JNA contributed to the conception and design of the study, background literature review, data acquisition and analysis, and manuscript preparation. PB contributed to the design of the study, data acquisition, and manuscript preparation. GY-KG contributed to data interpretation and revising the manuscript. CAM contributed to the conception and design of the study, data analysis, and manuscript preparation and revision.

Acknowledgements

The present study received funding received from University of Michigan International Institute, and Phi Kappa Phi, University of Michigan Chapter.

Conflicts of interest

The authors have no conflicts of interest.

References

- [1] Landry E, Vera F, Ruminjo J, et al. Profiles and experiences of women undergoing genital fistula repair: Findings from five countries. *Global Public Health*. 2013;8(8):926-942. doi:10.1080/17441692.2013.824018.
- [2] Cowgill KD, Bishop J, Norgaard AK, Rubens CE, Gravett MG. Obstetric fistula in low-resource countries: an under-valued and under-studied problem – systematic review of its incidence, prevalence, and association with stillbirth. *BMC Pregnancy and Childbirth*. 2015;15:193. doi:10.1186/s12884-015-0592-2.

[3] Uganda Bureau of Statistics (UBOS) and ICF International Inc. 2012. Uganda Demographic and Health Survey 2011. Kampala, Uganda: UBOS and Calverton, Maryland: ICF International Inc.

Table 1 Root causes and social consequences of birth injuries.

Factors	Examples provided by participants
Upstream factors identified as contributing to birth injury	
Delivering with a traditional birth attendant	“It’s not necessary to deliver in a hospital.” Quote from a patient aged 30 y with a fourth-degree AST (TSI 11 y)
Necessary to wait for a husband’s permission/economic input	Experienced a 9-h delay in hospital-transfer owing to having to wait for husband to arrive with funds. Patient aged 42 y with a VVF (TSI 4 mo)
Lack of transportation to facility	Began labor at midnight but had to delay seeking care until the morning when motorbike taxis would be available. Patient aged 30 y with a fourth-degree AST (TSI: 12 y)
Chastised for arriving at facility early, leading to future delays	At prior delivery, patient told she was too early. Consequently, during her next delivery she labored at home for 12 h before departing for the hospital. She delivered on route with a traditional birth attendant who did not recognize/disclose the injury. Patient aged 23 y with a fourth-degree AST (TSI 9 mo)
Referral/access challenges	There was no physician available at the first facility they presented at; consequently, they sought care at a government hospital. Patient aged 28 y with a VVF (TSI 2 mo)
Lack of recognition/explicit discussion of injury	After a facility delivery, the provider did not discuss the tear; as it was the patient’s first delivery, she thought it was normal and would heal. Patient aged 35 y with a fourth-degree AST (TSI 17 y)
Social consequences of birth injury	
Not allowed to prepare food	Owing to being perceived as unclean, the patient’s children must prepare food, although she is allowed to eat with the family. Patient aged 28 y with a VVF (TSI 2 mo)
Interferes with work	“I do not bring produce to market anymore” [because of fear of leaking in public]. Quote from a patient aged 33 y with a VVF (TSI 5 y) When she bends over to pick something up in front of her students she cannot control passing gas, which embarrasses her greatly. Patient

Social isolation

aged 30 y with a fourth-degree tear (TSI 11 y)

At the market where she sells fruit, people point and laugh, making fun of her wet skirts and odor. Patient aged 48 y with a VVF (TSI 26 y)

Patient avoids visiting friends because she is afraid of people looking after her to see if her skirts are wet. Patient aged 33 y with VVF (TSI 5 y)

Patient avoids community gatherings because she fears leaking gas and/or stool. Patient aged 30 y with a fourth-degree AST (TSI 11 y)

Limited sexual intercourse

The patient's husband has stopped coming to her bed and she is worried that he will leave her if this continues. Patient aged 23 y with a fourth-degree AST (TSI 9 mo)

Fear of being left by partner

The patient's husband keeps another woman and has been sleeping at her home more and more since the injury. She is worried he will leave her if the fistula is not repaired. Patient aged 28 y with a VVF (TSI 2 mo)

Abbreviations: AST, anal-sphincter tear; TSI, time since injury; VVF, vesicovaginal fistula.

Author Manuscript