Table S3. The Relation between Blood Pressure in Pregnancy and Hypertensive Status at 7-15 Years Follow-up using Unweighted Data.

	Prehypertensive at Follow-Up (n=162)			Hypertensive at Follow-Up (n=139)		
	Model 1: Unadjusted OR (95% CI)	Model 2: Adjusted OR ^a (95% CI)	Model 3: Model 2 + BMI Adjusted OR ^{a,b} (95% CI)	Model 1: Unadjusted OR (95% CI)	Model 2: Adjusted OR ^a (95% CI)	Model 3: Model 2 + BMI Adjusted OR ^{a,b} (95% CI)
Section A: Blood pressure in pregnancy						
Normotensive (n=190)	1 [Referent]	1 [Referent]	1 [Referent]	1 [Referent]	1 [Referent]	1 [Referent]
Moderately-elevated BP (n=399)	1.0 (0.7, 1.5)	1.0 (0.6, 1.5)	0.9 (0.6, 1.4)	2.5 (1.5, 4.3)	2.8 (1.5, 5.1)	2.5 (1.4, 4.7)
Hypertensive disorder (HDP; n=78)	2.8 (1.5, 5.6)	3.0 (1.4, 6.2)	2.8 (1.3, 5.9)	10.1 (4.9, 20.9)	14.1 (6.1, 32.7)	13.3 (5.7, 31.2)
Section B: Blood pressure in pregnancy by type of elevation ^c						
Normotensive (n=190)	1 [Referent]	1 [Referent]	1 [Referent]	1 [Referent]	1 [Referent]	1 [Referent]
Moderately-elevated, systolic only (n=156)	0.9 (0.5, 1.5)	0.9 (0.5, 1.5)	0.8 (0.5, 1.4)	2.2 (1.2, 4.1)	2.5 (1.2, 4.9)	2.3 (1.1, 4.6)
Moderately-elevated, both systolic and diastolic (n=241)	1.1 (0.7, 1.7)	1.0 (0.6, 1.7)	0.9 (0.6, 1.5)	2.8 (1.6, 4.9)	3.1 (1.6, 5.9)	2.8 (1.4, 5.4)
HDP (n=78)	2.8 (1.5, 5.6)	3.0 (1.5, 6.3)	2.8 (1.3, 5.9)	10.1 (4.9, 20.9)	14.4 (6.2, 33.3)	13.5 (5.8, 31.7)
Section C: Blood pressure in pregnancy by timing of elevation						
Normotensive (n=190)	1 [Referent]	1 [Referent]	1 [Referent]	1 [Referent]	1 [Referent]	1 [Referent]
Moderately-elevated, early (<20 wk gestation; n=97)	1.1 (0.6, 2.1)	1.0 (0.5, 1.9)	0.9 (0.5, 1.8)	3.6 (1.8, 7.1)	3.4 (1.5, 7.4)	3.2 (1.4, 7.0)
Moderately-elevated, late (≥20 wk gestation; n=302)	1.0 (0.6, 1.5)	1.0 (0.6, 1.5)	0.9 (0.5, 1.4)	2.2 (1.3, 3.9)	2.6 (1.4, 4.9)	2.4 (1.3, 4.5)
HDP (n=78)	2.8 (1.5, 5.6)	3.0 (1.4, 6.3)	2.8 (1.3, 5.9)	10.1 (4.9, 20.9)	14.4 (6.2, 33.4)	13.6 (5.8, 32.0)

Abbreviations: BMI, body mass index; BP, blood pressure; CI, confidence interval; HDP, hypertensive disorder of pregnancy; OR, odds ratio; POUCH, Pregnancy Outcomes and Community Health.

^a Adjusted for race/ethnicity; pre-pregnancy BMI; age, parity, and Medicaid at pregnancy; and interval between birth and follow-up.

^b Adjusted for all covariates listed above plus BMI at follow-up.

^c N=665 for these models, which exclude those with moderately-elevated diastolic BP only (n=2).