

# Hospital Performance on Percutaneous Coronary Intervention Process and Outcomes Measures

Philip W. Chui, MD; Craig S. Parzynski, MS; Brahmajee K. Nallamothu, MD; Frederick A. Masoudi, MD; Harlan M. Krumholz, MD, SM; Jephtha P. Curtis, MD

**Background**—The Physician Consortium for Performance Improvement recently proposed percutaneous coronary intervention (PCI)-specific process measures. However, information about hospital performance on these measures and the association of PCI process and outcomes measures are not available.

**Methods and Results**—We linked the National Cardiovascular Data Registry (NCDR) CathPCI Registry with Medicare claims data to assess hospital performance on established PCI process measures (aspirin, thienopyridines, and statins on discharge; door-to-balloon time; and referral to cardiac rehabilitation), newly proposed PCI process measures (documentation of contrast dose, glomerular filtration rate, and PCI indication; appropriate indication for elective PCI; and use of embolic protection device), and a composite of all process measures. We calculated weighted pair-wise correlations between each set of process metrics and performed weighted correlation analyses to assess the association between composite measure performance with corresponding 30-day risk-standardized mortality and readmission rates. We reported the variance in risk-standardized 30-day outcome rates explained by process measures. We analyzed 1 268 860 PCIs from 1331 hospitals. For many process measures, median hospital performance exceeded 90%. We found strong correlations between medication-specific process measures ( $P < 0.01$ ) and weak correlations between hospital performance on the newly proposed and established process measures. The composite process measure explained only 1.3% and 2.0% of the observed variation in mortality and readmission rates, respectively.

**Conclusions**—Hospital performance on many PCI-specific process measures demonstrated little opportunity for improvement and explained only a small percentage of hospital variation in 30-day outcomes. Efforts to measure and improve hospital quality for PCI patients should focus on both process and outcome measures. (*J Am Heart Assoc.* 2017;6:e004276. DOI: 10.1161/JAHA.116.004276.)

**Key Words:** outcomes • percutaneous coronary interventions • process measures • readmissions

The past decade has seen a dramatic increase in efforts to measure and report the quality of care delivered to patients undergoing percutaneous coronary intervention (PCI).<sup>1,2</sup> A number of PCI-related process measures already exist,<sup>3</sup> and there are ongoing efforts both to expand the number of process measures and implement outcomes measures to characterize the quality of care for patients undergoing PCI. Specifically, the Physician Consortium for Performance Improvement (PCPI) in partnership with

professional societies including American College of Cardiology (ACC), American Heart Association, Society for Cardiovascular Angiography and Interventions (SCAI) have recently proposed a set of 11 PCI-related process measures.<sup>4</sup> In addition, the ACC has developed risk-standardized measures of hospital 30-day mortality and readmission following PCI.<sup>5,6</sup> However, we have little information as to how hospitals currently perform on these measures and whether process and outcomes measures capture distinct or overlapping

From the Department of Internal Medicine, University of California Irvine School of Medicine, Orange, CA (P.W.C.); Center for Outcomes Research and Evaluation, Yale-New Haven Hospital, New Haven, CT (C.S.P., H.M.K., J.P.C.); Center for Clinical Management Research, Ann Arbor VA Medical Center (B.K.N.) and Department of Internal Medicine (B.K.N.), University of Michigan Medical School, Ann Arbor, MI; Department of Medicine, University of Colorado Anschutz Medical Campus, Aurora, CO (F.A.M.); Department of Health Policy and Management, Yale School of Public Health, New Haven, CT (H.M.K.); Section of Cardiovascular Medicine, Department of Internal Medicine, Yale University School of Medicine, New Haven, CT (H.M.K., J.P.C.).

This manuscript was presented, in part, as 2 oral presentations at the American Heart Association Scientific Sessions, November 15–19, 2014 in Chicago, IL.

**Correspondence to:** Jephtha P. Curtis, MD, 1 Church St, Suite 200, New Haven, CT 06510. E-mail: jephtha.curtis@yale.edu

Received November 8, 2016; accepted December 28, 2016.

© 2017 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

domains of quality. Every performance measure carries an opportunity cost in terms of the resources needed to collect the data and the efforts required to improve performance.<sup>7–10</sup> Expanding the portfolio of measures may be warranted if a new measure provides a more-comprehensive assessment of hospital quality. However, the extent to which the new measures achieve this goal has not been demonstrated.

To date, no study has examined hospital performance on PCI process and outcomes measures. To address this gap in knowledge, we used data from the ACC's National Cardiovascular Data Registry (NCDR) CathPCI Registry to describe variation in hospital performance and examine the extent to which hospital performance on PCI-related measures are correlated. Specifically, we identified the association of hospital performance on existing process measures with the PCPI's proposed process measures. We then examined whether hospital quality, as determined by process measures performance, was correlated with hospital performance on 30-day mortality and readmission.

## Methods

### Data Sources

With more than 1600 participating hospitals, the NCDR CathPCI registry, cosponsored by the ACC and the SCAI, is the largest registry of elective and emergency PCIs in the United States.<sup>11</sup> The registry collects data on patient demographics, procedural and clinical variables, and in-hospital outcomes using standardized definitions.<sup>12</sup> For this study, we used registry data reflecting PCIs performed between January 1, 2010 and December 31, 2011. To calculate the risk-standardized 30-day mortality and readmission rates, we linked registry data with corresponding Medicare claims data using direct patient identifiers, including name, date of birth, and Social Security number. Information regarding 30-day patient readmissions and mortality were obtained using Medicare's Inpatient and Outpatient Standard Analytical Files and enrollment database. Among patients with more than 1 PCI performed during a hospitalization, we only included information from the initial procedure.

### Study Design and Population

We performed a cross-sectional analysis of all hospitals in the registry that reported at least 25 PCI procedures during the study time period. As a result, we excluded 122 low-volume hospitals, leaving a total of 1331 hospitals for analysis. To ensure that our estimates of hospital performance on specific measures were reliable, we further required that each hospital have at least 25 procedures for each individual measure.

Accordingly, the number of hospitals included in the calculation of each measure varies.

### Process Measures

We classified PCI process metrics into existing National Quality Forum (NQF)-endorsed PCI-related process measures and the newer set of measures proposed by the PCPI. The NQF-endorsed measures included the following: aspirin at discharge; thienopyridines at discharge; statins at discharge; door-to-balloon time under 90 minutes for patients presenting to the emergency department or under 120 minutes for patients transferred to a facility, and referral to cardiac rehabilitation after PCI. Among the 11 new process measures proposed by the PCPI in 2013, we were able to calculate the following using data elements currently collected by the registry: comprehensive documentation of criteria needed to determine procedural appropriateness; appropriate indications for elective PCI; use of embolic protection devices in saphenous vein bypass grafts; documentation of contrast dose; documentation of a preprocedural assessment of renal function (glomerular filtration rate [GFR] calculation); optimal postprocedural medical therapy (defined as aspirin, P2Y<sub>12</sub> inhibitors, and statins for all patients upon discharge unless otherwise contraindicated); and referral to cardiac rehabilitation. There were several measures proposed by the PCPI that could not be assessed given the available elements in the registry: documentation of radiation dose and whether or not a patient's ability to tolerate and adhere to dual antiplatelet therapy had been evaluated. Furthermore, we did not consider physician and hospital PCI volume as potential measures, given that all hospitals participating in the NCDR registry routinely receive information about procedural volumes.

For each measure, we identified whether patients were eligible for that metric and aggregated patient-level results to calculate the hospital performance in the indicated performance measure. Measure-specific inclusion and exclusion criteria were applied to each case to ensure that the population used to define performance was appropriate. For door-to-balloon times, we used different thresholds for ST-segment elevation myocardial infarction (STEMI) patients who presented through the emergency department ( $\leq 90$  minutes) and those transferred from another acute care facility ( $\leq 120$  minutes).<sup>13</sup> For the documentation of indications for PCI process measure, we classified a procedure as correctly documented based on recommendations from the PCPI, which included the following: documentation of priority of diagnoses (acute coronary syndrome vs elective); presence and severity of angina symptoms; use of antianginal medical therapies within 2 weeks before the procedure; significance of angiographic stenosis on coronary angiography for treated lesion; and presence, results, and timing of noninvasive stress test, fractional flow reserve, or

intravascular ultrasound. For the measure of the proportion of elective PCIs considered appropriate, we defined it in a manner consistent with the PCPI guidelines, which is the sum of the total number of appropriate and uncertain cases (as opposed to inappropriate or unmappable) among all non-acute coronary syndrome PCIs defined in a manner consistent with 2012 appropriate usage criteria.<sup>14</sup> Finally, we created composite measures defined as the total number of process measures patients received over the total number of eligible performance measures for patients treated at that hospital. The first composite measure was restricted to the current NQF-endorsed PCI process measures, and the second composite measure included both the current and proposed PCPI measures.<sup>10</sup>

## Outcome Measures

We calculated hospital-specific risk standardized 30-day readmission and mortality rates in a manner consistent with NQF-approved mortality and readmission measures.<sup>15,16</sup> Specifically, for mortality, and consistent with the NQF-approved measures, we used separate models to calculate hospitals' risk-standardized 30-day mortality for (1) patients with STEMI or cardiogenic shock and (2) patients without STEMI and without cardiogenic shock.<sup>16</sup> All outcomes models use hierarchical logistic regression, which takes into account clustering of patients within hospitals and use preprocedural clinical characteristics of patients for risk adjustment.

## Statistical Analysis

Crude rates, defined as the number of times a specific process measure was performed on a patient over the total number of eligible patients at that hospital, were calculated for all process measures. To analyze the association between hospital performance on current and emerging process metrics, we used the hospital performance estimates for each process measure to calculate a set of pair-wise correlations.

In our analysis of the relationship between hospital performance on current and emerging process measures with performance on outcome metrics, we used correlation analyses to determine the association of hospital risk-standardized 30-day mortality and readmission rates with corresponding hospital composite score estimates of both the emerging and existing process measures. We repeated our analyses, first limiting the composite measure to current process measures, and again limiting to medication-specific measures. In each analysis, we calculated both correlation coefficients and the proportion of the hospital-specific variation in risk-standardized outcomes explained by performance on the composite measures. This variation is the

square of the correlation coefficient and is calculated as a percentage. We performed secondary analyses to assess the robustness of our findings, restricting the calculation of hospital performance on process measures to Medicare beneficiaries.

Analyses were conducted using SAS software (version 9.3; SAS Institute Inc., Cary, NC). Given that different numbers of patients have eligibility for the process measures at each hospital, analyses were weighted by the total number of patients from that hospital who were included in the calculation of the specific process measure. For each correlation, we tested the null hypothesis that there is no correlation between measures, adjusting *P* values for multiple comparisons using the Sidak correction.<sup>17</sup> All analyses were conducted with an overall family-wise error rate of 0.05. The Yale University Human Investigations Committee approved analyses of this limited NCDR data set.


## Results

### Hospital Performance on Process and Outcomes Measures

A total of 1 268 860 PCI procedures performed at 1331 hospitals between January 2010 and December 2011 met criteria for inclusion. Hospital performance on each of the current process measures, proposed PCPI process measures, and the composite measure are shown in Figure 1. Among the current process measures, there was relatively little variation in hospital performance on the discharge medications, with median hospital performance above 90%. We observed a larger gap in performance with regard to timely primary PCI (median, 78.9%; interquartile range [IQR], 71.2–85.1%) and referral to cardiac rehabilitation (median, 60.7%; IQR, 18.4–87.2%). Among the PCPI proposed measures, hospital performance on measures of documentation was uniformly high: contrast dose (median, 99.7%; IQR, 98.3–100%); GFR (median, 97.0%; IQR, 92.8–98.8%); and PCI indications (median, 99.8%; IQR, 99.2–100%). Larger variation was observed for use of embolic protection devices (median, 16.5%; IQR, 3.6–33.9%) and the proportion of non-acute coronary syndrome PCIs considered appropriate or uncertain (median, 50.0%; IQR, 38.3–61.2%). Variation in hospital 30-day mortality and readmission rates was modest (Table 1).

### Correlation of Existing Process Measures

We found moderate-to-strong correlations (correlation coefficient,  $>0.40$ ;  $P<0.05$ ) between the discharge medication-related process measures, particularly between aspirin and thienopyridines at discharge (Table 2). None of the discharge medication-related process measures for medications were



**Figure 1.** Hospital performance on proposed PCI process measures. Hospital performance on many of the process measures, including appropriate medications at discharge as well as documentation of contrast dose and GFR documentation, was close to having process metric performance nearing 100%. Hospital performance on referral to cardiac rehabilitation, use of embolic devices, proportion of PCIs with documentation of PCI eligibility, and proportion of elective PCIs consider appropriate or uncertain exhibit room for improvement. Central band represents median, box hinges represent the first and the third quintiles, and whiskers extend to the 5% and 95% percentile. DTB indicates door-to-balloon; GFR, glomerular filtration rate; NQF, National Quality Forum; PCI, percutaneous coronary intervention; PCPI, Physician Consortium for Performance Improvement.

significantly correlated with the proportion of patients who received timely primary PCI. Referral to cardiac rehabilitation was significantly correlated with other existing process measures (all correlation coefficients, >0.10;  $P < 0.05$ ). Referral to cardiac rehabilitation was strongly correlated with the overall composite measure (correlation coefficient, >0.90;  $P < 0.01$ ) and was responsible for most of the variation observed in the composite measure.

### Correlation of Existing and Emerging Process Measures

There were no significant correlations found among the PCPI's proposed process measures (all correlation coefficients, <0.10; all  $P > 0.05$ ; Table 2). However, both the use of embolic

devices in saphenous vein grafts and the proportion of appropriate elective PCIs were significantly correlated with the discharge medication process measures. Furthermore, all of the existing and emerging process measures, with the exception of documentation of contrast dose, were significantly correlated with the overall composite measure consisting of both existing and emerging process metrics.

### Correlations of Process Measures With Hospital-Level Outcome Measures

The overall composite measure was statistically significantly associated with both risk-standardized readmission rate and risk-standardized mortality rate (RSMR) for no STEMI/no shock patients, but it was not correlated with RSMR for

**Table 1.** Hospital Performance on 30-Day Outcomes in 2010–2011\*

| Variable   | N    | 5th Percentile | 25th Percentile | Median | 75th Percentile | 95th Percentile |
|--|------|----------------|-----------------|--------|-----------------|-----------------|
| Risk-standardized readmission rate                       | 1076 | 10.0           | 11.1            | 11.8   | 12.7            | 14.2            |
| Risk-standardized mortality rate (STEMI or shock)        | 743  | 9.5            | 10.9            | 12.1   | 13.4            | 16.1            |
| Risk-standardized mortality rate (no STEMI and no shock) | 1059 | 1.3            | 1.6             | 1.7    | 2.0             | 2.6             |

NSTEMI indicates non-ST segment elevation myocardial infarction; STEMI, ST-elevation myocardial infarction.  
 \*Hospitals were only considered eligible for each measure if they had more than 25 patients.

**Table 2.** Correlation Coefficients for Hospital Performance on PCI Process Measures\*†

|  | Aspirin at Discharge | Thienopyridines at Discharge | Statin at Discharge | Proportion DTB Time ≤90 Minutes | Referral to Cardiac Rehab | Documentation of Contrast Dose | Proportion of PCIs With Embolic Devices | Proportion of PCIs With GFR Documentation | Proportion of PCIs With Comprehensive Documentations for PCI | Proportion of Appropriate Elective PCIs Performed |
|--|----------------------|------------------------------|---------------------|---------------------------------|---------------------------|--------------------------------|---|---|--|---|
| Thienopyridines at discharge                                       | 0.713‡               |                              |                     |                                 |                           |                                |   |   |  |   |
| Statin at discharge  | 0.597‡               | 0.486‡                       |                     |                                 |                           |                                |   |   |  |   |
| Proportion DTB time ≤90 and ≤120 minutes for transfers             | 0.063                | 0.070                        | 0.092               |                                 |                           |                                |   |   |  |   |
| Referral to cardiac rehab  | 0.194‡               | 0.139‡                       | 0.181‡              | 0.111‡                          |                           |                                |   |   |  |   |
| Documentation of contrast dose                                     | 0.033                | 0.023                        | 0.000               | -0.002                          | 0.019                     |                                |   |   |  |   |
| Use of embolic device  | 0.247‡               | 0.182‡                       | 0.215‡              | -0.043                          | 0.125‡                    | -0.046                         |   |   |  |   |
| Proportion of PCIs with GFR documentation                          | 0.084                | -0.005                       | 0.114               | 0.081                           | 0.016                     | 0.055                          | 0.041                                   |   |  |   |
| Proportion of PCIs with comprehensive documentations of indication | 0.037                | 0.115‡                       | 0.074               | 0.087                           | 0.082                     | 0.032                          | 0.098                                   | 0.030                                     |  |   |
| Proportion of elective PCIs considered appropriate or uncertain‡§  | 0.149‡               | 0.202‡                       | 0.125‡              | 0.079                           | 0.018                     | 0.031                          | -0.009                                  | 0.052                                     | -0.041   |   |
| Overall proportion of existing process measures met                | 0.377‡               | 0.303‡                       | 0.384‡              | 0.151‡                          | 0.972‡                    |                                |   |   |  |   |
| Overall proportion of existing and emerging process measures met   | 0.397‡               | 0.338‡                       | 0.423‡              | 0.180‡                          | 0.909‡                    | 0.044                          | 0.206‡                                  | 0.154‡                                    | 0.303‡   | 0.205‡  |

DTB indicates door-to-balloon; GFR, glomerular filtration rate; PCI, percutaneous coronary intervention.

\*Hospitals with considered eligible only if they had more than 25 patients for each of the individual process measures.

†Weighted by number of eligible patients in each hospital.

‡Data points indicate significance ( $P < 0.01$ ) after adjusting for multiple comparisons.

§Not included in overall composite measure.

**Table 3.** Correlation Coefficients for 30-Day Risk-Standardized Readmission Rates and Mortality Rates With Hospital Performance on PCI Process Measures\*†

|   | Aspirin at Discharge | Thienopyridines at Discharge | Statin at Discharge | Proportion DTB Time ≤90 Minutes | Referral to Cardiac Rehab | Documentation of Contrast Dose | Proportion of PCIs With Embolic Devices | Proportion of PCIs With GFR Documentation | Proportion of PCIs With Comprehensive Documentations for PCI | Proportion of Appropriate Elective PCIs Performed‡ | Overall Proportion of Existing Process Measures Met | Overall Proportion of Existing and Emerging Process Measures Met |
|---|----------------------|------------------------------|---------------------|---------------------------------|---------------------------|--------------------------------|---|---|--|--|---|--|
| Risk-standardized readmission rate        | -0.120 <sup>§</sup>  | -0.131 <sup>§</sup>          | -0.139 <sup>§</sup> | -0.062                          | -0.100                    | 0.007                          | -0.048                                  | -0.032                                    | -0.059   | -0.047   | -0.128 <sup>§</sup>                                 | -0.132 <sup>§</sup>  |
| Risk-standardized mortality rate (STEMI)  | -0.135 <sup>§</sup>  | -0.153 <sup>§</sup>          | -0.116              | -0.069                          | -0.074                    | -0.003                         | -0.013                                  | -0.098                                    | 0.022  | 0.036  | -0.103  | -0.103   |
| Risk-standardized mortality rate (NSTEMI) | -0.223 <sup>§</sup>  | -0.143 <sup>§</sup>          | -0.240 <sup>§</sup> | 0.026                           | -0.063                    | 0.009                          | -0.134 <sup>§</sup>                     | -0.009                                    | -0.035   | 0.008  | -0.115 <sup>§</sup>                                 | -0.122 <sup>§</sup>  |

DTB indicates door-to-balloon; GFR, glomerular filtration rate; NSTEMI, non-ST segment elevation myocardial infarction; PCI, percutaneous coronary intervention; STEMI, ST-elevation myocardial infarction.

\*Hospitals with considered eligible only if they had more than 25 patients for each of the individual process measures.

†Weighted by number of eligible patients in each hospital.

‡Not included in overall composite measure.

§Data points indicate significance ( $P < 0.01$ ) after adjusting for multiple comparisons.

STEMI/shock patients (Table 3). Prescriptions of aspirin and thienopyridines at discharge were significantly correlated with all 3 outcomes measures. The individual process metrics explained 0.1% to 1.9% of hospital variation in 30-day risk-standardized readmission rate, 0% to 2.3% in 30-day RSMR for STEMI/shock patients, and 0% to 5.8% in 30-day RSMR for NSTEMI/no shock patients (Table 4). Hospital performance on the overall composite measure explained relatively little of the observed variation in 30-day outcomes—ranging from 0.7% for RSMR in STEMI patients to 2.0% in risk-standardized readmission rate. Hospital performance on outcomes measures was similar when grouped by quintiles of performance on the composite process measure (Figure 2). Secondary analyses demonstrated similar findings when we restricted the calculation of hospital process measures to include only Medicare beneficiaries and when we restricted analyses to the current process measures and the discharge medication-specific process measures.

### Discussion

In this cross-sectional analysis of the NCDR CathPCI registry, we found that the majority of hospitals performed well on existing PCI-related process measures. The additional PCI process measures proposed by PCPI were not strongly correlated with the existing process measures, but even among these, there was little variation among hospitals and thus limited opportunity for improvement. In addition, hospitals' performance on existing and proposed process measures were only weakly correlated with hospitals' 30-day risk-standardized mortality and readmission rates. Although the associations were often statistically significant, hospital performance on PCI-specific process metrics explained only between 0.0% and 5.8% of observed variation in risk standardized mortality rates and between 0.0% and 2.0% of risk-standardized readmission rates. These findings suggest that process and outcome measures capture complementary, and not overlapping, domains of quality.

With the exception of the medication-specific process measures, there is relatively little correlation between existing and proposed PCI process measures. This finding is consistent with previous studies, suggesting that distinct strategies are needed to improve performance across different assessments of hospital quality.<sup>10,18–20</sup> However, our findings raise questions as to whether there is enough of a gap in current performance to justify further investment in the proposed PCPI metrics. We found that hospital performance on many of the proposed PCPI measures is generally high, with the large majority of hospitals successfully meeting these metrics more than 90% of the time. The measurement and reporting of process measures carry opportunity cost, and implementation of the proposed process measures may have a limited

**Table 4.** Percent Variance in Hospital-Level 30-Day Outcome Measures for PCI Process Measures\*

| Variable   | RSRR % | RSMR % (STEMI) | RSMR % (NSTEMI) |
|--|--------|----------------|-----------------|
| Aspirin at discharge   | 1.5    | 1.8            | 5.0             |
| Thienopyridines at discharge                                     | 1.7    | 2.3            | 2.0             |
| Statin at discharge  | 1.9    | 1.3            | 5.8             |
| Timely primary PCI   | 0.4    | 0.5            | 0.1             |
| Referral to cardiac rehab  | 1.0    | 0.6            | 0.4             |
| Documentation of contrast dose                                   | 0.0    | 0.0            | 0.0             |
| Use of embolic device  | 0.2    | 0.0            | 1.8             |
| Documentation of PCIs with GFR documentation                     | 0.1    | 1.0            | 0.0             |
| Documentation of PCI indications                                 | 0.4    | 0.0            | 0.1             |
| Proportion of appropriate elective PCIs performed                | 0.2    | 0.1            | 0.1             |
| Overall proportion of existing and emerging process measures met | 2.0    | 0.7            | 1.3             |

GFR indicates glomerular filtration rate; NSTEMI, non-ST segment elevation myocardial infarction; PCI, percutaneous coronary intervention; RSMR, risk-standardized mortality rate; RSRR, risk-standardized readmission rate; STEMI, ST-elevation myocardial infarction.

\*Hospitals were considered eligible only if they had more than 25 patients for each of the individual process measures.

potential to improve patient outcomes. Our findings highlight the difficulty of identifying novel process measures that identify practice variations that are associated with patient outcomes. In the absence of novel PCI-related process measures, it may be worth focusing quality measurement efforts on expanding the portfolio of outcomes measures, such as rates of bleeding, acute kidney injury, and patient-report health status following PCI.<sup>21–23</sup>


With an increasing number of performance metrics, NQF has recommended consideration of composite measures to provide a more-comprehensive picture of quality.<sup>24</sup> Several composite measures already exist in the NQF's portfolio of endorsed measures for other areas of focus, such as acute myocardial infarction and congestive heart failure.<sup>24,25</sup> Given the increased number of PCI-related process measures, there may be advantages to creating a composite measure.<sup>26</sup> Nevertheless, we found that much of the variation in the composite measures reflected variations in referral to cardiac rehabilitation, bringing into question the efficacy of a composite measure for PCI.

Our results indicate that there is variation in 30-day risk-standardized readmission and mortality rates across hospitals that perform PCI and demonstrate the continued opportunity

to improve the outcomes of patients undergoing these procedures. Although previous analyses have examined the association between acute myocardial infarction outcome metrics and process measures and others have identified specific clinical profiles and risk factors of PCI patients that predict outcomes, our study is the first to characterize the relatively weak relationship between PCI-specific process and PCI outcome measures.<sup>10,27,28</sup> Our findings are relevant given that there are ongoing efforts to increase quality of care specifically for PCI patients. Furthermore, our analysis of recent performance on the PCPI PCI process measures suggests that despite this effort, there is a continued need to identify impactful process measures and potentially to shift focus to outcomes measures to include not only mortality and readmissions, but also other relevant outcomes such as rates of acute kidney injury and bleeding.<sup>21,22</sup> The absence of a strong association does not necessarily indicate that there is no role for process measures in assessing the quality of care provided to PCI patients. Indeed, many of these measures have both face validity and substantial evidence supporting their impact on individual patients. Nevertheless, the fact that most process measures demonstrate little variation and are not strongly associated with outcomes suggest the need to identify additional care processes for which there is a sufficiently large gap in care to warrant their collection. Evidence-based processes, including the proposed PCPI measures, may be necessary, but not sufficient to drive improvements in the outcomes of PCI patients.

Our study highlights the complementary role of process and outcome measures in assessing hospital quality and illustrates that high performance alone on process measures does not guarantee optimal outcomes. In fact, one can argue that a hospital's ability to drive improvements in outcomes can be limited given the heterogeneity and number of factors influencing outcomes that are outside a hospital's control. On the other hand, there is growing evidence suggesting that there are implementable hospital strategies to improve quality of care that are associated with lower mortality and readmissions rates.

For example, previous studies have shown that improvements in hospital systems such as organizational culture, including interdisciplinary rounding during hospitalization and at discharge, and optimization of patient care transitions were associated with improved 30-day outcomes.<sup>29–31</sup> Similarly, other qualitative studies have shown that high-performing hospitals have specific organizational strategies and enabling structures that distinguish them, including: active communication and coordination among care givers; senior management-level engagement and support; and an organizational commitment to developing and maintaining a focus on delivering high-quality care.<sup>32,33</sup> These strategies emphasize that efforts to improve outcomes need to be multifaceted,



**Figure 2.** Risk-standardized outcomes based on performance on percutaneous coronary intervention (PCI) process metrics. Box-and-whiskers plot of hospital performance on 30-day risk-standardized readmissions and mortality rates in STEMI/cardiogenic shock (A) and non-STEMI/no cardiogenic shock (B) patients as stratified by quintiles of hospital performance on overall proportion of PCI process measures met. There is minimal variation in hospital performance on readmission (C) and mortality rates in relationship to respective quintiles of hospital performance on overall PCI process measures met. Central band represents median, box hinges represent the first and the third quintiles, and whiskers extend to 1.5 times the interquartile range. Diamonds represent the means and circles represent outlier hospitals. STEMI indicates ST-elevation myocardial infarction.

involving a level of complexity that may not be captured by well-described processes such as discharge medications. Developing effective ways to promote the uptake of these strategies and structures at PCI hospitals will be needed to improve the outcomes of PCI patients and reduce variation in PCI outcomes across hospitals.

There are several limitations to our study that warrant consideration. First, our patient population was derived from a single registry, and our results may not be generalizable to all PCI centers in the United States. However, the registry captures over 95% of PCI procedures performed in the United States, and we believe our data sample is likely representative of the US experience. In addition, given the data elements available in the registry, we were not able to characterize hospital performance on all of PCPI's proposed process measures, and it is possible

that the additional measures or a composite measure reflecting all existing and proposed PCI process measures would be more strongly associated with 30-day outcome measures. Furthermore, our outcomes data were restricted to Medicare beneficiaries and may not be representative of the US population overall.

In summary, hospital performance on current and emerging PCI-metrics only explain a small amount of the variation in 30-day risk-standardized mortality and readmission rates. This fact highlights that these 3 sets of markers are all capturing distinct aspects of hospital quality. Additional efforts are needed to better characterize how hospitals can utilize these distinct markers of quality to improve hospital performance.

The CathPCI Registry is an initiative of the American College of Cardiology with partnering support from The Society for Cardiovascular Angiography and Interventions.



## Sources of Funding

This work was supported by grant U01 HL105270-03 (Center for Cardiovascular Outcomes Research at Yale University) from the National Heart, Lung, and Blood Institute (Bethesda, MD). This research was also supported by the American College of Cardiology's National Cardiovascular Data Registry (NCDR). The views expressed in this article represent those of the authors and do not necessarily represent the official views of the NCDR or its associated professional societies identified at CVQuality.ACC.org/NCDR.

## Disclosures

Dr Masoudi has a contract with the American College of Cardiology for his role as Senior Medical Officer, NCDR. Dr Curtis and Mr Parzynski receive salary support from the American College of Cardiology, NCDR. Dr Krumholz is a recipient of research agreements from Medtronic and from Johnson & Johnson (Janssen), through Yale University, to develop methods of clinical trial data sharing; is the recipient of a grant from the Food and Drug Administration and Medtronic to develop methods for post-market surveillance of medical devices; chairs a cardiac scientific advisory board for UnitedHealth; is a participant/participant representative of the IBM Watson Health Life Sciences Board; is a member of the Advisory Board for Element Science; and is the founder of Hugo, a personal health information platform. Drs Krumholz and Curtis work under contract with the Centers for Medicare & Medicaid Services to develop and maintain performance measures that are used for public reporting. No other authors report disclosures.

## References

- Klein LW, Ho KKL, Singh M, Anderson HW, Hillegass WB, Uretsky BF, Chambers C, Rao SV, Reilly J, Weiner BH, Kern M, Bailey S. Quality assessment and improvement in interventional cardiology: a Position Statement of the Society of Cardiovascular Angiography and Interventions, Part II: public reporting and risk adjustment. *Catheter Cardiovasc Interv*. 2011;78:493–502.
- Joynt KE, Blumenthal DM, Orav J, Resnic FS, Jha AK. Association of public reporting for percutaneous coronary intervention with utilization and outcomes among Medicare beneficiaries with acute myocardial infarction. *JAMA*. 2012;308:1460–1468.
- Table of cardiovascular project: NQF-Endorsed maintenance standards under review. National Quality Forum. Available at: [http://www.qualityforum.org/Setting\\_Priorities/Improving\\_Healthcare\\_Quality.aspx](http://www.qualityforum.org/Setting_Priorities/Improving_Healthcare_Quality.aspx). Accessed September 16, 2014.
- Nallamothu BK, Tommaso CL, Anderson HV, Anderson JL, Cleveland JC, Dudley A, Duffy PL, Faxon DP, Gurm HS, Hamilton LA, Jensen NC, Josephson RA, Malenka DJ, Mani CV, McCabe KW, Mortimer JD, Patel MR, Persell SD, Rumsfeld JS, Shunk KA, Smith SC, Stanko SJ, Watts B. ACC/AHA/SCAI/AMA-Convended PCPI/NCOA 2013 performance measures for adults undergoing percutaneous coronary intervention: a report of the American College of Cardiology/American Heart Association task force on performance measures, the Society for Cardiovascular Angiography and Interventions, the American Medical Association-Convended Physician Consortium for Performance Improvement, and the National Committee for Quality Assurance. *J Am Coll Cardiol*. 2014;63:722–745.
- American College of Cardiology percutaneous coronary intervention (PCI) readmission measure. Medicare.gov: Hospital compare. Available at: <http://www.medicare.gov/hospitalcompare/PCIReadmission.html>. Accessed September 14, 2014.
- Krumholz HM, Keenan PS, Brush JE Jr, Bufalino VJ, Chernew ME, Epstein AJ, Heidenreich PA, Ho V, Masoudi FA, Matchar DB, Normand SL, Rumsfeld JS, Schuur JD, Smith SC Jr, Spertus JA, Walsh MN. Standards for measures used for public reporting of efficiency in health care: a scientific statement from the American Heart Association Interdisciplinary Council on Quality of Care and Outcomes Research and the American College of Cardiology Foundation. *J Am Coll Cardiol*. 2008;52:1518–1526.
- Hayward RA. Performance measurement in search of a path. *N Engl J Med*. 2014;356:951–953.
- Williams SC, Schmaltz SP, Morton DJ, Koss RG, Loeb JM. Quality of care in US hospitals as reflected by standardized measures, 2002–2004. *N Engl J Med*. 2005;353:255–264.
- Werner RM, Bradlow ET. Public reporting on hospital process improvements is linked to better patient outcomes. *Health Aff*. 2010;29:1319–1324.
- Bradley EH, Herrin J, Elbel B, McNamara RL, Magid DJ, Nallamothu BK, Wang Y, Normand ST, Spertus JA, Krumholz HM. Hospital quality for acute myocardial infarction: correlation among process measures and relationship with short-term mortality. *JAMA*. 2006;296:72–78.
- Cover story. CathPCI: ACC's Flagship Registry. National Cardiovascular Data Registry; 2013.
- Moussa I, Hermann A, Messenger JC, Dehmer GJ, Weaver WD, Rumsfeld JS, Masoudi FA. The NCDR CathPCI Registry: a US national perspective on care and outcomes for percutaneous coronary intervention. *Heart*. 2013;99:297–303.
- O'Gara PT, Kushner FG, Ascheim DD, Casey DE, Chung MK, de Lemos JA, Ettinger SM, Fang JC, Fesmire FM, Franklin BA, Granger CB, Krumholz HM, Linderbaum JA, Morrow DA, Newby LK, Ornato JP, Ou N, Radford MJ, Tamis-Holland JE, Tommaso CL, Tracy CM, Woo YJ, Zhao DX. 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction: a reporting of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2013;61:e78–e140.
- Patel MR, Dehmer GJ, Hirshfeld JW, Smith PK, Spertus JA. ACCF/SCAI/STS/AATS/AHA/ASNC/HFSA/SCCT 2012 appropriate use criteria for coronary revascularization focused update: a report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, Society for Cardiovascular Angiography and Interventions, Society of Thoracic Surgeons, American Association for Thoracic Surgery, American Heart Association, American Society of Nuclear Cardiology, and the Society of Cardiovascular Computed Tomography. *J Am Coll Cardiol*. 2012;59:857–881.
- Curtis JP, Drye EE, Duffy SC, Geary LL, Krumholz HM, Partovian SC, Wang Y. Hospital 30-day readmission following percutaneous coronary intervention measure: measure methodology report. Prepared for Centers for Medicare and Medicaid Services (CMS). New Haven, CT: Yale New Haven Health Services Corporation/Center for Outcomes Research and Evaluation; 2009.
- Curtis JP, Geary LL, Wang Y, Chen J, Drye EE, Grosso LM, Spertus JA, Rumsfeld JS, Weintraub WS, Masoudi FA, Brindis RG. Development of 2 registry-based risk models suitable for characterizing hospital performance on 30-day all-cause mortality rates among patients undergoing percutaneous coronary intervention. *Circ Cardiovasc Qual Outcomes*. 2012;5:628–637.
- Sidak Z. Rectangular confidence regions for the means of multivariate normal distributions. *J Am Stat Assoc*. 1967;62:626–633.
- Werner RM, Bradlow ET, Asch DA. Does hospital performance on process measures directly measure high quality care or is it a marker of unmeasured care? *Health Serv Res*. 2008;43:1464–1484.
- Werner RM, Bradlow ET. Relationship between Medicare's hospital compare performance measures and mortality rates. *JAMA*. 2006;296:2694–2702.
- Mant J. Process versus outcome indicators in the assessment of quality in health care. *Int J Qual Health Care*. 2001;13:475–480.
- Hess CN, Rao SV, McCoy LA, Neely ML, Singh M, Spertus JA, Krone RJ, Weaver WD, Peterson ED. Identification of hospital outliers in bleeding complications after percutaneous coronary intervention. *Circ Cardiovasc Qual Outcomes*. 2015;8:1–8.
- Tsai TT, Patel UD, Chang TI, Kennedy KF, Masoudi FA, Matheny ME, Kosiborod M, Amin AP, Messenger JC, Rumsfeld JS, Spertus JA. Contemporary incidence, predictors, and outcomes of acute kidney injury in patients undergoing percutaneous coronary interventions: insights from the NCDR Cath-PCI Registry. *JACC Cardiovasc Interv*. 2014;7:1–9.
- Pederson SS, Versteeg H, Denollet J, Cheng JM, Serruys PW, van Domburg RT. Patient-rated health status predicts prognosis following percutaneous coronary intervention with drug-eluting stent. *Qual Life Res*. 2011;20:559–567.
- National Quality Forum (NQF). *Composite Measure Evaluation Framework and National Voluntary Consensus Standards for Mortality and Safety—Composite Measures: A Consensus Report*. Washington, DC: NQF; 2009.

25. Agency for Healthcare Research and Quality (AHRQ). *AHRQ Quality Indicator: National Quality Forum (NQF) Endorsed Composite and Individual Measures*. Columbus, OH: AHRQ; 2011.
26. Peterson ED, DeLong ER, Masoudi FA, O'Brien SM, Peterson PN, Rumsfeld JS, Shahian DM, Shaw RE. ACCF/AHA 2010 position statement on composite measures for healthcare performance assessment: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Performance Measures (Writing Committee to Develop a Position Statement on Composite Measures). *J Am Coll Cardiol*. 2010; 55:1755–1766.
27. Peterson ED, Roe MT, Mulgund J, DeLong ER, Lytle BL, Brindis RG, Smith SC, Pollack CV, Newby LK, Harrington RA, Gibler WB. Association between hospital process performance and outcomes among patients with acute coronary syndromes. *JAMA*. 2006;295:1912–1920.
28. Yeh RW, Rosenfield K, Zelevinsky K, Mauri L, Sakhuja R, Shivapour DM, Lovett A, Weiner BH, Jacobs AK, Normand SL. Sources of hospital variation in short-term readmission rates after percutaneous coronary intervention. *Circ Cardiovasc Interv*. 2012;5:227–236.
29. Bradley EH, Curry L, Horwitz LI, Sipsma H, Wang Y, Walsh MN, Goldmann D, White N, Piña IL, Krumholz HM. Hospital strategies associated with 30-day readmission rates for patients with heart failure. *Circ Cardiovasc Qual Outcomes*. 2013;6:444–450.
30. Townsend-Gervis M, Cornell P, Vardaman JM. Interdisciplinary rounds and structured communication reduce readmissions and improve some patient outcomes. *West J Nurs Res*. 2014;36:917–928.
31. Kripalani S, Theobald CN, Anctil B, Vasilevskis EE. Reducing hospital admission: current strategies and future directions. *Annu Rev Med*. 2014; 65:471–485.
32. Curry LA, Spatz E, Cherlin E, Thompson JW, Berg D, Ting HH, Decker C, Krumholz HM, Bradley EH. What distinguishes top-performing hospitals in acute myocardial infarction mortality rates? A qualitative study. *Ann Intern Med*. 2011;154:384–390.
33. Curtis JP, Minges KE, Cherlin E, Elma MC, Bernheim SM, Messenger J, Ting HH, Berg D, Chen P. A qualitative study of the organizational strategies of high and low-performing PCI hospitals: insights from TOP PCI. Abstract: *Circulation*. 2013;128:A14758.