

Received Date: 16-Nov-2015

Revised Date: 21-Jan-2016

Accepted Date: 03-Feb-2016

Article Type: Original Article

Subject category: Resuscitation Science

Long-Term Outcomes Among Elderly Survivors of Out-of-Hospital Cardiac Arrest

Running Title: Out-of-Hospital Cardiac Arrest Outcomes in Survivors

Paul S. Chan, M.D., M.Sc., Bryan McNally, M.D., M.P.H., Brahmajee K. Nallamothu, M.D., M.P.H., Fengming Tang, B.S., Bradley G. Hammill, M.S., John A. Spertus, M.D., M.P.H., and Lesley H. Curtis, Ph.D.

From Saint Luke's Mid America Heart Institute, Kansas City, Missouri (P.S.C., F.T., J.A.S.) and the University of Missouri-Kansas City (P.S.C., J.A.S.); Department of Emergency Medicine, Emory University, and Rollins School of Public Health, Atlanta, GA (B.M.), the VA Health Services Research and Development Center of Excellence, VA Ann Arbor Healthcare System, Department of Internal Medicine and Center for Healthcare Outcomes and Policy, University of Michigan, Ann Arbor (B.K.N.); and the Duke Clinical Research Institute (B.G.H., L.H.C.) and Department of Medicine (L.H.C.), Duke University School of Medicine, Durham, North Carolina.

Address correspondence to Dr. Chan at the Mid America Heart Institute, 5th Floor, 4401 Wornall Rd., Kansas City, MO 64111, or at pchan@saint-lukes.org.

Subject Terms: Quality and Outcomes, Mortality/Survival, Cardiopulmonary Arrest

Manuscript Text Word Count: 3292 words

Abstract

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1177/1077551215581396](https://doi.org/10.1177/1077551215581396)

This article is protected by copyright. All rights reserved

Background: Most studies on out-of-hospital cardiac arrest have focused on immediate survival. However, little is known about long-term outcomes and resource use among survivors.

Methods and Results: Within the national CARES registry, we identified 16,206 adults 65 years or older with an out-of-hospital cardiac arrest between 2005 and 2010. Among 1127 patients who were discharged alive, we evaluated whether 1-year mortality, cumulative readmission incidence, and follow-up inpatient costs differed according to patients' race, sex, initial cardiac arrest rhythm, bystander delivery of cardiopulmonary resuscitation (CPR), discharge neurological status, and functional status (hospital discharge disposition). Overall 1-year mortality after hospital discharge was 31.8%. Among survivors, there were no long-term mortality differences by sex, race, or initial cardiac arrest rhythm, but worse functional status and severe neurological disability at discharge were associated with higher mortality. Moreover, compared with first responders, CPR delivered by bystanders was associated with 23% lower mortality (HR, 0.77 [0.58-1.02]). Besides mortality, 638 (56.6%) patients were readmitted within the first year, and the cumulative readmission incidence was 197 per 100 patient-years. Mean 1-year inpatient costs were \$23,765±\$41,002. Younger age, black race, severe neurological disability at discharge, and hospital disposition to a skilled nursing or rehabilitation facility, were each associated with higher 1-year inpatient costs (*P* for all <0.05).

Conclusion: Among elderly survivors of out-of-hospital cardiac arrest, nearly 1 in 3 patients die within the first year. Long-term mortality and inpatient costs differed substantially by certain demographic factors, whether CPR was initiated by a bystander, discharge neurological status, and hospital disposition.

Key Words: cardiac arrest; outcomes research; survival; cost

Although there are an estimated 350,000 out-of-hospital cardiac arrests annually in the U.S.,¹ little is known about long-term outcomes among those surviving to hospital discharge. This is because most prior studies of out-of-hospital cardiac arrest have focused on pre-hospital and in-hospital survival. Furthermore, the few studies which have examined long-term outcomes²⁻⁶ have primarily examined mortality, been restricted to cardiac arrests due to ventricular fibrillation, were conducted more than a decade ago, or typically involved a single region or hospital center, many with small sample sizes. As a result, prior studies among survivors may have limited generalizability and have not examined resource utilization.

Besides the need to quantify long-term mortality and costs among survivors of out-of-hospital cardiac arrest, it would be important to examine these outcomes in specific patient subgroups. For instance, although prior out-of-hospital cardiac arrest studies have found racial and sex differences for in-hospital survival,^{7,8} whether long-term mortality and costs also differ by race and sex among survivors is unknown. Although initiation of cardiopulmonary resuscitation (CPR) by a bystander has been linked to higher rates of in-hospital survival,⁹ determining whether it is associated with lower or higher mortality and costs among survivors has potential implications for current public campaigns to increase bystander CPR rates and life-years saved.¹⁰ Finally, as many survivors of out-of-hospital cardiac arrest have neurological and functional disability, defining long-term survival and costs by neurological disability and hospital disposition at discharge would better enable patients and physicians to use this prognostic information for shared decision-making.

Given these gaps in knowledge, we linked data from a large, national out-of-hospital cardiac arrest registry with Medicare claims files and examined long-term mortality, readmission incidence, and cumulative inpatient costs at 1 year among survivors who were discharged after an out-of-hospital cardiac arrest. We examined rates of these outcomes overall, as well as by race, sex, initial cardiac arrest rhythm, bystander administration of CPR, discharge neurological status, and hospital disposition.

Methods

Data Sources and Linkage

CARES is a large, prospective clinical registry of patients with out-of-hospital cardiac arrest in the U.S. Established in October of 2005 by the Centers for Disease Control and Emory University for public health surveillance and continuous quality improvement, the design of the registry has been previously described in detail.^{11,12} Briefly, all patients with a confirmed out-of-hospital cardiac arrest (defined as apnea, pulselessness and unresponsiveness for which cardiopulmonary resuscitation was initiated) of presumed cardiac etiology and for whom resuscitation is attempted are identified and followed, including those with termination of resuscitation prior to hospital arrival. Data are collected from three sources that together define the continuum of emergency cardiac care: 911 dispatch centers, EMS agencies, and receiving hospitals. Standardized international Utstein definitions for defining clinical variables and

outcomes are used to ensure uniformity.¹³ The completeness of data submitted to CARES is confirmed during routine data audits, wherein the number of cardiac arrest cases reported to CARES by each participating EMS agency is compared with the number of cardiac arrest cases in the agency's medical records. Finally, a CARES analyst reviews every record for completeness and accuracy.¹²

Based on prior work linking registries with Medicare files,^{14, 15} we linked CARES patient-level data from October, 1, 2005, through December 31, 2010, with Medicare inpatient files using 5 identifiers: dates of hospital admission, patient age and sex, admitting hospital (deidentified), and *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)* diagnosis and procedure codes. We selected Medicare records for the linkage if they included a primary or secondary diagnosis code for cardiac arrest (427.5), ventricular fibrillation (427.41), or ventricular flutter (427.42) or a procedure code for cardiopulmonary resuscitation (99.60), defibrillation (99.62), or closed chest massage (99.63). For each linked patient, we obtained Medicare denominator and inpatient files from 2005 through 2010.

Study Population

During the study period, a total of 31,881 patients 18 years or older with an out-of-hospital cardiac arrest not occurring in the presence of EMS personnel were enrolled within CARES (**Figure 1**). We excluded 15,673 patients younger than 65 years who would not be eligible for a match to Medicare files, leaving 16,208 Medicare age-eligible patients. Of these, 3931 survived to hospital admission (4014 had terminated resuscitations outside the hospital and 8263 in a hospital emergency room). Using the method described above, we linked 2206 (56.1%) of these hospitalized patients to Medicare claims data. The reasons for non-linkage to Medicare data occurred when a patient 1) was admitted to a non-Medicare hospital (e.g., Veterans Administration hospital), 2) had insurance other than fee-for-service Medicare, 3) was admitted to a hospital with few registry patients (thus precluding a unique match), or 4) lacked a qualifying *ICD-9-CM* diagnosis or procedure code for cardiac arrest in the Medicare files. Patients who were and were not linked to Medicare files had similar demographic and clinical characteristics (**Table S1**). Finally, as we were interested in examining outcomes among survivors, we excluded 1079 patients who died during the index hospitalization for their cardiac

arrest. The final study cohort comprised 1127 patients who survived to hospital discharge after an out-of-hospital cardiac arrest.

Study Outcomes

The outcomes of interest were long-term mortality, cumulative readmission rate, and follow-up inpatient costs at 1 year after discharge from the index hospitalization. Information on vital status was obtained from the Medicare denominator files and on readmissions and inpatient costs from the Medicare inpatient files.

Statistical Analysis

Baseline characteristics of the study cohort were described using proportions for categorical variables and means with standard deviations for continuous variables. We constructed survival curves using Kaplan-Meier estimates to determine unadjusted rates of mortality. We also computed cumulative readmission incidence rates at 1 year of follow-up. From these rates, the mean number of readmissions per patient-year of follow-up was determined.

Multivariable Cox proportional hazard models were constructed to determine predictors of 1-year mortality. All models were adjusted for age (65 to 74 years, 75 to 84 years, and ≥ 85 years), sex, race (white, black, and other), initial cardiac arrest rhythm, (ventricular fibrillation, pulseless ventricular tachycardia, asystole, pulseless electrical activity), location of arrest (private residence, public outdoor area, outpatient healthcare facility, and other), whether the arrest was witnessed, who initiated CPR (first responder [police, fire department staff], bystander, or EMS personnel), who first applied an automated external defibrillator, hospital disposition at discharge (home self-care, home with home health care, skilled nursing facility, inpatient rehabilitation care, hospice, and other) and discharge neurological status. The latter was assessed using the Cerebral Performance Category (CPC) scale, which classifies patients as having mild to no neurological disability, moderate disability, severe disability, or coma or vegetative state.¹⁶ Finally, we adjusted for the primary reason for the initial hospitalization (cardiac arrest, other cardiac, pulmonary, and other), which we determined from *ICD-9-CM* codes for the principal discharge diagnosis in the Medicare inpatient files.

From the model, we derived risk-adjusted hazard ratios for long-term mortality for the following prespecified subgroups: sex, race, initial cardiac arrest rhythm, who initiated CPR, discharge neurological status, and hospital disposition at discharge. In addition to looking at 1-year mortality, we examined whether subgroup differences persisted beyond 1 year and constructed similar models for 3-year mortality. Lastly, to place our findings in proper context, we compared mortality rates using Cox models between our cohort and two different Medicare cohorts who survived to discharge matched by age, sex, admitting hospital, and hospitalization year. The first was for any hospitalized Medicare patient whereas the second was for Medicare patients who required mechanical ventilation (*ICD-9-CM* procedure codes 96.70, 96.71 and 96.72) during the index hospitalization.

To examine inpatient resource use, we first identified each rehospitalization in the cohort from the linked Medicare inpatient files. Costs for each patient were determined by summing costs for each readmission. We then computed adjusted 1-year cost ratios for the aforementioned pre-specified subgroups. To accomplish this, since some patients had no follow-up inpatient costs, we constructed a two-part model conditional on patients having follow-up inpatient costs, comprised of (1) a logistic regression model predicting the probability of having any follow-up costs,¹⁷ and (2) a gamma regression model with a log link for the costs (for those patients with non-zero follow-up costs),¹⁸ with this model adjusted for the same variables as described for the model for mortality above. From the model, we calculated adjusted costs for each reference group (e.g., men) by performing 1000 bootstrap samples and computing the mean over these 1000 samples. Adjusted cost ratios and 95% confidence intervals for the comparator strata in each subgroup (e.g., women) were then derived by performing 1000 bootstrap samples, with the 2.5th and 97.5th percentile cost ratios defined as the 95% CI.¹⁹

Overall, rates of missing data were low, with a missing data rate of ~7% for both race and discharge neurological status. Patients with missing information on these variables were categorized as ‘unknown’ as a separate dummy variable in our models. For each analysis, we evaluated the null hypothesis at a 2-sided significance level of 0.05 and calculated 95% confidence intervals (CIs) using robust standard errors. All analyses were performed using SAS version 9.2 (SAS Institute, Cary, North Carolina) and R version 2.10.0 (R Foundation for Statistical Computing, Vienna, Austria).²⁰ The institutional review boards of the Duke University Health System and the Mid America Heart Institute approved the study, and the

requirement for informed consent was waived by the institutional review board from the Mid America Heart Institute.

Results

Of 1127 survivors of out-of-hospital cardiac arrest, 58.3% were men and 19.5% were of black race; mean age was 75.4 ± 7.5 years (**Table 1**). Nearly half (46.3%) of survivors had a cardiac arrest due to a shockable cardiac arrest rhythm of ventricular fibrillation or pulseless ventricular tachycardia. Three in five patients were at home at the time of cardiac arrest, 12% occurred in a nursing home facility, and 14.3% in a public area. Although 70.6% of survivors had an out-of-hospital cardiac arrest that was witnessed, cardiopulmonary resuscitation was begun by a non-medical bystander in only 34.3% of instances and bystander deployment of an automated external defibrillator was uncommon (6.2%). At hospital discharge, more than half (52.6%) were discharged home (most without requirement for home health care), 38.3% transitioned to an inpatient skilled nursing or rehabilitation facility, 8.0% went to hospice or another facility, and 1.2% to other non-home facilities.

Mortality

Overall mortality after hospital discharge was initially steep (12.7% at 30 days) and then rose more gradually, with a mortality rate of 31.8% at 1 year and 47.2% at 3 years (**Figure S1**). Among the prespecified subgroups, there were no differences in long-term mortality by race, sex, or initial cardiac arrest rhythm (**Table 2**). However, compared with a 1-year mortality rate of 11.8% among those discharged home without requirement of home health services, the risk of dying during the first year was 85% higher among those discharged home with a need for home health care and more than 4-fold higher among those requiring additional inpatient care at a skilled nursing or rehabilitation facility. Moreover, compared with a 1-year mortality rate of 21.4% among those with mild to no neurological disability at discharge, those with severe neurological disability had a two-fold increase in long-term mortality, even after adjusting for hospital disposition. Finally, compared with when CPR was initiated by first responders, CPR

initiated by bystanders (Hazard Ratio [HR], 0.77 [95% CI: 0.58-1.02]) and EMS personnel (HR, 0.57 [95% CI, 0.43-0.77]; P across groups <0.001) was associated with lower long-term mortality. Mortality results were similar when we repeated the analyses to examine predictors of 3-year mortality (**Table S2**). For both 1-year and 3-year mortality, there were no significant interactions between any of the subgroups of interest by age group or by sex.

When compared to a matched Medicare cohort of patients hospitalized for any reason, survivors of out-of-hospital cardiac arrest had higher 1-year (31.8% vs. 20.4%) and 3-year mortality (47.2% vs. 37.6%; HR 1.55 [95% CI: 1.35-1.77]; $P<0.001$) (**Figure 2**). In contrast, when compared to a matched Medicare cohort of patients who required mechanical ventilation for other reasons, survivors of out-of-hospital cardiac arrest had lower 1-year (31.8% vs. 45.6%) and 3-year mortality rates (47.2% vs. 60.7%; HR 0.68 [95% CI: 0.60-0.77]; $P<0.001$).

Readmission and Costs

Although 31.8% of patients died within the first year after discharge, there were a total of 2015 readmissions, yielding a 1-year cumulative incidence rate of 197 readmissions per 100 patient years (95% CI, 181 to 214). Notably, 638 (56.6%) patients were readmitted during the first year, and 279 (24.8%) were readmitted three or more times (**Figure 3**).

During the first year, the mean and median 1-year cost for readmissions for the whole cohort (including those who were not admitted) were \$23,765 \pm \$41,002 and \$7054 (interquartile range: \$0-\$30,751), respectively. There were no differences in inpatient costs by sex, initial cardiac arrest rhythm, or by who initiated CPR (**Table 3**). However, compared with whites (1-year mean inpatient costs of \$20,299), black patients incurred nearly twice that rate for inpatient costs during the first year (adjusted cost ratio, 1.95; 95% CI, 1.55-2.46; $P<0.001$). For patients discharged home, readmission costs for those discharged to an inpatient skilled nursing or rehabilitation care facility were more than 2-fold higher. Finally, patients with moderate to severe neurological disability at discharge had significantly higher 1-year inpatient costs.

Discussion

Among patients 65 years or older who survived an out-of-hospital cardiac arrest, 1-year mortality was 32%, with the most vulnerable period during the first month after discharge, wherein 40% of deaths during the first year occurred. Despite this, long-term mortality for

survivors of out-of-hospital cardiac arrest was substantially lower than for patients who required mechanical ventilation for other reasons prior to discharge. Although there were no racial or sex differences in long-term mortality, mortality differed markedly by neurological status at discharge and hospital disposition. Moreover, CPR initiated by bystanders was associated with lower mortality, and the mortality differences among each of these subgroups persisted at 3 years. Readmission during the first year was common, with mean 1-year inpatient costs of ~\$23,000, and costs differed by race, hospital discharge disposition, and neurological status at discharge. Collectively, our findings highlight that survivors of out-of-hospital cardiac arrest have significant mortality and morbidity risks after hospital discharge and these risks differed based on certain demographic factors, whether CPR was initiated by a bystander, and neurological and functional status at discharge.

Until recently, there has been limited information on the long-term outcomes of survivors of out-of-hospital cardiac arrest. In general, prior studies have been largely restricted to patients with ventricular fibrillation or from communities with highly organized EMS systems. One study from Olmsted County of 79 cardiac arrest survivors with ventricular fibrillation found that the expected 5-year mortality rate was 21%.³ Another study from Seattle—a metropolitan region with high rates of bystander CPR and a well-organized EMS system—reported a 1-year mortality rate of 18%.⁴ A third retrospective study from the Netherlands two decades ago reported a 1-year mortality rate of only 12% among 441 survivors of out-of-hospital cardiac arrest⁵ while a more recent study of 95 cardiac arrest survivors from Copenhagen found a 1-year mortality rate of 13%.⁶ A fifth study exists for 61 cardiac arrest survivors from over 3 decades ago.² Our study was able to build upon this prior literature by examining outcomes across multiple communities throughout the U.S, many of which do not have EMS systems as robust as those in Seattle and Denmark. We evaluated not only mortality but also rates of readmission and inpatient resource use—which, to our knowledge, has not been previously described—and we were able to evaluate predictors of these outcomes given our sample size.

We observed several predictors of 1-year mortality and costs. Although men and patients of black race with out-of-hospital cardiac arrest are known to have higher rates of in-hospital mortality than women and whites, respectively^{7,8}, we found no differences by race or sex in long-term mortality among cardiac arrest survivors, although black survivors had higher inpatient costs than white survivors. Patients with severe neurological disability had both higher

mortality (confirming the results of a recent study²¹) and inpatient costs at 1 year. This suggests that renewed efforts are needed to treat both the heart and the brain during acute resuscitation care, as significant neurological disability in a cardiac arrest survivor is not only associated with lower quality of life but also substantially higher post-discharge morbidity and mortality.

A strong and powerful predictor of long-term mortality was functional status, as assessed by hospital disposition in our study. Patients who were able to be discharged independently to their homes had a mortality rate during the first year of only 11.8%, or half the cohort average, while those discharged home with a need for home health services had nearly doubled this mortality rate. Most notably, patients who required further inpatient care at a skilled nursing facility or rehabilitation site had a mortality rate was more than 4-fold higher than patients discharged independently to their homes.

Finally, we found that cardiac arrest survivors in whom CPR was initiated by a bystander or EMS personnel had lower mortality compared with those in whom CPR was initiated by first responders from police and fire departments. Although we did not have information on time to CPR (as many cases of cardiac arrest are unwitnessed), cardiac arrest patients treated by bystanders likely have shorter time intervals between the onset of cardiopulmonary arrest and when CPR is begun as compared with first responders. As a result, cardiac arrest survivors in whom CPR was initiated by a bystander are likely to have less neurological and functional disability at hospital discharge than those in whom CPR was not initiated by a bystander. Although the link between bystander CPR and lower long-term mortality among cardiac arrest survivors would seem to be intuitive, this association has been reported in only one prior study²² and supports ongoing efforts to broadly disseminate CPR instruction to improve rates of bystander CPR for out-of-hospital cardiac arrest.

Our study has some limitations. First, CARES is a quality-improvement registry. Although it collects data from a diverse group of EMS agencies, our study's overall rates for long-term mortality and inpatient costs in nonparticipating communities may differ, although we have no reason to believe that our subgroup analyses of predictors of these outcomes would be different. Second, we restricted the analysis to Medicare beneficiaries; outcomes of patients younger than 65 years may differ. Third, we excluded patients for whom a CARES record could not be linked to a Medicare hospitalization. This occurred because a patient was admitted to a federal hospital, did not have fee-for-service Medicare insurance, or lacked a qualifying ICD-9

diagnosis, or because there were too few cases admitted to a hospital to ensure a unique Medicare match. Nonetheless, excluded patients were similar to patients in the study cohort; therefore, their exclusion was unlikely to significantly bias the results. Third, the CARES registry only recently began collecting data on use of coronary angiography and targeted temperature management on out-of-hospital cardiac arrest patients who survived to hospital admission. These post-resuscitation factors may play a role in long-term outcomes among survivors, but we were unable to examine their impact as their inclusion into CARES occurred during the last year of this study sample. Finally, we did not have access to serial assessments of neurological status or quality of life after discharge to allow for a more refined understanding of the trajectory of health status among survivors, nor did we have information about cause of death.

In conclusion, we found that, among elderly survivors of out-of-hospital cardiac arrest, nearly 1 in 3 patients die within the first year and readmissions were common. There was no evidence for racial or sex disparities in survival, but long-term mortality differed by whether CPR was initiated by a bystander, patients' neurological status at discharge, and hospital disposition.

Funding Sources:

- Dr. Chan is supported by funding (K23HL102224 and R01HL123980) from the National Heart Lung and Blood Institute.
- CARES was funded by the Centers for Disease Control and Prevention from 2004-2012. The program is now supported through private funding from the American Red Cross, the Medtronic Foundation Heart Rescue Program, the American Heart Association, Zoll Corporation, and in-kind support from Emory University.

Disclosures:

None.

REFERENCES

1. Go AS, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, Borden WB, Bravata DM, Dai S, Ford ES, Fox CS, Franco S, Fullerton HJ, Gillespie C, Hailpern SM, Heit JA, Howard VJ, Huffman MD, Kissela BM, Kittner SJ, Lackland DT, Lichtman JH, Lisabeth LD, Magid D, Marcus GM, Marelli A, Matchar DB, McGuire DK, Mohler ER, Moy CS, Mussolino ME, Nichol G, Paynter NP, Schreiner PJ, Sorlie PD, Stein J, Turan TN, Virani SS, Wong ND, Woo D, Turner MB. Heart disease and stroke statistics--2013 update: a report from the American Heart Association. *Circulation*. 2013;127:e6-e245.
2. Myerburg RJ, Kessler KM, Estes D, Conde CA, Luceri RM, Zaman L, Kozlovskis PL, Castellanos A. Long-term survival after prehospital cardiac arrest: analysis of outcome during an 8 year study. *Circulation*. 1984;70:538-546.
3. Bunch TJ, White RD, Gersh BJ, Meverden RA, Hodge DO, Ballman KV, Hammill SC, Shen WK, Packer DL. Long-term outcomes of out-of-hospital cardiac arrest after successful early defibrillation. *N Engl J Med*. 2003;348:2626-2633.
4. Dumas F, Rea TD. Long-term prognosis following resuscitation from out-of-hospital cardiac arrest: role of aetiology and presenting arrest rhythm. *Resuscitation*. 2012;83:1001-1005.
5. Kuilman M, Bleeker JK, Hartman JA, Simoons ML. Long-term survival after out-of-hospital cardiac arrest: an 8-year follow-up. *Resuscitation*. 1999;41:25-31.
6. Holler NG, Mantoni T, Nielsen SL, Lippert F, Rasmussen LS. Long-term survival after out-of-hospital cardiac arrest. *Resuscitation*. 2007;75:23-28.
7. Johnson MA, Haukoos JS, Larabee TM, Daugherty S, Chan PS, McNally B, Sasson C. Females of childbearing age have a survival benefit after out-of-hospital cardiac arrest. *Resuscitation*. 2013;84:639-44.
8. Cowie MR, Fahrenbruch CE, Cobb LA, Hallstrom AP. Out-of-hospital cardiac arrest: racial differences in outcome in Seattle. *Am J Public Health*. 1993;83:955-959.
9. Sasson C, Rogers MA, Dahl J, Kellermann AL. Predictors of survival from out-of-hospital cardiac arrest: a systematic review and meta-analysis. *Circ Cardiovasc Qual Outcomes*. 2010;3:63-81.

10. Lindner T, Vossius C, Mathiesen WT, Soreide E. Life years saved, standardised mortality rates and causes of death after hospital discharge in out-of-hospital cardiac arrest survivors. *Resuscitation*. 2014;85:671-675.
11. McNally B, Stokes A, Crouch A, Kellermann AL. CARES: Cardiac Arrest Registry to Enhance Survival. *Ann Emerg Med*. 2009;54:674-683 e672.
12. McNally B, Robb R, Mehta M, Vellano K, Valderrama AL, Yoon PW, Sasson C, Crouch A, Perez AB, Merritt R, Kellermann A. Out-of-hospital cardiac arrest surveillance --- Cardiac Arrest Registry to Enhance Survival (CARES), United States, October 1, 2005--December 31, 2010. *MMWR Surveill Summ*. 2011;60:1-19.
13. Jacobs I, Nadkarni V, Bahr J, Berg RA, Billi JE, Bossaert L, Cassan P, Coovadia A, D'Este K, Finn J, Halperin H, Handley A, Herlitz J, Hickey R, Idris A, Kloeck W, Larkin GL, Mancini ME, Mason P, Mears G, Monsieurs K, Montgomery W, Morley P, Nichol G, Nolan J, Okada K, Perlman J, Shuster M, Steen PA, Sterz F, Tibballs J, Timerman S, Truitt T, Zideman D. Cardiac arrest and cardiopulmonary resuscitation outcome reports: update and simplification of the Utstein templates for resuscitation registries: a statement for healthcare professionals from a task force of the International Liaison Committee on Resuscitation (American Heart Association, European Resuscitation Council, Australian Resuscitation Council, New Zealand Resuscitation Council, Heart and Stroke Foundation of Canada, InterAmerican Heart Foundation, Resuscitation Councils of Southern Africa). *Circulation*. 2004;110:3385-3397.
14. Hammill BG, Hernandez AF, Peterson ED, Fonarow GC, Schulman KA, Curtis LH. Linking inpatient clinical registry data to Medicare claims data using indirect identifiers. *Am Heart J*. 2009;157:995-1000.
15. Chan PS, Nallamothu BK, Krumholz HK, Spertus JA, Li Y, Hammill BG, Curtis LH. Long-term outcomes of elderly survivors of in-hospital cardiac arrest. *N Engl J Med*. 2013;368:1019-1026.
16. Jennett B, Bond M. Assessment of outcome after severe brain damage. *Lancet*. 1975;1:480-484.
17. Heilbron DC. Zero-Altered And Other Regression Models For Count Data With Extra Zeros. *Biometrical Journal*. 1994;36:531-547.

18. Blough DK, Ramsey SD. Using Generalized Linear Models to Assess Medical Care Costs. *Health Services and Outcomes Research Methodology*. 2000;1:185-202.
19. Shao J, Tu D. *The jackknife and bootstrap*. New York: Springer Verlag. 1995.
20. R Development Core Team (2008). R: A language and environment for statistical computing. R Foundation for Statistical Computing V, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org>.
21. Phelps R, Dumas F, Maynard C, Silver J, Rea T. Cerebral Performance Category and long-term prognosis following out-of-hospital cardiac arrest. *Crit Care Med*. 2013;41:1252-1257.
22. Hasselqvist-Ax I, Riva G, Herlitz J, Rosenqvist M, Hollenberg J, Nordberg P, Ringh M, Jonsson M, Axelsson C, Lindqvist J, Karlsson T, Svensson L. Early cardiopulmonary resuscitation in out-of-hospital cardiac arrest. *N Engl J Med*. 2015;372:2307-2315.

Author Manuscript

Table 1. Characteristics of Study Cohort

	Status at 1 Year			P value
	All Patients N = 1127	Alive n = 777	Dead n = 350	
Age, mean \pm SD	75.4 \pm 7.5	74.6 \pm 7.2	77.1 \pm 7.7	<0.001
Age, years				<0.001
65 to 74	563 (50.0%)	425 (54.7%)	139 (39.7%)	
75 to 84	401 (35.6%)	268 (34.5%)	133 (38.0%)	
\geq 85	163 (14.5%)	84 (10.8%)	78 (22.3%)	
Men	657 (58.3%)	457 (58.8%)	200 (57.1%)	0.61
Race				0.27
White	828 (73.7%)	579 (74.8%)	249 (71.3%)	
Black	219 (19.5%)	141 (18.2%)	78 (22.3%)	
Other	76 (6.8%)	54 (7.0%)	22 (6.3%)	
Missing	4	3	1	
Initial cardiac arrest rhythm				0.18
Asystole	245 (21.7%)	165 (21.2%)	80 (22.9%)	
Pulseless electrical activity	268 (23.8%)	175 (22.5%)	93 (26.6%)	
Indeterminate unshockable rhythm	93 (8.3%)	61 (7.9%)	32 (9.1%)	
Ventricular fibrillation	367 (32.6%)	259 (33.3%)	108 (30.9%)	
VT or indeterminate shockable rhythm	154 (13.7%)	117 (15.1%)	37 (10.6%)	
Location of Arrest				0.08
Home	688 (61.0%)	465 (59.8%)	223 (63.7%)	
Nursing home	132 (11.7%)	85 (10.9%)	47 (13.4%)	
Public area	161 (14.3%)	123 (15.8%)	38 (10.9%)	
Hospital or healthcare facility	60 (5.3%)	39 (5.0%)	21 (6.0%)	
Other	86 (7.6%)	65 (8.4%)	21 (6.0%)	
Witnessed arrest	796 (70.6%)	554 (71.3%)	242 (69.1%)	0.47
CPR initiated by:				0.12
First Responder	333 (29.5%)	216 (27.8%)	117 (33.4%)	
Bystander	387 (34.3%)	268 (34.5%)	119 (34.0%)	
Responding EMS Personnel	407 (36.1%)	293 (37.7%)	114 (32.6%)	

Table 1. Characteristics of Study Cohort (continued)

	All Patients N = 1127	Status at 1 Year		P value
		Alive n = 777	Dead n = 350	
Use of public access AED	320 (28.4%)	232 (29.9%)	88 (25.1%)	0.10
Who first applied AED				0.10
First Responder	407 (36.1%)	273 (35.1%)	134 (38.3%)	
Responding EMS	650 (57.7%)	448 (57.7%)	202 (57.7%)	
Bystander	70 (6.2%)	56 (7.2%)	14 (4.0%)	
Principal discharge diagnosis				<0.001
Cardiac arrest	91 (8.1%)	63 (8.1%)	28 (8.0%)	
Acute myocardial infarction	159 (14.1%)	126 (16.2%)	33 (9.4%)	
Other cardiac diagnosis	241 (21.4%)	163 (21.0%)	78 (22.3%)	
Pulmonary diagnosis	220 (19.5%)	119 (15.3%)	101 (28.9)	
Infection and other diagnoses	416 (36.9%)	305 (39.3%)	110 (31.4%)	
Discharge neurological status				<0.001
Mild to no disability (CPC score 1)	555 (52.8%)	444 (60.5%)	111 (34.9%)	
Moderate disability (CPC score 2)	265 (25.2%)	204 (27.8%)	61 (19.2%)	
Severe disability (CPC score 3)	148 (14.1%)	57 (7.8%)	91 (28.6%)	
Coma or vegetative state (CPC score 4)	84 (8.0%)	29 (4.0%)	55 (17.3%)	
Missing	75	43	32	
Discharge destination				<0.001
Home self-care	430 (38.2%)	382 (49.2%)	48 (13.7%)	
Home with home health care	162 (14.4%)	131 (16.9%)	31 (8.9%)	
Skilled nursing or intermediate care facility	196 (17.4%)	111 (14.3%)	85 (24.3%)	
Rehabilitation center	235 (20.9%)	143 (18.4%)	92 (26.3%)	
Hospice	90 (8.0%)	4 (0.5%)	86 (24.6%)	
Other non-home facility	14 (1.2%)	6 (0.8%)	8 (2.3%)	

Abbreviations: AED, automated external defibrillator; CPC, cerebral performance category; CPR, cardiopulmonary resuscitation; EMS, emergency medical services; SD, standard deviation; VT, ventricular tachycardia

Table 2. One-Year Mortality for Pre-Specified Subgroups

Variable	Unadjusted 1-Year Mortality	Adjusted HR (95% CI)	P value
Sex			
Women	31.1%	Reference	Reference
Men	32.8%	1.03 (0.82-1.31)	0.79
Race			
White	30.7%	Reference	0.76
Black	36.7%	1.11 (0.84-1.46)	
Other	29.2%	1.01 (0.63-1.61)	
Initial cardiac arrest rhythm			
Asystole	33.1%	Reference	0.39
Pulseless electrical activity	35.5%	1.19 (0.87-1.63)	
Indeterminate unshockable rhythm	35.4%	1.59 (0.99-2.58)	
Ventricular fibrillation	30.2%	1.23 (0.88-1.70)	
VT or indeterminate shockable rhythm	25.0%	1.20 (0.72-2.01)	
Person initiating CPR			
First responder	36.2%	Reference	<0.001
Bystander	31.7%	0.76 (0.57-1.01)	
EMS personnel	28.6%	0.57 (0.43-0.77)	
Discharge neurological status (CPC score)			
Mild to no disability	21.4%	Reference	<0.001
Moderate disability	24.4%	0.71 (0.45-1.10)	
Severe disability	63.1%	2.08 (1.37-3.14)	
Coma or vegetative state	66.2%	2.09 (1.31-3.35)	

Hospital disposition

Home self-care	11.8%	Reference	<0.001
Home with home health care	19.5%	1.82 (1.15-2.88)	
Skilled nursing or intermediate care	44.8%	4.29 (2.92-6.29)	
Inpatient rehabilitation facility	40.0%	4.23 (2.92-6.12)	
Hospice	96.3%	45.1 (7.31-37.1)	

Abbreviations: CPC, cerebral performance category score; CPR, cardiopulmonary resuscitation; EMS, emergency medical services; HR, hazard ratio; VT, ventricular tachycardia

Table 3. Inpatient Costs at One Year, Overall and by Patient Subgroup

	Unadjusted 1-Year Costs	Adjusted Cost Ratio (95% CI)	P
ALL PATIENTS	\$23,765 ± \$41,003		
Sex			
Women	\$25,207	Reference	Reference
Men	\$22,086	0.97 (0.80-1.20)	0.38
Race			
White	\$20,299	Reference	Reference
Black	\$36,181	1.95 (1.55-2.45)	<0.001
Other	\$24,213	1.02 (0.66-1.50)	0.43
Person initiating CPR			
First responder	\$26,975	Reference	Reference
Bystander	\$21,557	0.87 (0.70-1.15)	0.82
EMS personnel	\$23,441	0.92 (0.72-1.23)	0.67
Initial cardiac arrest rhythm			
Asystole	\$25,854	Reference	Reference

Pulseless electrical activity	\$26,981	1.10 (0.82-1.55)	0.24
Indeterminate unshockable rhythm	\$23,950	0.88 (0.56-1.41)	0.30
Ventricular fibrillation	\$22,235	0.80 (0.59-1.11)	0.12
Pulseless ventricular tachycardia	\$18,516	0.63 (0.30-1.02)	0.06

Discharge neurological status (CPC score)

Mild to no disability	\$19,640	Reference	Reference
Moderate disability	\$31,332	1.27 (0.94-1.76)	0.07
Severe disability	\$43,641	1.54 (1.08-2.33)	0.01
Coma or vegetative state	\$32,107	1.54 (1.00-2.48)	0.03

Hospital disposition

Home	\$14,949	Reference	Reference
Skilled nursing or rehabilitation site	\$41,219	2.75 (2.23-3.33)	<0.001
Hospice	\$1,787	0.12 (0.03-0.26)	<0.001


Abbreviations: CPC, cerebral performance category score; CPR, cardiopulmonary resuscitation; EMS, emergency medical services

Figure Legends


Figure 1. Definition of the Study Cohort

Figure 2. Comparison of Long-Term Mortality with Matched Medicare Cohorts. Survivors of out-of-hospital cardiac arrest had higher mortality during follow-up than patients hospitalized for any reason but lower mortality than patients who required mechanical ventilation during the index hospitalization.

Figure 3. Cumulative Readmission Incidence and Frequency of Readmissions During the First Year. There were 197 readmissions per 100 patient years during the first year of follow-up, although 43% of cardiac arrest survivors were not readmitted during this time.




Author Manuscript



jah3_1396_f2.tif

Author Manuscript



Hospitalizations per Patient	N	% of Cohort
0	489	43.4%
1	206	18.3%
2	153	13.6%
3	86	7.6%
>3	193	17.1%

jah3_1396_f3.tif