

Pseudomonas aeruginosa defense systems against microbicidal oxidants

Bastian Groitl,^{1†} Jan-Ulrik Dahl,^{1†}
Jeremy W. Schroeder² and Ursula Jakob ^{1,3*}

¹Department of Molecular, Cellular, and Developmental Biology, University of Michigan, Ann Arbor, MI, USA.

²Department of Bacteriology, University of Wisconsin-Madison, Madison, WI, USA.

³Department of Biological Chemistry, University of Michigan Medical School, University of Michigan, Ann Arbor, MI, USA.

Summary

The most abundant oxidants controlling bacterial colonization on mucosal barrier epithelia are hypochlorous acid (HOCl), hypobromous acid (HOBr) and hypothiocyanous acid (HOSCN). All three oxidants are highly antimicrobial but little is known about their relative efficacies, their respective cellular targets, or what specific responses they elicit in bacteria. To address these important questions, we directly tested the individual oxidants on the virulent *Pseudomonas aeruginosa* strain PA14. We discovered that HOCl and HOBr work almost interchangeably, impacting non-growing bacterial cultures more significantly than actively growing bacteria, and eliciting similar stress responses, including the heat shock response. HOSCN treatment is distinctly different, affecting primarily actively growing PA14 and evoking stress responses suggestive of membrane damage. What all three oxidants have in common, however, is their ability to cause substantial protein aggregation. This effect became particularly obvious in strains lacking polyphosphate, a newly recognized chemical chaperone. Treatment of PA14 with the FDA-approved anti-inflammatory drug mesalamine, which has recently been shown to attenuate polyP production in a wide range of bacteria, effectively decreased the resistance of PA14 toward all three

oxidants, suggesting that we have discovered a novel, targetable defense system in *P. aeruginosa*.

Introduction

The natural ability of mammalian hosts to defend themselves against invading pathogens involves the production of antimicrobial oxidants, particularly hypochlorous acid (HOCl, bleach), hypobromous acid (HOBr) and hypothiocyanous acid (HOSCN) (Klebanoff *et al.*, 2013). These antimicrobial hypohalous acids (HOX) are produced by human haloperoxidases, which include (i) myeloperoxidases (MPO) in activated neutrophils, monocytes, and tissue macrophages at sites of inflammation (Davies, 2011); (ii) lactoperoxidases (LPO) and salivary peroxidases, that are secreted from salivary and other mucosal glands and play important roles in controlling the oral microflora (Ashby, 2008); (iii) eosinophil peroxidases in eosinophil granulocytes (EPO) (Wang and Slungaard, 2006) and (iv) gastric peroxidases (GPO) found in cells of the fundic region in the stomach (Das *et al.*, 1995). What all of these haloperoxidases have in common is their ability to catalyze the peroxide (H₂O₂)-mediated oxidation of halides (e.g., Cl⁻, Br⁻) and/or pseudohalides (e.g., SCN⁻), yielding the respective two-electron oxidants (Davies, 2011; Winterbourn *et al.*, 2016). The levels of oxidants produced depend on the concentrations of the respective anions and on the peroxidases catalyzing the reaction. Previous reports indicated that MPO in the plasma produces almost equal amounts of HOCl and HOSCN, and only a minor (~5%) amount of HOBr (van Dalen *et al.*, 1997). In contrast, LPO and GPO only oxidize SCN⁻ under physiological conditions, whereas EPO oxidizes primarily SCN⁻ and Br⁻ (van Dalen *et al.*, 1997; Hawkins *et al.*, 2009).

All three oxidants are effective antimicrobials and considered more bactericidal/bacteriostatic than H₂O₂ (Love *et al.*, 2016). HOCl, for instance, is known to rapidly oxidize a variety of amino acid side chains, causing extensive protein unfolding and aggregation (Winter *et al.*, 2008; Winterbourn, 2008; Dahl *et al.*, 2015). Not surprisingly, bacterial defense systems against HOCl include expression of transcription factors and genes involved in

Accepted 6 August, 2017. *For correspondence. E-mail ujakob@umich.edu; Tel. (+1) 734 615 1286; Fax (+1) 734 647 0884. †Both the authors contributed equally to this work.

detoxification and reduction of side chain modifications (Gebendorfer *et al.*, 2012; Gray *et al.*, 2013; Parker *et al.*, 2013), activation of stress specific molecular chaperones, such as Hsp33 and RidA that prevent protein aggregation (Winter *et al.*, 2008; Muller *et al.*, 2014), and conversion of ATP in inorganic polyphosphate, a chemical chaperone effective in stabilizing protein structures (Gray *et al.*, 2014; Gray and Jakob, 2015). Absence of any one of these systems causes a significant decrease in HOCl stress resistance, suggesting that protein unfolding is a major contributor to the bactericidal activity of HOCl. Much less is known about the effects of and response to HOBr, which is also a strong oxidant that readily brominates tyrosine, tryptophan and nucleic acids (Klebanoff, 2005; Davies *et al.*, 2008). Nevertheless, HOCl and HOBr are both associated with numerous diseases, including initiation and progression of various inflammatory diseases, such as atherosclerosis (Klebanoff, 2005; Davies *et al.*, 2008; Davies, 2011). HOSCN appears to react with great specificity with thiols (e.g., GSH; protein cysteines) and selenogroups (Skaff *et al.*, 2009, 2012), resulting in a number of reversible oxidation products, including sulfenyl thiocyanates, sulfenic acids or disulfides (Barrett *et al.*, 2012). High concentrations of HOSCN, however, result in non-reversible thiol products including sulfinic/sulfonic acids (Barrett and Hawkins, 2012). In contrast to HOCl/HOBr, the role of HOSCN in disease is still unclear and appears organism-specific, ranging from apparently protective functions in mammalian hosts to cytotoxic effects in bacteria (Pattison *et al.*, 2012).

It is largely unknown how the three oxidants compare in bactericidal efficacies, what their specific *in vivo* targets are, and whether bacteria contain universal or individualized defense systems against these three different stressors. Finding answers to these important questions is especially pressing for patients suffering from cystic fibrosis, one of the most common fatal hereditary diseases, affecting about 30,000 people in the US (Cohen and Prince, 2012). Caused by a defect in the cystic fibrosis transmembrane conductance regulator (CFTR), patients experience frequent and reoccurring lower respiratory tract infections, chronic inflammation and progressive tissue damage in the lungs (Rao and Grigg, 2006). Analysis of the bronchoalveolar lavage fluid of CF patients revealed substantially altered levels of HOCl and HOSCN, presumably due to a combination of increased MPO activity caused by sustained airway neutrophilia, and a lack of secreted SCN⁻ and glutathione (GSH), two CFTR-dependent compounds (Kogan *et al.*, 2003; Lorentzen *et al.*, 2011). HOCl levels were further increased in CF patients testing positive for *P. aeruginosa* infection, consistent with even higher numbers of

activated neutrophils and further increased MPO concentrations during infections.

In this study, we now directly compared the antimicrobial efficacy of HOCl, HOBr and HOSCN on the pathogenic *P. aeruginosa* strain PA14. We determined the effects that the three oxidants exert on PA14, and identified common and oxidant-specific response and defense systems by RNA_{seq} analysis. Despite their distinct reactivities, we discovered that all three oxidants cause extensive protein unfolding and aggregation. Stress-induced accumulation of polyphosphate appears to be the universal response to HOCl/HOBr and HOSCN treatment and serves as an effective strategy to prevent stress-induced protein aggregation and increase stress survival. Mesalamine, an FDA-approved drug recently discovered to reduce cellular polyP levels in a wide range of bacteria, including *P. aeruginosa*, effectively increases PA14's sensitivity towards the antimicrobial oxidants, making it a potential treatment option for chronic *P. aeruginosa* infections.

Results

The antimicrobial efficacies of HOCl, HOBr and HOSCN differ by growth conditions

Most growth media are rich in components that react with and potentially quench the effective concentration of oxidants like HOCl and HOBr, making direct comparisons in relative bactericidal efficacy challenging. As a first approach, we therefore decided to treat the bacteria with the respective oxidants in phosphate-buffered saline (PBS), which is known not to react with any of the oxidants. We grew *P. aeruginosa* PA14 to mid-logarithmic phase in MOPS-glucose media, switched the media to PBS and treated the bacteria with various concentrations of the respective oxidants for 30 min. Next, we quenched the oxidants, serially diluted and spotted the bacteria onto LB plates and monitored their survival after 14 h of incubation. We found that HOCl and HOBr showed very comparable bactericidal activity with 100–125 μ M of each oxidant being sufficient to reduce PA14 survival by about four orders of magnitude (Fig. 1A, Supporting Information Fig. S1). In contrast, HOSCN, when used at this concentration, reduced bacterial survival by less than one log, suggesting that HOSCN is much less bactericidal under these conditions (Fig. 1A, Supporting Information Fig. S1). Moreover, even at very high concentrations of HOSCN (i.e., 1 mM), a substantial fraction of bacteria still survived the treatment.

Surprisingly, we obtained very different results when we conducted the treatments in MOPS-glucose media. Under these conditions, HOBr was most effective in killing PA14, followed by HOSCN and HOCl (Fig. 1B, left




Fig. 1. Effects of HOCl, HOBr and HOSCN on the survival of non-growing or exponentially growing PA14. Wild-type PA14 was grown in MOPS-glucose media to mid-log phase ($A_{600} = 0.4\text{--}0.5$) and shifted to either (A) PBS buffer or (B) MOPS media in the absence or presence of glucose. The indicated concentrations of HOCl, HOBr or HOSCN were added. After 30 min incubation, excess HOX was quenched by the addition of 10 mM thiosulfate, and the cells were serially diluted with 0.85% NaCl, spot-titrated onto LB agar plates, and incubated o/n at 37°C. The experiments were repeated at least three independent times. Data shown in (A) represent means \pm s.e.m. A representative data set is shown in (B).

panel). Explanations for this result included the possibility that components of the MOPS-glucose medium differentially quench the respective oxidants, or that cells grown in MOPS-glucose media might react with oxidants in ways that generate reaction products of differing bactericidal activities (e.g., chloramines, bromamines). Alternatively, however, we also considered the possibility that the three oxidants might act differently on metabolically active and inactive bacteria. To distinguish between these possibilities, we cultivated PA14 in MOPS-glucose, washed the cells and, shortly before the stress treatment, shifted one part of the cells into fresh MOPS media with glucose (MOPS-glucose) to allow for continuous growth. The other part, we transferred into fresh MOPS media without glucose (MOPS), which resulted in an immediate growth arrest. We reasoned that this approach should allow us to keep the media composition constant while only varying the growth status of the bacteria. We found that HOSCN was substantially more

bactericidal on actively growing cells, while HOCl and HOBr were both more effective in killing non-growing *P. aeruginosa* (Fig. 1B, compare left and right panel). For instance, in the presence of 0.5 mM HOSCN, cell survival in MOPS-glucose media was reduced by 4 orders of magnitude while no substantial killing was observed when the treatment was conducted in MOPS media. In contrast, HOCl and HOBr caused significant killing non-growing bacteria at concentrations that posed little threat to actively metabolizing bacteria (Fig. 1B). Importantly, we confirmed that glucose does not quench HOCl, suggesting that it is indeed the growth status of the cells that affects the toxicity of the oxidants. These results are fully consistent with our previous experiments conducted in PBS buffer, which showed that both HOCl/HOBr are significantly more effective than HOSCN in killing non-growing *P. aeruginosa*. They furthermore suggest that the three oxidants work by different mechanisms and/or affect different *in vivo* targets.

PA14 transcriptional changes in response to HOCl, HOBr and HOSCN treatment

Adaptation of gene expression is key for bacteria to quickly respond to changes in their environment. As a first step towards obtaining insights into the physiological consequences of HOCl, HOBr or HOSCN treatment, we tested gene expression changes in PA14 treated with sublethal concentrations of the respective oxidants. Since we wanted to keep the cultivation conditions the same and directly compare the *in vivo* effects of the three oxidants, we had to first identify concentrations of each oxidant that cause comparable growth defects without killing the cells. We settled on 0.5 mM HOCl, 0.15 mM HOBr or 0.25 mM HOSCN, which triggered a 1–1.5 h growth arrest (Supporting Information Fig. S2A). We isolated total RNA from the treated cells and conducted RNA_{seq} analysis to monitor global changes in gene expression in PA14 in response to each of the three oxidants. All experiments were conducted in triplicates. For the transcriptome analysis, we included all genes annotated in the NCBI database and compared the expression values of the stress treated cells to non-stress treated controls. We set a false discovery rate (FDR) of <0.005 as a threshold for significance, and considered transcripts as upregulated when they showed a log₂ fold change of >1.5 and downregulated when they showed a log₂ fold change of <−1.5. A list of all expression data can be found in Supporting Information Table S1.

A multidimensional scaling plot of our data sets revealed that while gene expression changes in all three replicates treated with either HOCl or HOBr clustered in one dimension (Fig. 2A, green and red diamonds), treatment of PA14 with HOSCN led to distinctly different gene expression changes and clustered in two dimensions (Fig. 2A, purple diamonds). Closer analysis of all up- and downregulated genes in the three treatment groups revealed a >85% overlap between differentially expressed genes during HOBr and HOCl treatment while less than a 50% overlap existed between differentially expressed genes in PA14 treated with HOCl/HOBr and HOSCN (Fig. 2B). Closer analysis of the data revealed that all three stressors induced the expression of genes involved in oxidant detoxification (e.g., catalase, methionine sulfoxide reductase, alkyl hydroperoxide reductase) and restoration of redox homeostasis (e.g., glutathione peroxidase, glutathione reductase) (Fig. 2C and D; Supporting Information Table S1). Very similar results in regards to HOCl stress responses were previously reported for *Bacillus subtilis*, *P. aeruginosa* strain PAO1, *Bacillus cereus* ATCC14579 and *E. coli* MG1655 (Small *et al.*, 2007; Ceragioli *et al.*, 2010; Chi *et al.*, 2011; Gray *et al.*, 2013).

We next ran Fisher's Exact Tests on the 78 known regulons in PA14, and determined the FDR to identify differentially expressed genes that are statistically significantly enriched upon one or the other stress treatment (see *Experimental procedures* for details). Surprisingly, only 6 regulons showed an FDR <0.05 in stress-treated versus untreated PA14, and only one of these regulons, MexT, responded comparably to all three antimicrobials (Table 1; Supporting Information Fig. S2B). MexT is a redox-sensitive LysR-type transcriptional regulator that modulates the expression of virulence factors and significantly contributes to the increased antibiotic resistance in *P. aeruginosa* (Fargier *et al.*, 2012). It has recently been shown to induce the expression of the MexEF-OprN multidrug efflux system, which exports electrophilic metabolites to restore redox homeostasis in the cell (Fargier *et al.*, 2012). Our finding that all three oxidants induce the MexT regulon agrees with the redox-modifying properties of the three oxidants, and suggests that they activate MexT through thiol oxidation. Surprisingly, based on our FDR <0.05 criterium, we did not find the peroxide-sensitive OxyR regulon to be significantly induced by any of the three treatments (Supporting Information Fig. S2B), although OxyR has been previously shown to contribute to increased HOCl resistance in *E. coli* (Dukan and Touati, 1996).

The LasR regulon, which controls quorum sensing in *P. aeruginosa* was significantly down-regulated upon both HOCl and HOBr treatment but not by HOSCN. The remaining four regulons responded specifically to only one of the three oxidants; whereas both RhlR- and GacA regulons were significantly altered only upon HOBr treatment, Fur and PchR appeared to respond to HOCl treatment only (Table 1; Supporting Information Fig. S2B).

We also analyzed the expression levels of transcriptional regulators, whose regulons have not yet been identified in *P. aeruginosa* but which have previously shown to be HOCl-responsive in other organisms. Similar to our previous results, we found that HOCl, HOBr and HOSCN treatment elicits both overlapping and distinct responses in PA14. For instance, we found that the oxidative stress-sensing regulator OspR, which regulates glutathione peroxidase (Atichartpongkul *et al.*, 2016), was strongly induced upon treatment with any of the three antimicrobials (Supporting Information Table S1). Expression of NemR (PA14_36300), a known HOCl-specific repressor in *E. coli* (Gray *et al.*, 2013), was significantly upregulated in response to HOCl and HOSCN but not HOBr (Supporting Information Table S1). In contrast, the chlorine-specific transcription factor HypT/YjiE (PA14_71640) (Gebendorfer *et al.*, 2012) was significantly induced only by HOCl, while RclR (PA14_07340), a third HOCl-responsive transcriptional activator in *E. coli* (Parker *et al.*, 2013) was not

significantly up-regulated in response to any of the three oxidants (Supporting Information Table S1).

By far the most significant differences in the proportion of upregulated genes, however, were found in the functional categories of molecular chaperones (Fig. 2C and E). While a total of 12 and 9 heat shock genes (e.g., *ibpA*, *dnaK*, *groES*, *hslU/V*) were significantly upregulated in response to HOCl and HOBr respectively, only two (i.e., *grpE*, *hslV*) were upregulated in response to HOSCN (Fig. 2E). Moreover, the transcripts of both genes were about sixfold less abundant in HOSCN-treated cells as compared to HOCl-treated cells. The HOCl-results were consistent with our previous microarray data in *E. coli* (Gray *et al.*, 2013). Other functional categories that were specifically and significantly more upregulated in response to HOCl/HOBr as compared to HOSCN included genes involved in motility and attachment, chemotaxis and nucleotide biosynthesis (Fig. 2C). This result fully agrees with previous observations in *E. coli* MG1655 and *B. subtilis*, where exposure to sublethal HOCl concentrations caused the induction of motility and chemotaxis-related genes (Chi *et al.*, 2011; Gray *et al.*, 2013). Interchangeably, a much higher proportion of genes involved in antibiotic resistance or encoding membrane proteins were upregulated in response to HOSCN as compared to HOCl and HOBr, suggesting that HOSCN might target membrane proteins (Fig. 2C). These data demonstrate that HOCl and HOBr work quite similarly *in vivo*, and elicit distinct gene expression changes when compared with the effects of HOSCN treatment.


Effects of HOCl, HOBr and HOSCN on the PA14 proteome

HOCl has previously been shown to induce extensive protein unfolding and aggregation (Winter *et al.*, 2008). This effect of HOCl is presumably due to its ability to rapidly oxidize a variety of different amino acid side chains, thereby shifting the equilibrium of proteins more towards their aggregation-prone state (Winter *et al.*, 2008; Dahl *et al.*, 2015). HOCl-stress induced protein aggregation explains the upregulation of members of the heat shock regulon, whose expression is triggered by the accumulation of unfolded proteins (Arsene *et al.*, 2000). Since bacterial HOCl-stress resistance is increased by preventing HOCl-induced protein aggregation, we previously concluded that protein unfolding is one mechanism by which HOCl kills bacteria (Winter *et al.*, 2008). Based on the chemical similarity between HOCl and HOBr as well as the similarity by which cells react to the two stressors, it is likely that HOBr also triggers global protein aggregation. HOSCN, in contrast, is thought to be highly cysteine-specific, raising the

question whether it kills bacteria by different means (Skaff *et al.*, 2009; Barrett *et al.*, 2012; Chandler *et al.*, 2013).

To directly compare the extent and targets of protein aggregation during our various oxidative stress treatments, we therefore treated wild-type PA14 with our previously established sublethal concentrations of HOCl (0.5 mM), HOBr (0.15 mM) or HOSCN (0.25 mM). After 30 min of treatment, we lysed the cells and analyzed the aggregated proteins by SDS-PAGE. While treatment with HOCl and HOBr triggered similar protein aggregation, affecting a large number of different PA14 proteins, we found much less protein aggregation when cells were treated with growth-delaying concentrations (i.e., 0.25 mM) of HOSCN (Fig. 3A, Supporting Information Fig. S3). This result was fully consistent with our RNA-seq data that showed induction of the heat shock response when cells experienced sublethal doses of HOCl/HOBr but not HOSCN (Fig. 2E). Higher (i.e., lethal) concentrations of HOSCN, however, also triggered substantial protein aggregation, indicating that HOSCN either directly or indirectly exerts proteotoxic effects as well (Supporting Information Fig. S3).

Subsequent SILAC experiments on the protein aggregates on HOCl and HOSCN treated PA14 further confirmed these results and revealed a significantly increased aggregation of proteins in HOCl-treated bacteria as compared to HOSCN-treated PA14 (Fig. 3B). However, due to the fact that PA14 produces endogenous arginine, the incorporation rate of $^{13}\text{C}_6$ -Arg was very low (6–8%). This caused our H/L ratios to be quite small, and made direct comparisons of the respective H/L ratios in HOCl and HOSCN treated bacteria challenging. As a first approach, we therefore decided to use the H/L ratio of the over 30 identified ribosomal proteins found in both of our aggregate preparations as internal reference points (see Supporting Information Fig. S4 and Table S2, highlighted in orange). We then focused primarily on those annotated proteins, whose H/L ratio was substantially higher than that of ribosomal subunits. By using this analysis, we confirmed many of the proteins that have previously been identified to be aggregation prone in response to HOCl treatment (Winter *et al.*, 2008), including GapA, EF-Tu and ClpX (see Supporting Information Fig. S4 and Table S2, highlighted in blue). A few of these proteins were also identified among the aggregation-sensitive proteins in HOSCN-treated PA14. However, the H/L ratio was typically much lower. In addition, we found several proteins that appeared to be substantially more prone to HOSCN-mediated aggregation than to HOCl-mediated aggregation, suggesting that differences exist in the target proteins of the two oxidants. More detailed studies are needed in the future to reveal the features that make



proteins particularly sensitive towards HOCl vs. HOSCN treatment. A complete list of all proteins identified in the aggregates of HOCl and HOSCN-treated *P. aeruginosa* can be found in Supporting Information Table S2.

Polyphosphate – a universal oxidative stress defense system

Previous work in *E. coli* and *V. cholerae* revealed both protein and non-protein based chaperone strategies that

Fig. 2. Global gene expression changes in PA14 in response to HOCl, HOBr or HOSCN treatments. Exponentially growing PA14 wild-type cells ($A_{600} = 0.4\text{--}0.5$) were incubated with sublethal concentration of HOCl (0.5 mM), HOBr (0.15 mM) or HOSCN (0.25 mM) for 15 min. Incubation was stopped by the addition of ice-cold methanol. Reads were aligned to the *P. aeruginosa* UCBPP-PA14 reference genome (Accession number: NC_008463.1).

A. A multidimensional scaling plot revealed distinct clustering of transcriptomes between untreated PA14 (black) and PA14 treated with HOCl (green), HOBr (red) or HOSCN (purple).

B. Venn diagram shows the number of upregulated or downregulated genes upon treatment with HOCl (green), HOBr (red), or HOSCN (purple) and a potential overlap between treatments.

C. Proportion of upregulated genes with regards to functional groups defined in (Lee *et al.*, 2006).

D. Log₂-fold change in genes involved in redox homeostasis.

E. Log₂-fold change in heat shock genes. Each condition was tested in triplicate. Significance was defined by a false discovery rate of <0.005 and absolute values of log₂-fold change of >1.5.

protect bacteria against oxidative stress-mediated protein aggregation (Dahl *et al.*, 2015). These strategies include post-translational activation of the redox-regulated chaperone Hsp33 as well as the conversion of ATP into the chemical chaperone polyphosphate (polyP). Hsp33 is inactive under non-stress conditions but becomes rapidly activated as a molecular chaperone upon oxidative stress conditions that lead to widespread protein unfolding (Winter *et al.*, 2008). Accumulation of polyP, a universal polymer, is at least in part achieved by the oxidant-mediated inactivation of the polyP degrading enzyme exopolyphosphatase (PPX) (Gray *et al.*, 2014). Like activated Hsp33, polyP works as an effective protein-stabilizing chaperone, which binds unfolding proteins and prevents their *in vivo* aggregation. To test the extent to which Hsp33 and/or polyP participate in protecting *P. aeruginosa* against the three physiologically relevant oxidants, we constructed PA14 deletion mutants, lacking either the Hsp33 encoding gene *hsI/O*, the polyphosphate kinase (PPK) encoding gene *ppk*, or both *hsI/O* and *ppk* genes. As before, we exposed the cells to HOX-treatment in MOPS-glucose media, and either monitored growth in liquid culture after treatment (Fig. 4A) or their survival (Supporting Information Fig. S5). We found that the *hsI/O* deletion reproducibly increased the sensitivity of PA14 towards HOCl and HOBr treatment but not towards HOSCN stress (Fig. 4A; Supporting Information Fig. S5). This result was not completely unexpected given that 0.25 mM HOSCN did

not appear to cause significant protein unfolding in PA14 wild-type cells, and hence would be predicted to fail to trigger unfolding of Hsp33. Indeed, *in vitro* activation studies using recombinantly expressed PA14 Hsp33 revealed that both HOBr and HOCl rapidly activated the chaperone function of PA14 Hsp33 whereas HOSCN treatment had no activating effect (Fig. 4B).

In contrast to the *hsI/O* deletion, we found that the deletion of the *ppk* gene dramatically increased the sensitivity of PA14 towards all three oxidants, HOCl, HOBr and HOSCN. For instance, we observed no recovery upon treatment of the *ppk* mutant with 0.25 mM HOSCN within the timeframe of the experiment (Fig. 4A), and survival was decreased by at least 2 orders of magnitude compared to wild-type PA14 (Supporting Information Fig. S5). However, whereas additional deletion of *hsI/O* did not cause any further increase in the HOSCN sensitivity of PA14, deletion of *hsI/O* increased the sensitivity toward HOCl and HOBr, suggesting that polyP and Hsp33 participate in protecting PA14 against HOCl and HOBr stress (Fig. 4A, Supporting Information Fig. S5).

Analysis of the polyP levels in HOCl, HOBr and HOSCN-treated wild-type PA14 confirmed that all three treatments cause an increase in polyP level (Fig. 4C). Since the respective polyP levels were quite similar, however, these results also excluded the possibility that polyP's apparently more substantial influence on HOSCN resistance is due to higher amounts of polyP. Instead, our results suggest that polyP accumulation in wild-type PA14 is either sufficient to deal with HOSCN-mediated protein unfolding, explaining the lack of heat shock gene induction and failed activation of Hsp33, or that polyP prevents HOSCN-mediated damage of other cellular targets, such as DNA. In either case, our results suggested that we discovered a universal stress response system that protects PA14 against all three neutrophilic oxidants.

Absence of PolyP leads to massive protein aggregation in HOSCN-treated bacteria

To gain insights into the mechanism(s) by which polyP protects PA14 against neutrophil-derived oxidants, we

Table 1. Regulons determined to be significantly enriched for differentially expressed genes (FDR<0.05) using one-sided Fisher's Exact Tests.

| Regulon | FDR | | |
|-------------|----------------------|-----------------------|----------------------|
| | HOBr | HOCl | HOSCN |
| <i>mexT</i> | 1.8×10^{-6} | 1.1×10^{-9} | 5.8×10^{-4} |
| <i>lasR</i> | 1.6×10^{-8} | 1.6×10^{-12} | n.s. |
| <i>pchR</i> | n.s. | 3.0×10^{-4} | n.s. |
| <i>fur</i> | n.s. | 0.03 | n.s. |
| <i>rhIR</i> | 1.1×10^{-4} | n.s. | n.s. |
| <i>gacA</i> | 0.015 | n.s. | n.s. |




Fig. 3. Effects of HOCl or HOSCN on the aggregation propensity of the PA14 wild-type proteome. A SILAC experiment was performed to determine the distribution of aggregated proteins in PA14 upon treatment with either HOCl or HOSCN. PA14 wild-type cells were grown in MOPS-glucose medium supplemented with either 120 $\mu\text{g/ml}$ isotopically light [$^{12}\text{C}_6$]Arg or heavy [$^{13}\text{C}_6$]Arg. PA14 cultures in light [$^{12}\text{C}_6$]Arg were left untreated while cultures in heavy [$^{13}\text{C}_6$]Arg were exposed to either 0.5 mM HOCl or 0.25 mM HOSCN for 30 min. Cells were lysed, and the aggregated proteins were separated from the soluble fractions by centrifugation.

A. Insoluble proteins were analyzed by SDS-PAGE.

B. Equal volumes of insoluble proteins from light and heavy samples were mixed, and the proteins were identified by MS/MS analysis. A heat map was generated using the Hierarchical clustering function with Perseus. Row Trees were processed using the default settings with color red indicating those proteins that were enriched in the aggregates of HOCl (left) or HOSCN (right) treated cells.

compared protein aggregation in the *ppk* deletion strain in response to HOCl and HOSCN. We reasoned that if polyP protected PA14 against HOSCN-mediated protein damage, absence of polyP would cause a dramatic increase in protein aggregation. In contrast, no change in the aggregation pattern would be expected if polyP's effects were independent of protein unfolding. As before, we treated the *ppk* deletion strain with 0.25 mM HOSCN for 30-min. While this concentration did not cause any detectable protein aggregation in the wild-type strain (Fig. 3A and B, Supporting Information Fig. S3), in the *ppk* deletion strain it led to protein aggregation that was fully comparable to the massive protein aggregation triggered by a treatment with 0.5 mM HOCl (Fig. 5). In support of these results, we found that the *ppk* deletion strain now responds to HOSCN – treatment with the upregulation of the heat shock response, indicating that in the absence of polyP, cells try to compensate by increasing chaperone production (Supporting

Information Fig. S6). Moreover, and confirming our earlier SILAC results, we again noted that several proteins were differentially affected by HOSCN and HOCl treatment, indicating that the two oxidants have slightly different *in vivo* targets (Fig. 5, arrows). In summary, these results demonstrate that HOSCN has the capacity to cause widespread protein aggregation and suggest that in wild-type PA14, endogenous polyP levels are sufficient to protect proteins against HOSCN-mediated protein damage.

Mesalamine increases PA14 sensitivity toward all three oxidants

Our data revealed that accumulation of polyP serves as an efficient posttranslational stress response system that protects *Pseudomonas* against the deleterious effects of all three major neutrophilic oxidants. We recently identified mesalamine, the gold standard in




Fig. 4. Chaperones provide a defense mechanism against HOX-mediated killing. **A.** Logarithmically growing PA14 wild-type, *ppk*⁻, *hslO*⁻ or *ppk hslO*⁻ cells in MOPS-glucose medium ($A_{600} = 0.4\text{--}0.5$) were incubated with the indicated concentrations of either HOCl, HOBr or HOSCN. Growth was recorded every 30 min for 6 h post treatment. **B.** To test the effects of HOX on the *in vitro* activation of PA14-Hsp33, 12 μM chemically denatured citrate synthase was diluted 160-fold into 40 mM HEPES, pH 7.5 at 30°C in the presence and absence of PA14 Hsp33 that was oxidized with the indicated molar excess of HOX. Light scattering was monitored with excitation and emission wavelength set to 360 nm. The light scattering signal of citrate synthase incubated in the presence of 20-fold molar excess of HOBr-activated PA14-Hsp33 was set to 100% activity while the light scattering signal of CS in the absence of any Hsp33 was set to 0% chaperone activity. **C.** Exponentially growing PA14 wild-type and *ppk*⁻ cells ($A_{600} = 0.4\text{--}0.5$) were incubated with sublethal concentrations of either HOCl (0.5 mM), HOBr (0.15 mM) or HOSCN (0.25 mM) for 2.5 h. Then, polyP was extracted, quantified and the resulting amount was normalized to the respective protein amount in each sample. Experiments were performed independently at least three times. Data represent means \pm s.d.




Fig. 5. HOCl and HOSCN cause widespread protein aggregation in PA14 lacking polyP. Exponentially growing PA14 *ppk*⁻ cells in MOPS-glucose medium ($A_{600} = 0.4$ – 0.5) were either left untreated or incubated with HOCl (0.5 mM) or HOSCN (0.25 mM) for 30 min. Incubation was stopped by the addition of 10 mM thiosulfate. Cells were lysed and aggregated proteins were separated from the soluble fraction by centrifugation. The insoluble proteins were analyzed by SDS-PAGE. Arrows indicate changes in the aggregation pattern between HOCl- and HOSCN-treated samples. Experiments were performed independently at least three times.

treating patients with mild to moderate ulcerative colitis, as a potent inhibitor of bacterial polyP production in a wide range of bacteria, including *P. aeruginosa* PA14 (Dahl *et al.*, 2017). We now wondered whether mesalamine treatment could be used to sensitize PA14 toward the three physiological oxidants, making it potentially a powerful drug to combat *P. aeruginosa* infections within chronically inflamed environments. To test this idea, we treated logarithmically growing PA14 wild-type and *ppk* deletion mutants for 120 min with mesalamine before we exposed the cells to our 30-min HOCl, HOBr or HOSCN stress. As before, we monitored and compared cell survival. As shown in Fig. 6, mesalamine-treated

wild-type PA14 was almost as sensitive towards all three oxidants as the *ppk* deletion strain. Consistent with our previous findings (Dahl *et al.*, 2017), mesalamine did not alter the sensitivity of the *ppk* deletion strain. These results demonstrate that a decrease in polyphosphate levels sensitizes *P. aeruginosa* toward HOCl, HOBr and HOSCN stress, and suggest that mesalamine might be an appropriate option in the treatment of PA14 infections.

Discussion

The accumulation of reactive oxygen and chlorine species (RO/CS) is highly toxic, causing membrane alterations, protein unfolding and DNA damage. It is therefore not surprising that organisms developed systems that utilize increased RO/CS production and secretion as a strategy to defend themselves against their enemies. One example for such a system comes in the form of the mammalian innate immune defense, where neutrophils produce and release high levels of HOCl, HOBr and HOSCN to kill invading pathogens and control bacterial colonization (Davies *et al.*, 2008). To investigate what mechanisms bacteria use for their defense, we compared the effects and consequences of HOCl, HOBr and HOSCN treatment on *P. aeruginosa*, a microorganism known to persist within chronically inflamed environments that are enriched for neutrophil-derived oxidants (Rada, 2017). We reasoned that by identifying and subsequently targeting such bacteria-specific defense mechanisms, we would have the potential to increase the efficacy of the host to kill colonizing bacteria. This might potentially provide a novel approach to combat chronic *P. aeruginosa* infections.

We determined the bacteriotoxic concentrations for HOCl, HOBr and HOSCN, and compared their bactericidal effects on both actively growing and dormant cells. We found that HOCl and HOBr are most toxic for non-growing bacteria, cultivated in either PBS or MOPS buffer. These results are consistent with the high reactivity of these oxidants, and suggest that some of the organic compounds that are produced and secreted by actively metabolizing bacteria likely react with and partially detoxify the oxidants. In contrast, HOSCN was significantly more bactericidal for actively growing PA14 than for non-metabolizing cells, suggesting that HOSCN treatment affects cellular systems involved in metabolism and/or growth. Previous studies conducted in bacteria and mammalian cell cultures suggested several mechanisms by which HOSCN might affect metabolism (Love *et al.*, 2016). For instance, redox proteomic studies revealed active site cysteines in glycolytic enzymes as one of the major *in vivo* oxidation targets of HOSCN,




Fig. 6. Mesalamine-treatment increases HOX sensitivity of wild-type PA14 by targeting polyP homeostasis. Exponentially growing PA14 wild-type and *ppk*⁻ cells in MOPS-glucose medium ($A_{600} = 0.4\text{--}0.5$) were either left untreated or pretreated with 0.5 mM mesalamine for 120 min. Then, treatment with the indicated concentrations of HOCl, HOBr or HOSCN was conducted as before. After 30 min of incubation, excess HOX was quenched by the addition of 10 mM thiosulfate. Cells were serially diluted with 0.85% NaCl, spot-titrated onto LB agar plates and incubated overnight at 37°C. The experiments were repeated at least three independent times.

together with proteins involved in redox signaling, structure maintenance, and protein folding (Love *et al.*, 2016). Moreover, HOSCN has been shown to inhibit glucose and amino acid uptake in a broad spectrum of bacteria, most likely by targeting membrane-bound transport proteins for these nutrients (Chandler and Day, 2015). These results are consistent with our RNA_{seq} data, which revealed significant HOSCN-specific upregulation of genes encoding for membrane proteins and transporters. They furthermore revealed that HOCl/HOBr and HOSCN exert distinctly different effects on bacteria and, when produced in combination, as is the case in healthy individuals, should be effective in targeting both actively growing and dormant bacteria.

Analysis of the defense strategies in *P. aeruginosa* revealed at least one cellular system, which significantly promotes bacterial resistance towards all three

oxidants. We found that in response to HOCl/HOBr and HOSCN, PA14 produces high levels of polyphosphate, which confers considerable resistance against all three oxidants. PolyP is a universally conserved, energy-rich polymer, which was recently shown to bind unfolded proteins and prevent their non-specific aggregation, suggesting it has chaperone-like activities (Gray *et al.*, 2014). At first, this result was surprising since HOSCN treatment, in contrast to HOCl, did not cause any detectable protein aggregation or upregulation of the heat shock response in wild-type PA14, suggesting that HOSCN does not affect protein homeostasis. However, when we conducted these experiments in strains lacking PPK, the kinase responsible for polyP synthesis, we found that HOSCN treatment triggered massive amounts of protein aggregation, and led to increased expression of heat-shock regulated genes. Since wild-

type PA14 produced approximately the same levels of polyP in response to HOCl/HOBr and HOSCN treatment, we concluded that polyP production is more effective in protecting PA14 against HOSCN-mediated protein unfolding as compared to HOCl-mediated unfolding. These results make sense when one considers the reactivity of HOCl/HOBr and HOSCN. HOCl/HOBr are highly reactive toward a number of different amino acid side chains, whose oxidation rapidly cause protein unfolding both *in vitro* and *in vivo* (Winter *et al.*, 2008). In contrast, HOSCN, which is highly thiol specific (Skaff *et al.*, 2009), will presumably react first with surface-exposed cysteine residues, whose oxidation is unlikely to cause rapid protein unfolding. Cellular levels of polyP increase in both circumstances since polyP accumulation is in part mediated by the oxidative modification of a surface-exposed cysteine in the exopolyphosphatase PPX (Gray *et al.*, 2014). It is therefore possible that polyP production in response to HOCl treatment is simply not fast enough to compete with the sudden onset of protein unfolding, a conclusion that would also explain why cells require the extremely fast activation of the redox-regulated chaperone Hsp33 for additional protection. HOSCN might either work much slower, and hence give polyP the time to accumulate and stabilize unfolding proteins, or might affect only a few crucially important redox-sensitive proteins, such as cysteine containing chaperones (Winter *et al.*, 2005), whose inactivation would trigger more widespread protein aggregation. Either of these scenarios is consistent with our finding that HOCl and HOSCN target a set of only partially overlapping cellular proteins for aggregation.

In summary, our results suggest that we have identified a protection mechanism that would become crucial for PA14 when present in an environment of chronic inflammation, such as it is found in the airways of patients suffering from cystic fibrosis. Successful clearance of chronic *P. aeruginosa* infection is often hampered by the high antibiotic resistance of *P. aeruginosa* (Lister *et al.*, 2009). We have now discovered that mesalamine, which decreases endogenous polyP levels in PA14, substantially increases the sensitivity of PA14 to neutrophilic oxidants. The fact that PA14 does not require polyP for growth under non-stress conditions likely reduces their chance to develop resistance. Our studies therefore serve as a first step towards alternative treatment strategies aimed to improve the ability of the endogenous host defense systems to kill stress-resistant microbes.

Experimental procedures

Bacterial strains, plasmids and primers

All strains, plasmids and primers used in this study are listed in Supporting Information Table S3. A clean deletion

of *hslO*, *ppk* or *hslO/ppk* in *P. aeruginosa* strain PA14 was generated as previously described (Hmelo *et al.*, 2015). In brief, a 500 bp upstream and a 500 bp downstream region of the PA14 *hslO* and *ppk* gene were amplified by PCR, using the respective primer pairs (Supporting Information Table S3). Sowing overlap extension PCR was used to amplify the constructs *hslO*updwSOE and *ppk*updwSOE. The plasmids pEX18Gm*hslO*updwSOE and pEX18Gm*ppk*updwSOE were generated by integrating the constructs into the multicloning-site (MCS) of the allelic exchange vector pEX18Gm using *Bam*HI and *Hind*III restriction sites (Supporting Information Table S3). Plasmids were verified by PCR using MCS specific primers M13F-21 and M13R. For subsequent puddle mating with the respective PA14 strain, the plasmid constructs were transformed into the donor strain *E. coli* S17.1 (λ pir⁺) using standard chemical transformation. Transformed *E. coli* cells were plated onto LB agar plates supplemented with 10 μ g ml⁻¹ gentamicin and grown overnight at 37°C. The same day, an overnight culture of the recipient strain PA14 was prepared. The following day, an equal volume of fresh LB was added to the PA14 overnight culture, and the PA14 strain was incubated at 42°C for 3 h and until the *E. coli* donor strain had reached $A_{600} = 0.5$ – 0.6 . 1.5 ml of the *E. coli* donor strain and 0.5 ml of the PA14 recipient strain were mixed and centrifuged for 5 min at 10,000 *g* at room temperature. Cells were pelleted, resuspended in 50 μ l LB medium and transferred onto an LB plate and overnight incubation at 30°C. The bacterial lawn was scraped from the plate and resuspended in 1 ml of sterile PBS. Then 10 μ l, 100 μ l and 500 μ l aliquots of the resuspension were plated onto Vogel-Bonner Minimal Medium plates supplemented with 60 μ g ml⁻¹ gentamicin and the plates were incubated for 48 h at 37°C. For the counter-selection, an isolated colony from the VBMM plate was picked, streaked onto no-salt LB-plates supplemented with 15% (w/v) sucrose, and incubated for 36–48 h at 30°C. Colonies were PCR verified to confirm the deletion of the PA14 *hslO* and *ppk* genes using the respective sequencing primers (Supporting Information Table S3).

Preparation of HOCl, HOBr and HOSCN stock solutions

HOCl was prepared by diluting concentrated NaOCl into 40 mM potassium phosphate buffer (KPi), pH 7.5, and the concentration was determined in 10 mM NaOH using $\epsilon_{292 \text{ nm}} 350 \text{ M}^{-1} \text{ cm}^{-1}$ (Morris, 1966). HOBr was generated by mixing equal volumes of 40 mM HOCl and 45 mM KBr, and the concentration was determined in 10 mM NaOH using $\epsilon_{329 \text{ nm}} 332 \text{ M}^{-1} \text{ cm}^{-1}$ (Hawkins *et al.*, 2009). HOSCN was generated enzymatically as described previously with modifications (5). In brief, 26 mM NaSCN and 200 units ml⁻¹ lactoperoxidase (Sigma) were added to 2 mM aliquots of H₂O₂ in 40 mM KPi buffer, pH 7.5 over 5 min, followed by the addition of 500 units ml⁻¹ catalase. The solution was centrifuged at 14,000 *g* for 7.5 min at 4°C using a 10-kDa cutoff filter to remove proteins. The concentration of HOSCN was determined by monitoring the loss of signal at 412 nm ($\epsilon_{412 \text{ nm}} 14,150 \text{ M}^{-1} \text{ cm}^{-1}$), which occurs when HOSCN interacts with 5-thio-2-nitrobenzoic acid (Chandler *et al.*, 2013).

Growth conditions and bacterial survival assays

Unless specified otherwise, all strains were grown aerobically at 37°C in lysogenic broth (LB) or potassium morpholinopropanesulfonate (MOPS) minimal medium (Teknova, Inc.) containing 0.2% glucose, 1.32 mM potassium phosphate and 10 µM thiamine. For oxidative stress treatment, PA14 wild-type and the respective mutant strains were grown until $A_{600} = 0.4$ – 0.5 was reached. Then, the cells were washed with and resuspended in pre-warmed (i) phosphate buffer saline (PBS), pH 7.5, (ii) MOPS-glucose media or (iii) MOPS media, and treated with the indicated concentrations of HOCl, HOBr and HOSCN at either 30°C (PBS) or 37°C (MOPS-glucose; MOPS) respectively. Growth was recorded every 30–60 min for 10 h by measuring the optical density at 600 nm. To test for cell survival, 0.5 ml of bacteria were harvested by centrifugation at the indicated time points after normalizing them to their A_{600} , washed with MOPS medium containing 10 mM $\text{Na}_2\text{S}_2\text{O}_3$ to quench excess HOX, diluted into 0.85% NaCl and spotted onto LB agar using a Precision XS Microplate Sample Processor (Bio-Tek). The LB plates were incubated overnight at 37°C. To test the effects of mesalamine (Sigma Aldrich) treatment on the oxidative stress resistance of PA14 wild-type and *ppk* deletion strain, 500 µM mesalamine dissolved in MOPS media was added to the cultures 120 min before the stress treatment. Stress treatment with the respective oxidants and survival analysis was conducted as described above.

RNA_{Seq} library construction and sequence analysis

Wild-type *P. aeruginosa* PA14 was cultivated in MOPS-glucose medium at 37°C until OD_{600} of 0.4–0.5 was reached. Then, cells were treated with either 0.5 mM HOCl, 0.15 mM HOBr or 0.25 mM HOSCN. Cells equivalent to 1 ml with an $A_{600} = 0.4$ were harvested either before (control) or 15 min after the treatment onto 1 ml of ice-cold methanol (−80°C) to stop transcription. After centrifugation and removal of the supernatant, total RNA was prepared from three biological replicates of untreated and HOX-treated PA14 using the Ambion RiboPure-Bacteria Kit (Thermo Fisher Scientific) according to the manufacturer's instructions. The samples were DNase I treated, followed by depletion of rRNA using the Illumina Ribo Zero Kit (Illumina) for Gram-negative bacteria. Fifty base single end sequencing was performed on an Illumina HiSeq 2500 by the University of Michigan DNA Sequencing Core. Reads were aligned to the *P. aeruginosa* UCBPP-PA14 reference genome (Accession number: NC_008463.1) using bwa v. 0.7.8-r455 (Li and Durbin, 2009). Alignment files were generated using samtools (Li *et al.*, 2009). Subsequent analyses were performed using the statistical software package R. The R packages edgeR and limma were used for read count normalization and determining differential expression respectively (Robinson *et al.*, 2010; Ritchie *et al.*, 2015). The false discovery rate (fdr) was calculated via the Benjamini-Hochberg procedure (Benjamini and Hochberg, 1995). The threshold chosen for calling a gene differentially expressed was $\text{fdr} \leq 0.005$ and an absolute value of $\log_2(\text{fold change}) \geq 1.5$. Enrichment of differentially expressed

genes in each of 78 regulons was tested using one-sided Fisher's Exact Tests. *P*-values were adjusted for false discovery rate using the method of Benjamini and Hochberg (Benjamini and Hochberg, 1995). Information on *P. aeruginosa* regulons was taken from (Galan-Vasquez *et al.*, 2011), with additional information on the OxyR regulon supplemented from a recently published study (Wei *et al.*, 2012). Regulons determined to be significantly enriched for differentially expressed genes ($\text{FDR} < 0.05$) were presented as heatmaps using the heatmap function from the R package NMF. $\log_2(\text{Fold change})$ values of each treatment group compared to the untreated control group were used for clustering genes by their Euclidean distance with complete linkage.

In vivo aggregation assays

Preparation of cellular insoluble protein fractions was conducted according to (Tomoyasu *et al.*, 2001) with slight modifications. In brief, wild-type *P. aeruginosa* PA14 and mutant strains were grown in MOPS-glucose medium at 37°C to an $\text{OD}_{600} \sim 0.4$ – 0.5 and treated with the indicated concentrations of HOCl, HOBr and HOSCN for 30 min. Cells equivalent to 4 ml of $\text{OD}_{600} = 1$ were harvested by centrifugation and resuspended in 40 µl Buffer A (10 mM KPi, pH 6.5, 1 mM EDTA, 20% [w/v] sucrose, 1 mg/ml lysozyme, 50 U/ml benzonase), followed by a 30 min incubation on ice. The samples were then frozen at −80°C. After thawing the samples on ice and addition of 160 µl Buffer B (10 mM KPi, pH 6.5, 1 mM EDTA), ~ 100 µl 0.5 mm glass beads (BioSpec Products) were added, and the microfuge tubes were shaken for 30 min at 1400 rpm, 8°C, to lyse the cells completely. The samples were centrifuged for 20 min at $16,100 \times g$ and 4°C to separate protein aggregates from the soluble fraction. The pellets were washed once with Buffer B, subsequently with Buffer C (Buffer B plus 2% Nonidet P-40 [ICN Biomedicals]) to dissolve membrane proteins, and finally with Buffer B. The insoluble fractions were resuspended in SDS-Laemmli buffer and visualized by SDS-PAGE.

Analysis of in vivo protein aggregation upon HOX treatment and identification of protein aggregates by SILAC

Wild-type PA14 was cultivated in MOPS-glucose medium supplemented with either 120 µg/ml heavy [$^{13}\text{C}_6$]Arg or the isotopically light [$^{12}\text{C}_6$]Arg at 37°C until OD_{600} of 0.4–0.5 was reached. Then, cells were treated with either 0.5 mM HOCl or 0.25 mM HOSCN for 30 min. Cells equivalent to 4 ml of $\text{OD}_{600} = 1$ were harvested by centrifugation and resuspended in 40 µl Buffer A (10 mM KPi, pH 6.5, 1 mM EDTA, 20% [w/v] sucrose, 1 mg/ml lysozyme, 50 U/ml benzonase), followed by a 30 min incubation on ice. Preparation of cellular insoluble protein fraction was performed as described in the previous section. Aggregated proteins were dissolved in DAB buffer (200 mM Tris/HCl, pH 7.5, 6 M Urea, 10 mM EDTA, 0.5% SDS) and mixed in a 1:1 ratio. Protein mixtures were then processed by SDS-PAGE using a 10% Bis-Tris NuPAGE gel (Invitrogen). Protein

bands were cut out and in-gel digestion was processed on each using a robot (ProGest, DigiLab) with the following protocol: (i) washed with 25 mM ammonium bicarbonate followed by acetonitrile; (ii) reduced with 10 mM dithiothreitol at 60 °C followed by alkylation with 50 mM iodoacetamide at room temperature; (iii) digested with trypsin (Promega) at 37 °C for 4 h and quenched with formic acid, lyophilized and reconstituted in 0.1% trifluoroacetic acid. Samples were analyzed by nano-LC-MS/MS (MS Bioworks). Half of each digested sample was analyzed by nano LC-MS/MS with a Waters NanoAcquity HPLC system interfaced to a ThermoFisher Fusion Lumos. Peptides were loaded on a trapping column and eluted over a 75 µm analytical column at 350 nL/min; both columns were packed with Luna C18 resin (Phenomenex). A 2 h gradient was employed. The mass spectrometer was operated in data-dependent mode, with the Orbitrap operating at 60,000 FWHM and 17,500 FWHM for MS and MS/MS respectively. The instrument was run with a 3-s cycle for MS and MS/MS.

SILAC data were processed with MaxQuant 1.5.8.3 (Cox and Mann, 2008). Second peptide and Match between runs were enabled otherwise the default parameters were used. The multiplicity was set to 2, with Arg6 configured as the heavy label. Data were searched against UniProt *Pseudomonas aeruginosa* UCBPP-PA14 (2/14/2017). The MaxQuant output was further processed using Perseus. This involved filtering out contaminant and reverse protein identifications. The heat map in Fig. 3B was generated using the Hierarchical clustering function with Perseus, Row Trees were processed using the default settings.

Protein expression, purification and in vitro oxidation of PA14 Hsp33

The *hslO* gene from genomic *P. aeruginosa* PA14 was cloned into a pET11a vector using *Bam*HI and *Hind*III restriction sites. The plasmid was transformed into the *E. coli* expression strain BL21 (DE3) *hslO::kan*. PA14 Hsp33 was expressed and purified as described for *E. coli* Hsp33 (Graumann *et al.*, 2001). Reduced, chaperone-inactive PA14 Hsp33 (PA14-Hsp33_{red}) and HOX-treated PA14-Hsp33 (PA14 Hsp33_{HOX}) were prepared as previously described for *E. coli* Hsp33 using the indicated concentrations of HOCl, HOBr or HOSCN (Winter *et al.*, 2008).

Chaperone activity assays

Chaperone activity measurements with chemically denatured citrate synthase from porcine heart (CS, Sigma-Aldrich) were conducted as previously described (Ilbert *et al.*, 2007). In short, CS was denatured to a final concentration of 12 µM in 6.0 M Gdn·HCl, 40 mM HEPES, pH 7.5 overnight at room temperature. To initiate aggregation of CS, the unfolded enzyme was diluted 1:160 into 40 mM HEPES, pH 7.5 at 30 °C in the absence or presence of various concentrations of reduced or oxidized Hsp33. Light scattering was monitored using a Hitachi F4500 fluorimeter equipped with a thermostated cell holder and stirrer. Excitation and emission wavelengths were set to 360 nm, and the excitation and emission slit widths were set to 2.5 nm.

Extraction and quantification of polyphosphate (polyP)

Changes in intracellular polyP levels upon treatment with HOCl, HOBr or HOSCN were quantified as previously described (Dahl *et al.*, 2017) with minor modifications. PA14 wild-type and PA14 *ppk* strains were grown aerobically in 50 ml MOPS glucose medium and split into 10 ml cultures in new flasks when the culture reached an optical density at 600 nm of 0.4–0.5. Then, 0.5 mM HOCl, 0.15 mM HOBr or 0.25 mM HOSCN were added and cells sufficient to yield 200 µg total cellular protein were harvested by centrifugation after incubation for 2.5 h and lysed by incubation for 10 min at 95 °C in 0.25 ml GITC Lysis Buffer (4 M guanidine isothiocyanate, 50 mM Tris-HCl, pH 7.0). Protein content of each sample was determined using the Bradford assay (Bio-Rad). PolyP was extracted by sequential addition of 15 µl 10% sodium dodecyl sulfate, 0.5 ml 95% ethanol and 5 µl glassmilk (0.1 g/ml acid-washed silicon dioxide in GITC Lysis Buffer). This mixture was applied (1 min @ 3,000 × g) to silica membrane spin columns (Econospin™, Epoch Life Science, Inc.) and rinsed twice with 0.75 ml NW Buffer (5 mM Tris-HCl, pH 7.5, 50 mM NaCl, 5 mM EDTA, 50% v/v ethanol). After centrifuging once more to dry the membrane, polyP was eluted by adding 50 µl 50 mM Tris-HCl, pH 8, incubating for 15 min at room temperature and centrifuging to collect eluate. PolyP extracts were incubated for 1 h at 37 °C with 250 µM highly-pure ADP (Cell Technology), 50 mM HEPES (pH 7.5), 50 mM ammonium sulfate, 5 mM MgCl₂ in the presence or absence of 50 nM *E. coli* PPK. The resulting ATP was quantified with QuantiLum® Recombinant Luciferase (Promega). The reactions contained 50 mM Tricine buffer, pH 7.8, 10 mM MgSO₄, 0.2 mM EDTA, 0.2 mM sodium azide, 1 mM DTT, 100 µM luciferin, 25 nM luciferase and measurements were conducted in a FLUOstar Omega microplate reader (BMG Labtech). Values obtained from samples incubated in the absence of PPK were subtracted from the values of samples incubated in the presence of PPK and normalized to the values obtained for the untreated samples of each strain in order to calculate the fold-change in polyP accumulation in each sample.

Quantitative real-time PCR

Gene expression analysis by RT-PCR was performed upon HOSCN treatment of PA14 wild-type and PA14 *ppk* strains. Briefly, cells were grown aerobically to mid-log phase in MOPS glucose at 37 °C. HOSCN was added to a final concentration of 0.25 mM and incubated for 20 min. RNA was prepared using the NucleoSpin RNA kit (Macherey&Nagel) and DNA-free™ kit (Ambion). PrimeScript 1st strand cDNA Synthesis Kit (Takara) was used to generate cDNA, and RT-PCRs were set up with SYBR^(R) GreenER™ qRT-PCR mix (Invitrogen) and a Mastercycler^(R) realplex2 real-time PCR system (Eppendorf). Expression ratios were calculated compared with expression of each gene in untreated PA14 and PA14 *ppk* cultures respectively, by the $\Delta\Delta$ CT method (Pfaffl, 2001) and normalized to expression of *rrsD* (encoding 16S rRNA), expression of which did not change under the conditions tested. Primers used for RT-PCR analysis are listed in Supporting Information Table S3.

Acknowledgements

We thank Dao Ngyuen (McGill University) for providing us with protocols, plasmids and strains to construct the PA14 mutant strains. Wild-type PA14 was kindly provided by B. Boles. We are grateful to M. Gray for helpful discussions. We thank the University of Michigan DNA Sequencing Core for RNA sample processing and sequencing. The authors declare no conflict of interest. This work was funded by the National Institute of Health grants GM065318 and GM116582 to U.J.; J.-U. D. was supported by a postdoctoral fellowship from the Deutsche Forschungsgemeinschaft (DA 1697/1-1).

Author contributions

B.G. and J.-U.D. designed and carried out experiments. J.S. evaluated the RNA_{seq} analysis. B.G., J.-U.D. and U.J. conceived the study, interpreted the results, and wrote the paper.

References

- Arsene, F., Tomoyasu, T., and Bukau, B. (2000) The heat shock response of *Escherichia coli*. *Int J Food Microbiol* **55**: 3–9.
- Ashby, M.T. (2008) Inorganic chemistry of defensive peroxidases in the human oral cavity. *J Dent Res* **87**: 900–914.
- Atichartpongkul, S., Vattanaviboon, P., Wisitkamol, R., Jaroensuk, J., Mongkolsuk, S., and Fuangthong, M. (2016) Regulation of organic hydroperoxide stress response by two OhrR homologs in *Pseudomonas aeruginosa*. *PLoS One* **11**: e0161982.
- Barrett, T.J., and Hawkins, C.L. (2012) Hypothiocyanous acid: benign or deadly? *Chem Res Toxicol* **25**: 263–273.
- Barrett, T.J., Pattison, D.I., Leonard, S.E., Carroll, K.S., Davies, M.J., and Hawkins, C.L. (2012) Inactivation of thiol-dependent enzymes by hypothiocyanous acid: role of sulfenyl thiocyanate and sulfenic acid intermediates. *Free Radic Biol Med* **52**: 1075–1085.
- Benjamini, Y., and Hochberg, Y. (1995) Controlling the false discovery rate - a practical and powerful approach to multiple testing. *J Roy Stat Soc B Met* **57**: 289–300.
- Ceragioli, M., Mols, M., Moezelaar, R., Ghelardi, E., Senesi, S., and Abee, T. (2010) Comparative transcriptomic and phenotypic analysis of the responses of *Bacillus cereus* to various disinfectant treatments. *Appl Environ Microbiol* **76**: 3352–3360.
- Chandler, J.D., and Day, B.J. (2015) Biochemical mechanisms and therapeutic potential of pseudohalide thiocyanate in human health. *Free Radic Res* **49**: 695–710.
- Chandler, J.D., Nichols, D.P., Nick, J.A., Hondal, R.J., and Day, B.J. (2013) Selective metabolism of hypothiocyanous acid by mammalian thioredoxin reductase promotes lung innate immunity and antioxidant defense. *J Biol Chem* **288**: 18421–18428.
- Chi, B.K., Gronau, K., Mader, U., Hessling, B., Becher, D., and Antelmann, H. (2011) S-bacillithiolation protects against hypochlorite stress in *Bacillus subtilis* as revealed by transcriptomics and redox proteomics. *Mol Cell Proteomics* **10**: M111 009506.
- Cohen, T.S., and Prince, A. (2012) Cystic fibrosis: a mucosal immunodeficiency syndrome. *Nat Med* **18**: 509–519.
- Cox, J., and Mann, M. (2008) MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat Biotechnol* **26**: 1367–1372.
- Dahl, J.U., Gray, M.J., Bazopoulou, D., Beaufay, F., Lempart, J., Koenigsnecht, M.J., et al. (2017) The anti-inflammatory drug mesalamine targets bacterial polyphosphate accumulation. *Nat Microbiol* **2**: 16267.
- Dahl, J.U., Gray, M.J., and Jakob, U. (2015) Protein quality control under oxidative stress conditions. *J Mol Biol* **427**: 1549–1563.
- Das, D., De, P.K., and Banerjee, R.K. (1995) Thiocyanate, a plausible physiological electron donor of gastric peroxidase. *Biochem J* **305**: 59–64.
- Davies, M.J. (2011) Myeloperoxidase-derived oxidation: mechanisms of biological damage and its prevention. *J Clin Biochem Nutr* **48**: 8–19.
- Davies, M.J., Hawkins, C.L., Pattison, D.I., and Rees, M.D. (2008) Mammalian heme peroxidases: from molecular mechanisms to health implications. *Antioxid Redox Signal* **10**: 1199–1234.
- Dukan, S., and Touati, D. (1996) Hypochlorous acid stress in *Escherichia coli*: resistance, DNA damage, and comparison with hydrogen peroxide stress. *J Bacteriol* **178**: 6145–6150.
- Fargier, E., Mac Aogain, M., Mooij, M.J., Woods, D.F., Morrissey, J.P., Dobson, A.D., et al. (2012) MexT functions as a redox-responsive regulator modulating disulfide stress resistance in *Pseudomonas aeruginosa*. *J Bacteriol* **194**: 3502–3511.
- Galan-Vasquez, E., Luna, B., and Martinez-Antonio, A. (2011) The regulatory network of *Pseudomonas aeruginosa*. *Microb Inform Exp* **1**: 3.
- Gebendorfer, K.M., Drazic, A., Le, Y., Gundlach, J., Bepperling, A., Kastenmuller, A., et al. (2012) Identification of a hypochlorite-specific transcription factor from *Escherichia coli*. *J Biol Chem* **287**: 6892–6903.
- Graumann, J., Lilie, H., Tang, X., Tucker, K.A., Hoffmann, J.H., Vijayalakshmi, J., et al. (2001) Activation of the redox-regulated molecular chaperone Hsp33—a two-step mechanism. *Structure* **9**: 377–387.
- Gray, M.J., and Jakob, U. (2015) Oxidative stress protection by polyphosphate—new roles for an old player. *Curr Opin Microbiol* **24**: 1–6.
- Gray, M.J., Wholey, W.Y., Parker, B.W., Kim, M., and Jakob, U. (2013) NemR is a bleach-sensing transcription factor. *J Biol Chem* **288**: 13789–13798.
- Gray, M.J., Wholey, W.Y., Wagner, N.O., Cremers, C.M., Mueller-Schickert, A., Hock, N.T., et al. (2014) Polyphosphate is a primordial chaperone. *Mol Cell* **53**: 689–699.
- Hawkins, C.L., Morgan, P.E., and Davies, M.J. (2009) Quantification of protein modification by oxidants. *Free Radic Biol Med* **46**: 965–988.
- Hmelo, L.R., Borlee, B.R., Almlad, H., Love, M.E., Randall, T.E., Tseng, B.S., et al. (2015) Precision-engineering the *Pseudomonas aeruginosa* genome with two-step allelic exchange. *Nat Protoc* **10**: 1820–1841.

- Ilbert, M., Horst, J., Ahrens, S., Winter, J., Graf, P.C., Lilie, H., and Jakob, U. (2007) The redox-switch domain of Hsp33 functions as dual stress sensor. *Nat Struct Mol Biol* **14**: 556–563.
- Klebanoff, S.J. (2005) Myeloperoxidase: friend and foe. *J Leukoc Biol* **77**: 598–625.
- Klebanoff, S.J., Kettle, A.J., Rosen, H., Winterbourn, C.C., and Nauseef, W.M. (2013) Myeloperoxidase: a front-line defender against phagocytosed microorganisms. *J Leukoc Biol* **93**: 185–198.
- Kogan, I., Ramjeesingh, M., Li, C., Kidd, J.F., Wang, Y., Leslie, E.M., Cole, S.P., and Bear, C.E. (2003) CFTR directly mediates nucleotide-regulated glutathione flux. *EMBO J* **22**: 1981–1989.
- Lee, D.G., Urbach, J.M., Wu, G., Liberati, N.T., Feinbaum, R.L., Miyata, S., *et al.* (2006) Genomic analysis reveals that *Pseudomonas aeruginosa* virulence is combinatorial. *Genome Biol* **7**: R90.
- Li, H., and Durbin, R. (2009) Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**: 1754–1760.
- Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., *et al.* (2009) The sequence alignment/map format and SAMtools. *Bioinformatics* **25**: 2078–2079.
- Lister, P.D., Wolter, D.J., and Hanson, N.D. (2009) Antibacterial-resistant *Pseudomonas aeruginosa*: clinical impact and complex regulation of chromosomally encoded resistance mechanisms. *Clin Microbiol Rev* **22**: 582–610.
- Lorentzen, D., Durairaj, L., Pezzulo, A.A., Nakano, Y., Launspach, J., Stoltz, D.A., *et al.* (2011) Concentration of the antibacterial precursor thiocyanate in cystic fibrosis airway secretions. *Free Radic Biol Med* **50**: 1144–1150.
- Love, D.T., Barrett, T.J., White, M.Y., Cordwell, S.J., Davies, M.J., and Hawkins, C.L. (2016) Cellular targets of the myeloperoxidase-derived oxidant hypothiocyanous acid (HOSCN) and its role in the inhibition of glycolysis in macrophages. *Free Radic Biol Med* **94**: 88–98.
- Morris, J.C. (1966) The acid ionization constant of HOCl from 5 to 35°. *J Phys Chem* **70**: 3798–3805.
- Muller, A., Langklotz, S., Lupilova, N., Kuhlmann, K., Bandow, J.E., and Leichert, L.I. (2014) Activation of RidA chaperone function by N-chlorination. *Nat Commun* **5**: 5804.
- Parker, B.W., Schwessinger, E.A., Jakob, U., and Gray, M.J. (2013) The RclR protein is a reactive chlorine-specific transcription factor in *Escherichia coli*. *J Biol Chem* **288**: 32574–32584.
- Pattison, D.I., Davies, M.J., and Hawkins, C.L. (2012) Reactions and reactivity of myeloperoxidase-derived oxidants: differential biological effects of hypochlorous and hypothiocyanous acids. *Free Radic Res* **46**: 975–995.
- Pfaffl, M.W. (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res* **29**: e45.
- Rada, B. (2017) Interactions between neutrophils and *Pseudomonas aeruginosa* in cystic fibrosis. *Pathogens* **6**: 10. doi: 10.3390/pathogens6010010.
- Rao, S., and Grigg, J. (2006) New insights into pulmonary inflammation in cystic fibrosis. *Arch Dis Child* **91**: 786–788.
- Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015) limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* **43**: e47.
- Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010) edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**: 139–140.
- Skaff, O., Pattison, D.I., and Davies, M.J. (2009) Hypothiocyanous acid reactivity with low-molecular-mass and protein thiols: absolute rate constants and assessment of biological relevance. *Biochem J* **422**: 111–117.
- Skaff, O., Pattison, D.I., Morgan, P.E., Bachana, R., Jain, V.K., Priyadarsini, K.I., and Davies, M.J. (2012) Selenium-containing amino acids are targets for myeloperoxidase-derived hypothiocyanous acid: determination of absolute rate constants and implications for biological damage. *Biochem J* **441**: 305–316.
- Small, D.A., Chang, W., Toghrol, F., and Bentley, W.E. (2007) Toxicogenomic analysis of sodium hypochlorite antimicrobial mechanisms in *Pseudomonas aeruginosa*. *Appl Microbiol Biotechnol* **74**: 176–185.
- Tomoyasu, T., Mogk, A., Langen, H., Goloubinoff, P., and Bukau, B. (2001) Genetic dissection of the roles of chaperones and proteases in protein folding and degradation in the *Escherichia coli* cytosol. *Mol Microbiol* **40**: 397–413.
- van Dalen, C.J., Whitehouse, M.W., Winterbourn, C.C., and Kettle, A.J. (1997) Thiocyanate and chloride as competing substrates for myeloperoxidase. *Biochem J* **327**: 487–492.
- Wang, J., and Slungaard, A. (2006) Role of eosinophil peroxidase in host defense and disease pathology. *Arch Biochem Biophys* **445**: 256–260.
- Wei, Q., Minh, P.N., Dotsch, A., Hildebrand, F., Panmanee, W., Elfarash, A., *et al.* (2012) Global regulation of gene expression by OxyR in an important human opportunistic pathogen. *Nucleic Acids Res* **40**: 4320–4333.
- Winter, J., Ilbert, M., Graf, P.C., Ozcelik, D., and Jakob, U. (2008) Bleach activates a redox-regulated chaperone by oxidative protein unfolding. *Cell* **135**: 691–701.
- Winter, J., Linke, K., Jatzek, A., and Jakob, U. (2005) Severe oxidative stress causes inactivation of DnaK and activation of the redox-regulated chaperone Hsp33. *Mol Cell* **17**: 381–392.
- Winterbourn, C.C. (2008) Reconciling the chemistry and biology of reactive oxygen species. *Nat Chem Biol* **4**: 278–286.
- Winterbourn, C.C., Kettle, A.J., and Hampton, M.B. (2016) Reactive oxygen species and neutrophil function. *Annu Rev Biochem* **85**: 765–792.

Supporting information

Additional supporting information may be found in the online version of this article at the publisher's web-site.