

**Systolic Blood Pressure and Cardiovascular Outcomes in Heart Failure with Preserved
Ejection Fraction: An Analysis of the TOPCAT Trial**

Senthil Selvaraj, MD, MA; Brian Claggett, PhD; Sanjiv J. Shah, MD; Inder Anand, MD, FRCP, DPhil (Oxon); Jean L. Rouleau, MD; Akshay S. Desai, MD, MPH; Eldrin F. Lewis, MD, MPH; Bertram Pitt, MD; Nancy K. Sweitzer, MD; Marc A. Pfeffer, MD, PhD; Scott D. Solomon, MD.

From the Division of Cardiology, Department of Medicine, Hospital of the University of Pennsylvania, Philadelphia, PA (S.S.); Division of Cardiology, Department of Medicine, Brigham and Women's Hospital, Boston, MA (B.C., A.S.D., E.F.L., M.A.P., S.D.S.); the Division of Cardiology, Department of Medicine, Northwestern University Feinberg School of Medicine, Chicago, IL (S.J.S.); VA Medical Center and University of Minnesota, Minneapolis, MN (I.S.A); Department of Medicine, Montreal Heart Institute, University of Montreal, Quebec, Canada (J.L.R.); University of Michigan, Ann Arbor, MI (B.P.); and the Sarver Heart Center, University of Arizona College of Medicine, Tucson (N.K.S.).

Running title: Systolic Blood Pressure in HFpEF

Word count: 2900; Tables: 4; Figures: 2 (+2 Supplemental)

Address for correspondence:

Scott D. Solomon, MD
Brigham and Women's Hospital
75 Francis Street
Boston, MA 02115
Phone: 857.307.1954
Fax: 857.307.1944
E-mail: ssolomon@rics.bwh.harvard.edu

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1002/ejhf.1060](https://doi.org/10.1002/ejhf.1060)

Sponsor: National Institutes of Health, National Heart, Lung, and Blood Institute (NHLBI), Bethesda, MD (N01 HC45207)

ClinicalTrials.gov Identifier: NCT00094302

Author Manuscript

ABSTRACT (word count = 230)

Background: Recent guidelines have advocated for stricter systolic blood pressure (SBP) control in Heart Failure with Preserved Ejection Fraction (HFpEF), though data regarding the optimal SBP in HFpEF is sparse.

Methods and Results: We analyzed participants from the Americas from the Treatment of Preserved Cardiac Function Heart Failure with an Aldosterone Antagonist (TOPCAT) study with available baseline and 8-week visit SBP data (N=1645). We related baseline SBP to several efficacy and safety outcomes. To determine whether BP lowering was responsible for the potential beneficial effects of spironolactone observed in the Americas, we assessed the randomized treatment adjusting for baseline and change in 8-week SBP. The average age was 71.7 ± 9.7 years, 50% were women, and 79% were white. Patients in the lowest baseline SBP quartile were less often female, more often white, had lower body-mass index, lower baseline diastolic BP and pulse pressure, and more often had atrial fibrillation. After multivariable adjustment, there was no relationship observed between baseline SBP quartiles and any outcome. Spironolactone reduced SBP by 4.4 ± 0.6 mmHg compared to placebo (and consistently across baseline SBP quartiles). There was minimal change in the treatment effect for all outcomes after adjusting for baseline SBP and 8-week change in SBP.

Conclusions: No relationship was observed between baseline SBP quartiles and outcomes in TOPCAT. The anti-hypertensive effects of spironolactone did not account for the potential benefit in cardiovascular outcomes in the Americas.

Clinical Trials Registration Information: ClinicalTrials.gov identifier: NCT00094302

(<http://www.clinicaltrials.gov/ct2/show/NCT00094302>)

Keywords: heart failure with preserved ejection fraction; heart failure hospitalization; spironolactone; blood pressure

Author Manuscript

INTRODUCTION

Thus far, no interventions have been shown to reduce mortality in patients with heart failure with preserved ejection fraction (HFpEF), and therefore treatment has mainly focused on management of comorbidities.¹⁻³ Though expert opinion suggests management of systemic hypertension remains a cornerstone of therapy, and reduction of systolic blood pressure (SBP) has reduced cardiovascular events in the general population, there is a paucity of evidence to suggest the same effect in HFpEF. For example, while there are several clinical trials of BP lowering therapies in HFpEF,^{1,2,4} a dedicated trial of BP targets powered for cardiovascular events in HFpEF has not been performed. The relationship of SBP to cardiovascular events, likewise, has not been studied in stable HFpEF outpatients.⁵ The Systolic Blood Pressure Intervention Trial (SPRINT) demonstrated that intensive versus standard BP control improved cardiovascular outcomes, including incident HF, but this trial excluded patients with HF.⁶ Despite limited data, an SBP goal of less than 130 mmHg was recently recommended for HFpEF patients.⁷

Hypertension is a very common comorbidity in HFpEF, and the extent to which SBP control may influence clinical outcomes remains unclear. Control of BP in patients with HFpEF may reduce cardiovascular events by several mechanisms including improving hemodynamics, diastolic function, ventricular-arterial coupling, and left ventricular (LV) hypertrophy.⁸ In the Americas, spironolactone reduced the primary and several secondary endpoints in the Treatment

of Preserved Cardiac Function Heart Failure with an Aldosterone Antagonist (TOPCAT) study.⁹
¹⁰ Whether these effects were mediated through SBP reduction is unknown.

In this study, we first assessed the prognostic role of SBP in patients with HFpEF enrolled in the Americas in TOPCAT. Subsequently, we studied whether BP lowering effect of spironolactone was responsible for the potential beneficial effects observed in adverse cardiovascular event reduction in the Americas.

METHODS

TOPCAT study design and objectives

The design of the TOPCAT study have been described in detail previously.¹⁰ Briefly, TOPCAT was a multi-center, international, randomized, double blind, placebo-controlled trial of spironolactone in adults with HFpEF recruited from over 270 clinical sites. The trial was funded by the National Heart, Lung, and Blood Institute as a contract with the Brigham and Women's Hospital (Clinical Coordinating Center) and the New England Research Institute (Data Coordinating Center). Enrollment began in August 2006 and ended in January 2012, and the primary results of the trial were published in April 2014¹¹ (mean follow-up was 3.5 years). The primary aim of the TOPCAT study was to determine whether treatment with spironolactone, compared to placebo, can produce a clinically meaningful reduction in the composite outcome of cardiovascular mortality, aborted cardiac arrest, or HF hospitalization in adults with symptomatic HF and documented LVEF \leq 45%. All study participants provided written informed consent.

Inclusion criteria for TOPCAT were as follows: age \geq 50 years; diagnosis of HF based on at least one HF symptom at the time of study screening and at least one HF sign within the 12 months prior to screening; LVEF \geq 45% (per local reading); at least 1 HF hospitalization in the 12 months prior to study screening *or* B-type natriuretic peptide (BNP) $>$ 100 pg/ml or N-terminal pro-BNP $>$ 360 pg/ml (in the absence of an alternative explanation for elevated natriuretic peptide level) within the 60 days prior to screening; serum potassium $<$ 5.0 mmol/L prior to randomization.¹⁰ In addition, SBP was required to be controlled ($<$ 140 mmHg or 140-160 mmHg if the patient is taking at least 3 antihypertensive medications to control blood pressure). There were multiple exclusion criteria for TOPCAT, as detailed previously.¹⁰ Examples of exclusion criteria include severe systemic illness with a life expectancy of less than 3 years, severe chronic kidney disease, a history of significant hyperkalemia, known intolerance to aldosterone antagonists, and recent myocardial infarction, coronary artery bypass grafting, or percutaneous coronary intervention.

For the present study, we excluded participants 1) patients from Russia and Georgia (N=1678), given the significant regional differences previously described,⁹ and missing systolic BP at baseline or the 8-week visit (N=122). All hospitalizations were adjudicated by a clinical end-point committee at Brigham and Women's Hospital, blinded to study-drug assignments, according to pre-specified criteria.¹⁰ The primary endpoint of the study was the time to death from cardiovascular causes, aborted cardiac arrest, or HF hospitalization. Secondary endpoints included cardiovascular mortality, HF hospitalization, and recurrent HF hospitalization. Safety

outcomes included hyperkalemia (any serum potassium > 5.5 mEq/L), hypokalemia (any serum potassium < 3.5 mEq/L), and doubling of serum creatinine.

Statistical analysis

Baseline characteristics grouped by quartiles of baseline SBP were described using means±SD and medians and interquartile ranges or percentages as appropriate for the levels of measurement and distributions of the variables. We used quartiles of SBP rather than clinical cutoffs in order to avoid digit preference for BP readings ending in “0”. The SBP quartiles were compared using ANOVA for continuous variables and chi-squared tests (or Fisher’s exact test when appropriate) for categorical variables.

The association between SBP quartiles and the efficacy and safety outcomes were assessed using crude and multivariable-adjusted Cox regression. In a complementary analysis using restricted cubic splines, we examined the association between SBP and all outcomes. Four knots placed at the 5th, 35th, 65th, and 95th percentiles were used for all efficacy outcomes, while the safety outcomes were analyzed linearly. For recurrent HF hospitalization, we used negative binomial regression. Covariates were chosen based upon a combination of clinical relevance and previous prognostic implication in TOPCAT.¹² Multivariable models adjusted for New York Heart Association class, diabetes status, creatinine, heart rate, age, sex, race, smoking status, atrial fibrillation, peripheral arterial disease, ejection fraction, number of anti-hypertensive medications, and assignment to spironolactone vs. placebo.

We next determined the placebo-adjusted change in BP from baseline to the 8-week visit (the time at which maximal BP change occurred in the trial, as shown in **Figure 1**). Subsequent analyses were landmarked starting from the 8-week visit. To assess whether the treatment effect was independent of baseline SBP, an interaction term between treatment and continuous SBP was tested. Finally, to assess whether the change in SBP accounted for the beneficial effects of spironolactone, seen in the Americas, we generated Cox models assessing the relationship between treatment assignment and outcomes adjusting for baseline SBP and change in SBP between baseline and the 8-week visit. Analyses were performed using STATA version 12, and a two-sided p-value < 0.05 was considered statistically significant.

RESULTS

Baseline characteristics

Table 1 lists the baseline characteristics of the study population, stratified by quartiles of SBP. Of the initial study population of 1767 participants in the Americas, 1645 participants met the inclusion criteria for the present analysis. The mean baseline BP was 126±15 / 71±11 mmHg. The average age was 71.7±9.7 years, 50% were women, and 79% were white. Participants in the lowest quartile (SBP<118 mmHg) were less often female, more often white, had lower body mass index, lower baseline diastolic BP and pulse pressure, less frequently had hypertension and diabetes mellitus, and more often had atrial fibrillation (p<0.05 for all comparisons). These

participants were also less likely to use BP lowering therapy and more often had evidence of LV hypertrophy on electrocardiogram ($p<0.05$)

Table 2 shows event rates and crude and multivariable-adjusted hazard ratios for all outcomes, stratified by SBP quartiles (quartile 1 designated as the referent quartile). There was no relationship between SBP quartile and any outcome after multivariable adjustment. **Figure 2** shows event rates for both safety and efficacy outcomes by SBP quartile, stratified by randomization group, and demonstrates consistent effects of spironolactone across SBP quartiles. In a complementary analysis, **Supplementary Figures 1 and 2** shows the relationship between SBP and several efficacy and safety outcomes after multivariable adjustment. The relationship between the efficacy outcomes was non-linear ($p<0.05$ for non-linearity), whereas the relationship was linear for the safety endpoints ($p>0.05$ for non-linearity). A J-shaped relationship was observed between SBP as a continuous variable and the primary outcome and cardiovascular death ($p<0.05$ for overall relationship). The nadir risk appeared to occur around a SBP of 135 mmHg with the risk increasing below and above that cutoff. There was no relationship between SBP and any of the safety outcomes, though there was a trend toward an increased risk of doubling of serum creatinine with higher SBP ($p=0.056$)

In the overall population, spironolactone significantly reduced the systolic BP soon after start of the therapy (**Figure 1**) compared to placebo. By the 8-week visit, the SBP was -4.4 ± 0.6 mmHg lower than the placebo group (**Table 3**). The SBP lower remained lower in the spironolactone compared to placebo group by roughly the same amount through the 60 months

duration of the study (**Figure 1**). Interestingly, as seen in **Table 3**, spironolactone had a similar BP lowering effect in the 4 SBP baseline quartiles (P for interaction=0.30).

To determine whether the potential beneficial effects of spironolactone were mediated by BP reduction, we performed Cox regression (or negative binomial regression for recurrent HF hospitalization) adjusting for SBP and change in BP by the 8-week visit. As shown in **Table 4**, adjustment had little effect on the hazard ratios (or incidence rate ratio) for the composite endpoint, cardiovascular mortality, HF hospitalization, and recurrent HF hospitalization. There were no significant interactions between treatment group and baseline SBP or change in SBP (from baseline to 8-week visit) for any of the 4 efficacy outcomes (p for interaction >0.05 for all analyses).

DISCUSSION

In an analysis of the TOPCAT trial restricted to the Americas, there was no relationship between SBP quartiles and efficacy outcomes after multivariable adjustment. Additionally, the beneficial effects of spironolactone in the Americas were independent of baseline SBP, and the BP reduction by spironolactone accounted for only a small proportion of the beneficial effects observed in the trial. Our analysis questions the prognostic value of SBP, as well as the utility of SBP lowering, in HFpEF.

Our results are similar to findings observed in HF with reduced EF. Vasodilators, except for hydralazine and nitrates, consistently reduce SBP across the baseline SBP range.¹³⁻¹⁵ Patients

with the lowest baseline SBP assigned to hydralazine/nitrates showed a less significant decrease in SBP, which may be related to accompanied improved forward flow. In addition, similar to these studies, the beneficial effects of the therapies tested were independent of baseline SBP or change in SBP.

The J-shaped relationship observed in this analysis of HFpEF patients has been observed in several other high risk patient populations, including those with coronary artery disease and diabetes as well as previous stroke.^{16, 17} Additionally, in a post-hoc analysis of the Prospective comparison of ARNI with ACEI to Determine Impact on Global Mortality and morbidity in Heart Failure (PARADIGM-HF) trial, a randomized trial of patients with HF and reduced EF, the relationship between SBP and several outcomes was also noted to be J-shaped.¹⁸ It is difficult to know, however, whether the relationship is truly causal or confounded. For instance, low SBP in HFpEF may signify a sicker patient population. In addition, a low SBP may reflect low stroke volume from atrial fibrillation or perhaps a small LV cavity due to significant hypertrophy. Consistent with this notion, the lowest quartile SBP group had a higher frequency of atrial fibrillation and demonstrated lower pulse pressure, a surrogate of stroke volume.

As a result of the SPRINT Trial, professional society guidelines have recently recommended an SBP goal <130 mmHg, a deviation from the previously defined SBP goal of <140 mmHg.⁷ Notably, however, the SPRINT trial excluded patients with prevalent HF. While our primary analysis did not show significant relationships between SBP quartiles and adverse events, a complementary analysis using restricted cubic splines showed that a baseline SBP of

approximately 135 mmHg represented the lowest risk for adverse events. While it is impossible to suggest 135 mmHg as the goal of therapy given the non-randomized nature of this analysis, our results raise the question of the optimal SBP goal in HFpEF. Reducing SBP to <130 mmHg would likely require a significant increase in the use of anti-hypertensive therapy, which increases the risks associated with the anti-hypertensive therapy itself as well as polypharmacy. Therefore, a randomized trial of achieved SBP targets in HFpEF may be warranted.

We found that spironolactone modestly reduced BP in HFpEF. Interestingly, the amount of SBP reduction was less than that observed ALDO-DHF, another study of spironolactone in HFpEF [placebo-adjusted reduction 8 (95% CI 5-11) mmHg].¹⁹ This may be related to the differences in exclusion criteria. For instance, ALDO-DHF did not have similarly strict inclusion criteria for SBP as in TOPCAT, and therefore some patients may have had more significant hyperaldosteronism that was responsive to mineralocorticoid inhibition. The magnitude of BP reduction is similar to perindopril in HFpEF⁴, but less potent than angiotensin receptor neprilysin inhibition.²⁰ Furthermore, the SBP reduction in TOPCAT accounted for only a very small proportion of the adverse event reduction. The potential benefits of spironolactone in HFpEF are therefore multifactorial and go beyond reducing SBP. As an extension of this principle, in the echocardiographic substudy of TOPCAT, spironolactone did not significantly decrease LV mass index,²¹ which may partially underpin why SBP reduction in TOPCAT by spironolactone didn't relate to improved outcomes. Spironolactone, in other studies, decreases myocardial fibrosis, improves endothelial function and vascular compliance, and even reduces oxidative stress in the

failing heart.¹⁰ Accordingly, comorbidities have been suggested to induce a pro-inflammatory state in HFpEF, which induces reactive oxygen species from coronary microvascular endothelial cells. These events, in turn, limit nitric oxide bioavailability and decrease protein kinase G activity, accelerating LV remodeling and increasing collagen production, which worsens diastolic function.²² Thus, alternative biologic mechanisms may explain the potential beneficial effects of spironolactone and why SBP control may not be the primary driver of events in HFpEF.

Though hypertension is common in HFpEF and may drive a significant number of the pathophysiological disturbances, reducing BP when already reasonably controlled (SBP was <140 mmHg in the vast majority of patients by trial design) may not afford additional benefit. Our results are concordant with a study of angiotensin receptor neprilysin inhibition in HFpEF using surrogate endpoints, whereby improvement in several functional, laboratory, and echocardiographic parameters was also independent of change in SBP.²⁰

There are some limitations of the study. We used in office BP measures, while ambulatory BP monitoring may provide a more accurate assessment of the true treatment effect of spironolactone. In addition, because BP was well controlled in the trial, as an entry criterion, our results cannot be extrapolated to HFpEF patients with poorly controlled BP. Finally, we may have been underpowered to detect relationships between SBP and some of the less frequent efficacy or safety outcomes. Our analysis was restricted to the Americas due to regional

variation, but the majority of the events occurred in the Americas.⁹ Therefore, limitation of the cohort to the Americas likely resulted in only a small loss of statistical power.

In summary, in patients with HFpEF, we did not observe significant relationships between SBP quartiles and the vast majority of the efficacy and safety outcomes. In addition, spironolactone consistently reduced SBP across baseline SBP quartiles, and BP reduction explained only a small proportion of the effect of spironolactone on several cardiovascular outcomes. The potential benefits of spironolactone in HFpEF may thus be mediated by other mechanisms. Future clinical trials of SBP targets in HFpEF may be warranted.

DISCLOSURES

Dr Rouleau is a consultant for Novartis, Bayer and Astra-Zeneca. Dr Pfeffer has received consulting fees from Aastrom, Abbott Vascular, Amgen, Cerenis, Concert, Daiichi Sankyo, Fibrogen, Genzyme, GlaxoSmithKline, Hamilton Health Sciences, Medtronic, Merck, Novo Nordisk, Roche, Salix, Sanderling, Sanofi Aventis, Serono, Servier, and Teva, as well as research grants from New England Research Institute via subcontract from the National Institutes of Health, Amgen, Celladon, Novartis, and Sanofi-Aventis. The Brigham and Women's Hospital has patents for the use of inhibitors of the renin-angiotensin system in selected survivors of myocardial infarction with Novartis Pharmaceuticals on which Dr Pfeffer is a coinventor. His share of the licensing agreement is irrevocably transferred to charity. Dr Desai has received consulting fees from Novartis, Boston Scientific, Reata, Cardiomems, 5 am Ventures, Intel Corp, Coverys, and Relypsa, as well as research grants from AtCor Medical to support the Vascular

Stiffness Ancillary Study to the TOPCAT trial, for which he is listed as principal investigator. Dr Lewis has received research grants from the National Heart, Lung, and Blood Institute, Novartis, and Sanofi Aventis. Dr S. Shah has received research grants from the American Heart Association, National Institutes of Health, Actelion, AstraZeneca, Corvia, and Novartis, and consulting fees from Actelion, Amgen, AstraZeneca, Bayer, Boehringer-Ingelheim, Cardiora, Eisai, Gilead, Ironwood, Merck, MyoKardia, Novartis, Pfizer, Sanofi, and United Therapeutics. Dr Sweitzer has received research grants from the National Institutes of Health. Dr Pitt reports receiving consulting fees from Amorcyte, AstraZeneca, Aurasense, Bayer, BG Medicine, Gambro, Johnson & Johnson, Mesoblast, Novartis, Pfizer, Relypsa, and Takeda; receiving research grant support from Forest Laboratories; and holding stock in Aurasense, Relypsa, BG Medicine, and Aurasense. Dr Pitt also reports a pending patent related to site-specific delivery of eplerenone to the myocardium. Dr Solomon has received consulting fees from Novartis and Bayer and research grants from the National Heart, Lung, and Blood Institute. The other authors report no conflicts.

FUNDING SOURCE

This work was funded by the National Heart, Lung, and Blood Institute, National Institutes of Health, contract HHSN268200425207C. The content of this article does not necessarily represent the views of the National Heart, Lung, and Blood Institute or of the Department of Health and Human Services.

REFERENCES

1. Yusuf S, Pfeffer MA, Swedberg K, Granger CB, Held P, McMurray JJ, Michelson EL, Olofsson B, Ostergren J, Investigators C, Committees. Effects of candesartan in patients with chronic heart failure and preserved left-ventricular ejection fraction: the CHARM-Preserved Trial. *Lancet* 2003;**362**(9386):777-81.
2. Massie BM, Carson PE, McMurray JJ, Komajda M, McKelvie R, Zile MR, Anderson S, Donovan M, Iverson E, Staiger C, Ptaszynska A, Investigators IP. Irbesartan in patients with heart failure and preserved ejection fraction. *N Engl J Med* 2008;**359**(23):2456-67.
3. Pitt B, Pfeffer MA, Assmann SF, Boineau R, Anand IS, Claggett B, Clausell N, Desai AS, Diaz R, Fleg JL, Gordeev I, Harty B, Heitner JF, Kenwood CT, Lewis EF, O'Meara E, Probstfield JL, Shaburishvili T, Shah SJ, Solomon SD, Sweitzer NK, Yang S, McKinlay SM, Investigators T. Spironolactone for heart failure with preserved ejection fraction. *N Engl J Med* 2014;**370**(15):1383-92.
4. Cleland JG, Tendera M, Adamus J, Freemantle N, Polonski L, Taylor J, Investigators P-C. The perindopril in elderly people with chronic heart failure (PEP-CHF) study. *Eur Heart J* 2006;**27**(19):2338-45.
5. Shah SJ, Kitzman DW, Borlaug BA, van Heerebeek L, Zile MR, Kass DA, Paulus WJ. Phenotype-Specific Treatment of Heart Failure With Preserved Ejection Fraction: A Multiorgan Roadmap. *Circulation* 2016;**134**(1):73-90.

6. Group SR, Wright JT, Jr., Williamson JD, Whelton PK, Snyder JK, Sink KM, Rocco MV, Reboussin DM, Rahman M, Oparil S, Lewis CE, Kimmel PL, Johnson KC, Goff DC, Jr., Fine LJ, Cutler JA, Cushman WC, Cheung AK, Ambrosius WT. A Randomized Trial of Intensive versus Standard Blood-Pressure Control. *N Engl J Med* 2015;**373**(22):2103-16.
7. Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE, Jr., Colvin MM, Drazner MH, Filippatos GS, Fonarow GC, Givertz MM, Hollenberg SM, Lindenfeld J, Masoudi FA, McBride PE, Peterson PN, Stevenson LW, Westlake C. 2017 ACC/AHA/HFSA Focused Update of the 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *Circulation* 2017.
8. Lam CS, Shah AM, Borlaug BA, Cheng S, Verma A, Izzo J, Oparil S, Aurigemma GP, Thomas JD, Pitt B, Zile MR, Solomon SD. Effect of antihypertensive therapy on ventricular-arterial mechanics, coupling, and efficiency. *Eur Heart J* 2013;**34**(9):676-83.
9. Pfeffer MA, Claggett B, Assmann SF, Boineau R, Anand IS, Clausell N, Desai AS, Diaz R, Fleg JL, Gordeev I, Heitner JF, Lewis EF, O'Meara E, Rouleau JL, Probstfield JL, Shaburishvili T, Shah SJ, Solomon SD, Sweitzer NK, McKinlay SM, Pitt B. Regional variation in patients and outcomes in the Treatment of Preserved Cardiac Function Heart Failure With an Aldosterone Antagonist (TOPCAT) trial. *Circulation* 2015;**131**(1):34-42.
10. Desai AS, Lewis EF, Li R, Solomon SD, Assmann SF, Boineau R, Clausell N, Diaz R, Fleg JL, Gordeev I, McKinlay S, O'Meara E, Shaburishvili T, Pitt B, Pfeffer MA. Rationale and

design of the treatment of preserved cardiac function heart failure with an aldosterone antagonist trial: a randomized, controlled study of spironolactone in patients with symptomatic heart failure and preserved ejection fraction. *Am Heart J* 2011;**162**(6):966-972 e10.

11. Pfeffer MA, Pitt B, McKinlay SM. Spironolactone for heart failure with preserved ejection fraction. *N Engl J Med* 2014;**371**(2):181-2.
12. Solomon SD, Claggett B, Lewis EF, Desai A, Anand I, Sweitzer NK, O'Meara E, Shah SJ, McKinlay S, Fleg JL, Sopko G, Pitt B, Pfeffer MA, Investigators T. Influence of ejection fraction on outcomes and efficacy of spironolactone in patients with heart failure with preserved ejection fraction. *Eur Heart J* 2016;**37**(5):455-62.
13. Anand IS, Rector TS, Kuskowski M, Thomas S, Holwerda NJ, Cohn JN. Effect of baseline and changes in systolic blood pressure over time on the effectiveness of valsartan in the Valsartan Heart Failure Trial. *Circ Heart Fail* 2008;**1**(1):34-42.
14. Meredith PA, Ostergren J, Anand I, Puu M, Solomon SD, Michelson EL, Olofsson B, Granger CB, Yusuf S, Swedberg K, Pfeffer MA, McMurray JJ. Clinical outcomes according to baseline blood pressure in patients with a low ejection fraction in the CHARM (Candesartan in Heart Failure: Assessment of Reduction in Mortality and Morbidity) Program. *J Am Coll Cardiol* 2008;**52**(24):2000-7.
15. Anand IS, Tam SW, Rector TS, Taylor AL, Sabolinski ML, Archambault WT, Adams KF, Olukotun AY, Worcel M, Cohn JN. Influence of blood pressure on the effectiveness of a

fixed-dose combination of isosorbide dinitrate and hydralazine in the African-American Heart Failure Trial. *J Am Coll Cardiol* 2007;**49**(1):32-9.

16. Cooper-DeHoff RM, Gong Y, Handberg EM, Bavry AA, Denardo SJ, Bakris GL, Pepine CJ. Tight blood pressure control and cardiovascular outcomes among hypertensive patients with diabetes and coronary artery disease. *JAMA* 2010;**304**(1):61-8.

17. Ovbiagele B, Diener HC, Yusuf S, Martin RH, Cotton D, Vinisko R, Donnan GA, Bath PM, Investigators P. Level of systolic blood pressure within the normal range and risk of recurrent stroke. *JAMA* 2011;**306**(19):2137-44.

18. Bohm M, Young R, Jhund PS, Solomon SD, Gong J, Lefkowitz MP, Rizkala AR, Rouleau JL, Shi VC, Swedberg K, Zile MR, Packer M, McMurray JJV. Systolic blood pressure, cardiovascular outcomes and efficacy and safety of sacubitril/valsartan (LCZ696) in patients with chronic heart failure and reduced ejection fraction: results from PARADIGM-HF. *Eur Heart J* 2017;**38**(15):1132-1143.

19. Edelmann F, Wachter R, Schmidt AG, Kraigher-Krainer E, Colantonio C, Kamke W, Duvinage A, Stahrenberg R, Durstewitz K, Loffler M, Dungen HD, Tschope C, Herrmann-Lingen C, Halle M, Hasenfuss G, Gelbrich G, Pieske B, Aldo DHFI. Effect of spironolactone on diastolic function and exercise capacity in patients with heart failure with preserved ejection fraction: the Aldo-DHF randomized controlled trial. *JAMA* 2013;**309**(8):781-91.

20. Jhund PS, Claggett B, Packer M, Zile MR, Voors AA, Pieske B, Lefkowitz M, Shi V, Bransford T, McMurray JJ, Solomon SD. Independence of the blood pressure lowering effect

and efficacy of the angiotensin receptor neprilysin inhibitor, LCZ696, in patients with heart failure with preserved ejection fraction: an analysis of the PARAMOUNT trial. *Eur J Heart Fail* 2014;**16**(6):671-7.

21. Shah AM, Claggett B, Sweitzer NK, Shah SJ, Deswal A, Anand IS, Fleg JL, Pitt B, Pfeffer MA, Solomon SD. Prognostic Importance of Changes in Cardiac Structure and Function in Heart Failure With Preserved Ejection Fraction and the Impact of Spironolactone. *Circ Heart Fail* 2015;**8**(6):1052-8.

22. Paulus WJ, Tschope C. A novel paradigm for heart failure with preserved ejection fraction: comorbidities drive myocardial dysfunction and remodeling through coronary microvascular endothelial inflammation. *J Am Coll Cardiol* 2013;**62**(4):263-71.

FIGURE LEGENDS

Figure 1:

Title: *Average Systolic Blood Pressure during Follow-up by Treatment Arm.*

Caption: Average systolic blood pressure decreased to a maximum at the 8-week (2 month) visit in the spironolactone arm. Bars represent standard deviation.

Figure 2:

Title: *Event Rates by Systolic Blood Pressure Quartile.*

Caption: Event rates are displayed for safety and efficacy outcomes by systolic blood pressure quartile and stratified by assignment to spironolactone or placebo. Spironolactone demonstrated a consistent effect across the SBP quartiles.

TABLE 1. Baseline Clinical Characteristics by Systolic Blood Pressure Quartile

	SBP<118 mmHg N=403	118dSBP<128 mmHg N=373	128dSBP<138 mmHg N=403	SBPe138 mmHg N=466	P-value
SBP (mmHg)	107 ± 7	122 ± 3	132 ± 3	146 ± 7	
DBP (mmHg)	64 ± 9	70 ± 10	74 ± 10	77 ± 11	<0.001
Pulse pressure (mmHg)	43 ± 9	52 ± 9	58 ± 10	68 ± 13	<0.001
Randomization to spironolactone, n (%)	202 (50.1%)	191 (51.2%)	207 (51.4%)	225 (48.3%)	0.79
Age, years	71 ± 10	72 ± 9	72 ± 9	71 ± 10	0.41
Female, n (%)	186 (46.2%)	173 (46.4%)	192 (47.6%)	271 (58.2%)	<0.001
White race, n (%)	335 (83.1%)	307 (82.3%)	328 (81.4%)	332 (71.2%)	<0.001
NYHA Class III or IV, n (%)	143 (35.6%)	135 (36.3%)	125 (31.0%)	159 (34.1%)	0.41
Enrollment through HF hospitalization stratum, n (%)	225 (55.8%)	200 (53.6%)	208 (51.6%)	266 (57.1%)	0.39
Physical Characteristics					
Body mass index (kg/m ²)	32.8 ± 8.6	33.6 ± 8.6	34.0 ± 7.7	34.7 ± 8.1	0.008
Heart rate (beats/min)	69 ± 11	70 ± 11	69 ± 11	68 ± 12	0.2
Comorbidities, n (%)					
Hypertension	336 (83.4%)	320 (85.8%)	370 (92.0%)	450 (96.6%)	<0.001

Atrial fibrillation	199 (49.4%)	184 (49.3%)	175 (43.5%)	146 (31.3%)	<0.001
Diabetes mellitus	165 (40.9%)	161 (43.2%)	171 (42.5%)	235 (50.4%)	0.022
Myocardial Infarction	90 (22.3%)	74 (19.8%)	82 (20.4%)	95 (20.4%)	0.83
COPD	74 (18.4%)	51 (13.7%)	57 (14.2%)	88 (18.9%)	0.08
Asthma	46 (11.4%)	49 (13.1%)	35 (8.7%)	50 (10.7%)	0.26
Stroke	36 (8.9%)	26 (7.0%)	36 (9.0%)	49 (10.5%)	0.36
Peripheral arterial disease	52 (12.9%)	43 (11.5%)	43 (10.7%)	59 (12.7%)	0.75
Current smoker	23 (5.7%)	27 (7.2%)	23 (5.7%)	28 (6.0%)	0.79
Medication Use, n (%)					
ACE-I and/or ARB	309 (76.7%)	279 (75.0%)	307 (76.2%)	404 (86.7%)	<0.001
Beta-blocker	328 (81.4%)	309 (83.1%)	302 (74.9%)	356 (76.4%)	0.012
Calcium channel blocker	117 (29.0%)	125 (33.6%)	155 (38.5%)	237 (50.9%)	<0.001
Diuretic	355 (88.1%)	330 (88.7%)	348 (86.4%)	430 (92.3%)	0.039
Other anti-hypertensive medication	54 (13.4%)	49 (13.2%)	60 (14.9%)	106 (22.7%)	<0.001
Anti-hypertensive medications (n)	3.2 ± 1.2	3.1 ± 1.2	3.1 ± 1.2	3.6 ± 1.2	<0.001
Laboratory Testing					

Estimated glomerular filtration rate (mL/min/1.78 m ²)	60 ± 21	60 ± 20	60 ± 19	61 ± 20	0.68
Hemoglobin (mg/dL)	12.8 ± 1.7	13.0 ± 1.7	13.0 ± 1.7	12.8 ± 1.6	0.11
Electrocardiographic and Imaging Data					
Ejection fraction (%)	58 ± 8	58 ± 8	58 ± 8	59 ± 8	0.05
ECG Left ventricular hypertrophy (%)	27 (8.6%)	24 (8.1%)	47 (15.8%)	56 (16.6%)	<0.001
ECG atrial fibrillation (%)	132 (42.0%)	114 (38.3%)	105 (35.2%)	74 (22.0%)	<0.001

NYHA, New York Heart Association; SBP, systolic blood pressure; DBP, diastolic blood pressure; COPD, chronic obstructive pulmonary disease; ACE-I, angiotensin-converting-enzyme inhibitor; ARB, angiotensin II receptor blocker; HF, heart failure.

TABLE 2. Event Rates and Crude and Adjusted Hazard Ratios for Efficacy and Safety Outcomes by Systolic Blood Pressure Quartile

Efficacy outcomes, n (%)	SBP<118 mmhg N=403	118dSBP<128 mmHg N=373	128dSBP<138 mmHg N=403	SBPe138 mmHg N=466
Composite endpoint				
<ul style="list-style-type: none"> Event rate (per 100 person-years) 	13.0 (10.9, 15.4)	11.0 (9.1, 13.3)	9.6 (8.0, 11.6)	10.3 (8.6, 12.2)
<ul style="list-style-type: none"> Crude model HR (95% CI) 	ref	0.86 (0.66, 1.11)	0.75 (0.58, 0.96)	0.80 (0.62, 1.02)
<ul style="list-style-type: none"> Multivariable adjusted model HR (95% CI) 	ref	0.91 (0.70, 1.17)	0.84 (0.65, 1.07)	0.83 (0.65, 1.07)
Cardiovascular mortality				
<ul style="list-style-type: none"> Event rate (per 100 person-years) 	5.0 (3.8, 6.4)	4.7 (3.6, 6.1)	3.4 (2.6, 4.6)	3.3 (2.5, 4.4)
<ul style="list-style-type: none"> Crude model HR (95% CI) 	ref	0.94 (0.65, 1.37)	0.69 (0.46, 1.01)	0.67 (0.45, 0.98)
<ul style="list-style-type: none"> Multivariable adjusted model HR (95% CI) 	ref	0.95 (0.65, 1.39)	0.75 (0.51, 1.12)	0.71 (0.48, 1.06)
HF hospitalization				
<ul style="list-style-type: none"> Event rate (per 100 person-years) 	9.8 (8.1, 12.0)	7.9 (6.3, 9.9)	7.4 (6.0, 9.2)	8.7 (7.2, 10.5)
<ul style="list-style-type: none"> Crude model HR (95% CI) 	ref	0.81 (0.60, 1.09)	0.77 (0.58, 1.03)	0.88 (0.67, 1.16)

CI)				
• Multivariable adjusted model HR (95% CI)	ref	0.86 (0.64, 1.16)	0.88 (0.66, 1.18)	0.93 (0.70, 1.23)
Recurrent HF hospitalization				
• Event rate (per 100 person-years)	16.1 (14.0, 18.5)	14.6 (12.5, 17.0)	13.4 (11.6, 15.6)	15.2 (13.3, 17.4)
• Crude model HR (95% CI)	ref	0.92 (0.63, 1.34)	0.84 (0.58, 1.23)	0.91 (0.63, 1.30)
• Multivariable adjusted model HR (95% CI)	ref	0.81 (0.57, 1.16)	0.88 (0.62, 1.26)	0.83 (0.59, 1.18)
Hyperkalemia				
• Event rate (per 100 person-years)	6.8 (5.4, 8.6)	6.3 (4.9, 8.1)	5.9 (4.7, 7.5)	5.8 (4.6, 7.3)
• Crude model HR (95% CI)	ref	0.92 (0.65, 1.30)	0.90 (0.64, 1.26)	0.87 (0.62, 1.20)
• Multivariable adjusted model HR (95% CI)	ref	0.90 (0.64, 1.28)	0.91 (0.65, 1.28)	0.86 (0.61, 1.21)
Hypokalemia				
• Event rate (per 100 person-years)	6.4 (5.1, 8.2)	7.4 (5.9, 9.4)	6.5 (5.2, 8.2)	6.8 (5.5, 8.4)
• Crude model HR (95% CI)	ref	1.15 (0.82, 1.61)	1.05 (0.75, 1.46)	1.08 (0.78, 1.48)
• Multivariable adjusted model HR (95% CI)	ref	1.18 (0.84, 1.64)	1.10 (0.79, 1.54)	1.09 (0.78, 1.52)

Doubling of creatinine				
• Event rate (per 100 person-years)	4.9 (3.8, 6.4)	5.6 (4.4, 7.3)	3.9 (2.9, 5.2)	7.0 (5.7, 8.6)
• Crude model HR (95% CI)	ref	1.15 (0.79, 1.67)	0.80 (0.54, 1.18)	1.42 (1.01, 1.99)
• Multivariable adjusted model HR (95% CI)	ref	1.19 (0.82, 1.72)	0.86 (0.58, 1.28)	1.29 (0.91, 1.83)

TABLE 3. Placebo-adjusted Change in Systolic Blood Pressure at the 8 Week Visit by Treatment Arm

Baseline Systolic Blood Pressure Group	Treatment Effect of Spironolactone vs. Placebo	
	Change in SBP (95% CI)	P-value
All Patients	-4.4 (-6.2, -2.7)	<0.0001
SBP < 118 mmHg	-3.1 (-6.4, 0.2)	0.068
118 mmHg d SBP <1 28 mmHg	-6.2 (-9.3, -3.2)	0.0001
128 mmHg d SBP < 138 mmHg	-4.1 (-7.1, -1.0)	0.001
SBP e 138 mmHg	-5.2 (-8.6, -1.8)	0.003

Table 4. Effect of Change in Systolic Blood Pressure on the Efficacy of Spironolactone in Reducing Outcomes

Efficacy outcomes	Unadjusted Hazard Ratio Spironolactone vs. Placebo (95% CI)	P-value	Multivariable-Adjusted Hazard Spironolactone vs. Placebo (95% CI)*	P-value
Composite endpoint	0.82 (0.68, 0.99)	0.043	0.85 (0.70, 1.02)	0.08
Cardiovascular mortality	0.74 (0.56, 0.98)	0.037	0.73 (0.55, 0.96)	0.025
HF hospitalization	0.82 (0.66, 1.02)	0.07	0.85 (0.68, 1.06)	0.14
Recurrent HF hospitalization [^]	0.73 (0.56, 0.96)	0.022	0.76 (0.58, 0.99)	0.044

CI, confidence interval; HF, heart failure.

Analyses were landmarked from the 8-week visit.

*Adjusted for baseline systolic blood pressure and change in systolic blood pressure at the 8-week visit.

[^]Presented as incidence rate ratio (95% confidence interval).

Figure 1.




Figure 2.

