

**TITLE**

Differences in the Neighborhood Retail Food Environment and Obesity Among US Children and Adolescents by SNAP Participation

**AUTHORS**

Mary T. Gorski Findling,<sup>1</sup> Julia A. Wolfson,<sup>2</sup> Eric B. Rimm,<sup>3</sup> Sara N. Bleich<sup>4</sup>

**AFFILIATION**

<sup>1</sup> Interfaculty Initiative in Health Policy, Harvard University, Cambridge, Massachusetts, USA

<sup>2</sup> Departments of Health Management and Policy and Nutritional Sciences, University of Michigan School of Public Health, Ann Arbor, Michigan, USA

<sup>3</sup> Departments of Epidemiology and Nutrition, Harvard T.H. Chan School of Public Health, Boston, Massachusetts and Channing Division of Network Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA

<sup>4</sup> Department of Health Policy and Management, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA

**KEY WORDS**

Food environment; childhood obesity; Supplemental Nutrition Assistance Program

**RUNNING TITLE**

Neighborhood Food Environment & Childhood Obesity

**CONTACT INFORMATION**

Mary T. Gorski Findling, ScM

Interfaculty Initiative in Health Policy

Harvard University

14 Story Street

4<sup>th</sup> Floor

Cambridge, MA 02138

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of record](#). Please cite this article as [doi:10.1002/oby.22184](https://doi.org/10.1002/oby.22184).

Email: mgorski@mail.harvard.edu

Phone: 248-930-4586

Fax: 617-432-0092

### **WORD COUNT**

4019

### **FUNDING**

This study was funded by a pilot grant from the Hopkins Population Center, which is supported by a parent grant from the National Institute of Child Health and Human Development at NIH (R24HD042854). The funding source had no role in the design and conduct of the study, collection, management, analysis and interpretation of the data and preparation, review, or approval of the manuscript. M.T. Gorski Findling was supported by grant number T32HS000055 from the Agency for Healthcare Research and Quality. The content of this article is solely the responsibility of the authors and does not necessarily represent the official views of the Agency for Healthcare Research and Quality.

### **DISCLOSURE**

The authors declared no conflict of interest.

Author

## STUDY IMPORTANCE QUESTIONS

### What is already known about this subject?

- Some prior research suggests that the food retail environment may contribute to obesity risk among children and adolescents.
- The most consistent findings among children and adolescents are from studies where the availability of convenience stores and fast food restaurants are associated with higher BMI, and the availability of supermarkets is associated with lower BMI.
- However, several well-designed studies in both children and adults found no relationship between neighborhood food environment and BMI.

### What does this study add?

- To our knowledge, this is the first national study on the relationship between comprehensive neighborhood retail food access and obesity among all children and adolescents ages 2-18.
- This study addresses a key unanswered question of whether the relationship between children's food access and body mass index differs between children who live in households participating in the Supplemental Nutrition Assistance Program compared to those who do not, while also accounting for household food purchasing patterns.

## ABSTRACT

### Objective

To understand the association between children's neighborhood food access and overweight/obesity in a national sample of U.S. households, and whether it differs by Supplemental Nutrition Assistance Program (SNAP) participation or household purchases.

### Methods

Data were from the National Household Food Acquisition and Purchase Survey (2012–2013; n=3,748 children ages 2-18). We used logistic regression to examine associations between neighborhood retail food access ( $\leq 1$ -mile from home), food purchases (including sugary beverages), and overweight/obesity, stratified by SNAP status (1,720 participants, 453 eligible non-participants, 1,575 SNAP ineligible). Store types included: supermarkets/grocery, combination grocery/other (independent drug, dollar, and general stores), convenience, fast food, and non-fast food restaurants.

### Results

Odds of childhood overweight/obesity (OR [95% CI]) were higher with greater access to combination grocery/other stores overall (1.10 [1.03, 1.17]) and for children in SNAP (1.14 [1.05, 1.24]). Eligible non-SNAP children had higher odds of overweight/obesity with greater access to convenience stores (1.11 [1.04, 1.18]). The average child lived in a household with 6.3% of total spending at food outlets on sugary beverages (SNAP: 8.3%, eligible non-SNAP: 7.7%, SNAP ineligible: 5.5%).

### Conclusions

Greater neighborhood access to combination grocery/other stores is associated with higher obesity prevalence for children overall and those in SNAP.

## Introduction

Overweight and obesity affect one in five children in the U.S.,<sup>1</sup> disproportionately impacting racial and ethnic minorities, low-income children, and those living in southeastern states.<sup>2</sup> Increasingly, public health and policy efforts have focused on the neighborhood food environment as a key contributor to poor diet and high body mass index (BMI).<sup>3-5</sup>

Prior research suggests that the food retail environment may contribute to obesity risk.<sup>3, 6</sup> The most consistent findings among children and adolescents are from studies where the availability of convenience stores<sup>7-10</sup> and fast food restaurants<sup>11, 12</sup> are associated with higher BMI, and the availability of supermarkets is associated with lower BMI.<sup>7, 12, 13</sup> However, several well-designed studies in both children and adults found no relationship between neighborhood food environment and BMI.<sup>14-17</sup> In addition, a key limitation of past work is that few studies consider all types of food outlet access simultaneously.<sup>10</sup> To our knowledge, there have been no national studies on the relationship between comprehensive neighborhood retail food access and obesity among all children ages 2-18.

The relationship between the food environment and obesity may be especially important for low-income children, who often have limited access to healthy, affordable food.<sup>6, 18</sup> Research among low-income children shows inconsistent relationships between access to food outlets and BMI.<sup>19, 20</sup> Other research has shown some differences in households' food purchasing patterns by participation in the federal Supplemental Nutrition Assistance Program (SNAP), which is designed to support nutrition for low-income households through food assistance.<sup>21, 22</sup> SNAP households spend less per capita on food overall each week and have different food acquisition patterns than non-SNAP households,<sup>23</sup> including a higher percentage of food spending on sugary beverages.<sup>22</sup> In addition, the absolute dollar amount of sugary

beverage spending using SNAP benefits is higher than spending on sugary beverages without SNAP benefits.<sup>24</sup> Sugary beverages are a major source of empty calories in U.S. children's diets,<sup>25</sup> and children's consumption has been consistently linked with excess calorie intake and weight gain.<sup>26, 27</sup> Since almost half of all SNAP recipients (47%) are children,<sup>21</sup> we sought to answer whether SNAP eligibility and participation modifies the relationship between children's food access and weight.

Our study addressed key gaps in the literature by examining the relationship between neighborhood retail food outlets (number and type) and childhood obesity prevalence, stratified by SNAP participation. We additionally examined whether household and adolescent (ages 11-18) spending at specific types of food outlets, or on specific foods (sugary beverages), explained this relationship.

## Methods

Data were obtained from the Food Acquisition and Purchase Survey (FoodAPS), collected by the United States Department of Agriculture (USDA) in 2012-2013 and released in 2015. FoodAPS is the first nationally representative dataset of all households, SNAP households, and SNAP-eligible households in the U.S. that contains comprehensive data on food purchases and acquisitions.<sup>28</sup> Data were collected from in-person interviews, telephone interviews, survey booklets, scanners, and receipts over a seven-day period for all individuals (N=14,317) in 4,826 surveyed households. All household members 11 years and older were asked to report all weekly food purchases and free acquisitions, including events at home and away from home, by scanning barcodes for packaged foods, submitting receipts for stores and restaurants, and filling out daily food booklets. The primary respondent in each household reported food acquisitions for children under 11.

Children ages 2-18 were included in this analysis, excluding those missing data on BMI or SNAP status (n=413). The final analytic sample included 3,748 children from 1,942 households, including 1,720 children from 850 households participating in SNAP.

### *Dependent Variable*

Childhood overweight/obesity was the dependent variable, measured through self-reported height and weight of the child by each household's primary respondent, and defined as BMI  $\geq$ 85<sup>th</sup> percentile compared to age- and sex-specific Centers for Disease Control and Prevention 2000 growth charts. Because parent-reported child height and weight tends to result in an overestimation of childhood overweight/obesity prevalence for young children and overweight/obesity estimates were higher for children ages 2-5 years in FoodAPS compared to NHANES,<sup>29-31</sup> we conducted sensitivity analyses to account for known biases, including adjusting estimates using NHANES measured height/weight data<sup>29</sup> and restricting analyses to only children ages 6-18.<sup>30</sup>

### *Independent Variable*

The independent variable of primary interest was neighborhood food access, defined as the number of food outlets (stores and restaurants) within 1 mile (straight-line distance) of the child's home, approximating walking distance. This measure follows prior studies and USDA's definition of "low access" neighborhoods.<sup>14, 32, 33</sup> Access was reported for the following mutually exclusive food outlet types identified by USDA: superstores, supermarkets, and large/medium grocery stores; combination grocery

and other stores (including independent drug stores, dollar stores, and general stores); convenience stores; fast food restaurants, and non-fast food restaurants. Appendix A includes details on store types and definitions.

Alternate neighborhood food access measures tested in sensitivity analyses were food exposure (county-level density of different store types per 1,000 people), store type of closest SNAP retailer, number of household visits per week (by store type), and alternate access distances of 0.25, 0.5, and 2 miles.

### *Covariates*

Analyses were stratified by SNAP participation, defined for each child as at least one household member reporting that they currently participate in SNAP. For this analysis, we divided children in non-SNAP households into income-eligible ( $\leq 130\%$  FPL) and income-ineligible ( $>130\%$  FPL) categories.

Household food purchases and acquisitions were a key covariate used in this analysis, measured through a combination of event-level food acquisition data (i.e., each recorded visit to a location where food was purchased or acquired for free is an ‘event’) and item-level food acquisition data (i.e., an itemized list of every food and beverage acquired at each event). We created several measures of food acquisitions and purchases, with full definitions in Appendix B. These include the mean number of weekly food acquisition events per household for each child, mean weekly spending at food outlets per household member, and the percent of weekly spending at food outlets by store type for each child. To capture a measure of spending on unhealthy items, we also examined the mean of each child’s household sugary beverage spending as a percentage of total weekly food spending (mean of ratios). We also examined



individual purchases of adolescents ages 11-18, to understand their acquisitions separately from their households and to potentially account for behavioral effects.

Other demographic covariates included children's individual characteristics (age, gender, race/ethnicity), household characteristics (primary respondent's BMI, marital status, education, and work status, self-reported household income as a percent of the federal poverty level, food security based on USDA's 30-day Adult Food Security Scale, household participation in the Special Supplemental Nutrition Program for Women, Infants, and Children (WIC), urban or rural area of residence, and household vehicle access), and neighborhood characteristics (closest SNAP-authorized retailer by store type, county-level poverty rate of households). Household income and county-level poverty allowed examination of SNAP participants along the income gradient and whether the income at the area of residence mattered.

#### *Data Analysis*

First, we calculated group differences on demographic characteristics using chi-squared tests for categorical variables and regression for continuous variables, with post-hoc pairwise comparison of means. We then calculated unadjusted differences in children's food access and household food purchases and acquisitions using the same methods. Then, we estimated the association between food access and childhood obesity, overall and by SNAP status, with binary logit models that controlled for individual, household, and neighborhood covariates. All analyses included the use of sample weights provided by USDA to account for unequal probability of selection from oversampling, non-response, and stratified multistage probability sample design. Standard errors have been estimated to account for the complex sample design of the survey. Data analysis was performed using STATA v.14 (StataCorp LP, College Station, TX). As a secondary analysis, we examined child overweight/obesity by race/ethnicity, gender, and age (2-5 years, 6-10 years, 11-18 years) to understand heterogeneity among different subgroups.

For sensitivity analyses, we used multivariate ordinary least squares regression to analyze the relationship between food access and BMI percentile. To examine whether the 1-mile access measure was robust to alternate specifications, we conducted separate regressions for distances of 0.25, 0.5, and 2 miles, and we ran analyses restricted to children living in urban areas (a 1-mile radius may be inappropriate for rural children). We also included additional controls such as distance to the nearest store, county-level store density per 1,000 residents, county-level store totals, number of household events per week by store type, and month of the year to adjust for seasonal differences in purchasing patterns. We also ran analyses restricted to children in households where the primary respondent reported living at this address for 2+ years (since 2009), to examine the relationship between food access and childhood obesity only in a group with potentially more consistent recent neighborhood exposures and food access. To ensure that data were missing at random, we also examined whether children excluded from the sample due to missing BMI data differed from included children by demographic characteristics.

## Results

Table 1 displays the key individual, household, and neighborhood demographic characteristics of children in the FoodAPS sample, by children's household SNAP participation. The study sample included 1,720 children living in households participating in SNAP (SNAP children hereafter), 453 children living in households eligible for, but not participating in SNAP (non-SNAP children), and 1,575 children living in households ineligible for SNAP (SNAP-ineligible children). Children in SNAP had lower reported overweight/obesity prevalence (43.9%) than eligible non-SNAP children (50.0%,  $p=0.04$ ), but higher overweight/obesity prevalence than SNAP-ineligible children (33.8%,  $p<0.01$ ). Children in the three groups also differed in several household demographic characteristics, including the primary respondent's

marital status, education, work status, income, and WIC participation. SNAP and eligible non-SNAP children were more likely to live in urban areas (76.9% and 80.6%) than SNAP-ineligible children (61.1%,  $p<0.01$ ), and they were more likely to have convenience stores as their closest neighborhood SNAP-authorized retailer ( $p<0.01$ ). SNAP children also lived in counties with higher household poverty rates (15.5%) than eligible non-SNAP children (14.3%,  $p=0.03$ ) and SNAP-ineligible (12.8%,  $p<0.01$ ) children.

Table 2 shows unadjusted measures of food access and acquisitions/purchases by household SNAP participation. SNAP-ineligible children had fewer retail food outlets of all types within 1 mile of their home compared to SNAP and eligible non-SNAP children. SNAP-ineligible children also had different household food purchasing and acquisition patterns than SNAP and eligible non-SNAP children. SNAP-ineligible children had higher mean weekly household spending at food outlets per household member (\$58.87) compared to SNAP children (\$37.41,  $p<0.01$ ) and eligible non-SNAP children (\$31.51,  $p<0.01$ ). SNAP-ineligible children also lived in households with different weekly food expenditures by store type than SNAP children (e.g., higher percentage of overall spending at restaurants and lower percentage of overall spending at supermarkets, combination grocery and other stores, and convenience stores). For adolescent spending, we found that among adolescents with any food spending ( $N=800$ ), there were not significant differences in mean weekly spending among SNAP, eligible non-SNAP, and SNAP-ineligible adolescents (data not shown).

For sugary beverage spending, the average child lived in a household with 6.3% of total spending at food outlets on sugary beverages (SNAP: 8.3%, eligible non-SNAP: 7.7%, SNAP ineligible: 5.5%) (see Table S1). At convenience stores, 26.1% of the average child's total household food spending was on sugary beverages (SNAP: 29.8%, eligible non-SNAP: 15.5%, SNAP ineligible: 25.6%). At combination grocery/other stores, 13.3% of the average child's total household food spending was on sugary beverages

(SNAP: 18.4%, eligible non-SNAP: 17.7%, SNAP ineligible: 10.4%). Among adolescents with food spending, the average adolescent spent 10.0% of their individual weekly spending on sugary beverages (SNAP: 18.1%, eligible non-SNAP: 7.6%, SNAP ineligible: 8.9%). While the average child and adolescent in SNAP households spent a higher percentage of their food budget on sugary beverages than non-SNAP households, absolute dollars spent on sugary beverages was similar across groups (see Table S2).

When we estimated the association between food access (the number of each type of food retail outlet within 1 mile of the child's home) and childhood overweight/obesity, controlling for individual, household, and neighborhood-level covariates, the odds of a child being overweight/obese increased with each additional combination grocery/other store within 1 mile of a child's home (OR: 1.10, 95% CI: 1.03, 1.17) (shown in Table 3). Access to other store types was not significantly associated with childhood obesity in overall models. Children in SNAP households had higher odds of overweight/obesity with greater access to combination grocery/other stores (OR: 1.14, 95% CI: 1.05, 1.24). Eligible non-SNAP children had higher odds of overweight/obesity with greater access to convenience stores (OR: 1.11, 95% CI: 1.04, 1.18). Table S3 shows the full overall model, including all covariates.

Results for combination grocery/other stores remained significant when BMI percentile was used as a continuous outcome measure (combination grocery/other store  $\hat{\beta}$ =1.40, 95% CI: 0.62, 2.18). Combination grocery/other store access remained significant for children in SNAP (OR: 1.24, 95% CI: 1.04, 1.48) within a 0.5 mile radius and for eligible non-SNAP children (OR: 2.30, 95% CI: 1.08, 4.90) within a 0.25 mile radius. Results for convenience stores remained significant for eligible non-SNAP children (OR: 1.24, 95% CI: 1.053, 1.50) within 0.5 mile radius. There were no other statistically significant

relationships between any store types and overweight/obesity when access was set at 0.25 miles or 2 miles.

For adolescents ages 11-18, greater access to combination grocery/other stores was associated with higher overall odds for adolescent overweight/obesity, even after adjusting for adolescent spending at different food outlets and spending on sugary beverages (OR: 1.12, 95% CI: 1.02, 1.22). Access to combination grocery/other stores within 1 mile was associated with obesity for children living in urban areas (OR: 1.11, 95% CI: 1.04, 1.19) and for non-Hispanic black children (OR: 1.33, 95% CI: 1.04, 1.70), but not for children living in rural areas (OR: 1.25, 95% CI: 0.76, 2.07), non-Hispanic white children (1.07, 95% CI: 0.95, 1.20), or Hispanic children (OR: 1.09, 95% CI: 0.97, 1.21).

There was no significant relationship between any store type and overweight/obesity among 2-5 year old children, the group most likely to be affected by errors in parent-reported BMI.<sup>29-31</sup> The relationship between combination grocery/other stores and odds of overweight/obesity for all children remained significant after adjusting overweight/obesity prevalence for children 2-5 using NHANES measured height/weight data (adjusted OR: 1.10, 95% CI: 1.02, 1.18). Combination grocery/other store access also remained significant when children 2-5 were excluded from analysis (N=2,803 children 6-18 years; OR: 1.09, 95% CI: 1.02, 1.18).

For children living in their current house for at least two years, odds of overweight/obesity was associated with access within 1 mile to combination grocery/other stores (N=2,221; OR: 1.11, 95% CI: 1.03, 1.20), but this was not the case for children living at their current address for less than two years. Alternate access measures of food exposure (county density, number of stores in county, and closest SNAP retailer) were not associated with child overweight/obesity. Results were robust when these variables were

included as covariates and when we controlled for percent of free acquisitions and survey month (not included in final models).

## **Discussion**

We analyzed a unique national sample of U.S. children with rich information on their neighborhood retail food environment and household food spending to examine the relationship between food access and childhood obesity, and whether it differed by SNAP participation and household purchases. We found that access to combination grocery/other stores (which includes independent drug stores, dollar stores, and general stores) is associated with higher childhood obesity prevalence overall, and this association is particularly strong for children living in households participating in SNAP. We also found that greater access to convenience stores was associated with higher overweight/obesity prevalence for eligible non-SNAP children. Main results were robust to alternate model specifications, and there was a stronger association between combination grocery/other store access and overweight/obesity for adolescents ages 11–18, children living in urban areas, and non-Hispanic black children.

To our knowledge, previous studies have not included combination grocery/other stores as a unique store type when examining the food retail environment. Store type definitions, combined with the population studied (all children 2-18 vs. restricted child ages), access boundaries set (0.25, 0.5, 1, and 2 miles vs. other radii), and children's household geographic location (nationally representative vs. concentrated in specific states or cities) are all plausible explanations for why our results do not replicate previous studies finding positive associations between fast food restaurants and BMI and negative associations between supermarkets and BMI. It is likely that our finding among combination grocery/other stores is most consistent with previous study findings on associations between convenience stores and higher BMI,

since the definition of combination grocery/other stores includes some store types previous studies may have considered convenience (e.g., independent drug stores, dollar stores). Different classifications make comparisons difficult between studies, given the heterogeneity in store types. Distinguishing among different store types is important for future studies, since combination grocery/other stores make up 22% of all SNAP-authorized retailers in the U.S., while convenience stores make up 45%.<sup>34</sup> Together, these two store types account for more than 10% of all redeemed SNAP benefits and 5–9% of all children's total household weekly spending at food outlets, making them a substantial contributor to the food environment. Other studies have examined these store types as small grocery stores or non-traditional food stores (including corner/small grocery stores, gas stations, pharmacies, and dollar stores), working under the hypothesis that high availability of unhealthy food at the neighborhood level may lead to less healthy, higher-calorie diets in those neighborhoods.<sup>35, 36</sup> Consistent with our results, previous studies have found that small grocery store availability is associated with higher adult BMI,<sup>37, 38</sup> and that small grocery/non-traditional food retailers sell a high proportion of unhealthy options (e.g., in one sample, 0% of dollar stores and only 23% of pharmacies sold fresh vegetables).<sup>36</sup>

Our results also showed that among those who shopped at combination grocery/other stores and convenience stores, the average child's household spent 13% of weekly combination grocery store spending and 26% of weekly convenience store spending on sugary beverages. These results demonstrate significant household purchases of empty calories from these limited store types. Sugary beverage spending was also significant among all adolescents, and especially SNAP adolescents. Since there are often high amounts of sugar-sweetened beverages<sup>39</sup> and low amounts of few fruits/vegetables or whole grains purchased at small grocery and non-traditional food stores,<sup>40</sup> adolescents in particular may be incentivized to reduce sugary beverage purchases by providing easily understandable caloric information on prominent display at the point of purchase (e.g., signs on beverage cases with sugary beverage calories presented as physical activity equivalents).<sup>41</sup> In addition, practices such as offering a greater variety and

more shelf space for fruits/vegetables, having produce visible at the store entrance, and improving the ratio of shelf space for healthy versus less healthy items is associated with higher fruit/vegetable purchases at these store types.<sup>40</sup> This suggests that modifying store practices and offerings without displacing these store types may have a positive impact on children's diets and BMI.

This study makes several contributions to existing literature on food access and childhood obesity. Our results indicate a consistent association between some measures of the food environment (access to combination grocery/other stores) and children's weight, even after controlling for important covariates. We also found an association between convenience store access and overweight/obesity for eligible non-SNAP children. However, it is unclear which specific food outlets drive these relationships, as SNAP-authorized combination grocery/other stores and convenience stores are heterogeneous in their availability and stock of food items. For example, grocery stores and stores in higher income neighborhoods tend to offer a greater variety of healthful foods, while convenience/small food stores and stores in lower income neighborhoods tend to offer less healthful options.<sup>42</sup> Our robust finding of the positive relationship between number of combination grocery/other stores and childhood obesity may also reflect the types of neighborhoods which are characterized by these types of outlets (e.g., lower-income SNAP and eligible non-SNAP children are more likely to live in urban environments with a greater number of food outlets of all types compared to higher-income SNAP ineligible children). In other words, combination grocery/other stores may serve as a proxy for other unmeasured factors in the neighborhood environment which contribute to obesity risk. Consistent with this, previous research has found that socioeconomically disadvantaged and minority neighborhoods may draw less healthy food outlets—including fast food restaurants and smaller grocery stores—over time.<sup>43</sup> Future research should explore this possibility. In addition, our results suggest that further research is needed to better understand observed patterns in sugary beverage purchases among SNAP participants.



### *Limitations*

There are several important limitations of this study, including the cross-sectional nature of the data, which limits our ability to draw causal conclusions about the food environment and childhood obesity. Access measures tested may not account for children's neighborhood food environments in all settings (e.g., sprawling or rural areas). Specifically, our sensitivity analysis found that the 1-mile access measure was significant for urban but not rural children, which may reduce the generalizability of our results to children living in rural areas. In addition, there are known biases using parent-reported child height and weight, particularly for height of young children (2-5 years). However, our results were robust to sensitivity analyses that adjusted for these biases. In addition, 411 children were excluded because of missing BMI data. Excluded children were not statistically different than included children on most demographic variables (e.g., race/ethnicity, SNAP participation, income, food access). However, excluded children were statistically more likely to be younger, female, have the primary respondent report a lower BMI, and have a different SNAP-authorized retailer closest to their home compared to included children.

Data were limited to household purchases and acquisitions and did not include individual purchases of children under age 11, children's overall diet patterns, or children's actual consumption of the acquired foods; these factors are likely important unobserved variables in the relationship between the food environment and childhood obesity. This data also does not include children's physical activity levels or sedentary behaviors, other important correlates of obesity. As a proxy, we examined adolescent food purchases and acquisitions to investigate whether detailed individual-level food acquisitions attenuated the relationship between food access and obesity prevalence, and results remained robust to their inclusion.

This analysis is also subject to measurement error, as exposure and access measures may not be complex enough to account for how food purchasing decisions (and subsequently, consumption and consumption patterns) are made. Although USDA excluded non-food items when possible, calculations using only event-level acquisitions may include spending on non-food items for some events (e.g., supermarket spending may include non-grocery items), resulting in a likely overestimate of actual food spending. Calculations using item-level acquisitions (e.g. reported sugary beverage spending) are likely to underestimate actual food spending, as some event-level acquisitions were missing item-level product information. In addition, there may be unobservable variables that account both for increased childhood obesity prevalence and changes to the neighborhood food environment. This analysis also does not address the impact of relative food prices or policies on childhood obesity prevalence, and future research should examine their effects, while accounting for neighborhood characteristics.

## **Conclusion**

This study found a positive and consistent relationship between combination grocery/other store access and overweight/obesity prevalence in a national sample of children, particularly for low-income children living in households participating in SNAP. The findings were stronger among adolescent children and non-Hispanic black children. Higher income children living in SNAP-ineligible households live in different retail food environments than SNAP and eligible non-SNAP children, and they have different household food and sugary beverage purchasing and acquisition patterns. The presence of some types of food outlets (i.e. those with limited healthy grocery options) may serve as a proxy for obesity-promoting environments for some children. They also raise questions of whether policies aiming to improve healthy food availability in retail food outlets may improve dietary choices or health outcomes for disadvantaged

children and families. For example, USDA recently developed new rules to increase the availability of healthy options for SNAP participants by requiring SNAP-authorized retailers to stock a larger inventory and variety of healthy foods.<sup>44</sup> However, the implementation of these rules has been delayed indefinitely. Longitudinal and experimental research is needed to examine how these types of changes in food access—both at the neighborhood level and within certain store types—influence BMI in children over time.

Author Manuscript

## References

1. Ogden C, Carroll M, Lawman H, et al. Trends in obesity prevalence among children and adolescents in the United States, 1988-1994 through 2013-2014. *JAMA* 2016;315(21):2292-2299.
2. Wang Y. Disparities in pediatric obesity in the United States. *Adv Nutr* 2011;2:23-31.
3. Odoms-Young A, Singleton C, Springfield S, McNabb L, Thompson T. Retail environments as a venue for obesity prevention. *Curr Obes Rep* 2016;5(2):184-91.
4. Gordon-Larsen P. Food availability/convenience and obesity. *Adv Nutr* 2014;5(6):809-817.
5. U.S. Department of Health and Human Services, Administrative for Children & Families, Office of Community Services. "Healthy Food Financing Initiative." Accessed online September 6, 2017: <https://www.acf.hhs.gov/ocs/programs/community-economic-development/healthy-food-financing>.
6. Larsen N, Story M, Nelson M. Neighborhood environments: disparities in access to healthy foods in the U.S. *Am J Prev Med* 2009;36(1):74-81.
7. Powell L, Auld M, Chaloupka F, O'Malley P, Johnston L. Associations between access to food stores and adolescent body mass index. *Am J Prev Med* 2007;33(4 Suppl):S301-7.
8. Leung C, Laraia B, Kelly M, Nickleach D, Adler N, Kushi L, et al. The influence of neighborhood food stores on change in young girls' body mass index. *Am J Prev Med* 2011;41(1):43-51.
9. Laska M, Hearst M, Forsyth A, Pasch K, Lytle L. Neighbourhood food environments: are they associated with adolescent dietary intake, food purchases and weight status? *Pub Health Nutrition* 2010;13(11):1757-1763.
10. Ohri-Vachaspati P, Lloyd K, Delia D, Tulloch D, Yedidia M. A closer examination of the relationship between children's weight status and the food and physical activity environment. *Preventive Med* 2013;57(3):162-7.
11. Currie J, Della Vigna S, Moretti E, Pathania V. The effect of fast food restaurants on obesity and weight gain. *American Economic Journal: Economic Policy* 2010;2:32-63.
12. Chen H, Wang Y. Changes in the neighborhood food store environment and children's body mass index at prepuberty in the United States. *J Adolesc Health* 2016;58(1):111-8.
13. Liu G, Wilson J, Qi R, Ying J. Green neighborhoods, food retail and childhood overweight: differences by population density. *Am J Health Promot* 2007;21(4 Suppl):317-325.
14. An R, Sturm R. School and residential neighborhood food environment and diet among California youth. *Am J Prev Med* 2012;42(2):129-135.
15. Meija N, Lightstone A, Basurto-Davila R, Morales D, Sturm R. Neighborhood food environment, diet, and obesity among Los Angeles County adults, 2011. *Prev Chron Disease* 2015;12:E143.
16. Cummins S, Flint E, Matthews S. New neighborhood grocery store increased awareness of food access but did not alter dietary habits or obesity. *Health Affairs (Millwood)* 2014;33(2):283-291.

7. Lee H. The role of local food availability in explaining obesity risk among young school-aged children. *Soc Sci Med* 2012;74(8):1193-1203.
8. Lovasi G, Hutson M, Guerra M, Neckerman K. Built environments and obesity in disadvantaged populations. *Epidemiol Rev* 2009;31:7-20.
9. Feichtner L, Block J, Duncan D, et al. Proximity to supermarkets associated with higher body mass index among overweight and obese preschool-age children. *Prev Med* 2013;56(3-4):218-21.
0. Chaparro M, Whaley S, Crespi C, et al. Influences of the neighborhood food environment on adiposity of low-income preschool-aged children in Los Angeles County: a longitudinal study. *J Epidemiol Community Health* 2014;68(11):1027-1033.
1. U.S. Department of Agriculture, Food and Nutrition Service. Building a Healthy America: A Profile of the Supplemental Nutrition Assistance Program. Office of Research and Analysis, April 2012. Accessed online September 02, 2017: <https://fns-prod.azureedge.net/sites/default/files/BuildingHealthyAmerica.pdf>.
2. Food and Nutrition Service, U.S. Department of Agriculture. Foods Typically Purchased by Supplemental Nutrition Assistance Program (SNAP) Households. November 2016. Accessed online September 2, 2017: <https://fns-prod.azureedge.net/sites/default/files/ops/SNAPFoodsTypicallyPurchased-Summary.pdf>.
3. Todd J, Scharadin B. Where Households Get Food in a Typical Week: Findings From USDA's FoodAPS, EIB-156: U.S. Department of Agriculture, Economic Research Service. July 2016.
4. Franckle R, Moran A, Hou T, et al. Transactions at a Northeastern Supermarket Chain: Differences by Supplemental Nutrition Assistance Program Use. *Am J Prev Med* 2017;53(4):e131-e138.
5. Bleich S, Vercammen K, Koma J, Zhonghe L. Trends in Beverage Consumption Among Children and Adults, 2003-2014. *Obesity (Silver Spring)* 2018; 26(2):432-441.
6. Vartanian L, Schwartz M, Brownell K. Effects of Soft Drink Consumption on Nutrition and Health: A Systematic Review and Meta-analysis. *Amer J Pub Health* 2007;97(4):667-675.
7. Malik V, Schulze M, Hu F. Intake of Sugar-Sweetened Beverages and Weight Gain: A Systematic Review. *Am J Clin Nutr* 2006;84(2):274-288.
8. U.S. Department of Agriculture, Economic Research Service. FoodAPS National Household Food Acquisition and Purchase Survey: Overview. 2016. Accessed September 2, 2017 from: <http://www.ers.usda.gov/data-products/foodaps-national-household-food-acquisition-and-purchase-survey.aspx>. Accessed September 01, 2017.
9. Jo Y. The Differences in Characteristics Among Households With and Without Obese Children: Findings from USDA's FoodAPS, EIB-179. U.S. Department of Agriculture, Economic Research Service. September 2017. Accessed online January 5, 2018 from: <https://www.ers.usda.gov/webdocs/publications/85028/eib-179.pdf?v=42989>.

0. Akinbami L, Ogden C. Childhood Overweight Prevalence in the United States: The Impact of Parent-Reported Height and Weight. *Obesity* 2009;17(8):1574-1580.
1. Weden M, Brownell M, Rendall C, Lau M, Nazarov Z. Parent-Reported Height and Weight as Sources of Bias in Survey Estimates of Childhood Obesity. *Am J Epidemiology* 2013;178(3):461-473.
2. Hattori A, An R, Sturm R. Neighborhood food outlets, diet, and obesity among California adults, 2007 and 2009. *Prev Chron Disease* 2013;10:e35.
3. U.S. Department of Agriculture, Economic Research Service. "Food Access Research Atlas: Definitions." Last updated May 22, 2017. Accessed online June 6, 2017 from: <https://www.ers.usda.gov/data-products/food-access-research-atlas/documentation/#definitions>.
4. U.S. Department of Agriculture Food and Nutrition Service. Retailer Management Year End Summaries. Fiscal Year 2016 At a Glance. Accessed online August 20, 2017: <https://www.fns.usda.gov/sites/default/files/snap/2016-SNAP-Retailer-Management-Year-End-Summary.pdf>.
5. Farley T, Rice J, Bodor J, Cohen D, Bluthenthal R, Rose D. Measuring the food environment: shelf space of fruits, vegetables, and snack foods in stores. *J Urban Health* 2009;86(5):672–682.
6. Caspi C, Pelletier J, Harnack L, Erikson D, Laska M. Differences in healthy food supply and stocking practices between small grocery stores, gas-marts, pharmacies, and dollar stores. *Pub Health Nutrition* 2016;19(3):540-547.
7. Rummo P, Guilkey D, Ng S, et al. Does unmeasured confounding influence associations between the retail food environment and body mass index over time? The Coronary Artery Risk Development in Young Adults (CARDIA) study. *Int J Epidemiol* 2017. doi: 10.1093/ije/dyx070.
8. Gibson D. The neighborhood food environment and adult weight status: estimates from longitudinal data. *Am J Public Health* 2011;101(1):71-78.
9. Caspi C, Lenk K, Pelletier J, Barnes T, Harnack L, Erickson D, et al. Food and beverage purchases in corner stores, gas-markets, pharmacies, and dollar stores. *Pub Health Nutrition* 2016:1-11.
0. Caspi C, Lenk K, Pelletier J, et al. Association between store food environment and customer purchases in small grocery stores, gas-markets, pharmacies, and dollar stores. *Int J Behav Nutr Phys Act* 2017;14(1):76.
1. Bleich S, Barry C, Gary-Webb T, Herring B. Reducing Sugar-Sweetened Beverage Consumption by Providing Caloric Information: How Black Adolescents Alter Their Purchases and Whether the Effects Persist. *Am J Public Health* 2014;104(12):2417-2424.
2. Glanz K, Sallis J, Saelens B, et al. Nutrition environment measures survey in stores (NEMS-S): development and evaluation. *Am J Prev Med* 2007;32:282–289.

3. Rummo P, Guilkey D, Ng S, Popkin B, Evenson K, Gordon-Larsen P. Beyond supermarkets: food outlet location in four U.S. cities over time. *Am J Prev Med* 2017;52(3):300-310.
4. U.S. Department of Agriculture. Enhancing retailer standards in the Supplemental Nutrition Assistance Program (SNAP). Accessed online August 21, 2017: <https://www.fns.usda.gov/snap/enhancing-retailer-standards-supplemental-nutrition-assistance-program-snap>.

Author Manuscript

**TABLE 1. Children's (ages 2-18) Individual, household, and neighborhood characteristics in the Food Acquisition and Purchase Survey, 2012-2013<sup>d</sup> by SNAP status**

	SNAP	Eligible Non-SNAP (≤130% FPL)	SNAP Ineligible (>130% FPL)
<b>Sample size, n</b>			
Children ages 2-18	1720	453	1575
Households with children ages 2-18	850	217	875
<b>INDIVIDUAL CHARACTERISTICS</b>			
<b>Child Age, m</b>	9.4 <sup>bc</sup>	10.5 <sup>a</sup>	10.3 <sup>a</sup>
<b>Child Gender, n(%)</b>			
Male	890 (51.5)	215 (46.0)	779 (49.7)
Female	830 (48.5)	238 (54.0)	796 (50.3)
<b>Child Race/Ethnicity, n</b>			
White, non-Hispanic	646 (32.0) <sup>bc</sup>	168 (33.5) <sup>ac</sup>	888 (65.0) <sup>ab</sup>
Hispanic	598 (31.6)	176 (41.2)	395 (16.9)
Black, non-Hispanic	369 (29.7)	76 (19.6)	155 (9.8)
Other, non-Hispanic	107 (6.7)	33 (5.7)	137 (8.3)
<b>Child BMI,<sup>e</sup> n(%)</b>			
Healthy Weight	940 (56.1) <sup>bc</sup>	237 (50.0) <sup>a</sup>	982 (66.2) <sup>a</sup>
Overweight/Obese	780 (43.9)	216 (50.0)	593 (33.8)
<b>HOUSEHOLD CHARACTERISTICS<sup>f</sup></b>			
<b>Household size, m</b>	4.9 <sup>c</sup>	5.1 <sup>c</sup>	4.4 <sup>ab</sup>
<b>Primary Respondent BMI, m</b>	30.5 <sup>c</sup>	28.9	27.3 <sup>a</sup>
<b>Primary Respondent Marital Status</b>			
Married	602 (30.7) <sup>bc</sup>	260 (56.9) <sup>ac</sup>	1173 (75.6) <sup>ab</sup>
Not Married	1118 (69.3)	193 (43.1)	402 (24.4)
<b>Primary Respondent Education</b>			
Less than high school	554 (32.1) <sup>bc</sup>	146 (36.2) <sup>ac</sup>	179 (4.9) <sup>ab</sup>
High school	486 (25.6)	138 (26.4)	364 (20.0)
Some college	560 (34.8)	106 (26.1)	581 (37.7)
College or more	118 (7.5)	62 (11.4)	451 (37.4)
<b>Primary Respondent Work Status</b>			
Working	647 (40.4) <sup>bc</sup>	209 (50.0) <sup>ac</sup>	1062 (70.1) <sup>ab</sup>
Not working <sup>g</sup>	1073 (59.6)	244 (50.0)	513 (29.9)
<b>Income (% FPL), m</b>	111.8% <sup>bc</sup>	87.2% <sup>ac</sup>	385.2% <sup>ab</sup>
<b>Food Security</b>			
High (secure)	480 (32.3) <sup>bc</sup>	141 (32.9) <sup>ac</sup>	928 (69.0) <sup>ab</sup>
Marginal (secure)	452 (22.5)	90 (18.9)	364 (21.1)
Low/very low (insecure)	788 (45.1)	222 (48.2)	283 (9.9)
<b>WIC participation</b>	469 (26.8) <sup>bc</sup>	128 (22.8) <sup>ac</sup>	152 (5.4) <sup>ab</sup>
<b>Area of residence<sup>h</sup></b>			
Urban	1309 (76.9) <sup>c</sup>	356 (80.6) <sup>c</sup>	1098 (61.1) <sup>ab</sup>
Rural	411 (23.2)	97 (19.4)	477 (38.9)
<b>Household Vehicle Access<sup>i</sup></b>			
Owns/leases vehicle	1303 (75.0) <sup>bc</sup>	407 (88.0) <sup>ac</sup>	1534 (97.3) <sup>ab</sup>
<b>NEIGHBORHOOD CHARACTERISTICS</b>			
<b>Closest SNAP retailer, n(%)</b>			
Convenience Store	939 (53.5) <sup>c</sup>	234 (55.6)	654 (37.7) <sup>ab</sup>



Superstore, supermarket, grocery store	400 (24.5)	106 (24.8)	473 (34.6)
Combo grocery or other Store	282 (16.1)	92 (15.8)	362 (23.6)
Specialty store <sup>j</sup>	99 (6.0)	21 (3.7)	86 (4.1)
<b>County-level poverty rate of households, m</b>	15.5% <sup>bc</sup>	14.3% <sup>ac</sup>	12.8% <sup>ab</sup>

Notes: \*P-value for difference is based on chi-squared test for categorical variables and regression (P>F) for continuous variables with post-hoc t-tests; P-values significant at p<0.05 level. Column frequencies may not sum to the full sample due to missingness.

BMI= Body Mass Index, measured in kg/m<sup>2</sup>.

<sup>a</sup> Significantly different from SNAP, p<0.05

<sup>b</sup> Significantly different from Eligible non-SNAP ≤130% FPL, p<0.05

<sup>c</sup> Significantly different from SNAP Ineligible >130% FPL, p<0.05

<sup>d</sup> Percentage of US population estimated with survey weights to adjust for unequal probability of sampling. All percentages shown are weighted column percentages

<sup>e</sup> Weight categories based on BMI for sex and age, where healthy weight is <85<sup>th</sup> percentile, overweight/obese is ≥85<sup>th</sup> percentile

<sup>f</sup> 1,102 out of 1,942 households in this sample have at least two children between the ages of 2-18 years old; 80.5% of children in this sample live in households where the primary respondent is female; 19.5% live in households where the primary respondent is male

<sup>g</sup> Not working includes those looking for work

<sup>h</sup> Urban and rural are defined in the Census Bureau's urbanized area definitions, where rural areas are sparsely populated areas with fewer than 2,500 people, and urban areas are areas with more than 2,500 people. A census tract is urban if the geographic centroid of the tract is in an area with more than 2,500 people; all other tracts are rural. 77.3%, 81.8%, and 80.7% of SNAP, SNAP-eligible, and SNAP-ineligible children live in households in metropolitan areas (county-level metropolitan designation; metropolitan statistical areas have at least 1 urbanized area of 50,000 or more population, plus adjacent territory that has a high degree of social and economic integration with the core measured by community ties)

<sup>i</sup> 98.6% of children in this sample live in households that either own or lease a vehicle, or have access to a car when one is needed for food shopping (only 1.5% of children live in households that reported issues accessing a vehicle)

<sup>j</sup> Specialty stores includes fruit/vegetable, meat/poultry, bread/bakery, seafood specialty stores and non-profit cooperatives, delivery route, direct marketing & farmer's markets

Author

**TABLE 2. Children's food access and household food purchases and acquisitions, 2012-2013<sup>d</sup> FoodAPS Survey, by SNAP status**

	SNAP N=1720	Eligible Non-SNAP (≤130% FPL) N=453	SNAP Ineligible (>130% FPL) N=1575
Mean number of stores within 1 mile of home			
Superstores, Supermarkets, Grocery stores	3.4 <sup>c</sup>	3.2 <sup>c</sup>	1.9 <sup>ab</sup>
Combination grocery and other stores	2.4 <sup>c</sup>	2.5 <sup>c</sup>	1.5 <sup>ab</sup>
Convenience stores	5.8 <sup>c</sup>	5.5 <sup>c</sup>	2.6 <sup>ab</sup>
Fast food restaurants	5.6 <sup>c</sup>	6.7 <sup>c</sup>	4.1 <sup>ab</sup>
Non-fast food restaurants	23.9 <sup>c</sup>	24.2 <sup>c</sup>	17.8 <sup>ab</sup>
Mean number of food acquisition events <sup>e</sup> per week per household	14.7 <sup>c</sup>	15.6	17.6 <sup>a</sup>
Mean weekly spending at food outlets per household member <sup>f</sup>	\$37.41 <sup>bc</sup>	\$31.51 <sup>ac</sup>	\$58.87 <sup>ab</sup>
Percent of household weekly spending at food outlets for the average child, by store type <sup>g</sup>	N=1636 100%	N=441 100%	N=1559 100%
Superstores, Supermarkets, Grocery stores	58.0% <sup>c</sup>	50.2%	49.3% <sup>a</sup>
Restaurants	24.4% <sup>c</sup>	26.7%	29.3% <sup>a</sup>
Combination grocery and other stores	4.9% <sup>c</sup>	5.3%	2.9% <sup>a</sup>
Other (Miscellaneous) <sup>h</sup>	4.6%	4.7%	5.8%
Convenience stores	4.3% <sup>bc</sup>	2.1% <sup>a</sup>	2.2% <sup>a</sup>
Club stores	2.4% <sup>c</sup>	8.7%	6.4% <sup>a</sup>
Family/friends, school, work	1.4% <sup>c</sup>	2.2%	4.1% <sup>a</sup>

Notes: P-value for difference based on regression (P>F) for continuous variables with post-hoc t-tests; P-values significant at p<0.05 level.

<sup>a</sup> Significantly different from SNAP, p<0.05

<sup>b</sup> Significantly different from Eligible non-SNAP ≤130% FPL, p<0.05

<sup>c</sup> Significantly different from SNAP Ineligible >130% FPL, p<0.05

<sup>d</sup> Percentage of US population estimated with survey weights to adjust for unequal probability of sampling. All percentages shown are weighted column percentages

<sup>e</sup> Food acquisition events=number of visits to each place per week where food was purchased or acquired for free. See Appendix B for full details on how this measure was calculated.

<sup>f</sup> Event-level weekly spending may include spending on non-food items for some events (e.g, supermarket spending may include non-grocery items); includes children living in households with \$0 weekly spending. See Appendix B for full details on how this measure was calculated.

<sup>g</sup> Percent of weekly food spending excludes free events and excludes 112 children in households with \$0 weekly spending. See Appendix B for full details on how this measure was calculated.

<sup>h</sup> Other (Miscellaneous) includes several miscellaneous food outlets, including vending machines, non-food retailers, travel places (airports, hotels, truck stops), gardens, hospitals and institutions, multiple places, and unknown

**TABLE 3: Estimated Odds of Childhood Overweight or Obesity, by Household Food Access and SNAP Status, FoodAPS Survey, 2012-2013**

	Overall (All Children) N=3,742		SNAP N=1,715		Eligible Non-SNAP (≤130% FPL) N=452		SNAP Ineligible (>130% FPL) N=1,575	
	OR	CI	OR	CI	OR	CI	OR	CI
Food access (number of stores within 1 mile of child’s home), by outlet type								
Superstore, supermarket, grocery	0.98	0.95, 1.02	0.99	0.95, 1.04	1.00	0.92, 1.10	0.98	0.92, 1.04
Combination grocery store/other	1.10*	1.03, 1.17	1.14*	1.05, 1.24	0.99	0.76, 1.29	1.10	0.99, 1.24
Convenience stores	1.01	0.97, 1.04	0.97	0.93, 1.02	1.11*	1.04, 1.18	1.02	0.97, 1.07
Fast Food restaurants	0.99	0.96, 1.02	0.96	0.91, 1.01	1.05	0.97, 1.13	0.99	0.94, 1.05
Non-Fast Food restaurants	1.00	0.99, 1.00	1.00	1.00, 1.01	0.98*	0.97, 0.995	0.99	0.99, 1.00

\*Statistically significant at p<0.05. All models adjust for child’s age, gender, race/ethnicity, primary respondent BMI, primary respondent marital status, primary respondent education, primary respondent work status, household income, food security, WIC participation, rural v. urban household location, vehicle access, closest SNAP retailer, county-level poverty rate, amount of weekly spending at food outlets per household member, number of weekly food acquisitions visits to each store type, and % of household spending by store type (includes children living in households with no spending). The Overall Model also adjusts for SNAP participation. Models did not adjust for household size, as it is correlated with household income as a percent of the Federal Poverty Level.

Author Manuscript  
 Copyright Clearance Center  
 www.copyright.com

**TABLE S1. Proportion of Sugary Beverage Purchases for the Average Child and Adolescent, 2012-2013 FoodAPS Survey,<sup>d</sup> Overall and by SNAP status**

	N	Total (All Children)	SNAP	Eligible Non-SNAP (≤130% FPL)	SNAP Ineligible (>130% FPL)
<b>CHILDREN'S HOUSEHOLD SUGARY BEVERAGE SPENDING</b>					
<b>Percent household weekly sugary beverage spending for the average child</b>					
Overall <sup>e</sup>	3627	6.3%	8.3% <sup>c</sup>	7.7% <sup>c</sup>	5.5% <sup>ab</sup>
At convenience stores <sup>f</sup>	1336	26.1%	29.8% <sup>b</sup>	15.5% <sup>ac</sup>	25.6% <sup>b</sup>
At family/friend events, school, or work <sup>f</sup>	1040	7.9%	18.2% <sup>c</sup>	16.9%	5.9% <sup>a</sup>
At combination grocery and other stores <sup>f</sup>	973	13.3%	18.4% <sup>c</sup>	17.7%	10.4% <sup>a</sup>
At superstores, supermarkets, and grocery Stores <sup>f</sup>	3276	6.7%	9.0%	9.1% <sup>c</sup>	5.8% <sup>b</sup>
At club stores <sup>f</sup>	356	3.5%	7.7%	13.1%	2.1%
At restaurants <sup>f</sup>	2960	4.8%	5.4%	4.8%	4.6%
At other venues (miscellaneous) <sup>fg</sup>	1321	13.3%	17.3%	13.0%	12.6%
<b>ADOLESCENT SUGARY BEVERAGE SPENDING</b>					
<b>Percent weekly sugary beverage spending for the average adolescent<sup>i</sup></b>	800	10.0%	18.1% <sup>bc</sup>	7.6% <sup>a</sup>	8.9% <sup>a</sup>

Notes: P-value for difference based on regression (P>F) for continuous variables with post-hoc t-tests; P-values significant at p<0.05 level.

<sup>a</sup> Significantly different from SNAP, p<0.05

<sup>b</sup> Significantly different from Eligible non-SNAP ≤130% FPL, p<0.05

<sup>c</sup> Significantly different from SNAP Ineligible >130% FPL, p<0.05

<sup>d</sup> Percentage of US population estimated with survey weights to adjust for unequal probability of sampling. All percentages shown are weighted column percentages

<sup>e</sup> Percent of household food spending on SSBs calculated for each child with item-level spending on foods. Overall percent excludes 121 children in households with \$0 in itemized food spending. See Appendix B for full details on how this measure was calculated.

<sup>f</sup> Percent of household food spending on SSBs calculated for each child includes item-level spending on foods by store type. Store type percentages exclude children in households with \$0 in itemized spending at that store type (e.g., for Combination grocery and other store spending, percent only calculated among the N=973 children living in households with any itemized purchases from combination grocery stores/other over the survey period (excludes those living in households with no spending at this store type). See Appendix B for full details on how this measure was calculated.

<sup>g</sup> Other (Miscellaneous) includes several miscellaneous food outlets, including vending machines, non-food retailers, travel places (airports, hotels, truck stops), gardens, hospitals and institutions, multiple places, and unknown

<sup>i</sup> Percent of adolescent spending on SSBs includes item-level spending on SSBs; percent excludes adolescents with \$0 in itemized food spending. See Appendix B for full details on how this measure was calculated.

**TABLE S2. Absolute Spending on Sugary Beverages Among Households and Adolescents, 2012-2013 FoodAPS Survey,<sup>d</sup> Overall and by SNAP status**

	N	SNAP	Eligible Non-SNAP ( $\leq 130\%$ FPL)	SNAP Ineligible ( $>130\%$ FPL)
<b>CHILDREN'S HOUSEHOLD SUGARY BEVERAGE SPENDING</b>				
Mean of all weekly itemized spending on food outlets per household member	3748	\$30.76 <sup>bc</sup>	\$25.17 <sup>ac</sup>	\$45.51 <sup>ab</sup>
Mean weekly spending on sugary beverages per household member <sup>e</sup>	3748	\$2.05	\$1.80 <sup>c</sup>	\$2.31 <sup>b</sup>
<b>ADOLESCENT SUGARY BEVERAGE SPENDING</b>				
Mean of all weekly itemized spending on food outlets for adolescents with $> \$0$ in itemized food spending	800	\$13.33	\$13.93	\$15.49
Mean weekly spending on sugary beverages per adolescent for adolescents with $> \$0$ in itemized food spending	800	\$1.44	\$1.12	\$1.01

Notes: P-value for difference based on regression ( $P > F$ ) for continuous variables with post-hoc t-tests; P-values significant at  $p < 0.05$  level.

<sup>a</sup> Significantly different from SNAP,  $p < 0.05$

<sup>b</sup> Significantly different from Eligible non-SNAP  $\leq 130\%$  FPL,  $p < 0.05$

<sup>c</sup> Significantly different from SNAP Ineligible  $> 130\%$  FPL,  $p < 0.05$

<sup>d</sup> Percentage of US population estimated with survey weights to adjust for unequal probability of sampling. All percentages shown are weighted column percentages

<sup>e</sup> Mean weekly spending per hh member for all children in the sample ( $N=3748$ ), including those living in households with  $\$0$  in food spending over the survey period

Author Manuscript

**TABLE S3: Full Model Estimating the Odds of Childhood Overweight or Obesity, by Household Food Access and SNAP Status, FoodAPS Survey, 2012-2013**

	OR	Overall (All Children) N=3,742 CI
Food access (number of stores within 1 mile of child's home), by outlet type		
Superstore, supermarket, grocery	0.98	0.95, 1.02
Combination grocery store/other	1.10*	1.03, 1.17
Convenience stores	1.01	0.97, 1.04
Fast Food restaurants	0.99	0.96, 1.02
Non-Fast Food restaurants	1.00	0.99, 1.00
<b>Covariates</b>		
<b>Child Age</b>	0.94*	(0.92, 0.96)
<b>Child Gender</b>		
Male	[Ref]	--
Female	0.71*	(0.56, 0.91)
<b>Child Race/Ethnicity</b>		
White, non-Hispanic	[Ref]	--
Hispanic	1.14	(0.78, 1.66)
Black, non-Hispanic	1.20	(0.87, 1.66)
Other	0.92	(0.65, 1.30)
<b>Primary Respondent BMI</b>	1.03*	(1.01, 1.05)
<b>Primary Respondent Marital Status</b>		
Not Married	[Ref]	--
Married	0.72*	(0.56, 0.94)
<b>Primary Respondent Education</b>	0.83*	(0.73, 0.95)
<b>Primary Respondent Work Status</b>		
Not Working	[Ref]	--
Working	0.91	(0.69, 1.21)
<b>Income (% FPL)</b>	1.00	(1.00, 1.00)
<b>Food Security</b>		
High (secure)	[Ref]	--
Marginal (secure)	1.31	(0.89, 1.92)
Low/very low (insecure)	1.04	(0.71, 1.52)
<b>SNAP Participation</b>	0.79	(0.62, 1.01)
<b>WIC Participation</b>	0.99	(0.78, 1.28)
<b>Area of residence</b>		
Urban	[Ref]	--
Rural	0.89	(0.63, 1.25)
<b>Household owns/leases vehicle</b>	0.84	(0.32, 2.19)
<b>Closest SNAP retailer</b>		
Convenience Store	[Ref]	--
Superstore, supermarket, grocery store	1.33*	(1.02, 1.73)
Combo grocery or other Store	1.18	(0.77, 1.84)
Specialty store	1.72*	(1.03, 2.86)
<b>County-level poverty rate of households</b>	1.03*	(1.01, 1.06)
<b>Amount of weekly spending at food outlets per</b>	1.00	(1.00, 1.01)

**household member****Number of household food acquisition events per week,  
by store type**

Superstores, Supermarkets, Grocery stores	0.99	(0.93, 1.05)
Restaurants	1.02	(0.97, 1.06)
Combination grocery/other stores	0.94	(0.83, 1.07)
Convenience stores	1.07	(0.99, 1.16)
Club stores	0.71*	(0.52, 0.98)
Family/friends, school, work	1.00	(0.99, 1.02)
Other (Miscellaneous)	0.92*	(0.86, 0.98)

**Percent of household weekly spending at food outlets,  
by store type**

Superstores, Supermarkets, Grocery stores	1.00	(0.99, 1.01)
Restaurants	1.00	(0.99, 1.01)
Combination grocery and other stores	1.01	(1.00, 1.02)
Convenience stores	1.00	(0.98, 1.01)
Club stores	1.02*	(1.01, 1.03)
Family/friends, school, work	1.00	(0.98, 1.01)
Other (Miscellaneous)	1.00	(0.99, 1.01)

---

\*Statistically significant at  $p < 0.05$ . Overall model adjusts for child's age, gender, race/ethnicity, primary respondent BMI, primary respondent marital status, primary respondent education, primary respondent work status, household income, food security, SNAP participation, WIC participation, rural v. urban household location, vehicle access, closest SNAP retailer, county-level poverty rate, amount of weekly spending at food outlets per household member, number of weekly food acquisitions visits to each store type, and % of household spending by store type (includes children living in households with no spending). Models did not adjust for household size, as it is correlated with household income as a percent of the Federal Poverty Level.

Author Manuscript

**APPENDIX A. ABBREVIATED USDA STORE TYPE DEFINITIONS, FoodAPS Survey, 2012-2013**

- (1) Supermarkets and grocery stores—includes SNAP-authorized superstores, “big box” stores, supermarkets, large and medium grocery stores
- (2) Combination grocery and other stores—includes SNAP-authorized combination grocery and other stores where the primary business is sale of general merchandise, but also sells a variety of food products (e.g., independent drug stores, dollar stores, general stores)
- (3) Convenience stores—SNAP-authorized convenience stores, where the primary business is retail sale of canned goods, dairy, pre-packaged meat, and other grocery items in limited amounts
- (4) Fast food restaurants—identified by USDA using InfoUSA and a Wikipedia list of global and U.S. fast food restaurant chains
- (5) Non-fast food restaurants—all other restaurants identified by USDA through InfoUSA

Author Manuscript



**APPENDIX B. DEFINITIONS FOR MEASURES USED FROM FOODAPS**

- **Table 2: Mean number of food acquisition events per week per household** was measured using event data and calculated for each child by averaging the sum of their household members' individual food purchases and acquisitions over the week.
- **Table 2: Mean weekly spending at food outlets per household member (overall)** was measured using event data and calculated by summing each household members' individual purchases over the week, then dividing total household purchases by household size, with the mean calculated for each child. This measure includes households with \$0 in weekly spending (n=112 children).
- **Table 2: Percent of household weekly spending at food outlets for the average child, by store type** was calculated for each child by dividing the total weekly household spending at each of the following store types over total weekly household spending: supermarkets, superstores, grocery stores; restaurants; combination grocery and other stores; convenience stores; club stores; family, school, and work; and other venues (miscellaneous). We then calculated the mean of these percentages across all children (average of ratios). This measure excludes free acquisitions and households with \$0 in weekly spending and includes more granular classifications of place types (including non-food outlets such as family and friend events) than the predefined food retailer access measures provided by USDA.
- **Table S1: To capture a measure of spending on unhealthy items, the percentage of household weekly food spending on sugary beverages for the average child, overall and by store type**, was measured using both event- and item-level data.
  - **Overall:** Total itemized spending at all food outlets was calculated for each child by summing all household members' total item-level food purchases over the week for each child, and total sugary beverage spending at all food outlets was calculated by summing all household members' item-level purchases of sugary beverages (defined by USDA as sweetened beverages—includes non-diet soft drinks, fruit drinks, sport and energy drinks,

and nutritional beverages, but not flavored milk) over the week for each child. Then, we calculated the percent of sugary beverage spending for each child by dividing their household total spending on sugary beverages by household total item-level spending. We then calculated the mean of these percentages across all children (average of ratios). This measure excludes 148 children living in households with \$0 in itemized food spending. This procedure was also used to calculate the percent of adolescents' overall food spending on sugary beverages among adolescents with any food spending (n=800) in Table S1.

- **By Store Type: Percent of sugary beverage spending** was then calculated by store type. For example, the percent of households' weekly convenience store food spending on sugary beverages was calculated for each child by dividing the sum of their household members' total sugary beverage purchases at convenience stores by the sum of their household members' total item-level food purchases at convenience stores. We then calculated the mean of these percentages across all children (average of ratios). This measure excludes children living in households with \$0 in itemized food spending at convenience stores over the survey period.

Author Manuscript