

Article type : Original Article

## **Phosphodiesterase 4 Inhibition Reduces Lung Fibrosis Following Targeted Type II Alveolar Epithelial Cell Injury**

Thomas H. Sisson<sup>1</sup>, Paul J. Christensen<sup>2</sup>, Yo Muraki<sup>3</sup>, Anthony J. Dils<sup>1</sup>, Lauren Chibucos<sup>1</sup>, Natalya Subbotina<sup>1</sup>, Kimio Tohyama<sup>3</sup>, Jeffrey C. Horowitz<sup>1</sup>, Takanori Matsuo<sup>3</sup>, Marc Bailie<sup>4</sup>, Sham Nikam<sup>3</sup>, and Masatoshi Hazama<sup>3</sup>

- 1) Pulmonary and Critical Care Division, Department of Internal Medicine, University of Michigan Medical Center. Ann Arbor, MI.
- 2) Division of Pulmonary & Critical Care Medicine, Department of Internal Medicine, William Beaumont Medical Center, Troy, MI.
- 3) Takeda Pharmaceutical Company Limited, Fujisawa, Japan.
- 4) In Vivo Facility. Dept. of Pharmacology and Toxicology, Michigan State University, East Lansing, MI.

### **Address all Correspondence to:**

Thomas H. Sisson, M.D.;  
Division of Pulmonary and Critical Care Medicine  
Department of Internal Medicine  
University of Michigan Medical Center  
6301c MSRB3

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.14814/phy2.13753](https://doi.org/10.14814/phy2.13753)

This article is protected by copyright. All rights reserved

1150 W. Medical Center Drive, Ann Arbor, MI 48109-0600 U.S.A

Phone: 734-763-8008

FAX: 734-764-4556

E-mail: [tsisson@med.umich.edu](mailto:tsisson@med.umich.edu)

**Running Head:** PDE4 inhibition and Lung Fibrosis

**Key Words:** Pulmonary, Collagen, Fibroblast, Epithelium, cAMP

## **Abstract**

Fibrosis of the lung constitutes a major clinical challenge and novel therapies are required to alleviate the associated morbidity and mortality. Investigating the anti-fibrotic efficacy of drugs that are already in clinical practice offers an efficient strategy to identify new therapies. The phosphodiesterase 4 (PDE4) inhibitors, approved for the treatment of chronic obstructive pulmonary disease, harbor therapeutic potential for pulmonary fibrosis by augmenting the activity of endogenous anti-fibrotic mediators that signal through cyclic AMP. In the present study, we tested the efficacy of several PDE4 inhibitors including a novel compound (Compound 1) in a murine model of lung fibrosis that results from a targeted type II alveolar epithelial cell injury. We also compared the anti-fibrotic activity of PDE4 inhibition to the two therapies that are FDA-approved for idiopathic pulmonary fibrosis (pirfenidone and nintedanib). We found that both preventative (day 0-21) and therapeutic (day 11-21) dosing regimens of the PDE4 inhibitors significantly ameliorated the weight loss and lung collagen accumulation that are the sequelae of targeted epithelial cell damage. In a therapeutic protocol, the reduction in lung fibrosis with PDE4 inhibitor administration was equivalent to pirfenidone and nintedanib. Treatment with this class of drugs also resulted in a decrease in plasma surfactant protein D concentration, a reduction in the plasma levels of several chemokines implicated in lung fibrosis, and an in vitro inhibition of fibroblast pro-fibrotic gene expression. These results motivate further investigation of PDE4 inhibition as a treatment for patients with fibrotic lung disease.

## Introduction:

Fibrosis of the lung interstitium occurs in response to multiple insults. Inhalation of organic and inorganic dusts and systemic exposure to specific medications including chemotherapy can all induce alveolar scarring. Lung fibrosis is also associated with a subset of rheumatologic diseases including systemic sclerosis and rheumatoid arthritis. In idiopathic pulmonary fibrosis (IPF), interstitial scarring occurs in the absence of an identifiable injury, although type II alveolar epithelial cell specific gene mutations are found in a subset of patients with both sporadic and familial forms of this disease (1, 2). All of these disorders that result in fibrosis of the lung interstitium are associated with significant morbidity and mortality. However, IPF arguably represents the greatest therapeutic challenge as this disease portends a median survival of only 3 years, and less than 20% of patients with this disease live beyond eight years of their initial diagnosis (2). The recent FDA approval of two new therapies for the treatment of IPF (pirfenidone and nintedanib) has resulted in guarded optimism, but there continues to be a desperate need for additional anti-fibrotic medications with increased efficacy and fewer side effects (3, 4).

One approach to identify novel anti-fibrotic therapies is to investigate the efficacy of currently FDA approved agents that target specific pro-fibrotic mechanisms. A significant advantage of this approach to repurpose existing compounds is the potential for rapid translation into patients because of the existing pharmacokinetic and pharmacodynamic data and safety studies in humans. As an example, phosphodiesterase 4 (PDE4) inhibitors including roflumilast, which is approved for the treatment of chronic obstructive pulmonary disease (COPD), hold promise as anti-fibrotic agents. This class of drugs blocks the degradation of cAMP and thereby augments the activity of anti-fibrotic mediators that signal through G protein-coupled receptors including prostaglandin E<sub>2</sub> (PGE<sub>2</sub>), prostacyclin, and adenosine. PGE<sub>2</sub> exhibits a multitude of anti-fibrotic effects including inhibiting fibroblast activation, increasing fibroblast susceptibility to apoptosis, and maintaining alveolar epithelial cell integrity (5-8). The anti-fibrotic potential of PDE4 inhibitors is supported by several *in vitro* studies. Specifically, the PDE4 inhibitor piclamilast was found to blunt transforming growth factor-beta (TGF- $\beta$ )-mediated fibroblast to myofibroblast differentiation, an effect that was accentuated by the addition of exogenous PGE<sub>2</sub> (9). In a separate *in vitro* study, roflumilast inhibited both TGF- $\beta$  stimulated

fibroblast contraction of 3-dimensional collagen gels and fibroblast chemotaxis toward fibronectin (10). Importantly, roflumilast also increased PGE<sub>2</sub> release by fibroblasts. PDE4 inhibitors have also been found to mitigate bleomycin induced fibrosis in vivo. Specifically, the initiation of roflumilast treatment on day 0 and day 7 following bleomycin injury reduced day 14 and day 21 lung fibrosis respectively (11). In a separate study, roflumilast also reversed the metabolic alterations of bleomycin injury when administered in a preventative regimen (12).

The aforementioned studies supporting the anti-fibrotic potential of PDE4 inhibitors led us to further investigate the efficacy of this class of medications in a murine model of lung fibrosis that results from a targeted insult to the type II alveolar epithelium (13). We hypothesized that both preventative and therapeutic PDE4 inhibitor administration protocols would mitigate pulmonary fibrosis in this model. Type II alveolar epithelial cell injury-induced fibrosis differs from bleomycin in that the inflammatory response is modest (marked by the selective accumulation of Ly-6c high monocytes and exudate macrophages) and the resultant fibrosis is more diffuse (13-15). PDE4 inhibitors including piclamilast, roflumilast and a novel agent (Compound 1) were tested and the efficacy of PDE4 inhibition was compared to pirfenidone and nintedanib. The effect of PDE4 inhibition on lung chemokine levels, whole lung fibrogenic gene expression, and plasma surfactant protein D (SP-D) levels (as an indicator of epithelial cell integrity) was also measured (1).

We found that both prophylactic and therapeutic administration of PDE4 inhibitors including the novel agent Compound 1 significantly reduced lung fibrosis following type II AEC injury with an efficacy comparable to pirfenidone and nintedanib. The protection against lung fibrosis was associated with diminished weight loss and a significant decline in plasma SP-D levels. These findings highlight the antifibrotic potential of PDE4 inhibition, and with roflumilast already FDA approved for the treatment of COPD and additional novel agents on the horizon, our data motivate the clinical investigation of this class of drugs in patients with IPF and other fibrotic lung diseases.

## **Methods:**

Animals: All animal experiments were performed in accordance with institutional guidelines set forth by the University Committee on the Use and Care of Animals at the University of Michigan. Transgenic mice expressing the diphtheria toxin receptor (DTR) under the control of the murine SPC promoter were generated in our laboratory on a C57BL/6 background (designated DTR+

mice) (13). The presence of the DTR was detected using PCR for the SPC-DTR transgenic construct as previously described (13). Control mice included littermates of SPC-DTR transgenic mice that were PCR-negative for the transgenic construct and C57BL/6 mice purchased from Jackson Laboratories (Bar Harbor, ME). These control animals are designated DTR- and prior studies revealed that neither of these strains of mice develop significant fibrosis in response to diphtheria toxin (DT) administration.

Drug concentration and cAMP levels: Roflumilast (5 mg/kg) and Compound 1 (10 mg/kg) were administered to C57BL/6N mice (Charles River Laboratories Japan, Inc., Japan) which were then sacrificed at 0.5, 1, 2, 4, 8 and 24 hours after the dosing. Blood samples were collected and centrifuged to obtain plasma specimens. Plasma concentrations of roflumilast, the active metabolite of roflumilast N-oxide and Compound 1 were measured using liquid chromatography/tandem mass spectrometry. Bronchoalveolar Lavage Fluid (BALF) was collected by instilling 3 aliquots of 0.5 mL 0.9% saline into the lungs. BALF was centrifuged to remove cells, the supernatant was deproteinized with trichloroacetic acid, and washed by diethylether. The aqueous layer was freeze-dried and dissolved by lysis buffer enclosed in cAMP XP Assay kit (Cell Signaling Technology Inc., MA, USA). cAMP content was measured in accordance with the manufacturer's instruction, and, total PDE4 inhibitory activities (tPDE4i) were calculated by following equation:

$$tPDE4i = \sum \frac{(C_n \text{ or } AUC_n) \times fu_n}{IC50_n}$$

where C<sub>n</sub> and AUC<sub>n</sub> represent the plasma concentration and the AUC of active pharmacological ingredient (API) respectively, fu is the unbound fraction in plasma, IC<sub>50</sub> is the concentration resulting in 50% inhibition in an in vitro assay and n represents the number of APIs(18). Of note, roflumilast has two APIs and Compound 1 has one API.

Preparation of test compounds: All drugs were suspended in 0.5% methylcellulose and administered by oral gavage. Roflumilast (1.0 and 5.0 mg/kg), piclamilast (30 mg/kg) and Compound 1 (N-[Amino(dimethylamino)methylidene]-4-[(3aS,9bR)-8-ethoxy-7-methoxy-1,3,3a,9b-tetrahydrofuro[3,4-c]isoquinolin-5-yl]benzamide at 1.0, 5.0, and 10.0 mg/kg) were given once daily by gavage. Pirfenidone was dosed at 100 mg/kg three times daily (16) and nintedanib was dosed at 100 mg/kg twice daily (17) by gavage.

Diphtheria Toxin Administration and Experimental Design: Six to ten week old mice were intraperitoneally injected with DT (Sigma Chemical, St. Louis, MO) once daily for 14 days at a dose of 12.5 µg/kg in 100 µl of PBS. Control groups received intraperitoneal PBS alone. Prophylactic dosing of the experimental treatments was initiated on day 1 and continued daily at the indicated doses throughout the 21 day study protocol. Therapeutic dosing was initiated on day 11 and continued daily at the indicated doses for 10 days. Mice were weighed daily and, on day 21 of the study protocol (7 days after the last dose of DT), blood and lungs were harvested 1 hour after the last drug administration. Lungs were either sectioned for histology or homogenized for hydroxyproline analysis and whole lung gene expression. Plasma was separated from whole blood and analyzed for chemokine and SP-D levels.

Hydroxyproline assay: Hydroxyproline content of both lungs was measured as previously reported (13).

Lung histology: The left lung was inflation-fixed at 25 cm H<sub>2</sub>O pressure with 10% neutral-buffered formalin, removed en bloc, further fixed in 10% neutral-buffered formalin overnight, and then paraffin embedded. Five micron sections were stained using picosirius red.

Plasma SP-D levels: Plasma SP-D levels were measured using SP-D ELISA Kit (Yamasa Corp., Japan). Samples were diluted using the provided buffer and measurements were made in accordance with the manufacturer's protocol.

Plasma Chemokine analysis: Plasma chemokine concentrations were measured using the Milliplex MAP mouse cytokine/chemokine magnetic bead panel kit (Merck Millipore Corp., Germany) according to the manufacturer's instructions.

In vitro fibroblast studies: For cAMP assay, WI-38 human lung fibroblasts were suspended in E-MEM (Thermo Fisher Scientific Inc., MA,USA) containing 10% heat-inactivated fetal bovine serum (FBS) and seeded on 24-well plates at 50,000 cells/500 µL/well. Twenty-four hours after seeding, cells were treated with Compound 1 (1×10<sup>-10</sup> M - 1×10<sup>-5</sup> M) dissolved in HBSS (Thermo Fisher Scientific Inc., MA,USA) containing 0.1 % bovine serum albumin (BSA) and 5 mM HEPES. After 30 min, cells were stimulated by 1 µM forskolin for 30 min and then lysed.

cAMP was measured in accordance with the cAMP XP Assay kit protocol (Cell Signaling Technology Inc., MA, USA).

To analyze fibroblast gene expression, WI-38 human lung fibroblasts were suspended in E-MEM (Thermo Fisher Scientific, MA, USA) containing 10% heat-inactivated FBS and seeded on 24 well plates at 50,000 cells/500  $\mu$ L/well. After seeding, cells were cultured in E-MEM containing 0.5% FBS for 24 hours. The cultures were then treated with Compound 1 ( $1 \times 10^{-10}$  M -  $1 \times 10^{-5}$  M) for 1 hour followed by TGF- $\beta$  (3 ng/mL) and forskolin (1  $\mu$ M) for 24 hours. Total RNA was extracted from cell lysate using RNeasy 96 Kit (QIAGEN, Germany). cDNA was amplified using High-Capacity cDNA Reverse Transcription Kit (Thermo Fisher Scientific, MA, USA), and target gene mRNA was measured using TaqMan PCR (7900HT Thermo Fisher Scientific, MA, USA). The target gene expression level was normalized by mRNA expression of Glyceraldehyde 3-phosphate dehydrogenase (GAPDH). TaqMan gene expression assays (Thermo Fisher Scientific, MA, USA) were used to measure gene expression. Assay ID of human GAPDH, type-1 collagen, fibronectin, connective tissue growth factor and plasminogen activator inhibitor-1 were Hs02758991g1, Hs00164004m1, Hs00365052m1 Hs01026927g1 and Hs00167155m1, respectively.

In vitro A549 cell studies: For cAMP assays, A549 human lung epithelial cells were suspended in Ham's F-12K (Thermo Fisher Scientific Inc., MA, USA) containing 10% heat-inactivated FBS and seeded on 24 well plates as 50,000 cells/500  $\mu$ L/well for 24 hours. Cells were treated with Compound 1 ( $1 \times 10^{-10}$  M -  $1 \times 10^{-5}$  M) dissolved in HBSS (Thermo Fisher Scientific Inc., MA, USA) containing 0.1% bovine serum albumin (BSA) and 5 mM HEPES for 30 minutes and stimulated with 10  $\mu$ M forskolin and 10  $\mu$ M prostaglandin E2 for 30 min. Cells were lysed and cAMP content in cell lysates and supernatants were measured using cAMP XP Assay kit (Cell signaling technology, MA, USA).

Quantitative RT-PCR on lung homogenates: RNA was isolated from the left lung using the RNeasy Plus Mini Kit (Qiagen) and first-strand cDNA was synthesized using SuperScriptIII (Invitrogen, Carlsbad, CA). Specific qPCRs with SYBR Green-based detection using an MX 3000P system (Stratagene, La Jolla, CA) were conducted for forty cycles (94  $^{\circ}$ C for 15 seconds followed by 60  $^{\circ}$ C for 30 seconds and 72  $^{\circ}$ C for 30 seconds) using each cDNA template. The mRNA levels were normalized with the expression of GAPDH using the following formula:

%GAPDH expression =  $100 / 2^{-\Delta\Delta CT}$ . Primer pairs for murine collagen1a1, tumor necrosis factor  $\alpha$ , fibronectin, connective tissue growth factor, and plasminogen activator inhibitor 1 were purchased from Applied Biosystems.

**Statistical Analysis.** All data were expressed as mean  $\pm$  SEM. Data were evaluated by unpaired Student's t-test (for comparison between two samples), a two-tailed William's test, one-way ANOVA with a Tukey multiple comparisons test or a two-way ANOVA with a Tukey multiple comparisons test. A statistically significant difference was accepted at  $p < 0.05$ .

## **Results:**

### **Pharmacokinetics of the phosphodiesterase 4 inhibitors:**

Prior to testing the efficacy of PDE4 inhibitors in our murine model of lung fibrosis, we assessed the pharmacokinetics of Compound 1 with our planned dosing regimen in uninjured wild-type mice. Compound 1 is a novel PDE4 inhibitor with a different structural scaffold than Roflumilast and Piclamilast, and in contrast to Piclamilast and Roflumilast which both inhibit CYP3A4, Compound 1 has no CYP inhibitory activity. For Compound 1, a single dose of (10.0 mg/kg) was administered by oral gavage at time 0. Plasma was then collected at 0.5, 1.0, 2.0, 4.0, 8.0 and 24.0 hours. Bronchoalveolar lavage fluid (BALF) was collected simultaneously and analyzed for cAMP levels. As demonstrated in **Figure 1A**, the 10.0 mg/kg dose of Compound 1 resulted in a peak plasma concentration at 0.5 hours of 1585.4 tPDE4i. The plasma level then decreased over the next 8 hours to a level of 96.1, and this low PDE4i of drug persisted at 24 hours. The BALF concentration of cAMP correlated with the drug level of the PDE4 inhibitor with a peak level of 27.4 pmol/mg protein occurring at the 0.5 hour time point (up from a basal concentration of 7.9 pmol/mg protein) (**Figure 1B**). By 8 hours, the BALF concentration of cAMP had returned to basal levels. **Figure 1C** demonstrates the linear increase in the BALF concentration of cAMP in direct relation to the increasing drug level of Compound 1 suggesting that BALF cAMP concentration can be used as a marker of drug effect in the target tissue.

Similar pharmacokinetics were observed following a single administration of roflumilast at 5.0 mg/kg (data not shown). The peak concentration was measured at 0.5 hours and returned to baseline at 8 hours. As was the case with Compound 1, the BALF concentration of cAMP correlated with the drug level of the PDE4 inhibitor with a peak level of 33.3 pmol/mg protein occurring at the 0.5 hour time point (up from a basal concentration of 10.9 pmol/mg protein) and returning near baseline by 8 hours (data not shown).



### **Prophylactic administration of piclamilast and roflumilast in the type II AEC injury model of lung fibrosis:**

To determine if PDE4 inhibition would mitigate pulmonary fibrosis that results from targeted type II AEC injury, we treated DTR+ mice with daily doses of intraperitoneal DT for 14 days. In previous studies, we demonstrated that this dosing regimen induced a progressive weight loss and lung fibrosis that was readily detected by day 21, one week after the last DT injection (13). To assess the efficacy of PDE4 inhibition in this model, we initially administered a prophylactic dosing regimen of piclamilast (30 mg/kg daily) or roflumilast (1.0 and 5.0 mg/kg) via oral gavage beginning on day 0 and continuing through day 21 of the model. By treating throughout the 21 day experimental time course, we sought to optimize our ability to detect a significant effect of PDE4 inhibition in our fibrosis model. We found that treatment of DTR+ mice with either piclamilast or roflumilast blunted the decline in body weight due to DT administration (**Figure 2A**). By day 11, the protection against weight loss afforded by these two drugs was statistically significantly different when compared to the vehicle-treated DTR+-DT injured mice. Treatment with either drug (and for roflumilast, both doses) resulted in a significant reduction in lung collagen content as measured by hydroxyproline compared to DT-injured DTR+ mice that were treated with vehicle (**Figure 2B**). The resulting level of lung hydroxyproline following PDE4 inhibition was similar to the uninjured control groups (i.e. DTR- mice treated with DT and DTR+ mice treated with PBS).

### **Prophylactic administration of Compound 1 and piclamilast in the type II AEC injury model of lung fibrosis:**

After confirming the efficacy of prophylactic PDE4 inhibition with piclamilast and roflumilast in reducing fibrosis following targeted type II AEC injury, we next assessed whether the novel PDE4 inhibitor Compound 1 would also protect against fibrosis in our murine model. Compound 1 was administered daily at doses of 1.0 mg/kg, 5.0 mg/kg and 10.0 mg/kg from day 0 through day 21. The efficacy of this molecule was compared to daily treatment with piclamilast (30 mg/kg). We found that all doses of Compound 1 lessened the DT-injury induced weight loss in DTR+ mice, and the protection afforded by this agent (at each dose) was comparable to piclamilast (**Figure 3A**). By day 15, the weight losses in the groups of DTR+-DT injured mice receiving the three doses of Compound 1 were significantly less than in the vehicle-treated DTR+-DT injured group. All doses of Compound 1 also significantly reduced the day 21 lung collagen content when compared to DT-injured DTR+ mice that were treated with

vehicle (**Figure 3B**). Also, the levels of lung hydroxyproline in the 3 dose groups of Compound 1 were not statistically different from the group that was treated with 30 mg/kg of piclamilast.

### **Therapeutic administration of Compound 1 and roflumilast in the type II AEC injury model of lung fibrosis:**

We next tested whether PDE4 inhibition would also diminish lung fibrosis when delivered in a therapeutic protocol. For these experiments, DTR+ mice were administered daily doses of DT from day 1 through day 14. From day 11 through day 21, subsets of mice were treated with Compound 1 at doses of 1.0 mg/kg or 5.0 mg/kg, roflumilast at 5.0 mg/kg, or vehicle. The time point of treatment initiation was chosen based on our prior studies which demonstrated that the onset of fibrosis was detectable around day 11 and that 10 days of treatment was sufficient to detect a therapeutic effect of a previously studied anti-fibrotic drug (19). Weight loss was tracked throughout the study, and the severity of lung fibrosis was assessed on day 21 by lung hydroxyproline concentration and picosirius red-stained histologic sections. At the time of treatment initiation on day 11, the mean body weight was equivalent in all DT-treated DTR+ cohorts. Treatment with the PDE4 inhibitors limited the further decline in body weight that was observed in the vehicle treated group from day 11 through day 21 (**Figure 4A**,  $p \leq 0.05$  on day 21 for Compound 1 (5.0 mg/kg) and Roflumilast (5.0 mg/kg)-treated mice compared to the vehicle treated group). In the 5.0 mg/kg doses of Compound 1 and roflumilast groups, the PDE4 inhibitor-mediated protection against weight loss was associated with a significant reduction in lung collagen content (**Figure 4B**). Of note, the 1.0 mg/kg therapeutic treatment dose of Compound 1 did not significantly reduce lung collagen content compared to the group that received vehicle, suggesting that this dose is below the efficacy threshold. The therapeutic administration of Compound 1 at 5.0 mg/kg also substantially reduced picosirius red staining (specific for collagen) and alveolar wall thickening characteristic of the histologic abnormalities observed in DT-injured DTR+ mice on day 21 (**Figure 4C**).

### **Compound 1 increases cAMP levels in A549 cells and reduces plasma surfactant protein D levels:**

After establishing both prophylactic and therapeutic efficacy of PDE4 inhibition in mitigating type II AEC injury-induced lung fibrosis, we performed studies to gain insight into the mechanisms by which this class of drugs exerts its anti-fibrotic effect. We first examined whether alveolar epithelial cells were responsive to PDE4 inhibition by measuring the level of cAMP following Compound 1 treatment. For these studies, A549 cells, a malignant human

alveolar epithelial cell line, were treated with increasing doses of Compound 1 (in the presence of 10  $\mu$ M PGE2 and 10  $\mu$ M forskolin to stimulate cAMP production). As demonstrated in **Figure 5A** and **Figure 5B**, Compound 1 induced a dose dependent increase in cAMP concentrations in both cell lysates and conditioned media. At lower concentrations of intracellular cAMP (below 20 pmol/mg protein), a linear relationship existed between intracellular and extracellular levels of this mediator (**Figure 5C**). After confirming activity in A549 cells, we examined whether Compound 1 (1.0 mg/kg, 5.0 mg/kg and 10.0 mg/kg) or piclamilast (30 mg/kg) modulated alveolar epithelial cell integrity following targeted type II AEC injury by measuring day 21 plasma surfactant protein D (SP-D) levels. Using the prophylactic treatment protocol, we found that all three doses of Compound 1 and the 30 mg/kg dose of piclamilast significantly reduced plasma SP-D compared to DT-injured DTR+ mice treated with vehicle (**Figure 6A**). Of note, there was a weak but statistically significant correlation between the plasma SP-D level at day 21 and lung collagen content in individual mice ( $r^2 = 18.7\%$ ,  $p=0.0053$ ) (**Figure 6B**).

#### **Compound 1 treatment decreases the plasma level of select chemokines:**

As another potential mechanism by which PDE4 inhibition reduces lung fibrosis in the setting of targeted type II AEC injury, we determined whether Compound 1 altered plasma levels of a select subset of chemokines. For these studies DT-injected DTR mice were treated with a therapeutic administration of Compound 1 at 5.0 mg/kg. Plasma was collected on day 21, and levels of eotaxin (CCL11), IP-10 (CXCL10), LIX (CXCL5) and Rantes (CCL5) were measured by bead-based multi-analyte profiling. We found that PDE4 inhibition with Compound 1 reduced the concentration of all of these mediators with the greatest effect on CXCL5 and CCL5 (**Figure 7**).

#### **Compound 1 treatment decreases fibrogenic gene expression in fibroblasts in vitro and blunts TNF $\alpha$ expression in vivo:**

Fibroblasts and myofibroblasts are considered the primary effector cells in tissue fibrosis. These cells are the primary source of key extracellular matrix constituents including the fibrillar collagens (including collagen 1) and fibronectin (Fn). These cells also secrete other profibrotic molecules including plasminogen activator inhibitor 1 (PAI-1) and connective tissue growth factor (CTGF). Prior studies have demonstrated that PDE4 inhibitors impede certain fibroblast functions including 3-dimensional gel contraction, migration, and myofibroblast differentiation (9-10). To assess whether PDE4 inhibition also influenced fibroblast expression of pro-fibrotic mediators, WI-38 human lung fibroblasts were treated with increasing doses of

Compound 1 in the presence of forskolin (1.0  $\mu$ M) and transforming growth factor- $\beta$  (TGF- $\beta$ , 3.0 ng/ml). We then measured the expression levels of Fn, collagen 1 $\alpha$ 1, PAI-1, and CTGF. Exposure of WI-38 cells to Compound 1 in the presence of forskolin increased the level of intracellular cAMP in a dose dependent manner (**Figure 8**). This increase in intracellular cAMP reduced the TGF- $\beta$  stimulated expression of Fn, collagen 1 $\alpha$ 1 chain, PAI-1, and CTGF in a dose-dependent manner (**Figure 8**).

To determine if treatment with Compound 1 also inhibited profibrotic gene expression in vivo, DT-injured DTR+ mice were treated with 5.0 mg/kg of Compound 1 from day 11 through day 21. On day 21, RNA was purified from lung tissue, and message levels for murine TNF $\alpha$ , collagen 1 $\alpha$ 1, PAI-1, CTGF and Fn were assessed by quantitative RT-PCR. We found that only TNF $\alpha$  expression was significantly reduced at the day 21 time point (**Figure 9B**). The mean expression of collagen 1 $\alpha$ 1, CTGF and Fn were also decreased, but the reduction did not achieve statistical significance from the vehicle treated DT-injured DTR+ group (**Figure 9A, C, D**).

#### **The therapeutic efficacy of PDE4 inhibition versus pirfenidone and nintedanib:**

After establishing a therapeutic benefit of roflumilast and Compound 1 in reducing type II AEC injury-induced fibrosis, we next compared the efficacy of PDE4 inhibition to treatment with the two FDA approved drugs for IPF (pirfenidone and nintedanib). DT was administered for 14 days according to our standard protocol. Beginning on day 11, subsets of DT-injured DTR+ mice were treated with vehicle, roflumilast (5 mg/kg once daily), pirfenidone (100 mg/kg 3 times daily), or nintedanib (100 mg/kg twice daily). For pirfenidone and nintedanib, doses were chosen based on PK profiles and published data in the bleomycin model. The mice were weighed daily, and on day 21, lungs were harvested for lung collagen content. Consistent with our prior results, we found that roflumilast blunted the weight loss induced by DT administration in the DTR+ animals. Pirfenidone also reduced weight loss compared to the vehicle treated group. In contrast, therapeutic administration of nintedanib accentuated weight loss in this fibrosis model (**Figure 10A**,  $p \leq 0.05$  on day 20-21 for roflumilast and pirfenidone-treated mice compared to the nintedanib-treated group). Measurements of lung hydroxyproline revealed that all three drugs effectively reduced type II AEC-injury induced fibrosis, and there was no difference in the anti-fibrotic efficacy of these agents at the given doses (**Figure 10B**).

## Discussion:

Diseases that are characterized by tissue fibrosis constitute major clinical challenges. Therefore, the approval of the first two drugs for patients with IPF, the prototypic scarring disorder of the lung, created excitement in the pulmonary community (20, 21). Despite this recent advance, there is still a desperate need for additional well-tolerated and efficacious treatments, and exploring the anti-fibrotic effect of already approved drugs with acceptable safety profiles offers an efficient approach to meet this demand. In this regard, we investigated the antifibrotic activity of both prophylactic and therapeutic PDE4 inhibitor administration in a type II AEC injury model of lung fibrosis. We found that 3 different members of this drug class including roflumilast (which is FDA approved for the treatment of COPD) and the novel agent Compound 1. The efficacy of Compound 1 in reducing lung fibrosis and blunting the associated weight loss was similar to roflumilast and piclamilast. Although roflumilast is known to show a body weight lowering effect in the clinical setting, treatment with the tested PDE4 inhibitors has a beneficial effect on weight loss in our fibrosis model compared to nintedanib and, less so, pirfenidone. Importantly, the antifibrotic activity of PDE4 inhibition in this model was equivalent to these two FDA-approved therapies for IPF.

A large number of potential antifibrotic therapies have been shown to reduce fibrosis following bleomycin-induced lung injury (22). Based on their promising effect in this rodent model, a subset of these agents has been tested in well-designed clinical trials with frustrating results. This lack of translation has led the fibrosis research community to question the utility of the bleomycin model in predicting the efficacy of prospective drugs. It should be noted, however, that the anti-fibrotic benefit of the majority of these drugs in the bleomycin model was established with preventative/prophylactic dosing. Because bleomycin induces inflammation which is a key element of the lung injury in this model, prophylactic administration of any drug that alters this inflammatory response is likely to impact fibrosis even if the agent lacks specific anti-fibrotic activity.

In light of this limitation of bleomycin, we chose to employ our type II AEC targeted injury model of lung fibrosis to evaluate the efficacy of the PDE4 inhibitors. This model shares pathogenic features with the human disease. For example, histologic studies consistently demonstrate type II cell defects including hyperplasia and apoptosis in patients with IPF (23-25). Also, AEC-specific gene mutations including surfactant proteins A2 and C are linked to familial lung fibrosis (26, 27). The limited and specific inflammatory response that occurs in this model may also better mimic the human disease (15). To further enhance the translatability of our

findings, we tested the efficacy of the PDE4 inhibitors not only in a preventative treatment regimen but also with therapeutic dosing (i.e. day 11-21). Although the potent effect of the PDE4 inhibitors in mitigating type II AEC injury-induced fibrosis should motivate the further development of this class of drugs for patients with IPF, we acknowledge that the predictive value of our model of fibrosis requires validation. It should also be mentioned that the two approved therapies for IPF (nintedanib and pirfenidone) comprise a small group of drugs that, when administered therapeutically, reduced bleomycin-induced fibrosis, suggesting that results from the bleomycin model, in the right context, can also guide clinical research decisions (28, 29). Furthermore, nintedanib and pirfenidone significantly reduce fibrosis in both the bleomycin and type II AEC injury models, supporting the notion that efficacy in several distinct models of fibrosis helps predict translatability to human disease.

Multiple anti-fibrotic mediators including PGE<sub>2</sub>, prostacyclin, and adenosine signal through cAMP, motivating our interest in evaluating the efficacy of PDE4 inhibitors in our animal model of fibrosis. Amongst the pleiotropic anti-fibrotic effects of PGE<sub>2</sub>, it has been shown to protect the lung epithelium. For example, prior studies reveal that PGE<sub>2</sub> inhibits Fas ligand induced apoptosis of type II AECs isolated from fibrotic lungs (8). This lipid mediator also promotes in vitro wound repair of airway epithelial cells (30). In our studies, treatment with Compound 1 effectively increased cAMP levels in an alveolar epithelial cell line, and we observed that the in vivo administration of this drug significantly reduced the plasma levels of SP-D following targeted type II AEC injury in mice. Of note, SP-D is increased in the serum of patients with IPF, and several studies indicate that this molecule, either alone or in conjunction with other proteins, functions as a diagnostic and prognostic biomarker in IPF (31-33). Although the mechanism by which serum levels of SP-D are increased in IPF and other lung fibrotic disorders is unknown, ongoing type II AEC injury in conjunction with hyperplasia likely contributes (34, 35). The PDE4 inhibitor-mediated decrease in the plasma concentration of SP-D after the cessation of DT supports an effect of this class of drugs on the alveolar epithelium as an element of their anti-fibrotic activity.

In addition to promoting the integrity of the alveolar epithelium, PGE<sub>2</sub> as well as prostacyclin analogs have been found to hinder a myriad of fibroblast functions. Specifically, PGE<sub>2</sub> and treprostinil inhibit TGF- $\beta$ -induced myofibroblast differentiation and promote the reversal of differentiated myofibroblasts to fibroblasts (36, 37, 38). In addition, PGE<sub>2</sub> impedes the proliferation and migration of fibroblasts and also increases their susceptibility to apoptosis (7, 8). In the setting of lung fibrosis, these anti-fibrotic activities of PGE<sub>2</sub> are mitigated by a down-regulation of its receptor in addition to an inhibition of its synthesis (39, 40). Through the

preservation of intracellular cAMP levels, PDE4 inhibitors have the capacity to augment the residual inhibitory actions of PGE<sub>2</sub> on fibroblasts, and indeed, prior studies have found that roflumilast slows migration and reduces 3-dimensional gel contraction by these cells (10). In the current study, we extend these findings by demonstrating that Compound 1-mediated PDE4 inhibition decreases in vitro fibroblast expression of several profibrotic mediators including Col 1a1, Fn and PAI-1. Of note, we did not find a significant PDE4 inhibitor-induced reduction in the expression of these same genes in vivo although we did observe a trend for decreased expression of Col 1a1, Fn. This lack of a statistically significant effect on pro-fibrotic gene expression may be a reflection of the chosen time point for these studies, as we have previously shown that lung collagen content does not appreciably increase beyond day 21 in the type II AEC injury model(13). Our findings might also be attributed to the dilution of a PDE4 inhibitor mediated effect in a specific cell population (i.e. fibroblast) resulting from the assessment of the whole lung RNA. We more likely would have detected a significant effect of PDE4 inhibition on profibrotic gene expression if measured specifically in fibroblasts on day 14 when collagen production is ongoing.

Beyond its effect of the alveolar epithelium and fibroblasts, it is also plausible that PDE4 inhibition limited fibrosis following type II AEC injury through an anti-inflammatory effect. Notably, our prior studies have implicated a specific monocyte/macrophage predominant inflammation in the development of type II AEC-injury induced fibrosis. In the current study, we found that Compound 1 treatment significantly reduced the plasma levels of a subset of chemokines that are upregulated in pulmonary fibrosis and/or are influenced by cAMP signaling. Specifically, CXCL5/LIX concentrations were found to be increased in the lavage fluid of systemic sclerosis patients with associated interstitial lung disease (41), and elevated levels of CCL5/RANTES were measured in the BALF of IPF patients (42). Furthermore, roflumilast treatment of lung fibroblasts has been found to decrease CCL5/RANTES and CXCL10/IP-10 expression(43). Finally, forskolin-mediated induction of cAMP levels was shown to inhibit CCL11/eotaxin production by bronchial epithelial cells (44). In addition to its effect on plasma chemokine levels, we also found that PDE4 inhibition decreased the day 21 lung expression levels of TNF $\alpha$ . Prior studies using adenoviral-mediated gene transfer suggest a direct role for this cytokine in fibrogenesis (45). On the other hand, the expression of TNF $\alpha$  in the alveolar compartment in the setting of a second hit (e.g. bleomycin) actually limited the fibrotic response (46). Furthermore, the intratracheal instillation of recombinant TNF $\alpha$  to bleomycin-injured mice with established fibrosis resulted in improvement in the severity of scarring (47). These disparate results suggest that the effect of TNF $\alpha$  on fibrosis is complex and may depend on the

timing and location of its expression. Ultimately, future studies are necessary to establish a causal link between the PDE4 inhibition-mediated reduction in cytokine/chemokine levels and the amelioration of fibrosis. However, the efficacy of both Compound 1 and roflumilast when initiated at a time point when the monocyte/macrophages have already maximally accrued suggests that the anti-fibrotic effect of PDE4 inhibition occurs predominantly through mechanisms other than blocking inflammation.

In conclusion, we have found that PDE4 inhibition significantly reduces lung fibrosis induced by targeted type II AEC injury. The antifibrotic activity of this class of drugs appears to be mediated through an array of cellular effects that promote the integrity of the alveolar epithelium and inhibit pro-fibrotic gene expression by lung fibroblasts. The equal potency of therapeutic PDE4 inhibition compared to the FDA approved drugs in a model of lung fibrosis that shares pathogenic features with the human disease serves as strong motivation to investigate the efficacy of roflumilast or Compound 1 in patients with IPF. The absence of CYP inhibitory activity of Compound 1 suggests that it will have fewer drug-drug interactions than roflumilast and may be a more attractive candidate for early phase clinical trials.

#### **Acknowledgments:**

This work was supported by grant funding from Takeda Pharmaceuticals and the NIH NHLBI: R01 HL078871 (THS) and R01 HL141195 (JCH).

#### **References:**

1. Hambly, N., Shimbori, C., and Kolb, M. 2015. Molecular classification of idiopathic pulmonary fibrosis: personalized medicine, genetics and biomarkers. *Respirology* 20:1010-1022.
2. Wolters, P.J., Collard, H.R., and Jones, K.D. 2014. Pathogenesis of idiopathic pulmonary fibrosis. *Annu Rev Pathol* 9:157-179.



3. Spagnolo, P., Wells, A.U., and Collard, H.R. 2015. Pharmacological treatment of idiopathic pulmonary fibrosis: an update. *Drug Discov Today* 20:514-524.
4. Raghu, G., Rochweg, B., Zhang, Y., Garcia, C.A., Azuma, A., Behr, J., Brozek, J.L., Collard, H.R., Cunningham, W., Homma, S., et al. 2015. An Official ATS/ERS/JRS/ALAT Clinical Practice Guideline: Treatment of Idiopathic Pulmonary Fibrosis. An Update of the 2011 Clinical Practice Guideline. *Am J Respir Crit Care Med* 192:e3-19.
5. Thomas, P.E., Peters-Golden, M., White, E.S., Thannickal, V.J., and Moore, B.B. 2007. PGE(2) inhibition of TGF-beta1-induced myofibroblast differentiation is Smad-independent but involves cell shape and adhesion-dependent signaling. *Am J Physiol Lung Cell Mol Physiol* 293:L417-428.
6. Penke, L.R., Huang, S.K., White, E.S., and Peters-Golden, M. 2014. Prostaglandin E2 inhibits alpha-smooth muscle actin transcription during myofibroblast differentiation via distinct mechanisms of modulation of serum response factor and myocardin-related transcription factor-A. *J Biol Chem* 289:17151-17162.
7. Huang, S.K., White, E.S., Wettlaufer, S.H., Grifka, H., Hogaboam, C.M., Thannickal, V.J., Horowitz, J.C., and Peters-Golden, M. 2009. Prostaglandin E(2) induces fibroblast apoptosis by modulating multiple survival pathways. *FASEB J* 23:4317-4326.
8. Maher, T.M., Evans, I.C., Bottoms, S.E., Mercer, P.F., Thorley, A.J., Nicholson, A.G., Laurent, G.J., Tetley, T.D., Chambers, R.C., and McAnulty, R.J. 2010. Diminished prostaglandin E2 contributes to the apoptosis paradox in idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 182:73-82.
9. Dunkern, T.R., Feurstein, D., Rossi, G.A., Sabatini, F., and Hatzelmann, A. 2007. Inhibition of TGF-beta induced lung fibroblast to myofibroblast conversion by phosphodiesterase inhibiting drugs and activators of soluble guanylyl cyclase. *Eur J Pharmacol* 572:12-22.
10. Togo, S., Liu, X., Wang, X., Sugiura, H., Kamio, K., Kawasaki, S., Kobayashi, T., Ertl, R.F., Ahn, Y., Holz, O., et al. 2009. PDE4 inhibitors roflumilast and rolipram augment PGE2 inhibition of TGF-beta1-stimulated fibroblasts. *Am J Physiol Lung Cell Mol Physiol* 296:L959-969.
11. Cortijo, J., Iranzo, A., Milara, X., Mata, M., Cerda-Nicolas, M., Ruiz-Sauri, A., Tenor, H., Hatzelmann, A., and Morcillo, E.J. 2009. Roflumilast, a phosphodiesterase 4 inhibitor, alleviates bleomycin-induced lung injury. *Br J Pharmacol* 156:534-544.
12. Milara, J., Morcillo, E., Monleon, D., Tenor, H., and Cortijo, J. 2015. Roflumilast Prevents the Metabolic Effects of Bleomycin-Induced Fibrosis in a Murine Model. *PLoS One* 10:e0133453.

13. Sisson, T.H., Mendez, M., Choi, K., Subbotina, N., Courey, A., Cunningham, A., Dave, A., Engelhardt, J.F., Liu, X., White, E.S., et al. 2010. Targeted injury of type II alveolar epithelial cells induces pulmonary fibrosis. *Am J Respir Crit Care Med* 181:254-263.
14. Osterholzer, J.J., Christensen, P.J., Lama, V., Horowitz, J.C., Hattori, N., Subbotina, N., Cunningham, A., Lin, Y., Murdock, B.J., Morey, R.E., et al. 2012. PAI-1 promotes the accumulation of exudate macrophages and worsens pulmonary fibrosis following type II alveolar epithelial cell injury. *J Pathol* 228:170-180.
15. Osterholzer, J.J., Olszewski, M.A., Murdock, B.J., Chen, G.H., Erb-Downward, J.R., Subbotina, N., Browning, K., Lin, Y., Morey, R.E., Dayrit, J.K., et al. 2013. Implicating exudate macrophages and Ly-6C(high) monocytes in CCR2-dependent lung fibrosis following gene-targeted alveolar injury. *J Immunol* 190:3447-3457.
16. Liu, Y., Lu, F., Kang, L., Wang, Z., and Wang, Y. 2017. Pirfenidone attenuates bleomycin-induced pulmonary fibrosis in mice by regulating Nrf2/Bach1 equilibrium. *BMC Pulm Med* 17:63.
17. Wollin, L., Maillet, I., Quesniaux, V., Holweg, A., and Ryffel, B. 2014. Antifibrotic and anti-inflammatory activity of the tyrosine kinase inhibitor nintedanib in experimental models of lung fibrosis. *J Pharmacol Exp Ther* 349:209-220.
18. Hermann, R., Nassr, N., Lahu, G., Peterfai, E., Knoerzer, D., Herzog, R., Zech, K., and de Mey, C. 2007. Steady-state pharmacokinetics of roflumilast and roflumilast N-oxide in patients with mild and moderate liver cirrhosis. *Clin Pharmacokinet* 46:403-416.
19. Sisson, T.H., Ajayi, I.O., Subbotina, N., Dodi, A.E., Rodansky, E.S., Chibucos, L.N., Kim, K.K., Keshamouni, V.G., White, E.S., Zhou, Y., et al. 2015. Inhibition of myocardin-related transcription factor/serum response factor signaling decreases lung fibrosis and promotes mesenchymal cell apoptosis. *Am J Pathol* 185:969-986.
20. Richeldi, L., du Bois, R.M., Raghu, G., Azuma, A., Brown, K.K., Costabel, U., Cottin, V., Flaherty, K.R., Hansell, D.M., Inoue, Y., et al. 2014. Efficacy and safety of nintedanib in idiopathic pulmonary fibrosis. *N Engl J Med* 370:2071-2082.
21. King, T.E., Jr., Bradford, W.Z., Castro-Bernardini, S., Fagan, E.A., Glaspole, I., Glassberg, M.K., Gorina, E., Hopkins, P.M., Kardatzke, D., Lancaster, L., et al. 2014. A phase 3 trial of pirfenidone in patients with idiopathic pulmonary fibrosis. *N Engl J Med* 370:2083-2092.
22. Moeller, A., Ask, K., Warburton, D., Gauldie, J., and Kolb, M. 2008. The bleomycin animal model: a useful tool to investigate treatment options for idiopathic pulmonary fibrosis? *Int J Biochem Cell Biol* 40:362-382.

23. Katzenstein, A.L. 1985. Pathogenesis of "fibrosis" in interstitial pneumonia: an electron microscopic study. *Hum Pathol* 16:1015-1024.
24. Kasper, M., and Haroske, G. 1996. Alterations in the alveolar epithelium after injury leading to pulmonary fibrosis. *Histol Histopathol* 11:463-483.
25. Kawanami, O., Ferrans, V.J., and Crystal, R.G. 1982. Structure of alveolar epithelial cells in patients with fibrotic lung disorders. *Lab Invest* 46:39-53.
26. Thomas, A.Q., Lane, K., Phillips, J., 3rd, Prince, M., Markin, C., Speer, M., Schwartz, D.A., Gaddipati, R., Marney, A., Johnson, J., et al. 2002. Heterozygosity for a surfactant protein C gene mutation associated with usual interstitial pneumonitis and cellular nonspecific interstitial pneumonitis in one kindred. *Am J Respir Crit Care Med* 165:1322-1328.
27. Wang, Y., Kuan, P.J., Xing, C., Cronkhite, J.T., Torres, F., Rosenblatt, R.L., DiMaio, J.M., Kinch, L.N., Grishin, N.V., and Garcia, C.K. 2009. Genetic defects in surfactant protein A2 are associated with pulmonary fibrosis and lung cancer. *Am J Hum Genet* 84:52-59.
28. Kakugawa, T., Mukae, H., Hayashi, T., Ishii, H., Abe, K., Fujii, T., Oku, H., Miyazaki, M., Kadota, J., and Kohno, S. 2004. Pirfenidone attenuates expression of HSP47 in murine bleomycin-induced pulmonary fibrosis. *Eur Respir J* 24:57-65.
29. Chaudhary, N.I., Roth, G.J., Hilberg, F., Muller-Quernheim, J., Prasse, A., Zissel, G., Schnapp, A., and Park, J.E. 2007. Inhibition of PDGF, VEGF and FGF signalling attenuates fibrosis. *Eur Respir J* 29:976-985.
30. Savla, U., Appel, H.J., Sporn, P.H., and Waters, C.M. 2001. Prostaglandin E(2) regulates wound closure in airway epithelium. *Am J Physiol Lung Cell Mol Physiol* 280:L421-431.
31. Takahashi, H., Fujishima, T., Koba, H., Murakami, S., Kurokawa, K., Shibuya, Y., Shiratori, M., Kuroki, Y., and Abe, S. 2000. Serum surfactant proteins A and D as prognostic factors in idiopathic pulmonary fibrosis and their relationship to disease extent. *Am J Respir Crit Care Med* 162:1109-1114.
32. Greene, K.E., King, T.E., Jr., Kuroki, Y., Bucher-Bartelson, B., Hunninghake, G.W., Newman, L.S., Nagae, H., and Mason, R.J. 2002. Serum surfactant proteins-A and -D as biomarkers in idiopathic pulmonary fibrosis. *Eur Respir J* 19:439-446.
33. White, E.S., Xia, M., Murray, S., Dyal, R., Flaherty, C.M., Flaherty, K.R., Moore, B.B., Cheng, L., Doyle, T.J., Villalba, J., et al. 2016. Plasma Surfactant Protein-D, Matrix Metalloproteinase-7, and Osteopontin Index Distinguishes Idiopathic Pulmonary Fibrosis From Other Idiopathic Interstitial Pneumonias. *Am J Respir Crit Care Med*.

34. Nishikiori, H., Chiba, H., Arika, S., Kuronuma, K., Otsuka, M., Shiratori, M., Ikeda, K., Watanabe, A., Kuroki, Y., and Takahashi, H. 2014. Distinct compartmentalization of SP-A and SP-D in the vasculature and lungs of patients with idiopathic pulmonary fibrosis. *BMC Pulm Med* 14:196.
35. Borensztajn, K., Crestani, B., and Kolb, M. 2013. Idiopathic pulmonary fibrosis: from epithelial injury to biomarkers--insights from the bench side. *Respiration* 86:441-452.
36. Epa, A.P., Thatcher, T.H., Pollock, S.J., Wahl, L.A., Lyda, E., Kottmann, R.M., Phipps, R.P., and Sime, P.J. 2015. Normal Human Lung Epithelial Cells Inhibit Transforming Growth Factor-beta Induced Myofibroblast Differentiation via Prostaglandin E2. *PLoS One* 10:e0135266.
37. Garrison, G., Huang, S.K., Okunishi, K., Scott, J.P., Kumar Penke, L.R., Scruggs, A.M., and Peters-Golden, M. 2013. Reversal of myofibroblast differentiation by prostaglandin E(2). *Am J Respir Cell Mol Biol* 48:550-558.
38. Corboz, M.R., Zhang, J., LaSala, D., DiPetrillo, K., Li, Z., Malinin, V., Brower, J., Kuehl, P.J., Barrett, T.E., Perkins, W.R., et al. 2018. Therapeutic administration of inhaled INS1009, a treprostinil prodrug formulation, inhibits bleomycin-induced pulmonary fibrosis in rats. *Pulm Pharmacol Ther* 49:95-103.
39. Huang, S.K., Fisher, A.S., Scruggs, A.M., White, E.S., Hogaboam, C.M., Richardson, B.C., and Peters-Golden, M. 2010. Hypermethylation of PTGER2 confers prostaglandin E2 resistance in fibrotic fibroblasts from humans and mice. *Am J Pathol* 177:2245-2255.
40. Gabasa, M., Royo, D., Molina-Molina, M., Roca-Ferrer, J., Pujols, L., Picado, C., Xaubet, A., and Pereda, J. 2013. Lung myofibroblasts are characterized by down-regulated cyclooxygenase-2 and its main metabolite, prostaglandin E2. *PLoS One* 8:e65445.
41. Hesselstrand, R., Wildt, M., Bozovic, G., Andersson-Sjoland, A., Andreasson, K., Scheja, A., Westergren-Thorsson, G., Bjermer, L., and Wuttge, D.M. 2013. Biomarkers from bronchoalveolar lavage fluid in systemic sclerosis patients with interstitial lung disease relate to severity of lung fibrosis. *Respir Med* 107:1079-1086.
42. Kodama, N., Yamaguchi, E., Hizawa, N., Furuya, K., Kojima, J., Oguri, M., Takahashi, T., and Kawakami, Y. 1998. Expression of RANTES by bronchoalveolar lavage cells in nonsmoking patients with interstitial lung diseases. *Am J Respir Cell Mol Biol* 18:526-531.
43. Tannheimer, S.L., Wright, C.D., and Salmon, M. 2012. Combination of roflumilast with a beta-2 adrenergic receptor agonist inhibits proinflammatory and profibrotic mediator release from human lung fibroblasts. *Respir Res* 13:28.

44. Chu, Y.T., Chang, T.T., Jong, Y.J., Kuo, P.L., Lee, H.M., Lee, M.S., Chang, H.W., and Hung, C.H. 2010. Suppressive effects of formoterol and salmeterol on eotaxin-1 in bronchial epithelial cells. *Pediatr Allergy Immunol* 21:345-352.
45. Sime, P.J., Marr, R.A., Gauldie, D., Xing, Z., Hewlett, B.R., Graham, F.L., and Gauldie, J. 1998. Transfer of tumor necrosis factor-alpha to rat lung induces severe pulmonary inflammation and patchy interstitial fibrogenesis with induction of transforming growth factor-beta1 and myofibroblasts. *Am J Pathol* 153:825-832.
46. Fujita, M., Shannon, J.M., Morikawa, O., Gauldie, J., Hara, N., and Mason, R.J. 2003. Overexpression of tumor necrosis factor-alpha diminishes pulmonary fibrosis induced by bleomycin or transforming growth factor-beta. *Am J Respir Cell Mol Biol* 29:669-676.
47. Redente, E.F., Keith, R.C., Janssen, W., Henson, P.M., Ortiz, L.A., Downey, G.P., Bratton, D.L., and Riches, D.W. 2014. Tumor necrosis factor-alpha accelerates the resolution of established pulmonary fibrosis in mice by targeting profibrotic lung macrophages. *Am J Respir Cell Mol Biol* 50:825-837.

#### Figure Legends:

**Figure 1:** The pharmacokinetics of Compound 1 administration. 10.0 mg/kg of Compound 1 was administered by oral gavage at time 0. Plasma and BALF were collected simultaneously from separate cohorts of mice at 0.5, 1, 2, 4, 8 and 24 hours after the dosing. The plasma concentration of drug over time (1A) and the BALF cAMP levels over time (1B) are reported. A linear relationship between plasma drug level and BALF cAMP concentration was observed (1C). Data is reported as mean  $\pm$  SD (n= 4/group).

**Figure 2:** Prophylactic administration of roflumilast and piclamilast to targeted type II AEC-injured mice reduces weight loss and ameliorates pulmonary fibrosis. DTR-expressing mice (DTR+) were administered daily I.P. PBS or DT from day 0 through Day 14. Subsets of the DTR+:DT-injured animals were treated by oral gavage once daily beginning on day 0 with vehicle, roflumilast (1.0 mg/kg or 5.0 mg/kg), or Piclamilast at 30 mg/kg. Mice were weighed daily (A) and on day 21, lungs were harvested and analyzed for total collagen content using an assay for hydroxyproline (B). Data is reported as mean  $\pm$  SEM (n = 10 per group). Weight curves are analyzed with a two-way ANOVA + Tukey multiple comparison tests (# p  $\leq$  0.05 for

DTR+:DT Piclamilast-treatment versus DTR+:DT vehicle treatment. ##  $p \leq 0.05$  DTR+:DT piclamilast and roflumilast (1.0 mg/kg or 5.0 mg/kg) treatment versus DTR+:DT vehicle treatment), and hydroxyproline is analyzed with a one-way ANOVA + Tukey multiple comparison tests.

**Figure 3:** Prophylactic administration of Compound 1 to targeted type II AEC-injured mice reduces weight loss and ameliorates pulmonary fibrosis. DTR-expressing mice (DTR+) were administered daily I.P. PBS or DT from day 0 through Day 14. Subsets of the DTR+:DT treated animals were treated by oral gavage once daily beginning on day 0 with vehicle, Compound 1 (1.0 mg/kg, 5.0 mg/kg, or 10.0 mg/kg) or Piclamilast at 30 mg/kg. Mice were weighed daily (A) and on day 21, lungs were harvested and analyzed for total collagen content using an assay for hydroxyproline (B). Data is reported as mean  $\pm$  SEM (n = 8 per group). Weight curves are analyzed with a two-way ANOVA + Tukey multiple comparison tests (#  $p \leq 0.05$  for DTR+:DT piclamilast and Compound 1-treatments (all doses) versus DTR+:DT vehicle treatment), and hydroxyproline is analyzed with a one-way ANOVA + Tukey multiple comparison tests.

**Figure 4:** Therapeutic administration of Compound 1 and roflumilast to targeted type II AEC-injured mice reduces weight loss and ameliorates pulmonary fibrosis. DTR-expressing mice (DTR+) were administered daily I.P. PBS or DT from day 0 through Day 14. Subsets of the DTR+:DT treated animals were treated by oral gavage once daily beginning on day 11 with vehicle, roflumilast (5.0 mg/kg), or Compound 1 (1.0 mg/kg or 5.0 mg/kg). Mice were weighed daily (A) and on day 21, lungs were harvested and analyzed for total collagen content using an assay for hydroxyproline (B) and by histopathology (C) with representative sections stained with picrosirius red (400x magnification). Data is reported as mean  $\pm$  SEM (n = 10-12 per group). Weight curves are analyzed with a two-way ANOVA + Tukey multiple comparison tests (#  $p \leq 0.05$  for DTR+:DT roflumilast and Compound 1 (5.0 mg/kg)-treatments versus DTR+:DT vehicle treatment), and hydroxyproline is analyzed with a one-way ANOVA + Tukey multiple comparison tests.

**Figure 5:** Compound 1 treatment of A549 human lung epithelial cell cultures increases intracellular and extracellular cAMP concentrations. A549 human lung epithelial cells were seeded in a 24 well plate for 24 hours. The cultures were treated with increasing concentrations

of Compound 1 ( $1 \times 10^{-10}$  M -  $1 \times 10^{-5}$ ) for 30 minutes and stimulated with 10  $\mu$ M forskolin and 10  $\mu$ M prostaglandin E2 for 30 minutes. cAMP concentrations were measured in cell lysates (A) and culture supernatants (B). The correlation between intracellular cAMP and cAMP in the culture supernatants is shown in (C). Data is reported as mean + SD (n = 3 per group).

**Figure 6:** Prophylactic treatment of targeted type II AEC injured mice with Compound 1 reduces plasma SP-D levels. DTR-expressing mice (DTR+) were administered daily I.P. PBS or DT from day 0 through Day 14. Subsets of the DTR+:DT treated animals were treated by oral gavage once daily beginning on day 0 with vehicle, Compound 1 (1.0 mg/kg, 5.0 mg/kg, or 10.0 mg/kg) or Piclamilast at 30 mg/kg. Plasma was collected on day 21 and analyzed for SP-D concentration (A). The plasma SP-D concentration was then correlated with lung collagen content (B). Data is reported as mean +SEM with individual data points shown (n = 8 per group). #:  $p \leq 0.05$  vs. Vehicle group by two-tailed Williams' test. \$:  $p \leq 0.05$  vs. Vehicle group by Student's t-test.

**Figure 7:** Therapeutic treatment of targeted type II AEC injured mice with Compound 1 reduces plasma cytokine levels. DTR-expressing mice (DTR+) were administered daily I.P. PBS or DT from day 0 through Day 14. Subsets of the DTR+:DT treated animals were treated by oral gavage once daily beginning on day 0 with vehicle or Compound 1 at 5.0 mg/kg. Plasma was collected on day 21 and analyzed for CCL11/eotaxin, CXCL10/IP-10, CXCL5/LIX, and CCL5/Rantes. Data is reported as mean  $\pm$  SD with individual data points shown (n = 13 for control, 22 for Compound 1 per group). #:  $p \leq 0.05$  vs. Control group by Student's t-test.

**Figure 8:** Compound 1 treatment of human lung fibroblasts cell cultures increases intracellular cAMP and decreases TGF- $\beta$ -mediated induction of fibrotic gene expression. WI-38 human lung fibroblasts were seeded on 24-well plates and treated with increasing concentrations of Compound 1 ( $1 \times 10^{-10}$  M -  $1 \times 10^{-5}$  M). After 30 min, cells were stimulated by 1  $\mu$ M forskolin for 30 minutes and cell lysates were analyzed for cAMP concentration (A). To analyze fibroblast gene expression, WI-38 human lung fibroblasts were cultured in media containing 0.5% FBS for 24 hours and treated with increasing concentrations of Compound 1 ( $1 \times 10^{-10}$  M -  $1 \times 10^{-5}$  M) for 1 hour followed by TGF- $\beta$  (3 ng/mL) and forskolin (1  $\mu$ M) for 24 hours. Total RNA was extracted

from cell lysates, cDNA was amplified, and target gene mRNA was measured. The target gene expression levels of Col1a1 (Type-1 collagen), Fn (Fibronectin), CTGF (connective tissue growth factor) and PAI-1 (plasminogen activator inhibitor-1) was normalized to Glyceraldehyde 3-phosphate dehydrogenase (GAPDH). Data is reported as mean  $\pm$  SD with individual data points shown (n = 3 per group). #;  $p \leq 0.05$  vs. forskolin group by two-tailed Williams' test.


**Figure 9:** Therapeutic administration of Compound 1 to targeted type II AEC injured mice reduces the expression of TNF $\alpha$  within the lung. DTR-expressing mice (DTR+) were administered daily I.P. PBS or DT from day 0 through Day 14. Subsets of the DTR+:DT treated animals were treated by oral gavage once daily beginning on day 11 with vehicle or Compound 1 at 5.0 mg/kg. On day 21, the left lung was harvested and homogenized, and total RNA was extracted. First-strand cDNA was synthesized and mRNA levels for Col1a1, Fibronectin, CTGF TNF $\alpha$  and PAI-1 (plasminogen activator inhibitor-1) were assessed using SYBR Green-based detection. The expression levels were normalized to GAPDH using the following formula: %GAPDH expression =  $100 / 2^{-\Delta\Delta CT}$ . Data is presented as an average  $\pm$  SEM (n = 9 per DTR+:DT vehicle and drug treated groups).

**Figure 10:** Roflumilast is equivalent to pirfenidone and nintedanib in ameliorating fibrosis following targeted type II AEC injury. DTR-expressing mice (DTR+) were administered daily I.P. PBS or DT from day 0 through Day 14. Subsets of the DTR+:DT treated animals were treated by oral gavage beginning on day 11 with vehicle, roflumilast at 5.0 mg/kg once daily, pirfenidone at 100.0 mg/kg three times daily, or nintedanib 100.0 mg/kg two times daily. Mice were weighed daily (A) and on day 21, lungs were harvested and analyzed for total collagen content using an assay for hydroxyproline (B). Data is reported as mean  $\pm$  SEM (n = 6-10 per group). Weight curves are analyzed with a two-way ANOVA + Tukey multiple comparison tests (#  $p \leq 0.05$  for DTR+:DT roflumilast and pirfenidone-treatment versus DTR+:DT nintedanib treatment, and hydroxyproline is analyzed with a one-way ANOVA + Tukey multiple comparison tests.




# Author Manuscript


Figure 1



**Figure 2**



**Figure 3**



**Figure 4**






Figure 4

Author Manuscript<sup>c</sup>




DTR:DT




DTR:DT:Cmp1

Figure 5

A cAMP in cell lysate



B cAMP in supernatant



C Correlation




Figure 6







Figure 7



**Figure 8**




Figure 9




Figure 10

