

Title: Allostatic Load: A Useful Concept for Advancing Nursing Research

Running head: Allostatic Load and Nursing Research

Authors: Marie-Anne S. Rosemberg, PhD, MN, RN
Assistant Professor
Department of Systems, Populations and Leadership
University of Michigan, School of Nursing
Email: sanon@umich.edu

Yang Li, RN, PhD Candidate
University of Michigan, School of Nursing
Email: lyx@umich.edu

Julia Seng, PhD, RN CNM, FAAN
Professor
Department of Systems, Populations and Leadership
University of Michigan, School of Nursing
Email: jseng@med.umich.edu

Corresponding Author: Marie-Anne S. Rosemberg, PhD, RN
400 North Ingalls, Room 3175, Ann Arbor, MI, 48109,
Email: sanon@umich.edu
Tel: (734)-647-0146 (office)

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/jocn.13753](https://doi.org/10.1111/jocn.13753)

This article is protected by copyright. All rights reserved

DR. MARIE-ANNE SANON ROSEMBERG (Orcid ID : 0000-0003-0536-2062)

Received Date : 30-Sep-2016

Revised Date : 19-Jan-2017

Accepted Date : 20-Jan-2017

Article type : Discursive Paper

ABSTRACT

Aims and objectives. To elucidate the historical development of the allostatic load concept, alongside its use in nursing research, and to explore how allostatic load has been investigated among two stress-vulnerable populations.

Background. “Stress” is a prominent term in understanding the development of disease. Allostatic load is among several approaches undertaken to quantify the magnitude of stress and understand how stress can affect health.

Method. We explored the advent of allostatic load including its antecedents, and consequences. We used an exemplar case to apply the concept. We reviewed studies that used allostatic load among workers and women of childbearing age.

Results. There remains a need to consolidate a common definition and operationalization of allostatic load. Despite this need for further work, allostatic load is a good fit for nursing science which focuses on the client, environment, and health. Only 12 studies explored allostatic load among workers (n=6) and women of childbearing age (n=6). In some studies allostatic load was used as a predictor while in others it was used as an outcome. None of the studies considered it as a mediator.

This article is protected by copyright. All rights reserved

Conclusions. The concept of allostatic load holds promise for nursing researchers to operationalize a holistic view of multiple stressors and to quantify their effects on health. Studies are needed to affirm the role of allostatic load as a potential mediator between multiple stressors and outcomes. Longitudinal studies are also needed to demonstrate a causal pathway from stressor exposure to tertiary outcomes such as chronic conditions and morbidity.

Relevance to clinical practice. Allostatic load is a useful concept for nurses working with stress-vulnerable populations. With the use of an interpretable allostatic load index, nurses will be able to intervene at various stages of the allostasis-adaptation process (stress-response) and adjust interventions accordingly.

Key words: allostatic load, concept analysis, stress, nursing research, workers, women of childbearing age, vulnerable populations, biomarkers

SUMMARY BOX

What does this paper contribute to the wider global clinical community?

- Nurse researchers and clinicians working with stress-vulnerable populations need clear points of intervention and ability to determine impact of the interventions in the short term.
- The concept of allostatic load is useful at broader levels, including further developing measurement of multiple levels of stress that could take structural inequalities, toxic and traumatic stress into account as predictors in all studies. This concept is also useful in advocating for policies that decrease stress and allostatic load as a way to build a culture of health and decrease burden, morbidity, and costs of stress-related disease.
- In the future, nursing is well-positioned to place the concept of allostatic support within the theoretical framework and operationalize it—in research and in practice.

AIMS

Nursing's meta-paradigm defines the basis of nursing's work as focused on the person, environment, health, and nursing (Fawcett 1984). Stressors are a ubiquitous

aspect of “environment”. Types of stressors that must be taken into account include those at each eco-social level which are cumulating and interacting, including social determinants, marginalized identities, socio-economics, interpersonal, and intra-personal factors (Juster *et al.* 2010). The literature is replete with evidence supporting a strong relationship between socio-environmental stressors and health outcomes (National Center for Health Statistics 2012, Cohen *et al.* 2013). Nursing care includes health promotion and risk reduction interventions that address this stress diathesis. These are necessarily complex because the stressors and adverse effects are complex while the clients are diverse. Nursing research could be enhanced by a theory that elucidates a mechanism for assessment of the near-term impact of our interventions (i.e., have a valid proxy endpoint) without having to await the long-term end-point of morbidity or mortality. The theory of allostasis, and the core concept of allostatic load are excellent candidates for this purpose. The aim of this discursive paper is to provide a concept analysis of “allostatic load” to advance its use in nursing research.

BACKGROUND

McEwen (1998) proposed the concept of allostatic load (AL) to explicate how chronic life stressors, including toxic and traumatic stress, impact individuals' health via the physiologic responses to such chronic stressors. The concept of AL has been applied in research across various disciplines and findings have generally confirmed that cumulative effects of social and environmental stressors increase the risks for physiologic dysregulation and ill-mental and physical health, especially among vulnerable groups (Seeman *et al.* 1997, McEwen 2000, McEwen 2004, Read & Grundy 2012, Seeman *et al.* 2014). The clinical value of this concept is that it could serve as a signal of health risk early enough (i.e., when the physiologic dysregulation is still sub-clinical) to lead to interventions that may prevent further deterioration of health and thus prevent future associated morbidity and mortality. For research purposes, it could also serve as a near-term proxy outcome or end-point in clinical health promotion and risk reduction studies (Juster *et al.* 2010). Despite its promise AL remains underutilized in nursing research.

There may be several reasons for underuse of this concept (McEwen & Wingfield 2010, Read & Grundy 2012). A recent systematic review by a nurse researcher exploring the concept of AL among 58 studies indicated the lack of homogeneity in the operationalization and measurement of AL (Beckie 2012). Beckie (2012) also noted inconsistencies in the study results. For example, some studies were able to show a strong relationship between socioeconomic status (SES) and AL biomarkers (Seeman *et al.* 2004), while others did not (Dowd & Goldman 2006). There were also variations in the selection and number of indicators used in operationalizing the concept, which usually are added together into an AL index (Juster *et al.* 2010). There's also a lack of consensus about how best to score the AL index (Beckie 2012). Thus, work remains to advance this promising concept into one that can be broadly and consistently used for health research.

DESIGN AND METHOD

In this discursive paper we extend Beckie's (2012) work. Having selected a concept and determined the aim, we will follow the rest of the steps outlined by Walker and Avant (2005). The work to identify uses of the concept and determine defining attributes has been accomplished by others (McEwen & Seeman 1999), but we will summarize. We will identify antecedents and consequences. The concept has already been situated in a theoretical framework (Beckie 2012), but we will illustrate this with a model case. We will discuss issues related to empirical referents. Finally, we will extend focus on cases by looking at two small sets of studies that used AL to research stress effects on health of workers and childbearing women.

Evolution of the concept

Homeostasis

In 1932, Cannon introduced the term homeostasis to describe the tight regulation of physiological and biochemical function (Cannon 1932). Homeostasis involves the maintenance of balance within the bodily systems. With self-correcting negative feedback actions, it could reduce variability and maintain constancy from those systems where invariability is a characteristic of a healthy system (Sterling & Eyer 1988). Those physiological systems work in concert to reestablish the body's initial conditions

whenever one or more of the systems exhibited perturbation. As such, homeostatic systems are those where a narrow physiological range is indicative of health and deviance from this range is an indication of pathology (e.g., an elevated body temperature [Carlson & Chamberlain 2005]). However, the homeostasis concept of stable states and feedback loops is insufficient to capture the complexity of physiologic systems or account for the variability of integrative networks of adaptation to environmental stressors (Goldberger *et al.* 2002). Hence the theory of allostasis was developed.

Allostasis

Allostasis refers to the mechanisms through which physiological systems adapt to a changing environment or to stressful challenges (Sterling & Eyer 1988, Karlamangla *et al.* 2002, McEwen 2002). The resting points change according to dynamic biological processes. Variability is a healthy adaptive mechanism in response to environmental demands, which is in contrast to homeostasis which supports stability (Carlson & Chamberlain 2005). Sterling and Eyer (1988) proposed allostasis as the process of achieving stability through change.

AL and Allostatic Overload

McEwen and Stellar (1993) elucidated the concept of AL as a multi-systemic approach to understand the cumulative effects of stress on health as the body responds to stressors that are chronic or severe enough to force adaptation. This central concept, AL, refers to the accumulation of wear and tear on interacting physiological systems from the adaptation process (McEwen & Seeman 1999). This physiologic wear and tear is a natural consequence or the price paid for the adaptation. Variation in AL among individuals may reflect individual differences in exposure to stressors and/or the ability to adapt to environments and challenges (Lipowicz *et al.* 2014). It may also reflect allostatic supports they find or receive, which can be multiple and synergistic (Friedman & McEwen 2004). In essence, given stressors and physiologic responses to them, if the allostatic load is not too high and adaptation occurs, health is likely to be maintained. If the allostatic load is too high (i.e., if there is allostatic overload), the price paid is dysregulation across multiple systems which leads to disease development. Read and

Grundy (2012) defined (high) AL as “a sub-clinical dysregulation state, resulting from the body’s response to stress” (p.1). Allostatic overload is a more extreme form of AL. Allostatic overload, which is AL at the pathological level, can result from sustained, severe, or repeated stress, the failure to habituate to repeated challenge, the inability to shut off allostatic responses, and inadequate allostatic responses (McEwen & Wingfield 2010).

Antecedents and consequences

Antecedents refer to the events or attributes that must precede the occurrence of a concept (Walker & Avant 2005). Antecedents that occur prior to AL include psychosocial factors (e.g. low socioeconomic status, stressful life events including trauma) and individual factors (e.g., posttraumatic stress, isolation, or maladaptive coping). These can be thought of as occurring at all eco-social levels (Bronfenbrenner & Morris 2006) and so would include structural inequalities and time-specific stressors, such as exposure to a disaster, war, or famine.

A variety of types of stressors have been considered as antecedents. For example, in the National Health and Nutrition Examination Surveys, the antecedent, race, was associated with AL. Blacks had higher AL index scores than did Whites at all ages (Geronimus *et al.* 2006). The differences in AL index scores increased with age. Black women, in particular, had higher AL index scores compared with either Black men or White women. Lower SES (lower education, occupational status, income) and greater social challenges (recent widowhood, high demands) have been reported to be highly correlated with higher AL (Weinstein *et al.* 2003). A study in alzheimer patients’ caregivers showed a greater number of negative life events related to higher AL (von Kanel *et al.* 2003). Among the elderly, ties with close friends and/or neighbors have been reported in relation to lower AL for both men and women (Seeman *et al.* 2004). Type A personality traits were associated with higher AL in a large sample (Sun *et al.* 2007). Inactivity and poor diet have also been reported in relation to higher AL (Juster *et al.* 2010).

Consequences refer to the events that result from the occurrence of a concept (Walker & Avant 2005). Consequences that could occur as a result of AL include the

leading causes of death for vulnerable populations whose stress levels are higher by definition. Consequences of AL include negative health outcomes such as cardio- and cerebro-vascular disease, cognitive deficits, weaker physical performance, depression, and premature mortality (Seplaki *et al.* 2004, Szanton *et al.* 2009). In the MacArthur Studies of Successful Aging, AL index score explained 35% of socioeconomic variation in mortality (Seeman *et al.* 2004).

Theoretical framework and model case

The British Economic and Social Research Council's National Centre for Research Methods Working Paper elaborated a conceptual framework with three stages in the allostasis-adaptation process (Read & Grundy 2012, p. 3). The primary mediators between stress and outcomes are neuroendocrine responses (i.e., stress regulation hormones such as cortisol, oxytocin, or catecholamines). Secondary outcomes are (potentially still sub-clinical) dysregulations that can be seen in immune, metabolic, cardiovascular, and anthropometric indicators (i.e., c-reactive protein, glucose, blood pressure, hip-waist ratio). Tertiary outcomes are the clinical manifestations that develop as dysregulations accumulate (i.e., poor subjective health, cognitive decline, disease states, and premature death [Read & Grundy 2012]). The premise and promise of AL is that a cumulative index that captures levels of multiple biomarker or anthropometric indicators can signal overload in time to provide a clinical response and prevent adverse outcomes.

A model case scenario can illustrate this allostasis-adaptation process (see Figure 1). Imagine first, a scenario where overload occurs. A 22-year-old Latina immigrant hotel housekeeper gave birth by cesarean to an infant with a genetic anomaly who was admitted to the NICU. The stressors are young age, immigration status, non-native English speaker, low wages, surgical delivery, and separation from the infant. Primary mediators involve elevated cortisol and dysregulated oxytocin. Secondary outcomes appear as sleep alterations with changes in insulin levels, altered immune and inflammatory responses, and mildly elevated blood pressure. The tertiary outcomes manifest as near- and long-term outcomes. In the near term, she experiences fatigue, weight gain, delayed bonding, and slow cesarean wound healing. In the longer term she

develops type 2 diabetes, depression, chronic pain from adhesions, and hypertension. Her adaptation to motherhood is not fulfilling to her, and she chooses not to risk having any additional children, especially given depression and early ill health. It would have been possible, however, to have allostasis and adaptation be the result without adverse outcomes. Imagine an alternative scenario in which her stress had been reduced with a regular Spanish interpreter in the NICU, ability to Skype with her mother in Mexico, adequate insurance to prevent worry, maternity leave to accommodate the slow recovery from surgery and the need to be with the infant, as well as nursing home visiting to support her maternal development, monitor her low mood, and provide health promotion interventions to support her in regaining healthy sleep and diet until her blood glucose and blood pressure normalize. Her adaptation to mothering a child with chronic health needs becomes a source of pride, and she channels some of her healthy energy into becoming an interpreter in the hospital's NICU.

Empirical referents

Empirical referents are the measurable ways to validate the occurrence of a concept (Walker & Avant 2005). To operationalize AL, measurement of biomarker and anthropometric indicators chosen to represent the primary mediator response and secondary outcomes are combined in a clinometric index—an AL-index score. When AL was first proposed, an AL index of 10 biomarkers was used to measure the concept. The 10 original AL measures included systolic blood pressure (SBP), diastolic blood pressure (DBP), waist-hip ratio (WHR), high density lipoprotein (HDL), total HDL/cholesterol ratio, total cholesterol, dehydroepiandrosterone sulfate (DHEA-S), urinary free cortisol, norepinephrine, and epinephrine (Seeman *et al.* 1997). However, studies have since used a variety of biomarkers, some using additional biomarkers (Goldman *et al.* 2006, Gleib *et al.* 2007, Bellingrath *et al.* 2009) and others using fewer (Gersten 2008, Loucks *et al.* 2008, Evans & Schamberg 2009) based on the outcomes of interest that they are trying to predict or feasibility.

In measuring AL, each of the different physiologic systems must be represented. However, there are inconsistencies in the combination of indicators included in AL index across studies (Beckie 2012, Read & Grundy 2012). Given the variation in which

indicators are included in the AL index, it is not surprising that there are inconsistencies in study findings.

Scoring is also a source of inconsistency. A summation approach is the most often used method to calculate the index score. Individuals are ranked by the indicator values, ranging between the highest and lowest risks (Seeman *et al.* 1997). Individuals in the highest risk quartile for each indicator are “1” and others are “0”. These are summed so that those with the highest sum score have the highest AL and are considered at risk for poor tertiary outcomes (Read & Grundy 2012). Thus, AL has been operationalized as a count-based multi-systemic index representing the sum of biomarkers and anthropometric indicators falling within a high-risk percentile based on the sample's distribution of values (Juster *et al.* 2010). This approach presents some unique challenges in that there is loss of granularity in the data reduction and the amount of risk depends on the nature of the sample itself. Alternative scoring approaches have been considered (Karlmanngla *et al.* 2002, Seplaki *et al.* 2005, Gruenewald *et al.* 2006) and more consensus may emerge in the near future.

Extending the analysis of AL concept from clinical case to research examples with two vulnerable populations

The model case scenario above was a clinical example at how the concept of AL could function within a single individual. In the near term, it is more likely that AL's usefulness in nursing will be as a concept deployed in research. So we'll extend the illustration of AL as a concept linking stress and adverse health outcomes by exploring studies reported about two stress-vulnerable populations, workers and women of childbearing age.

The workplace represents a major source of stress and thus can impact AL leading to poor health outcomes. First, work relates to SES, and lower SES has shown to be strongly correlated with high AL (Szanton *et al.* 2005). Second, work where employees experience high demand and low control (a phenomenon known as job strain), results in high stress (Karasek & Theorell 1992). Some workers are therefore at risk for high stress and poor health outcomes, making it important to explore AL among workers.

Trauma exposure and sequelae are strongly related to gender, and pregnancy is another stress exposure unique to females, making women another key group to study in relation to AL. McEwen and Seeman (1999) pointed out that early traumatic events combined with ongoing life stress could contribute to AL. In a study with a random sample of 1,442 subjects from the United States, 32.3% of women reported childhood sexual abuse (CSA), which was twice as much as in men (Briere & Elliott 2003). In a review of population-based studies, 10% to 69% of women reported intimate partner violence (Krug *et al.* 2002). The peak age of trauma exposure for women is in adolescence, so posttraumatic stress—where re-experiencing the trauma in memory, flashbacks, or nightmares is a hallmark of the disorder and an intra-personal chronic stress—is also prevalent early in the lifespan and during reproduction (Breslau 2002). There is growing evidence suggesting the causal links between adverse childhood events, adult traumatic events, traumatic stress sequelae, and physical and reproductive health outcomes (Felitti *et al.* 1998, Campbell 2002, Groer *et al.* 2016). Thus women of childbearing age who are experiencing traumatic stress are another priority population among which to explore the mediating role of AL on the association of stress with adverse health outcomes.

To find articles with which to explore these research “cases” to illustrate the AL model, we searched PubMed, Web of Science, CINAHL, PsycInfo, and Scopus databases using the keywords: AL and workers and AL and women of childbearing age. Our initial search yielded 207 articles across all the databases. We only included articles that were written in English and for which full texts were provided. We then limited selection to the handful that specifically used the theory of allostasis as their conceptual framework for the study in relation to workers (6 papers, Table 1) or women of childbearing age (6 papers, Table 2).

AL and workers

Six studies were identified exploring AL among workers. The studies were conducted in industrialized nations, included men and women, and focused primarily on industry employees. The studies took place in across several countries including the US, Sweden, Germany, and China. There were significant variations in the

operationalization of AL across the studies. The number of indicators used in the AL index ranged from 6 to 14 (Table 3). All the studies included primary and secondary indicators in the AL index, and all the studies reviewed used the high risk scoring approach (e.g., highest quartile for BMI or glucose, lowest quartile for HDL cholesterol or DHEA-S). In all of the studies, AL was used as a dependent variable (an outcome). None of the studies used AL as a mediator or an independent variable (a predictor). All of the studies found a positive relationship between the stressors (i.e. job strain, recovery from work stress) and AL index score. None of the studies explored tertiary outcomes such as chronic diseases, quality of life, or mortality.

AL and women of childbearing age

Two of the articles focused on pregnant women to examine the relationships between AL and adverse birth outcomes. All six studies included women of childbearing age, with the age ranging from 13 to 44 years. Four studies used primary and secondary outcome indicators in the AL index (Table 3). All of the studies except one used the upper and lower quartile scoring approach to AL. Wallace and Harville (2013) used the sum of the z-scores for the biomarkers. Again these six studies modeled either the contributing factors of AL (as a dependent variable) or adverse outcomes of AL (AL as a predictor). The predictors of AL included: PTSD, age, race, ethnicity, chronic work, financial and caregiving stress, lifestyle factors (e.g., exercise, alcohol consumption, and smoking), social adversity (e.g., parental illness or loss, residential instability, exposure to threat/violence), and occupational social class. Adverse outcomes of AL included: sense of coherence (Lindfors *et al.* 2006) and adverse birth outcomes (e.g., preterm birth, low birth weight [Wallace & Harville 2013]).

CONCLUSIONS

We aimed to extend analysis of the concept of AL to advance its use in nursing research since it is a concept well-suited to the nursing meta-paradigm and to health promotion and risk reduction intervention science. Important papers have recently reviewed and clearly delineated the theory (Read & Grundy 2012) and systematically reviewed AL research (Beckie 2012), and these reviews highlighted areas for further methodological work. It is apparent that we need more consistent operationalization in

terms of indicators to include and scoring methods to apply. From examining these two sets of study reports on different populations to serve as research cases to illustrate the concept in use, we see mostly affirmation for the proposition that stress leads to higher AL and for the proposition that higher AL leads to adverse outcomes. Further research is needed to test the entire model, with AL as the mediator between stress and adverse outcomes—ideally with prospective designs. If such theory-testing research validates AL as a mediator, that will serve as a more solid test of concept for AL to be used as a proximal endpoint for clinical research.

RELEVANCE TO CLINICAL PRACTICE

Nurse researchers and clinicians working with stress-vulnerable populations need clear points of intervention and ability to determine impact of the interventions in the short term. Health promotion and risk reduction research is usually behavioral and often depends on self-reported, near-term outcomes such as self-efficacy, intention, or observation of behavior. This is because it's notoriously difficult to demonstrate prevention of disease in the long-term. Practical considerations also foster adopting simple designs (i.e., one primary outcome) that cannot represent complexity that we know is operating, and often the outcome has to be a proximal end-point (e.g., change in mean blood pressure) because follow-up to the manifestation of disease (e.g., prevention of hypertension diagnosis) is not feasible. Clinical practice has constraints that are similar. Intervening early and seeing progress during the sometimes extended period of care needed to achieve the intervention goal are sustaining and reinforcing for both the nurse and the client. Being able to explain the concept of AL to clients, being able to run a panel of tests that yield an interpretable "AL-Index" result, and being able to adjust interventions in response would be useful.

The theory of allostasis is useful at broader levels as well, including population health and policy. Further developing measurement of multiple levels of stress could take what we know about structural inequalities, as well as toxic and traumatic stress, into account as predictors in all studies. Tailoring with additional measures for specific populations will also be needed (e.g., measures of job strain and work-life balance for studies of workers, and measures of pregnancy-specific stress for perinatal studies).

We could advocate for policies that decrease stress and AL as a way to build a culture of health and decrease burden, morbidity, and costs of stress-related disease.

Finally, the concept of AL has been paired with the concept of allostatic support (Friedman & McEwen 2004), but this twin concept has not been synthesized formally. Given the state of the science on AL, there may be higher priority tasks. However, the goal of the theory is to depict not only the scenario where dysregulation from overload leads to disease but also the scenario where allostasis leads to adaptation for optimal health under the circumstances. In the future, nursing is well-positioned to place the concept of allostatic support within the theoretical framework and operationalize it—in research and in practice.

References

- Beckie TM (2012) A systematic review of allostatic load, health, and health disparities. *Biological Research for Nursing* **14**, 311-346. doi: 10.1177/1099800412455688
- Bellingrath S, Weigl T & Kudielka BM (2009) Chronic work stress and exhaustion is associated with higher allostatic load in female school teachers. *Stress* **12**, 37-48. <http://dx.doi.org/10.1080/10253890802042041>
- Breslau N (2002) Gender differences in trauma and posttraumatic stress disorder. *Journal of Gender Specific Medicine* **5**, 34-40.
- Briere J & Elliott DM (2003) Prevalence and psychological sequelae of self-reported childhood physical and sexual abuse in a general population sample of men and women. *Child Abuse & Neglect* **27**, 1205-1222. <http://dx.doi.org/10.1016/j.chiabu.2003.09.008>
- Bronfenbrenner U & Morris PA (2006) The bioecological model of human development. In *Handbook of child psychology* 6th edn. (Damon W & Lerner R eds.), John Wiley & Sons, Hoboken, NJ, pp. 793-828. DOI: 10.1002/9780470147658.chpsy0114
- Campbell JC (2002) Health consequences of intimate partner violence. *The Lancet* **359**, 1331-1336. [http://dx.doi.org/10.1016/S0140-6736\(02\)08336-8](http://dx.doi.org/10.1016/S0140-6736(02)08336-8)
- Cannon WB (1932) *The wisdom of the body*. W W Norton & Co, New York, NY.

- Carlson ED & Chamberlain RM (2005) Allostatic load and health disparities: a theoretical orientation. *Research in Nursing & Health* **28**, 306-315. doi: 10.1002/nur.20084
- Cohen S, Evans GW, Stokols D & Krantz DS (2013) *Behavior, health, and environmental stress*. Springer Science & Business Media, New York, NY.
- de Castro AB, Voss JG, Ruppin A, Dominguez CF & Seixas NS (2010) Stressors among Latino day laborers. A pilot study examining allostatic load. *American Association of Health Nurses Journal* **58**, 185-196. doi: 10.3928/08910162-20100428-01
- Dowd JB & Goldman N (2006) Do biomarkers of stress mediate the relation between socioeconomic status and health? *Journal of Epidemiology & Community Health* **60**, 633-639. doi:10.1136/jech.2005.040816
- Evans GW & Schamberg MA (2009) Childhood poverty, chronic stress, and adult working memory. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 6545-6549. doi: 10.1073/pnas.0811910106
- Fawcett J (1984) The metaparadigm of nursing: present status and future refinements. *Journal of Nursing Scholarship* **16**, 84-89. doi: 10.1111/j.1547-5069.1984.tb01393.x
- Felitti VJ, Anda RF, Nordenberg D, Williamson DF, Spitz AM, Edwards V, Koss MP & Marks JS (1998) Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences (ACE) Study. *American Journal of Preventative Medicine* **14**, 245-258. [http://dx.doi.org/10.1016/S0749-3797\(98\)00017-8](http://dx.doi.org/10.1016/S0749-3797(98)00017-8)
- Friedman MJ & McEwen BS (2004) Posttraumatic stress disorder, allostatic load, and medical illness. In *Trauma and Health: Physical Health Consequences of Exposure to Extreme Stress*. (Schnurr P & Green L eds.) American Psychological Association, Washington, D.C. pp 157-188. <http://dx.doi.org/10.1037/10723-007>

- Geronimus AT, Hicken M, Keene D & Bound J (2006) "Weathering" and age patterns of allostatic load scores among blacks and whites in the United States. *American Journal of Public Health* **96**, 826-833. doi: 10.2105/AJPH.2004.060749
- Gersten O (2008) Neuroendocrine biomarkers, social relations, and the cumulative costs of stress in Taiwan. *Social Science & Medicine* **66**, 507-519. <http://dx.doi.org/10.1016/j.socscimed.2007.09.004>
- Glei DA, Goldman N, Chuang YL & Weinstein M (2007) Do chronic stressors lead to physiological dysregulation? Testing the theory of allostatic load. *Psychosomatic Medicine* **69**, 769-776. doi: 10.1097/PSY.0b013e318157cba6
- Goldberger AL, Peng CK & Lipsitz LA (2002) What is physiologic complexity and how does it change with aging and disease? *Neurobiology of Aging* **23**, 23-26. [http://dx.doi.org/10.1016/S0197-4580\(01\)00266-4](http://dx.doi.org/10.1016/S0197-4580(01)00266-4)
- Goldman N, Turra CM, Gleib DA, Lin YH & Weinstein M (2006) Physiological dysregulation and changes in health in an older population. *Experimental Gerontology* **41**, 862-870. <http://dx.doi.org/10.1016/j.exger.2006.06.050>
- Groer MW, Kostas-Polston EA, Dillahunt-Aspillaga C, Beckie TM, Johnson-Mallard V, Duffy A & Evans ME (2016) Allostatic perspectives in women veterans with a history of childhood sexual assault. *Biological Research for Nursing* **18**, 454-464. doi: 10.1177/1099800416638442
- Gruenewald TL, Seeman TE, Ryff CD, Karlamangla AS & Singer BH (2006) Combinations of biomarkers predictive of later life mortality. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 14158-14163. doi: 10.1073/pnas.0606215103
- Gustafsson PE, Janlert U, Theorell T, Westerlund H & Hammarström A (2012) Social and material adversity from adolescence to adulthood and allostatic load in middle-aged women and men: Results from the Northern Swedish Cohort. *Annals of Behavioral Medicine* **43**, 117-128. doi: 10.1007/s12160-011-9309-6
- Hasson D, Schwarz UV & Lindfors P (2009) Self-rated health and allostatic load in women working in two occupational sectors. *Journal of Health Psychology* **14**, 568-577. doi: 10.1177/1359105309103576

- Hux VJ, Catov JM & Roberts JM (2014) Allostatic load in women with a history of low birth weight infants: The national health and nutrition examination survey. *Journal of Womens Health* **23**, 1039-1045. doi: 10.1089/jwh.2013.4572
- Juster RP, McEwen BS & Lupien SJ (2010) Allostatic load biomarkers of chronic stress and impact on health and cognition. *Neuroscience & Biobehavioral Reviews* **35**, 2-16. <http://dx.doi.org/10.1016/j.neubiorev.2009.10.002>
- Karasek RA & Theorell T (1992) *Healthy work: stress, productivity, and the reconstruction of working life*. Basic Books, New York, NY.
- Karlamangla AS, Singer BH, McEwen BS, Rowe JW & Seeman TE (2002) Allostatic load as a predictor of functional decline MacArthur studies of successful aging. *Journal of Clinical Epidemiology* **55**, 696-710. [http://dx.doi.org/10.1016/S0895-4356\(02\)00399-2](http://dx.doi.org/10.1016/S0895-4356(02)00399-2)
- Krug EG, Dalhberg LL, Mercy JA, Zwi AB & Lozano R (2002) Sexual violence. In *World Report on Violence and Health*. pp. 147-181. World Health Organization, Geneva, Switzerland.
- Li W, Zhang JQ, Sun J, Ke JH, Dong ZY & Wang S (2007) Job stress related to glycolipid allostatic load, adiponectin and visfatin. *Stress & Health* **23**, 257-266. doi: 10.1002/smi.1145
- Lindfors P, Lundberg O & Lundberg U (2006) Allostatic load and clinical risk as related to sense of coherence in middle-aged women. *Psychosomatic Medicine* **68**, 801-807. doi: 10.1097/01.psy.0000232267.56605.22
- Lipowicz A, Szklarska A & Malina RM (2014) Allostatic load and socioeconomic status in Polish adult men. *Journal of Biosocial Science* **46**, 155-167. <http://dx.doi.org/10.1017/S0021932013000345>
- Loucks EB, Juster RP & Pruessner JC (2008) Neuroendocrine biomarkers, allostatic load, and the challenge of measurement: A commentary on Gersten. *Social Science & Medicine* **66**, 525-530. <http://dx.doi.org/10.1016/j.socscimed.2007.09.006>

- McEwen BS (1998) Stress, adaptation, and disease: Allostasis and allostatic load. *Annals of the New York Academy of Sciences* **840**, 33-44. doi: 10.1111/j.1749-6632.1998.tb09546.x
- McEwen BS (2000) Allostasis and allostatic load: Implications for neuropsychopharmacology. *Neuropsychopharmacology* **22**, 108-124. doi:10.1016/S0893-133X(99)00129-3
- McEwen BS (2002) Sex, stress and the hippocampus: Allostasis, allostatic load and the aging process. *Neurobiology of Aging* **23**, 921-939. [http://dx.doi.org/10.1016/S0197-4580\(02\)00027-1](http://dx.doi.org/10.1016/S0197-4580(02)00027-1)
- McEwen BS (2004) Protection and damage from acute and chronic stress: Allostasis and allostatic overload and relevance to the pathophysiology of psychiatric disorders. *Annals of the New York Academy of Sciences* **1032**, 1-7. doi: 10.1196/annals.1314.001
- McEwen BS & Stellar E (1993) Stress and the individual. Mechanisms leading to disease. *Archives of Internal Medicine* **153**, 2093-2101. doi: 10.1001/archinte.1993.00410180039004
- McEwen BS & Seeman T (1999) Protective and damaging effects of mediators of stress: Elaborating and testing the concepts of allostasis and allostatic load. *Socioeconomic Status and Health in Industrial Nations* **896**, 30-47. doi: 10.1111/j.1749-6632.1999.tb08103.x
- McEwen BS & Wingfield JC (2010) What is in a name? Integrating homeostasis, allostasis and stress. *Hormones and Behavior* **57**, 105-111. doi: 10.1016/j.yhbeh.2009.09.011
- Morrison S, Shenassa ED, Mendola P, Wu TT & Schoendorf K (2013) Allostatic load may not be associated with chronic stress in pregnant women, NHANES 1999-2006. *Annals of Epidemiology* **23**, 294-297. <http://dx.doi.org/10.1016/j.annepidem.2013.03.006>
- National Center for Health Statistics (2012) *Health, United States, 2011: With special feature on socioeconomic status and health*. Hyattsville, MD. <http://dx.doi.org/10.1037/e582072012-001>

- Read S & Grundy E (2012) Allostatic load: A challenge to measure multisystem physiological dysregulation. National Centre for Research Methods.
- Schnorpfeil P, Noll A, Schulze R, Ehler U, Frey K & Fischer JE (2003) Allostatic load and work conditions. *Social Science & Medicine* **57**, 647-656. [http://dx.doi.org/10.1016/S0277-9536\(02\)00407-0](http://dx.doi.org/10.1016/S0277-9536(02)00407-0)
- Seeman M, Stein Merkin S, Karlamangla A, Koretz B & Seeman T (2014) Social status and biological dysregulation: The "status syndrome" and allostatic load. *Social Science & Medicine* **118**, 143-151. <http://dx.doi.org/10.1016/j.socscimed.2014.08.002>
- Seeman TE, Singer BH, Rowe JW, Horwitz RI & McEwen BS (1997) Price of adaptation—Allostatic load and its health consequences. MacArthur studies of successful aging. *Archives of Internal Medicine* **157**, 2259-2268. doi: 10.1001/archinte.1997.00440400111013
- Seeman TE, Crimmins E, Huang MH, Singer B, Bucur A, Gruenewald T, Berkman LF & Reuben DB (2004) Cumulative biological risk and socio-economic differences in mortality: MacArthur studies of successful aging. *Social Science & Medicine* **58**, 1985-1997. [http://dx.doi.org/10.1016/S0277-9536\(03\)00402-7](http://dx.doi.org/10.1016/S0277-9536(03)00402-7)
- Seplaki CL, Goldman N, Weinstein M & Lin YH (2004) How are biomarkers related to physical and mental well-being? *Journals of Gerontology Series A Biological Sciences & Medical Sciences* **59**, 201-217. doi: 10.1093/gerona/59.3.B201
- Seplaki CL, Goldman N, Gleib D & Weinstein M (2005) A comparative analysis of measurement approaches for physiological dysregulation in an older population. *Experimental Gerontology* **40**, 438-449. <http://dx.doi.org/10.1016/j.exger.2005.03.002>
- Sterling P & Eyer J (1988) *Allostasis: A new paradigm to explain arousal pathology*. Wiley, New York, NY.
- Sun J, Wang S, Zhang JQ & Li W (2007) Assessing the cumulative effects of stress: The association between job stress and allostatic load in a large sample of Chinese employees. *Work & Stress: An International Journal of Work, Health & Organisations* **21**, 333-347. <http://dx.doi.org/10.1080/02678370701742748>

- Szanton SL, Gill JM & Allen JK (2005) Allostatic load: A mechanism of socioeconomic health disparities? *Biological Research for Nursing* **7**, 7-15. doi: 10.1177/1099800405278216
- Szanton SL, Allen JK, Seplaki CL, Bandeen-Roche K & Fried LP (2009) Allostatic load and frailty in the women's health and aging studies. *Biological Research for Nursing* **10**, 248-256. doi: 10.1177/1099800408323452
- von Kanel R, Dimsdale JE, Patterson TL & Grant I (2003) Acute procoagulant stress response as a dynamic measure of allostatic load in Alzheimer caregivers. *Annals of Behavioral Medicine* **26**, 42-48. doi: 10.1207/S15324796ABM2601_06
- von Thiele U, Lindfors P & Lundberg U (2006) Self-rated recovery from work stress and allostatic load in women. *Journal of Psychosomatic Research* **61**, 237-242. <http://dx.doi.org/10.1016/j.jpsychores.2006.01.015>
- Walker LO & Avant KC (2005) *Strategies for theory construction in nursing, 4th edn.* Pearson/Prentice Hall, Upper Saddle River, New Jersey.
- Wallace M, Harville E, Theall K, Webber L, Chen W & Berenson G (2013) Preconception biomarkers of allostatic load and racial disparities in adverse birth outcomes: The Bogalusa Heart Study. *Paediatric and Perinatal Epidemiology* **27**, 587-597. doi: 10.1111/ppe.12091
- Wallace ME & Harville EW (2013) Allostatic load and birth outcomes among White and Black women in New Orleans. *Maternal and Child Health Journal* **17**, 1025-1029. doi: 10.1007/s10995-012-1083-y
- Weinstein M, Goldman N, Hedley A, Yu-Hsuan L & Seeman T (2003) Social linkages to biological markers of health among the elderly. *Journal of Biosocial Science* **35**, 433-453. <http://dx.doi.org/10.1017/S0021932003004334>

Table 1. Allostatic load and workers

Reference (author/year)	Sample (workers vs. women)	Study Purpose	AL- Definition	AL Component	Scoring Method	Role of AL (mediator or?)	Study Findings	Notes
de Castro <i>et al.</i> 2010	N = 30 male Latino day laborers in Seattle Washington, US	This was a pilot study - cross sectional design. To evaluate the feasibility of conducting a research project focused on stressors (work related; economic; social) and AL among day laborers.	The physiologic effects of chronic stress	SBP, DBP, salivary cortisol, WHR; CRP; and BMI.	The sum of the number of biological measures on which each participant scored in the top quartile of risk for the total sample.	Outcome (dependent variable)	Average age was 46 years old and the average length of residency in the US was 12 years. 47% were married and the majority (97%) spoke Spanish and were from Mexico (77%). Workers who reported more stressors (work, economic, social) had higher AL physically and mentally.	Suggested need for studies to explore AL as a mediator between stressors and clinical health outcomes. Used the finger prick and sterile lancet to collect the blood for the CRP. Trust and collaboration between researchers and community partners are very important. They calculated AL score by summing the number of parameters in which levels were in

Reference (author/year)	Sample (workers vs. women)	Study Purpose	AL- Definition	AL Component	Scoring Method	Role of AL (mediator or?)	Study Findings	Notes
							They found no statistical significance between the groups compared to those with low AL who have been working as day laborers for just under 5 years.	the highest quartile (at or above the 75% quartile) and the parameters in the lowest quartile. AL scores can range from 0 to 6. The AL scores were then further dichotomized into two categories low and high.
von Thiele <i>et al.</i> 2006	N= 241 healthy women working in two public health care organizations in Stockholm, Sweden	This was a secondary data analysis. To investigate the relationships between self-rated recovery from work stress and biologic	A multisystems approach describes how daily stress relates to health and disease.	SBP, DBP, HR, HDL, LDL, LDL/HDL ratio, TC, TG, serum DHEA-S, glucose, HbA1c, prolactin, and WHR.	The sum of the number of biological measures on which each participant scored in the top quartile of risk for the total sample (except for	Outcome	No significant differences in the demographic characteristics of study participants. The fatigued women had an increased risk for high AL. In	Insufficient recovery from work stress may result in high AL. Acknowledged that the operationalization of AL differs between studies depending on the type and number of

Reference (author/year)	Sample (workers vs. women)	Study Purpose	AL- Definition	AL Component	Scoring Method	Role of AL (mediator or?)	Study Findings	Notes
		dysregulation in terms of AL and individual biomarkers, respectively, in healthy women within the public health care sector.			HDL, and DHEA-S, where inclusion into the lowest quartile constitutes risk).		contrast, there was no significant difference in individual biomarkers between recovered and non-recovered women.	biomarkers available
Sun <i>et al.</i> 2007	N= 1,219 healthy Chinese employees including both men (50%) and women (50%) from five industries in China.	This was a cross sectional study design. To determine the relationship between job strain and AL	1. The accumulation of wear and tear on physiological systems from the process of adaptation to chronic stress. 2. A possible biological warning	FIB, CRP, cortisol, adnephrin, BMI, WHR, SBP, DBP, HbA1c, IGR, TC/HDL, HDL, and TG.	The sum of the number of biological measures on which each participant scored in the top quartile of risk for the total sample (except for HDL in the lowest	Outcome	The average age was 38. About 70% had a college level education. High job strain had higher AL compared to low job strain. Positive association between AL and age ($r=.203$,	Excluded people already diagnosed with conditions such as hypertension and diabetes or taking medication for these conditions; also measured type A behavior which showed to be associated with higher job strain.

Reference (author/year)	Sample (workers vs. women)	Study Purpose	AL- Definition	AL Component	Scoring Method	Role of AL (mediator or?)	Study Findings	Notes
			system for health outcomes. 3. The cumulative physiological toll that may be exacted on the body through its attempts to adapt to life's demands.		quartile).		p<0.001), and positive relationship between AL and education level (lower education yielded to higher job strain) (r=-.122, p<0.001).	They also assessed for behavioral responses associated with stress such as smoking, alcohol intake, diet, and exercise
Schnorpfeil <i>et al.</i> 2003	N=324 employees from an airplane manufacturing plant in Germany	This was a cross sectional study design. To evaluate the relationship between objective health status and work characteristics in the workers.	A biological warning system	BMI, WHR, SBP, DBP, CRP, TNF- α , HDL, cholesterol, DHEA-S, HbA1c, urinary cortisol, epinephrine, norepinephrine, and albumin.	The sum of the number of biological measures on which each participant scored in the top quartile of risk for the total sample	Outcome	Older individuals and men had higher AL scores than younger participants and women. Job demand related significantly to AL.	

Reference (author/year)	Sample (workers vs. women)	Study Purpose	AL- Definition	AL Component	Scoring Method	Role of AL (mediator or?)	Study Findings	Notes
					(except for HDL, DHEA-S, and cholesterol in the lowest quartile).			
Hasson <i>et al.</i> 2009	Female employees from the health care sector (n=241) and IT/media sector (n=98), Stockholm, Sweden	To investigate how biological dysregulation, in terms of AL, relates to self-rated health (SRH) in women.	A multisystems approach describes how daily stress relates to health and disease.	SBP, DBP, HR, HDL, LDL, LDL/HDL ratio, TC, TG, serum DHEA-S, glucose, HbA1c, prolactin, and WHR.	The sum of the number of biological measures on which each participant scored in the top quartile of risk for the total sample (except for HDL and DHEA-S in the lowest quartile).	Outcome	A poor self-rated health, along with occupational sector, age and education, were significantly associated with a high AL, particularly for those working within the health care sector.	
Li <i>et al.</i> 2007	N=504 industrial	To evaluate the relationship	The price of allostasis	BMI, WHR, HbA1c, B-cell	The sum of the number	Outcome	50% were male. Most were	

Reference (author/year)	Sample (workers vs. women)	Study Purpose	AL- Definition	AL Component	Scoring Method	Role of AL (mediator or?)	Study Findings	Notes
	employees, China	between job stress and glycolipid metabolic change in healthy industrial employees in China		function, TC, TG, HDL, LDL, adiponectin, and visfatin.	of biological measures on which each participant scored in the top quartile of risk for the total sample (except for HDL, adiponectin, and B-cell function in the lowest quartile).		married (78%). About 70% had a college level education. High job stress was extremely correlated with increased glycolipid AL score. Low job control showed to be the main stressor impacting health outcomes in the participants.	

Note. AL, allostatic load; BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; DHEA-S, dehydroepiandrosterone sulfate; HDL, high-density lipoprotein; LDL, low-density lipoprotein; TC, total cholesterol; TG, triglyceride; FIB, fibrinogen; IGR, insulin glucose ratio; HbA1c, glycosylated hemoglobin; HR, heart rate; WHR, waist/hip ratio; CRP, C-reactive protein; TNF- α , tumor necrosis factor alpha.

Table 2. Allostatic load and women

Reference (author/year)	Sample (workers vs. women)	Study Purpose	AL-Definition	AL Component	Scoring Method	Role of AL (mediator or?)	Study Findings	Notes
Lindfors <i>et al.</i> 2006	N=200 women with no previously diagnosed pathology in Sweden	To investigate how physiologic dysregulation, in terms of AL and clinical risk, respectively, relates to sense of coherence (SOC) in women with no previously diagnosed pathology.	The dysregulation in multiple bodily systems in response to stress challenges over the life course.	Resting SBP and DBP, HDL, TC, HbA1c, WHR, and PEF.	The sum of the number of biological measures on which each participant scored in the top quartile of risk for the total sample (except for HDL and PEF in the lowest quartile).	Predictor	All women were 43 years old. 77.5% were married or living with a partner. 94.5% had children. 51.5% had a lower education. AL was found to predict future SOC, whereas clinical risk did not.	
Gustafsson <i>et al.</i> 2012	N= 394 women in Sweden	To examine whether social and material adversity over the life course is related to AL in	The cumulative dysregulations which eventually develop across multiple interconnected	SBP, DBP, BMI, waist circumference, fasting glucose, TC, HDL, TG, apolipoprotein	The sum of the number of biological measures which was divided into	Outcome	All women were 43 years of age. Social adversity accumulated over the life course was related to AL,	

Reference (author/year)	Sample (workers vs. women)	Study Purpose	AL-Definition	AL Component	Scoring Method	Role of AL (mediator or?)	Study Findings	Notes
		mid-adulthood.	physiological systems as a result of frequently repeated or chronic activation over the life course.	A1 and B, CRP, and diurnal salivary cortisol AUC	tertiles (coded 0, 1, 2), except for cortisol (coded symmetrically with sextile: 1 and 6=2, 2 and 5=1, 3 and 4=0) and HDL (coded inversely: 2, 1, 0).		independently of cumulative socioeconomic disadvantage. Moreover, social adversity in adolescence was related to AL, independently of cumulative socioeconomic disadvantage and also of later adversity exposure during adulthood.	
Morrison <i>et al.</i> 2013	N= 6,131 pregnant women (n=1,138) and nonpregnant (n=4,993)	To determine whether AL can be measured in a meaningful way during pregnancy.	The cumulative wear and tear on the body resulting from exposure to chronic stress.	SBP, DBP, 60-second pulse rate, homocysteine, CRP, serum albumin, HbA1c, HDL,	The sum of the number of biological measures on which each participant scored in the	Outcome	Women were aged 15-44. Among pregnant women, 56.4% were non-Hispanic white, 14.2% were non-Hispanic black,	

Reference (author/year)	Sample (workers vs. women)	Study Purpose	AL-Definition	AL Component	Scoring Method	Role of AL (mediator or?)	Study Findings	Notes
	women in the U.S.			TC, and creatinine.	top quartile of risk for the total sample (except for HDL and serum albumin in the lowest quartile).		16.1% were Mexican American, and 13.2% were other races; Among nonpregnant women, 66.1% were non-Hispanic white, 12.6% were non-Hispanic black, 9.4% were Mexican American, and 11.9% were other races. The distribution of each AL-related biomarker differed significantly between pregnant and nonpregnant women. Among nonpregnant women, higher AL	

Reference (author/year)	Sample (workers vs. women)	Study Purpose	AL-Definition	AL Component	Scoring Method	Role of AL (mediator or?)	Study Findings	Notes
							was found in women who are black, are older, and who have lower incomes). However, these associations were not seen in pregnant women.	
Wallace <i>et al.</i> 2013	N= 886 women from the Bogalusa Heart Study in the U.S	To examine more closely the relationships between AL, race, and adverse birth outcomes within the context of neighborhood- level poverty.	The dysregulation across the body's multiple physiological systems responsible for maintaining equilibrium when faced with physical or social challenges.	SBP, DBP, TC, HDL, LDL, TG, glucose, insulin, and waist circumference.	The sum of the number of biological measures on which each participant scored in the top quartile of risk for the total sample (except for HDL in the lowest quartile).	Predictor	Women were aged 13-41. 59.4% of women are white, and 40.6% of women are African American. African American women resided in more impoverished neighborhoods and had higher AL scores compared to whites; however,	

Reference (author/year)	Sample (workers vs. women)	Study Purpose	AL-Definition	AL Component	Scoring Method	Role of AL (mediator or?)	Study Findings	Notes
							AL was not associated with preterm birth or low birth weight.	
Wallace & Harville 2013	N= 123 women at 26-28 weeks gestation in the U.S	To identify associations between AL and birth outcomes and to assess differences in AL and its relation to birth outcomes between white and black women.	The wear and tear on the body that arises from chronic, prolonged, or persistent activation of allostatic effectors and a breakdown of the regulatory feedback mechanisms.	Cholesterol, cortisol, DHEA-S, HbA1c, and SBP.	The sum of z-scores for all five biological measures.	Predictor	Women were aged 20-35. All are white or African American. Black women had a lower AL index than white women. Gestational age was associated with AL. A significant interaction with age indicated that the effect wasn't as strong at higher maternal ages. There was no racial difference in the	

Reference (author/year)	Sample (workers vs. women)	Study Purpose	AL-Definition	AL Component	Scoring Method	Role of AL (mediator or?)	Study Findings	Notes
							effect of AL on birth outcomes.	
Hux <i>et al.</i> 2014	N= 877 women in the U.S	To determine whether past history of having had a pregnancy with low birth weight outcome is associated with higher AL.	The cumulative burden of chronic physiologic and psychological stress.	SBP, DBP, BMI, CRP, serum albumin, HbA1c, HDL, TC, and creatinine.	The sum of the number of biological measures on which each participant scored in the top quartile of risk for the total sample (except for HDL and serum albumin in the lowest quartile).	Predictor	Women were aged 17-35. 62.4% are white, 20.1% are Latino/Hispanic, 14.9% are Black, and 2.6% are other races. Women with history of small for gestational age or preterm birth had higher AL than did those with normal birth weight outcomes.	Since the study was poorly designed, it is difficult to determine whether differences in AL occur prior to or after pregnancy as a result of the adverse outcomes.

Note. AL, allostatic load; BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; DHEA-S, dehydroepiandrosterone sulfate; HDL, high-density lipoprotein; LDL, low-density lipoprotein; TC, total cholesterol; TG, triglyceride; FIB, fibrinogen; HbA1c, glycosylated hemoglobin; WHR, waist/hip ratio; CRP, C-reactive protein; AUC, area under curve; PEF, peak expiratory flow.

Table 3: Measures of AL across each studies

Reference (Author/year)	Measures of AL
de Castro <i>et al.</i> 2010	SBP, DBP, salivary cortisol, WHR; CRP; and BMI.
von Thiele <i>et al.</i> 2006	SBP, DBP, HR, HDL, LDL, LDL/HDL ratio, TC, TG, serum DHEA-S, glucose, HbA1c, prolactin, and WHR.
Sun <i>et al.</i> 2007	FIB, CRP, cortisol, adnephrin, BMI, WHR, SBP, DBP, HbA1c, IGR, TC/HDL, HDL, and TG.
Schnorpfeil <i>et al.</i> 2003	BMI, WHR, SBP, DBP, CRP, TNF- α , HDL, cholesterol, DHEA-S, HbA1c, urinary cortisol, epinephrine, norephinephrine, and albumin.
Hasson <i>et al.</i> 2009	SBP, DBP, HR, HDL, LDL, LDL/HDL ratio, TC, TG, serum DHEA-S, glucose, HbA1c, prolactin, and WHR.
Li <i>et al.</i> 2007	BMI, WHR, HbA1c, B-cell function, TC, TG, HDL, LDL, adiponectin, and visfatin.
Lindfors <i>et al.</i> 2006	Resting SBP and DBP, HDL, TC, HbA1c, WHR, and PEF
Gustafsson <i>et al.</i> 2012	SBP, DBP, BMI, waist circumference, fasting glucose, TC, HDL, TG, apolipoprotein A1 and B, CRP, and diurnal salivary cortisol AUC
Morrison <i>et al.</i> 2013	SBP, DBP, 60-second pulse rate, homocysteine, CRP, serum albumin, HbA1c, HDL, TC, and creatinine.
Wallace <i>et al.</i> 2013	SBP, DBP, TC, HDL, LDL, TG, glucose, insulin, and waist circumference.
Wallace & Harville 2013	Cholesterol, cortisol, DHEA-S, HbA1c, and SBP
Hux <i>et al.</i> 2014	SBP, DBP, BMI, CRP, serum albumin, HbA1c, HDL, TC, and creatinine.

Note. AL, allostatic load; BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; DHEA-S, dehydroepiandrosterone sulfate; HDL, high-density lipoprotein; LDL, low-density lipoprotein; TC, total cholesterol; TG, triglyceride; FIB, fibrinogen; IGR, insulin glucose ratio; HbA1c, glycosylated hemoglobin; HR, heart rate; WHR, waist/hip ratio; CRP, C-reactive protein; TNF- α , tumor necrosis factor alpha; AUC, area under curve.

Figure 1. Model Case Illustrating Read and Grundy's (2012) Allostatic-Adaptation Process

