

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29

Received Date:

Revised Date:

Accepted Date:

Article Type: Letters

**Elevated Atmospheric Concentrations of Carbon Dioxide Reduce Monarch
Tolerance and Increase Parasite Virulence by Altering the Medicinal Properties of
Milkweeds**

Leslie E. Decker:

Department of Ecology and Evolutionary Biology, University of Michigan, Biological
Sciences Building, 1105 North University Avenue, Ann Arbor, MI 48109-1085,
USA. lesldeck@umich.edu

Jacobus C. de Roode:

Biology Department, Emory University, Rollins 1113
O. Wayne Rollins Research Center, 1510 Clifton Road, Atlanta, GA 30322, USA.
jacobus.deroode@emory.edu

Mark D. Hunter:

Department of Ecology and Evolutionary Biology, University of Michigan, Biological
Sciences Building, 1105 North University Avenue, Ann Arbor, MI 48109-1085,
USA. mdhunter@umich.edu

Short Running Title: CO₂ alters tolerance and virulence

Key-words: anthropogenic, *Asclepias*, cardenolides, *Danaus plexippus*, global
environmental change, monarch butterfly, *Ophryocystis elektroscirrha*, host-parasite
interactions.

This is the author manuscript accepted for publication and has undergone full peer review
but has not been through the copyediting, typesetting, pagination and proofreading process,
which may lead to differences between this version and the [Version of Record](#). Please cite
this article as [doi: 10.1111/ele.13101](https://doi.org/10.1111/ele.13101)

This article is protected by copyright. All rights reserved

30

31 **Type of Article:** Letter

32 **Abstract Word Count:** 149

33 **Main Text Word Count:** 4,992

34 **Total References:** 86

35 **Total Figures:** 6

36 **Total Tables:** 0

37

38 **Corresponding Author:** Leslie E. Decker. Department of Ecology and Evolutionary
39 Biology, University of Michigan, Biological Sciences Building, 1105 North University
40 Avenue, Ann Arbor, MI 48109-1085, USA. lesldeck@umich.edu. Phone: 443-928-2357.

Author Manuscript

41 **Statement of Authorship:**

42 LED, MDH & JCdR designed the experiment. JCdR provided butterfly and parasite
43 materials along with disease protocols. LED & MDH collected and analyzed the data
44 with suggestions from JCdR. LED wrote the manuscript and all authors contributed
45 substantially to revisions.

46

47 **Data Accessibility:** Data will be made available in the Dryad Digital Repository and the
48 DOI will be provided, should the manuscript be accepted.

49

50 **Abstract**

51 Hosts combat their parasites using mechanisms of resistance and tolerance, which
52 together determine parasite virulence. Environmental factors, including diet, mediate the
53 impact of parasites on hosts, with diet providing nutritional and medicinal properties.
54 Here, we present the first evidence that ongoing environmental change decreases host
55 tolerance and increases parasite virulence through a loss of dietary medicinal quality.
56 Monarch butterflies use dietary toxins (cardenolides) to reduce the deleterious impacts of
57 a protozoan parasite. We fed monarch larvae foliage from four milkweed species grown
58 under either elevated or ambient CO₂, and measured changes in resistance, tolerance, and
59 virulence. The most high-cardenolide milkweed species lost its medicinal properties
60 under elevated CO₂; monarch tolerance to infection decreased, and parasite virulence
61 increased. Declines in medicinal quality were associated with declines in foliar
62 concentrations of lipophilic cardenolides. Our results emphasize that global
63 environmental change may influence parasite-host interactions through changes in the
64 medicinal properties of plants.

65

66 **Introduction**

67 When facing infection, hosts utilize two avenues of defense: resistance and tolerance
68 (Råberg *et al.* 2007; Best *et al.* 2008). Resistance reduces the probability and degree of
69 parasitic infection, and subsequent parasite replication (e.g. parasite fitness) (Best *et al.*
70 2008; Boots *et al.* 2009). In contrast, tolerance describes the ability of hosts to mitigate
71 the negative fitness impacts of infection for a given pathogen load (Råberg *et al.* 2009;
72 Kutzer & Armitage 2016a). While host resistance reduces parasite fitness by preventing
73 infection or lowering parasite replication, host tolerance does not reduce parasite fitness.
74 Therefore, the two defense strategies should engender different co-evolutionary outcomes
75 for host-parasite dynamics (Roy & Kirchner 2000; Restif & Koella 2004).

76
77 Resistance and tolerance evolve because of the inherent damage (virulence) that parasites
78 cause to their hosts. Virulence emerges from the interaction between parasite and host
79 genotype (Lambrechts *et al.* 2006; Råberg *et al.* 2007) and varies with host ecology and
80 condition (Thomas & Blanford 2003; Boots 2011; Howick & Lazzaro 2014). Together,
81 host resistance and tolerance influence the rate at which parasites replicate and damage
82 the host, governing the severity of virulence (de Roode *et al.* 2008a, b; Tao *et al.* 2015).

83
84 Understanding variation in resistance has long been a focus of disease ecology. Host
85 genotype, physiology and environment all contribute to parasite resistance (Lambrechts
86 *et al.* 2006; Wolinska & King 2009). In contrast, our understanding of host tolerance
87 derives largely from the study of pests that attack plants (*reviewed in* Baucom & De
88 Roode 2011). Recent work investigating host tolerance in animals has focused largely on
89 host genotype under laboratory conditions (Råberg *et al.* 2007; Rohr *et al.* 2010; Jackson
90 *et al.* 2014; Regoes *et al.* 2014). However, we miss important factors that contribute to
91 variation in tolerance (Lefèvre *et al.* 2011) by isolating host-parasite interactions from
92 surrounding the complex community of organisms or environmental conditions
93 (Sternberg *et al.* 2012; Hayward *et al.* 2014; Tao *et al.* 2015; Clough *et al.* 2016; Kutzer
94 & Armitage 2016b; Debes *et al.* 2017; Zeller & Koella 2017).

95

96 Global environmental change directly affects the ecology and evolution of host-parasite
97 interactions (Tylianakis *et al.* 2008; Altizer *et al.* 2013; Becker *et al.* 2015). Generally,
98 environmental change increases the distribution and prevalence of parasites in host
99 populations (Garamszegi 2011; Zamora-Vilchis *et al.* 2012). Host resistance can increase
100 or decrease in response to the direct effects of environmental change on parasite life cycle
101 and host physiology (Bruno *et al.* 2003; Adamo & Lovett 2011; Paull & Johnson 2014).
102 However, surprisingly little work has investigated how future environmental conditions
103 may impact host tolerance of disease (Franke *et al.* 2017).

104
105 Moreover, indirect effects of environmental change on host-parasite interactions remain
106 largely unexplored. Future environmental conditions will alter the composition and traits
107 of the surrounding community (Tylianakis *et al.* 2008; Gunderson *et al.* 2017), which
108 may lead to shifts in host resistance and tolerance, and parasite virulence. Phytophagous
109 insect-parasite systems are excellent models with which to study the indirect effects of
110 global change on host-parasite interactions in the context of their communities. Host-
111 plant quality mediates the impacts of parasites on phytophagous insects (Cory & Hoover
112 2006; Shikano 2017), with significant medicinal effects of plant secondary metabolites
113 (Felton & Duffey 1990; Hunter & Schultz 1993; Bernays & Singer 2005). Importantly,
114 plant nutritional and defensive chemistry vary in response to environmental change
115 (Bidart-Bouzat & Imeh-Nathaniel 2008). For instance, elevated concentrations of
116 atmospheric CO₂ increase foliar carbohydrates, reduce nitrogen concentrations, and
117 change secondary metabolite production (Hunter 2001; Robinson *et al.* 2012; Zavala *et al.*
118 2013). Such changes in host plant chemistry can alter herbivore performance against
119 parasites (Cory & Hoover 2006; Shikano *et al.* 2010; Lampert 2012). In essence,
120 environmental change alters plant quality, which can affect the interactions between
121 herbivores and their parasites.

122
123 Here, we assess the impact of a pervasive driver of environmental change, elevated
124 atmospheric CO₂ concentration (eCO₂), on the interaction between monarch butterflies,
125 *Danaus plexippus*, and their protozoan parasite, *Ophryocystis elektroscirrha* (Mclaughlin
126 *et al.* 1970). *Ophryocystis elektroscirrha* infection reduces adult monarch lifespan,

127 fecundity, and flight ability (Bradley & Altizer 2005; de Roode *et al.* 2008b, 2009).
128 Monarchs become infected by consuming dormant parasite spores on the surface of egg
129 chorea and leaf tissue. During monarch development, parasites replicate within the insect
130 and butterflies emerge covered in dormant parasite spores. As specialists, monarchs lay
131 eggs on milkweed, *Asclepias*, host-plants (Malcolm & Brower 1989), thereby
132 contaminating foliage and eggs with spores.

133

134 Certain milkweed species are strongly medicinal, reducing the probability of infection,
135 growth rate, and virulence of *O. elektroscirra* (de Roode *et al.* 2008a). The medicinal
136 qualities of milkweed are related to cardenolides, toxic steroids produced in a majority of
137 milkweed species (Gowler *et al.* 2015). Larvae that feed on plants with high cardenolide
138 concentrations, or a high diversity of lipophilic cardenolides, suffer lower rates of
139 infection, maintain higher fitness at a given parasite load, and produce fewer new
140 parasites (de Roode *et al.* 2011b; Sternberg *et al.* 2012; Gowler *et al.* 2015). Additionally,
141 high foliar nutrient concentrations can increase monarch tolerance to their parasites (Tao
142 *et al.* 2015). Thus, foliar cardenolides and nutrients combine to mediate the resistance and
143 tolerance of monarchs to their parasites. Recent work shows that eCO₂ causes decreases
144 in cardenolide concentrations, changes in the composition of cardenolides, and declines
145 in nutrient concentrations of milkweed (Vannette & Hunter 2011; Matiella 2012).
146 Therefore, increasing atmospheric CO₂ concentrations may influence milkweed-mediated
147 interactions between monarchs and their parasites

148

149 Together these data motivate the overarching question of this study: Will monarch
150 resistance and tolerance and mean *O. elektroscirra* virulence change with milkweed
151 phytochemistry under future atmospheric CO₂ concentrations? We performed a field
152 mesocosm experiment to explore how eCO₂ alters the foliar chemistry of four milkweed
153 species. We then measured the CO₂-mediated effects of altered food-plant chemistry on
154 three aspects of monarch and parasite performance: 1) the spore load of infected
155 monarchs; 2) the tolerance of monarchs, expressed as the rate of decline in lifespan with
156 increasing spore load; and 3) the virulence of *O. elektroscirra*, calculated as the decline
157 in adult monarch lifespan due to infection. We hypothesized that the presence of

158 lipophilic cardenolides, in conjunction with foliar nutrient quality, dictates the effects of
159 eCO₂ on monarch-parasite interactions.

160

161

162 **Materials and methods**

163 We performed the experiment in two temporal blocks during 2014 and 2015. General
164 experimental procedures were the same for both blocks, with minor differences noted
165 below. The experiment was fully factorial, with milkweed species, parasite treatment
166 (infected or uninfected), and CO₂ treatment (ambient or elevated) as main effects (Table
167 S1).

168

169 *Plant Materials*

170 We grew four species of milkweed under current (400 ppm, aCO₂) and future (760 ppm,
171 eCO₂) concentrations of atmospheric CO₂ at the University of Michigan Biological
172 Station (45.5587° N, 84.6776° W). Within this century, the concentration of atmospheric
173 CO₂ will likely exceed 700 ppm (Solomon *et al.* 2009). Thus, following previous studies
174 in this system (Vannette & Hunter 2011, 2014), we chose 760 ppm as our target future
175 CO₂ concentration. Plants grew in a mesocosm array of 40 chambers, 20 maintained at
176 aCO₂, and 20 at eCO₂ from dawn until dusk (Drake *et al.* 1989, Figure S1). We chose
177 milkweed species that differed in their cardenolide concentrations, on a gradient of anti-
178 parasitic effects from high to low: *A. curassavica* (high), *A. speciosa*, *A. syriaca* (both
179 medium) and *A. incarnata* (low) (Sternberg *et al.* 2012). All four milkweed species occur
180 in sympatry in North America (Woodson 1954; Malcolm & Zalucki 1996). Seeds were
181 obtained from commercial sources (Butterfly Encounters, CA in 2014 and Prairie Moon
182 Nurseries, MN in 2015). After six weeks of cold stratification (for all but tropical *A.*
183 *curassavica*), seeds were germinated and planted on 3-May-2014 and 5-May-2015 in
184 deepotsTM containing Metromix 360 and Osmocote 16:16:16 (N:P:K) controlled release
185 fertilizer. Seedlings were watered daily and kept in a greenhouse for two weeks following
186 germination to avoid frost damage. We transferred seedlings outside to their assigned
187 chambers on 25-May-2014 and 23-May-2015.

188

189 Monarch caterpillars can consume three entire plants as larvae. Due to space limitations
190 in 2014, only two plants of the assigned milkweed species and CO₂ treatment were
191 grown for each larva. This made for 16 experimental plants in total in each chamber (4
192 milkweed species x 2 parasite treatments x 2 plants/monarch). Once larvae had consumed
193 both assigned plants, they were fed cuttings from *A. tuberosa*, a milkweed with negligible
194 cardenolides. Inoculation with the parasite took place 3 days after hatching (see below).
195 Milkweed chemistry influences parasite infection success and severity just before and
196 during consumption of parasite spores on plant material (de Roode *et al.* 2011a).
197 Therefore, switching to an almost cardenolide-free host-plant just before pupation should
198 not affect monarch-parasite interactions. In 2015, each chamber held enough plants to
199 feed all larvae for their entire larval periods (4 milkweed species x 2 infection treatments
200 x 3 plants/ monarch = 24 plants/ chamber) with 20 aCO₂ and 20 eCO₂ chambers as
201 before.

202

203 CO₂ concentrations were monitored during daylight hours in all eCO₂ chambers and one
204 ambient chamber using a LI-COR 320 IRGA (LI-COR, Lincoln, NE, USA). The
205 concentrations of CO₂ were adjusted throughout the day to maintain target values in each
206 elevated chamber. Temperatures within the chambers were monitored using iButton
207 dataloggers (iButtonLink, Whitewater, WI, USA). Elevated CO₂ chambers averaged
208 21.03 (±0.03)°C, and aCO₂ chambers averaged 21.24 (± 0.04)°C which were roughly 2°C
209 higher than the outside average temperature of 18.93 (± 0.04)°C and fell well within those
210 temperatures experienced by monarchs in eastern North America. Plants were maintained
211 in their chambers for 61 days in 2014 and 42 days in 2015 before experimental trials
212 began.

213

214 *Monarch Sources and Rearing Methods*

215 Monarchs were F₁ offspring of eight (2014) and seven (2015) crosses between eastern
216 North American lineages. Monarch lineages are lab-bred full sib families, obtained by
217 outcrossing wild-collected monarchs. We crossed 7 lineages in 2014 and 6 lineages in
218 2015 to produce our different families. Because we collected new wild monarchs each
219 year (monarchs suffer from inbreeding depression and cannot be maintained as lab stocks

220 between years), we did not make the same crosses in both years. Darkened monarch eggs
221 (those about to hatch) were placed individually on milkweed cuttings taken from plants
222 grown in the array. Only one larva from each treatment (4 milkweed species x 2 levels of
223 parasite infection = 8 treatments) was reared on plants from each chamber (Table S1).
224 We kept larvae individually in 0.64L plastic containers under aCO₂ on foliage transferred
225 from the appropriate atmosphere to avoid any confounding direct effects of eCO₂ on
226 insect performance (such effects are negligible (Bale *et al.* 2002)).

227
228 Each year, we infected monarchs with a single parasite family cultured from spores
229 collected from an eastern North American butterfly. Hatchlings fed for three days on their
230 assigned leaf tissue before inoculation with *O. elektroscirra*. To infect larvae, 10
231 parasite spores were transferred to a 70.6 mm² leaf disk taken from each larva's assigned
232 host plant. The leaf disk was placed in a petri dish containing moist filter paper and the
233 assigned larva. Control larvae received spore-free leaf disks. Immediately after disks
234 were punched from plants, foliage was collected for chemical analysis (below). Petri
235 dishes containing disks and larvae were kept in an incubator held at 26°C with 16-hour
236 daylight. Once each larva had consumed its entire leaf disk (and therefore all spores) it
237 was returned to its assigned container and fed foliage *ad libitum* from its designated
238 plants until pupation.

239 240 *Measures of Monarch Performance*

241 Upon emergence, butterflies were transferred to individual glassine envelopes, stored at
242 14°C and inspected daily until death. Lifespan under these conditions correlates strongly
243 with monarch lifetime reproduction (fitness) (De Roode *et al.* 2008b). Parasite virulence
244 was measured as the magnitude reduction in lifespan of infected monarchs when
245 compared to control monarchs. After death, infection success and spore loads were
246 measured from adults following established methods (de Roode *et al.* 2008a, Table S2).
247 Wings were removed and each monarch body was placed in a scintillation vial with 5 mL
248 of deionized water. The mixture was vortexed for five minutes, and 10 µL aliquots were
249 deposited into 4 wells in a hemocytometer for counting. Spore load represents the inverse
250 of monarch resistance. Tolerance to *O. elektroscirra* was measured as the slope of the

251 relationship between spore load and lifespan, with a separate line (slope estimate) for
252 each milkweed species by CO₂ treatment.

253

254 *Plant Defense Measurements*

255 On the same day as inoculations, we sampled milkweed cardenolide and nutrient
256 concentrations using established methods (Zehnder & Hunter 2009; Tao & Hunter 2012).
257 Six 424 mm² leaf disks were taken, deposited in 1 mL methanol and stored at -10°C for
258 cardenolide analysis. Cardenolides were extracted, separated and quantified by reverse-
259 phase high-performance liquid chromatography (HPLC) on a Waters Acquity UPLC with
260 PDA detector (Waters Corporation, Milford, MA, USA) with 0.15mg/mL digitoxin
261 internal standard (Sigma Chemical Company, St. Louis, Missouri, USA). Peaks with
262 symmetrical absorbance between 217-222 nm were identified as cardenolides. Another
263 six disks were collected, weighed, dried, and reweighed to provide estimates of sample
264 dry mass. Remaining foliage from the two punched leaves was collected, oven dried, and
265 analyzed using a TruMac CN Analyzer (Leco Corporation, St. Joseph, MI) to estimate
266 foliar nitrogen (N) concentrations.

267

268

269 *Statistical Methods*

270 Analyses were carried out in R (version 3.3.2). In all of the linear mixed effects models
271 (lme4 package, LMMs) that follow, we included experimental year and chamber identity
272 as random effects, to account for unintended temporal and spatial variation. We also
273 included monarch family as a random effect in all models of monarch performance. For
274 all models we visually inspected homogeneity of variance to confirm model best fit
275 (Crawley 2012) and model comparisons were performed using likelihood ratio tests.

276

277 *Monarch performance*

278 Our analyses are limited to only those monarchs that survived to adulthood, including
279 successfully infected monarchs, and uninoculated (control) monarchs (Table S1 & S2,
280 Appendix S1) (Sternberg *et al.* 2012). Analyses of parasite burden were restricted to

281 infected monarchs, but analyses of tolerance included both infected and uninfected
282 monarchs.
283
284 To investigate effects of our treatments on monarch tolerance, we modeled adult lifespan
285 (square-root-transformed) as a function of spore load (square-root-transformed)
286 (Sternberg et al. 2012), including milkweed species and CO₂ treatment as fixed effects.
287 To investigate the effects of CO₂ treatment and milkweed species on parasite virulence,
288 we used monarch life span (square-root-transformed) as the dependent variable and
289 parasite treatment as a fixed effect. Finally, to estimate monarch resistance, we included
290 monarch spore load (square-root-transformed) as the dependent variable, with milkweed
291 species and CO₂ treatment as fixed effects. Analysis of monarch resistance was only
292 conducted on infected individuals. Due to the nature of the parasite life cycle (McLaughlin
293 et al. 1970), we cannot measure the spore load of larval monarchs that died before
294 adulthood because spores have not yet formed.
295
296 To test for a trade-off between host tolerance and resistance, we associated monarch
297 tolerance with resistance to *O. elektroscirra* using a linear regression (Råberg et al.
298 2009). We assessed any relationship between the 16 tolerance slope values of each
299 treatment group in each year and the mean resistance values (1/spore load) of those
300 treatment groups.
301
302 *Milkweed chemistry and elevated CO₂*
303 We explored the responses to our CO₂ treatments of foliar (a) cardenolide concentration
304 (log-transformed), (b) cardenolide diversity, (c) cardenolide polarity, and (d) N
305 concentration using LMMs. Cardenolide diversity was calculated using the Shannon
306 diversity index: $H = -\sum(P_i \log[P_i])$ where P_i is the relative amount of a cardenolide peak
307 compared to the total amount of cardenolides in an individual plant (Rasmann & Agrawal
308 2011). We excluded *A. incarnata* plants from the LMM exploring effects of CO₂
309 treatments on cardenolide diversity because only 2 individuals produced more than one
310 cardenolide peak. Following Rasmann & Agrawal (2011), we calculated an index of
311 cardenolide polarity $P = \sum(P_i RT_i)$, where RT_i is the retention time of the i th peak in the

312 individual. The polarity index values that result range from 0 (highly polar) to 1 (highly
313 lipophilic).

314

315 We also compared cardenolide composition among milkweed species and CO₂
316 treatments using permutational multivariate analysis of variance (PerMANOVA)
317 (Anderson 2001) following Bray-Curtis ordination. The analysis was performed using the
318 metaMDS function from the Adonis procedure of the Vegan package in R. We performed
319 Nonmetric Multidimensional Scaling (NMDS) (McCune & Grace 2002) reducing the
320 dimensions of the model with 999 permutations per model run and a maximum of 500
321 runs per dimension. We ultimately used a three-dimensional model in subsequent
322 analyses (model stress = 0.119) (McCune and Grace 2002).

323

324 Within the cardenolides monarchs ingest, two specific cardenolides (RT585 and RT650)
325 have been associated with the medicinal efficacy of milkweed species against *O.*
326 *elektroscirra* (de Roode *et al.* 2011b). In our experiment, we found these two
327 cardenolides only in the foliage of *A. curassavica*, with the exception of one *A. syriaca*
328 plant grown under aCO₂. Milkweed species are known to vary substantially in the types
329 of cardenolides they produce (Sternberg *et al.* 2012). Given the established importance of
330 these two compounds, and the losses that we observed in the medicinal activity of *A.*
331 *curassavica* under elevated CO₂ (see Results, below), we ran separate LMMs with each
332 of these cardenolides as dependent variables and CO₂ treatment as a fixed effect.

333

334 *Milkweed chemistry and monarch performance*

335 We observed significant effects of elevated CO₂ on monarch tolerance and parasite
336 virulence on only one milkweed species, *A. curassavica* (see Results). We did not
337 analyze resistance further because it did not vary with any of our treatments. We used
338 LMMs to assess any individual and interactive effects of *A. curassavica* traits on
339 monarch tolerance and parasite virulence. All analyses were restricted to those *A.*
340 *curassavica* plants for which we had measures of cardenolide diversity and
341 corresponding N data (N=77). We performed a PerMANOVA on the cardenolide
342 communities produced in *A. curassavica* alone. Additionally, we created a PCA of the

343 center-log-ratios of cardenolide concentrations and used PCA axes 1 and 2 in separate
344 LMMs to gauge the explanatory strength of the CO₂ treatment in determining monarch
345 tolerance.

346
347 We used Akaike's information criterion (AIC) scores to select chemical traits that were
348 associated with virulence or tolerance. We planned to add additional traits (and
349 interactions) to each model only if the AIC scores improved by two points (Burnham &
350 Anderson 2002, Table S6). However, in no case was more than one independent variable
351 included in any model.

352
353 *(i) Chemistry and Tolerance.* Using the procedure described above, we assessed
354 associations between *A. curassavica* traits and monarch tolerance (slope of fitness
355 decline) by investigating effects of spore load and plant traits on lifespan (square-root-
356 transformed).

357
358 *(ii) Chemistry and Virulence.* Likewise, we measured associations between individual *A.*
359 *curassavica* traits and parasite virulence by investigating the effects of infection
360 treatment and plant traits on lifespan (square-root-transformed).

361

362

363 **Results**

364 ***Monarch tolerance and parasite virulence***

365 Monarch tolerance to *O. elektroscirra* declined by 77% under eCO₂ for individuals
366 reared on the medicinal milkweed, *A. curassavica* (Fig. 1). The tolerance of monarchs
367 feeding on less medicinal milkweed species remained unchanged under eCO₂ (spore load
368 *species*CO₂: F_{3,315}= 4.50, p= 0.00415, Fig. 1, Table S3). These results suggest that
369 eCO₂ reduced the protective properties of *A. curassavica* to similar strength as *A.*
370 *incarnata* under eCO₂ (Fig. 1).

371

372 Consistent with effects on tolerance, the virulence of *O. elektroscirra* increased under
373 eCO₂ in those monarchs reared on *A. curassavica* and remained unchanged on the lower

374 cardenolide milkweed species (infection*species*CO₂: F_{3,308} = 4.44, p = 0.0045, Fig. 2a).
375 Essentially, eCO₂ made *A. curassavica* non-medicinal, magnifying the reduction in
376 fitness caused by infection to values similar to those of monarchs feeding on the other
377 three milkweed species. The magnitude of the reduction in lifespan between control and
378 infected monarchs feeding on *A. curassavica* increased from 1.8 days to 7.2 days under
379 eCO₂ (Fig. 2b). As expected, all infected monarchs had shorter lifespans than uninfected
380 monarchs (infection: F_{1,314} = 263.55, p < 0.0001, Fig. 2b).

381
382 In contrast to their effects on monarch tolerance and parasite virulence, we found no
383 effects of eCO₂ (F_{1,129} = 1.71, p = 0.1931), host plant species (F_{3,137} = 1.28, p = 0.2845), or
384 their interaction (F_{3,138} = 1.35, p = 0.2596) on monarch resistance to the parasite as
385 measured by spore load. Additionally, we found no tradeoff between monarch tolerance
386 and resistance to *O. elektroscirra* (F_{1,14} = 0.91, p = 0.3559).

387

388 ***Milkweed chemistry and elevated CO₂***

389 Foliar cardenolide concentrations were twelve times higher in *A. curassavica* than in *A.*
390 *syriaca* (milkweed species: F_{3,166} = 192.31, p < 0.0001, Fig. 3a). Cardenolide
391 concentrations declined under eCO₂ across all milkweed species (CO₂: F_{1,166} = 5.77,
392 p = 0.0174, Fig. 3a), and there was no interaction between milkweed species and CO₂
393 treatment (F_{1,166} = 0.48, p = 0.6963). The diversity of cardenolide molecular forms was four
394 times higher in *A. curassavica* than in the other milkweed species (milkweed species: F_{3,}
395 ₁₀₉ = 47.11, p < 0.0001, Fig. 3b) and declined under eCO₂ in all milkweed species but *A.*
396 *incarnata*, a species which rarely produces more than one cardenolide (CO₂: F_{1,33} = 5.63,
397 p = 0.02362). There was no interaction between milkweed species and CO₂ treatment on
398 cardenolide diversity (milkweed species*CO₂: F_{2,141} = 0.54, p = 0.58274). The average
399 polarity of *A. curassavica* and *A. speciosa* cardenolides was reduced by eCO₂ treatments,
400 while the average polarity of cardenolides increased in *A. syriaca* (species*CO₂: F_{3,}
401 ₁₅₃ = 2.99, p = 0.03281, Fig. 3c).

402

403 Across all four species, foliar N concentrations (plant nutritional quality estimate)
404 declined under eCO₂ (CO₂: F_{1,48} = 12.33, p = 0.00098, Fig. 3d). Milkweed species also

405 varied in their foliar N concentrations from 1.42% in *A. curassavica* to 1.14% in *A.*
406 *syriaca* ($F_{1,137} = 4.43$, $p = 0.0052$, Fig. 3d).

407

408 Beyond simple measures of cardenolide polarity and diversity, milkweed species differed
409 in the composition of foliar cardenolides (PerMANOVA; species: $F_{3, 171} = 20.02$, $p =$
410 0.001 , $R^2 = 0.26$, Fig. S2). The effects of CO₂ treatment on cardenolide community
411 composition varied among milkweed species (PerMANOVA; species*CO₂: $F_{1, 171} = 2.12$,
412 $p = 0.001$, $R^2 = 0.027$, Fig. S2). In *A. curassavica* alone, the communities of cardenolides
413 produced by individual plants differed between CO₂ treatments (CO₂: $F_{1, 76} = 2.80$, $p =$
414 0.03 , $R^2 = 0.036$, Fig. 4a). NMDS1 was associated with declines in the concentrations of
415 lipophilic cardenolides (decline in polarity index) ($p < 0.0001$, $R^2 = 0.47$, Fig. 4b).

416

417 Concentrations of both RT585 and RT650, the two cardenolides with established
418 medicinal activity (de Roode et al. 2011b), declined in *A. curassavica* under eCO₂.
419 Concentrations of RT585 declined by 25% under eCO₂ ($F_{1, 61} = 5.36$, $p = 0.02401$, Fig. 5a).
420 Far fewer *A. curassavica* individuals produced RT650 (N=6). Nonetheless, we detected a
421 marginally significant 65% decline in RT650 under eCO₂ ($F_{1, 4} = 5.92$, $p = 0.0717$, Fig.
422 5b).

423

424

425 ***A. curassavica* chemistry and monarch performance**

426 *Tolerance*

427 A significant interaction between spore load and a given plant trait on monarch lifespan
428 indicates a correlation between that trait and tolerance to *O. elektroscirra*. Monarch
429 tolerance varied with the expression of lipophilic cardenolides in leaves (spore load*
430 polarity: $F_{1, 70} = 4.10$, $p = 0.04678$, Fig. 6a). Interestingly, the model containing
431 cardenolide polarity fit the tolerance data better than did a model containing CO₂
432 treatment alone. Further, neither models containing the PCA axes of center-log-ratio-
433 transformed cardenolide concentrations fit the data better than a model with just CO₂
434 treatment (Table S5), supporting our findings that cardenolide traits are more important
435 in determining tolerance than concentration.

436

437 *Virulence*

438 A significant interaction between a plant trait and parasite treatment on monarch lifespan
439 indicates a relationship between that trait and parasite virulence. As with tolerance, there
440 was a positive relationship between the expression of lipophilic cardenolides and the
441 lifespan of infected individuals (declining virulence) but it was marginally significant
442 (parasite treatment*polarity: $F_{1, 70} = 3.26$, $p = 0.0753$, Fig. 6b).

443

444

445 **Discussion**

446 Monarch butterflies benefit from the medicinal properties of milkweeds when combating
447 their parasites (de Roode *et al.* 2008a, 2011a; Sternberg *et al.* 2012; Gowler *et al.* 2015),
448 and infected females can choose the most medicinal milkweeds for oviposition in
449 laboratory choice tests (Lefèvre *et al.* 2010, 2012). Here, we show that a medicinal
450 milkweed species, *A. curassavica*, loses its protective abilities under eCO₂. Our results
451 suggest that rising CO₂ will reduce the tolerance of monarch butterflies to their common
452 parasite, *Ophryocystis elektroscirrha*, and will increase parasite virulence. Ongoing
453 changes in water availability (Andrews 2015), temperature (Couture *et al.* 2015) and soil
454 nutrient loading (Zehnder & Hunter 2009; Tao *et al.* 2014) have already been shown to
455 influence the cardenolide chemistry of milkweeds, with consequences for parasite-
456 monarch interactions. Here, we add eCO₂ to the list of drivers that may alter monarch
457 parasite-host interactions in a changing world.

458

459 We observed the lowest tolerance values in those monarchs feeding on *A. syriaca* grown
460 under eCO₂ and the highest tolerance values in those monarchs feeding on *A. curassavica*
461 grown under aCO₂. However, monarchs feeding on the same species of milkweed that
462 once conveyed a tolerance advantage under aCO₂ (*A. curassavica*), experienced a 77%
463 reduction in tolerance under eCO₂. Parasites caused the most virulence when monarchs
464 fed on *A. incarnata* under aCO₂, reducing host lifespan by nearly 8 days (see Fig. 2b).
465 Parasites caused the least virulence in those monarchs feeding on *A. curassavica* grown
466 under aCO₂, reducing mean lifespan by only 2 days. Importantly, monarchs feeding on

467 this same species, *A. curassavica*, under eCO₂ experienced virulence of comparable
468 values to non-medicinal species like *A. incarnata*, suffering a reduction in lifespan of 7
469 days due to infection. These results are the first to show effects of environmental change
470 on host tolerance to parasites and parasite virulence resulting from indirect effects
471 mediated by community members.

472
473 A growing number of studies stress the importance of understanding the indirect
474 mechanisms by which disease will respond to changing environmental conditions
475 (Tylianakis *et al.* 2008; Altizer *et al.* 2013; Gunderson *et al.* 2017). Indirect effects of
476 environmental change on host-parasite interactions emerge from additional members of
477 ecological communities (Keesing *et al.* 2006; Wolinska & King 2009; Vuong *et al.* 2017).
478 Associated predators, competitors and symbionts are all subject to the effects of
479 environmental change, which may alter their interactions with host-parasite pairs (Ritchie
480 2006; Gherlenda *et al.* 2015). Here, we discover a previously unrecognized indirect
481 mechanism by which environmental change can act on disease: the loss of medicinal
482 compounds in host diet, contributing to reductions in host tolerance and increases in
483 parasite virulence.

484
485 Changes in host tolerance and parasite virulence under future environmental conditions
486 have important evolutionary implications. Theory predicts that reductions in resistance
487 will lesson antagonistic coevolution between host and parasite (Roy & Kirchner 2000;
488 Råberg *et al.* 2009; Rohr *et al.* 2010). However, we are less certain what changes in host
489 tolerance could mean for host-parasite dynamics (Best *et al.* 2008; Schneider & Ayres
490 2008). Because tolerance helps to maintain host fitness when infected, less tolerant hosts
491 should suffer shorter infections due to increased mortality, thereby potentially decreasing
492 transmission and the prevalence of parasites in the host population (Miller *et al.* 2006). In
493 our study, reduced tolerance was also accompanied by increased parasite virulence. In
494 some cases, increased in virulence may lead to local extinction (Kutzer & Armitage
495 2016a; Wilber *et al.* 2017). We expect parasites that cause higher virulence to be selected
496 against when host tolerance is also reduced because the risk of premature host mortality
497 is higher. Early host death reduces parasite fitness and thus, induces selection on parasite

498 virulence to decrease to a new optimum (Little *et al.* 2010). Given the reductions in host
499 tolerance and increases in early monarch death, we predict that future environmental
500 conditions may select for intrinsically less virulent parasites. Moreover, it is important to
501 note that even when parasites evolve lower levels of intrinsic virulence, the actual
502 virulence experienced by infected monarchs is likely to increase due to their reduced
503 plant-derived tolerance. Additionally, our study only investigated the response of two
504 parasite families (one in each year) to eCO₂ and milkweed species. To make better
505 inferences about the evolutionary trajectories of parasite virulence and host tolerance
506 under future conditions, further studies should examine the importance of parasite
507 families under eCO₂.

508

509 Our results add to a substantial body of work that emphasizes the role of environmental
510 factors in phytophagous host-parasite interactions (Cory & Hoover 2006; Myers & Cory
511 2016; Shikano 2017). The largest declines in tolerance and increases in virulence
512 occurred in monarchs feeding on *A. curassavica*, a species in which cardenolide
513 production declined by nearly 25% when grown under eCO₂. However, total cardenolide
514 concentrations did not correlate with changes in tolerance. Rather, reductions in
515 cardenolide concentration under eCO₂ occurred in concert with changes in cardenolide
516 community composition and declines in the expression of lipophilic cardenolides (Figs. 4,
517 5). Because the polarity of cardenolides partially determines their biological activity
518 (Agrawal *et al.* 2012), the loss of lipophilic cardenolides under eCO₂ compromises the
519 anti-parasitic properties of milkweed foliage. Infected monarchs that consume lipophilic
520 cardenolides live longer than do those infected monarchs consuming polar cardenolides
521 (Sternberg *et al.* 2012; Tao *et al.* 2016a). Previous work has shown that declines in the
522 concentrations of two key lipophilic cardenolides, RT585 and RT650, increase parasite
523 virulence (de Roode *et al.* 2011b). We also observed reductions in these two cardenolides
524 in our populations of *A. curassavica* that were exposed to eCO₂, which likely led to the
525 observed increases in parasite virulence. Interestingly, *A. syriaca* and *A. speciosa* did not
526 produce these compounds and were also not protective in our study. Perhaps this is
527 because different milkweed populations and individuals vary substantially in the
528 concentration and composition of cardenolides produced within species (Vannette &

529 Hunter 2011). *Asclepias syriaca*, specifically, is known to exhibit striking variation in
530 cardenolide concentrations among populations (Andrews 2015). It is also important to
531 note that the lifespan of uninfected monarchs feeding on *A. curassavica* increased under
532 eCO₂. This increase in the fitness of uninfected monarchs may illustrate the cost of
533 consuming toxic, lipophilic cardenolides (Sternberg *et al.* 2012; Tao *et al.* 2016b)
534 because the frequency of those potent compounds declined under eCO₂ as monarch
535 lifespan increased.

536

537 By demonstrating that tolerance to parasite infection can be altered by environmental
538 change, we reinforce the idea that tolerance is not solely determined by intrinsic host
539 factors but relies additionally on environmental conditions including interactions with
540 other community members. As the environmental factors that mediate disease change,
541 further empirical studies are sorely needed to explore the interplay between multiple
542 global change drivers and host-parasite interactions embedded within diverse ecological
543 communities.

544 **Acknowledgements**

545 Many thanks to H.B. Streit, A.R. Meier, K.C. Crocker, C.R. Chappell, M.O. Hemken, Y.
546 Yang, J. McMahon, J.J. Shi, R. Peterson, and J.D. Den Uyl, L. Tao, C.D. Gowler, A.A.
547 Pierce, and A.J. Mongue. This work was supported by National Science Foundation
548 grants DEB-1257160 and DEB-1256115 awarded to J.C.dR. and M.D.H., respectively.

549

550

551 **References**

552 Adamo, S.A. & Lovett, M.M. (2011). Some like it hot: the effects of climate change on
553 reproduction, immune function and disease resistance in the cricket *Gryllus texensis*.
554 *J. Exp. Biol.*, 214, 1997–2004.

555 Agrawal, A.A., Petschenka, G., Bingham, R.A., Weber, M.G. & Rasmann, S. (2012).
556 Toxic cardenolides: chemical ecology and coevolution of specialized plant-
557 herbivore interactions. *New Phytol.*, 194, 28–45.

558 Altizer, S., Ostfeld, R.S., Johnson, P.T.J., Kutz, S. & Harvell, C.D. (2013). Climate
559 change and infectious diseases: from evidence to a predictive framework. *Science*,

560 341, 514–9.

561 Anderson, M.J. (2001). A new method for non parametric multivariate analysis of
562 variance. *Austral Ecol.*, 26, 32–46.

563 Andrews, H. (2015). Changes in water availability and variability affect plant defenses
564 and herbivore responses in grassland forbs. University of Michigan.

565 Bale, J.S., Masters, G.J., Hodkinson, I.D., Awmack, C., Bezemer, T.M., Brown, V.K., *et*
566 *al.* (2002). Herbivory in global climate change research: direct effects of rising
567 temperature on insect herbivores. *Glob. Chang. Biol.*, 8, 1–16.

568 Baucom, R.S. & De Roode, J.C. (2011). Ecological immunology and tolerance in plants
569 and animals. *Funct. Ecol.*, 25, 18–28.

570 Becker, D.J., Streicker, D.G. & Altizer, S. (2015). Linking anthropogenic resources to
571 wildlife-pathogen dynamics: a review and meta-analysis. *Ecol. Lett.*, 1–13.

572 Bernays, E.A. & Singer, M.S. (2005). Taste alteration and endoparasites. *Nature*, 436,
573 476–476.

574 Best, A., White, A. & Boots, M. (2008). Maintenance of host variation in tolerance to
575 pathogens and parasites. *Proc. Natl. Acad. Sci. U. S. A.*, 105, 20786–20791.

576 Bidart-Bouzat, M.G. & Imeh-Nathaniel, A. (2008). Global change effects on plant
577 chemical defenses against insect herbivores. *J. Integr. Plant Biol.*, 50, 1339–54.

578 Boots, M. (2011). The Evolution of Resistance to a Parasite Is Determined by Resources.
579 *Am. Nat.*, 178, 214–220.

580 Boots, M., Best, A., Miller, M.R. & White, A. (2009). The role of ecological feedbacks in
581 the evolution of host defence: what does theory tell us? *Philos. Trans. R. Soc. Lond.*
582 *B. Biol. Sci.*, 364, 27–36.

583 Bradley, C.A. & Altizer, S. (2005). Parasites hinder monarch butterfly flight:
584 implications for disease spread in migratory hosts. *Ecol. Lett.*, 8, 290–300.

585 Bruno, J.F., Petes, L.E., Drew Harvell, C. & Hettinger, A. (2003). Nutrient enrichment
586 can increase the severity of coral diseases. *Ecol. Lett.*, 6, 1056–1061.

587 Burnham, K.P. & Anderson, D.R. (2002). *Model Selection and Multimodel Inference: A*
588 *Practical Information-Theoretic Approach. Ecol. Modell.* 2nd edn. Springer Science
589 & Business Media.

590 Clough, D., Prykhodko, O. & Råberg, L. (2016). Effects of protein malnutrition on

591 tolerance to helminth infection. *Biol. Lett.*, 12.

592 Cory, J.S. & Hoover, K. (2006). Plant-mediated effects in insect-pathogen interactions.
593 *Trends Ecol. Evol.*, 21, 278–86.

594 Couture, J.J., Serbin, S.P. & Townsend, P.A. (2015). Elevated temperature and periodic
595 water stress alter growth and quality of common milkweed (*Asclepias syriaca*) and
596 monarch (*Danaus plexippus*) larval performance. *Arthropod. Plant. Interact.*, 9,
597 149–161.

598 Crawley, M.J. (2012). *Statistical Modelling*. R B. John Wiley & Sons, Ltd, Chichester,
599 UK.

600 Debes, P.V., Gross, R. & Vasemägi, A. (2017). Quantitative Genetic Variation in, and
601 Environmental Effects on, Pathogen Resistance and Temperature-Dependent
602 Disease Severity in a Wild Trout. *Am. Nat.*, 190, 244–265.

603 Drake, B.G., Leadley, P.W., Arp, W.J., Nassiry, D. & Curtis, P.S. (1989). An Open Top
604 Chamber for Field Studies of Elevated Atmospheric CO₂ Concentration on
605 Saltmarsh Vegetation. *Funct. Ecol.*, 3, 363.

606 Felton, G.W. & Duffey, S.S. (1990). Inactivation of baculovirus by quinones formed in
607 insect-damaged plant tissues. *J. Chem. Ecol.*, 16, 1221–1236.

608 Franke, F., Armitage, S.A.O., Kutzer, M.A.M., Kurtz, J. & Scharsack, J.P. (2017).
609 Environmental temperature variation influences fitness trade-offs and tolerance in a
610 fish-tapeworm association. *Parasit. Vectors*, 10, 252.

611 Garamszegi, L.Z. (2011). Climate change increases the risk of malaria in birds. *Glob.*
612 *Chang. Biol.*, 17, 1751–1759.

613 Gherlenda, A.N., Haigh, A.M., Moore, B.D., Johnson, S.N. & Riegler, M. (2015).
614 Climate change, nutrition and immunity: Effects of elevated CO₂ and temperature
615 on the immune function of an insect herbivore. *J. Insect Physiol.*

616 Gowler, C.D., Leon, K.E., Hunter, M.D. & de Roode, J.C. (2015). Secondary Defense
617 Chemicals in Milkweed Reduce Parasite Infection in Monarch Butterflies, *Danaus*
618 *plexippus*. *J. Chem. Ecol.*, 41, 520–523.

619 Gunderson, A.R., Tsukimura, B. & Stillman, J.H. (2017). Indirect Effects of Global
620 Change: From Physiological and Behavioral Mechanisms to Ecological
621 Consequences. *Integr. Comp. Biol.*, 57, 48–54.

622 Hayward, A.D., Nussey, D.H., Wilson, A.J., Berenos, C., Pilkington, J.G., Watt, K.A., *et*
623 *al.* (2014). Natural Selection on Individual Variation in Tolerance of Gastrointestinal
624 Nematode Infection. *PLoS Biol.*, 12, 1–13.

625 Howick, V.M. & Lazzaro, B.P. (2014). Genotype and diet shape resistance and tolerance
626 across distinct phases of bacterial infection. *BMC Evol. Biol.*, 14, 56.

627 Hunter, M.D. (2001). Effects of elevated atmospheric carbon dioxide on insect-plant
628 interactions. *Agric. For. Entomol.*, 3, 153–159.

629 Hunter, M.D. & Schultz, J.C. (1993). Induced plant defenses breached? Phytochemical
630 induction protects an herbivore from disease. *Oecologia*, 94, 195–203.

631 Jackson, J.A., Hall, A.J., Friberg, I.M., Ralli, C., Lowe, A., Zawadzka, M., *et al.* (2014).
632 An immunological marker of tolerance to infection in wild rodents. *PLoS Biol.*, 12,
633 1–13.

634 Keesing, F., Holt, R.D. & Ostfeld, R.S. (2006). Effects of species diversity on disease
635 risk. *Ecol. Lett.*, 9, 485–98.

636 Kutzer, M.A.M. & Armitage, S.A.O. (2016a). Maximising fitness in the face of parasites:
637 a review of host tolerance. *Zoology*, 119, 281–289.

638 Kutzer, M.A.M. & Armitage, S.A.O. (2016b). The effect of diet and time after bacterial
639 infection on fecundity, resistance, and tolerance in *Drosophila melanogaster*. *Ecol.*
640 *Evol.*

641 Lambrechts, L., Fellous, S. & Koella, J.C. (2006). Coevolutionary interactions between
642 host and parasite genotypes. *Trends Parasitol.*, 22, 12–6.

643 Lampert, E. (2012). Influences of plant traits on immune responses of specialist and
644 generalist herbivores. *Insects*, 3, 573–592.

645 Lefèvre, T., Chiang, A., Kelavkar, M., Li, H., Li, J., de Castillejo, C.L.F., *et al.* (2012).
646 Behavioural resistance against a protozoan parasite in the monarch butterfly. *J. Anim.*
647 *Ecol.*, 81, 70–79.

648 Lefèvre, T., Oliver, L., Hunter, M.D. & De Roode, J.C. (2010). Evidence for trans-
649 generational medication in nature. *Ecol. Lett.*, 13, 1485–1493.

650 Lefèvre, T., Williams, A.J. & de Roode, J.C. (2011). Genetic variation in resistance, but
651 not tolerance, to a protozoan parasite in the monarch butterfly. *Proc. Biol. Sci.*, 278,
652 751–9.

653 Little, T.J., Shuker, D.M., Colegrave, N., Day, T. & Graham, A.L. (2010). The
654 Coevolution of Virulence: Tolerance in Perspective. *PLoS Pathog.*, 6, e1001006.

655 Malcolm, S.B. & Brower, L.P. (1989). Reviews Evolutionary and ecological implications
656 of cardenolide sequestration in the monarch butterfly. *Experientia*, 45, 284–295.

657 Malcolm, S.B. & Zalucki, M.P. (1996). Milkweed latex and cardenolide induction may
658 resolve the lethal plant defence paradox. *Entomol. Exp. Appl.*, 80, 193–196.

659 Matiella, T.J. (2012). The effects of carbon dioxide on three Species of milkweed
660 (*Asclepiadaceae*) and Monarch butterfly (*Danaus plexippus*) larva feeding
661 preference. Retrieved from ProQuest Diss. Theses. (Accession Order No. 3548664).

662 McCune, B. & Grace, J.B. (2002). *Analysis of ecological communities. Struct. Equ.*
663 *Model.*

664 Mclaughlin, R.E., Myers, J., Diw, E.R., Sem, A.R. & College, S. (1970). Monarch
665 Butterfly *Danaus plexippus* (L .) and the Florida Queen Butterfly *D . gilippus*
666 *berenicae* Cramerl. *J. Protozool.*, 17, 300–305.

667 Miller, M.R., White, A. & Boots, M. (2006). The evolution of parasites in response to
668 tolerance in their hosts: the good, the bad, and apparent commensalism. *Evolution*
669 (*N. Y.*), 60, 945–956.

670 Myers, J.H. & Cory, J.S. (2016). Ecology and evolution of pathogens in natural
671 populations of Lepidoptera. *Evol. Appl.*, 9, 231–247.

672 Paull, S.H. & Johnson, P.T.J. (2014). Experimental warming drives a seasonal shift in the
673 timing of host-parasite dynamics with consequences for disease risk. *Ecol. Lett.*

674 Råberg, L., Graham, A.L. & Read, A.F. (2009). Decomposing health: tolerance and
675 resistance to parasites in animals. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.*, 364, 37–
676 49.

677 Råberg, L., Sim, D. & Read, A.F. (2007). Disentangling Genetic Variation for Resistance
678 and Tolerance to Infectious Diseases in Animals. *Science*. 318, 1305–1309.

679 Rasmann, S. & Agrawal, A.A. (2011). Latitudinal patterns in plant defense: evolution of
680 cardenolides, their toxicity and induction following herbivory. *Ecol. Lett.*, 14, 476–
681 83.

682 Regoes, R.R., McLaren, P.J., Battagay, M., Bernasconi, E., Calmy, A., Günthard, H.F., *et*
683 *al.* (2014). Disentangling Human Tolerance and Resistance Against HIV. *PLoS Biol.*,

684 12.

685 Restif, O. & Koella, J.C. (2004). Concurrent evolution of resistance and tolerance to
686 pathogens. *Am. Nat.*, 164, E90–E102.

687 Ritchie, K.B. (2006). Regulation of microbial populations by coral surface mucus and
688 mucus-associated bacteria. *Mar. Ecol. Prog. Ser.*, 322, 1–14.

689 Robinson, E.A., Ryan, G.D. & Newman, J.A. (2012). A meta-analytical review of the
690 effects of elevated CO₂ on plant–arthropod interactions highlights the importance of
691 interacting environmental and biological variables. *New Phytol.*, 194, 321–336.

692 Rohr, J.R., Raffel, T.R. & Hall, C.A. (2010). Developmental variation in resistance and
693 tolerance in a multi-host-parasite system. *Funct. Ecol.*, 24, 1110–1121.

694 de Roode, J.C., de Castillejo, C.L.F., Faits, T. & Alizon, S. (2011a). Virulence evolution
695 in response to anti-infection resistance: toxic food plants can select for virulent
696 parasites of monarch butterflies. *J. Evol. Biol.*, 24, 712–22.

697 de Roode, J.C., Chi, J., Rarick, R.M. & Altizer, S. (2009). Strength in numbers: high
698 parasite burdens increase transmission of a protozoan parasite of monarch butterflies
699 (*Danaus plexippus*). *Oecologia*, 161, 67–75.

700 de Roode, J.C., Pedersen, A.B., Hunter, M.D. & Altizer, S. (2008a). Host plant species
701 affects virulence in monarch butterfly parasites. *J. Anim. Ecol.*, 77, 120–126.

702 de Roode, J.C., Rarick, R.M., Mongue, A.J., Gerardo, N.M. & Hunter, M.D. (2011b).
703 Aphids indirectly increase virulence and transmission potential of a monarch
704 butterfly parasite by reducing defensive chemistry of a shared food plant. *Ecol. Lett.*,
705 14, 453–61.

706 de Roode, J.C., Yates, A.J. & Altizer, S. (2008b). Virulence-transmission trade-offs and
707 population divergence in virulence in a naturally occurring butterfly parasite. *Proc.*
708 *Natl. Acad. Sci. U. S. A.*, 105, 7489–94.

709 Roy, B.A. & Kirchner, J.W. (2000). Evolutionary Dynamics of Pathogen Resistance and
710 Tolerance. *Evolution (N. Y.)*, 54, 176–90.

711 Schneider, D.S. & Ayres, J.S. (2008). Two ways to survive infection: what resistance and
712 tolerance can teach us about treating infectious diseases. *Nat. Rev. Immunol.*, 8,
713 889–895.

714 Shikano, I. (2017). Evolutionary Ecology of Multitrophic Interactions between Plants,

715 Insect Herbivores and Entomopathogens. *J. Chem. Ecol.*, 43, 586–598.

716 Shikano, I., Ericsson, J.D., Cory, J.S. & Myers, J.H. (2010). Indirect plant-mediated
717 effects on insect immunity and disease resistance in a tritrophic system. *Basic Appl.*
718 *Ecol.*, 11, 15–22.

719 Solomon, S., Plattner, G.-K., Knutti, R. & Friedlingstein, P. (2009). Irreversible climate
720 change due to carbon dioxide emissions. *Proc. Natl. Acad. Sci. U. S. A.*, 106, 1704–9.

721 Sternberg, E.D., Lefevre, T., Li, J., Lopez, C., Castillejo, F. De, Li, H., *et al.* (2012).
722 Food Plant-Derived Disease Tolerance and Resistance in a Natural Butterfly- Plant-
723 Parasite Interactions. *Evolution (N. Y.)*, 66, 3367–3377.

724 Tao, L., Ahmad, A., de Roode, J.C. & Hunter, M.D. (2016a). Arbuscular mycorrhizal
725 fungi affect plant tolerance and chemical defences to herbivory through different
726 mechanisms. *J. Ecol.*, 104, 561–571.

727 Tao, L., Berns, A.R. & Hunter, M.D. (2014). Why does a good thing become too much?
728 Interactions between foliar nutrients and toxins determine performance of an insect
729 herbivore. *Funct. Ecol.*, 28, 190–196.

730 Tao, L., Gowler, C.D., Ahmad, A., Hunter, M.D. & de Roode, J.C. (2015). Disease
731 ecology across soil boundaries: effects of below-ground fungi on above-ground
732 host–parasite interactions. *Proc. R. Soc. B Biol. Sci.*, 282, 20151993.

733 Tao, L., Hoang, K.M., Hunter, M.D. & de Roode, J.C. (2016b). Fitness costs of animal
734 medication: antiparasitic plant chemicals reduce fitness of monarch butterfly hosts. *J.*
735 *Anim. Ecol.*, 85, 1246–1254.

736 Tao, L. & Hunter, M.D. (2012). Does anthropogenic nitrogen deposition induce
737 phosphorus limitation in herbivorous insects? *Glob. Chang. Biol.*, 18, 1843–1853.

738 Thomas, M.B. & Blanford, S. (2003). Thermal biology in insect-parasite interactions.
739 *Trends Ecol. Evol.*, 18, 344–350.

740 Tylianakis, J.M., Didham, R.K., Bascompte, J. & Wardle, D.A. (2008). Global change
741 and species interactions in terrestrial ecosystems. *Ecol. Lett.*, 11, 1351–1363.

742 Vannette, R.L. & Hunter, M.D. (2011). Genetic variation in expression of defense
743 phenotype may mediate evolutionary adaptation of *Asclepias syriaca* to elevated
744 CO₂. *Glob. Chang. Biol.*, 17, 1277–1288.

745 Vannette, R.L. & Hunter, M.D. (2014). Genetic variation in plant below-ground response

746 to elevated CO₂ and two herbivore species. *Plant Soil*, 384, 303–314.

747 Vuong, H.B., Chiu, G.S., Smouse, P.E., Fonseca, D.M., Brisson, D., Morin, P.J., *et al.*

748 (2017). Influences of Host Community Characteristics on *Borrelia burgdorferi*

749 Infection Prevalence in Blacklegged Ticks. *PLoS One*, 12, 1–17.

750 Wilber, M.Q., Knapp, R.A., Toothman, M. & Briggs, C.J. (2017). Resistance , tolerance

751 and environmental transmission dynamics determine host extinction risk in a load-

752 dependent amphibian disease. *Ecol. Lett.*, 20, 1169–1181.

753 Wolinska, J. & King, K.C. (2009). Environment can alter selection in host-parasite

754 interactions. *Trends Parasitol.*, 25, 236–44.

755 Woodson, R.E. (1954). The North American species of *Asclepias L.* *Ann. Missouri Bot.*

756 *Gard.*, 41, 1–211.

757 Zamora-Vilchis, I., Williams, S.E. & Johnson, C.N. (2012). Environmental temperature

758 affects prevalence of blood parasites of birds on an elevation gradient: Implications

759 for disease in a warming climate. *PLoS One*, 7.

760 Zavala, J.A., Nability, P.D. & DeLucia, E.H. (2013). An emerging understanding of

761 mechanisms governing insect herbivory under elevated CO₂. *Annu. Rev. Entomol.*,

762 58, 79–97.

763 Zehnder, C.B. & Hunter, M.D. (2009). More is not necessarily better: the impact of

764 limiting and excessive nutrients on herbivore population growth rates. *Ecol.*

765 *Entomol.*, 34, 535–543.

766 Zeller, M. & Koella, J.C. (2017). The Role of the Environment in the Evolution of

767 Tolerance and Resistance to a Pathogen. *Am. Nat.*, 190, 389–397.

768 **Figure 1.** Monarch tolerance to *O. elektroscirra* infection as a function of milkweed
769 species and CO₂ treatment. Light gray lines and points correspond to tolerance slopes of
770 monarchs reared on plants grown under ambient CO₂ (400 ppm) and dark gray lines and
771 points correspond to tolerance slopes of monarchs reared on plants grown under elevated
772 CO₂ (760 ppm). Tolerance slopes are presented by milkweed species: CUR = *A.*
773 *curassavica*, SYR = *A. syriaca*, SPE = *A. speciosa*, INC = *A. incarnata*.

774
775 **Figure 2.** The virulence of *O. elektroscirra* parasites increases under elevated CO₂
776 when monarch larvae feed on *A. curassavica*. Virulence is measured as the magnitude of
777 the reduction in host fitness resulting from infection. In (a), points represent the
778 standardized difference (Hedge's $d \pm 95\%$ CI) in mean lifespan between uninfected and
779 infected monarchs fed different species of milkweed under ambient CO₂ (400 ppm, light
780 gray) and elevated CO₂ (760 ppm, dark gray). In (b), we show mean lifespan of parasite-
781 infected (dark gray bars) and uninfected (light gray bars) monarchs (± 1 SE) used to
782 calculate the Hedge's d values shown in (a). Longevities were transformed to
783 approximate normality of errors before statistical analyses but are presented here as
784 untransformed values for ease of interpretation. Milkweed species codes match those
785 presented above.

786
787 **Figure 3.** Effects of elevated CO₂ on foliar cardenolide concentrations (mg/g dry mass,
788 a), cardenolide diversity (b), cardenolide polarity index (c), and foliar nitrogen
789 concentration (%N) (d) of four milkweed species. Trait values were transformed to
790 approximate normality of errors before analyses but are presented here in their
791 untransformed values for ease of interpretation. Light gray bars represent plants grown
792 under ambient CO₂ and dark gray bars are those from elevated CO₂ (± 1 SE). Milkweed
793 species codes match those presented above.

794
795 **Figure 4.** *A. curassavica* plants differed in the composition of cardenolides that they
796 produced under the different CO₂ treatments. In (a) light gray points represent plants
797 grown under ambient CO₂ (400ppm) and dark gray points represent plants grown under

798 elevated CO₂ (760 ppm). In (b), we illustrate the negative association between NMDS
799 axis 1 and the occurrence of lipophilic cardenolides.

800

801 **Figure 5.** Effects of elevated CO₂ on the concentration of two medicinal cardenolides:
802 (a) RT585 and (b) RT650 in *A. curassavica*. Light gray bars represent foliar samples
803 taken from plants grown under ambient CO₂ and dark gray bars are those from elevated
804 CO₂ (± 1 SE).

805

806 **Figure 6.** The effects of cardenolide polarity on (a) monarch tolerance to infection by *O.*
807 *elektroskirra* and (b) the lifespan of infected and uninfected monarchs. A high polarity
808 index reflects greater expression of lipophilic cardenolides. The slopes of the lines in (a)
809 indicate monarch tolerance to infection, with steeper slopes representing lower tolerance.
810 For visual simplicity, we have binned butterflies in (a) by the polarity of the cardenolides
811 that they consumed as larvae. However, the analysis was performed with un-binned
812 polarity data, and binning was used purely as a simplified alternative to a 3D graph. We
813 present the square-root-transformed variables for ease of comparison with figure 1. In (b),
814 light gray points and lines indicate uninfected (Control) monarchs, and dark gray points
815 and lines indicate infected monarchs.

Figure 1.





Figure 2.

(a)



(b)




Figure 3.





Figure 4.

(a)



(b)




Figure 5.





Figure 6.

(a)



(b)

