

1

2 MISS TEAGAN GALE (Orcid ID : 0000-0003-0259-5071)

3 PROFESSOR ROB BROOKS (Orcid ID : 0000-0001-6926-0781)

4

5

6 Article type : Research Article

7

8

9 Section: Animal Growth and Development

10 Editor: Dr Christine Miller

11

12 **Perceived threats of infanticide reduce maternal allocation during lactation and lead to**
13 **elevated oxidative damage in offspring**

14

15 Teagan Gale¹, Michael G. Garratt², & Robert C. Brooks¹

16 ¹*Evolution and Ecology Research Centre, School of Biological, Earth and Environmental Sciences (BEES), the*
17 *University of New South Wales, High Street, Kensington, NSW 2052, Australia. 2**Department of Pathology,*

18 *University of Michigan Medical School, Ann Arbor, MI 48109, United States.*

19 **Corresponding Author:** Teagan Gale: t.gale@unsw.edu.au

20 **Abstract**

- 21 1. Maternal investment is costly to the mother but essential to offspring survival in
22 altricial species. Infanticide by novel males results in loss of maternal investment, and
23 maternal strategies have evolved to mitigate these losses. One such maternal strategy,
24 the Bruce effect, involves spontaneous abortion by females of some mammal species
25 when exposed to a novel male during pregnancy.
- 26 2. In mice, the Bruce effect only occurs during early pregnancy, but we have previously
27 found that female mice exposed to a novel male's scent in late pregnancy weaned
28 smaller offspring. Here we replicate that manipulation in order to resolve the cause of
29 the reduced weaning weight and subsequent effects on offspring fitness.

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/1365-2435.13146](https://doi.org/10.1111/1365-2435.13146)

This article is protected by copyright. All rights reserved

- 30 3. Females exposed to an unfamiliar male's scent in late pregnancy spent significantly
31 less time nursing their pups during lactation, suggesting that reduced maternal
32 allocation contributes to slower offspring growth. The offspring with a reduced
33 weaning weight exhibited catch-up growth and reached a normal weight at adulthood.
34 These offspring, however, were found to bear oxidative damage in adulthood,
35 revealing long-term effects on offspring condition.
- 36 4. We conclude that female mice strategically alter their investment in lactation in
37 relation to the likelihood of infanticide, but that this results in long term fitness costs
38 to their offspring.

39
40 **Running head:** Female mice reduce investment when they perceive a threat of infanticide

41 **Key words:** catch-up growth, infanticide, maternal effects, oxidative stress.

42 **Introduction**

43

44 In mammals, reproduction is characterised by large maternal investments in pregnancy, lactation
45 and other behaviours critical for offspring survival, growth and development (Wolff and Sherman
46 2008). This investment such as nutrition, warmth, and protection can be costly to the mother
47 (Bronson 1989; Clutton-Brock and Vincent 1991), and those costs are exacted in the currency of
48 residual reproductive value (König et al. 1988). Generally, the relationship between mother and
49 offspring is a cooperative interaction, with the mother benefitting from the success of her progeny.
50 However, with current and future reproduction known to trade-off (Reznick 1985; Stearns 1992;
51 Kirkwood and Austad 2000), a mother can also benefit from altering, and sometimes reducing, her
52 investment in current offspring according to environmental conditions, in order to optimize her
53 overall fitness.

54

55 In mice (as in most mammals) lactation is the most energetically demanding episode of
56 reproduction (Speakman 2008) and is a key time to modify investment in relation to contextual
57 conditions. Maternal investment in lactation is somewhat flexible in mice (König 1985; König and
58 Markl 1987), and offspring do not get more milk than corresponds to the mother's optimal
59 allocation during lactation, despite frequent suckling attempts (König and Markl 1987). Other
60 species, such as wild primates have been observed to alter maternal investment and wean offspring
61 prematurely when an unfamiliar male enters the natal group (Teichroeb and Sicotte 2008; Zhao et
62 al. 2011). Such flexibility may significantly enhance maternal fitness.

63

64 Females have also evolved other strategies to optimise their maternal investment when faced
65 with the prospect of infanticide. Infanticide represents one of the more extreme forms of
66 sexual conflict that results in a significant loss (and waste) of maternal investment for
67 females. Many mammalian species have been documented to commit infanticide, which is
68 thought to be an adaptive strategy for the perpetrator to enhance their reproductive success
69 (Agrell et al. 1998), provide nutritional benefits, increase access to limited resources,
70 increase reproductive opportunities, or ensure parental care is directed to the perpetrator's
71 own offspring (Ebensperger and Blumstein 2007). Social counter-tactics to infanticide may
72 include forming affiliative relationships with adult males (Smuts 1985; Palombit et al. 1997;
73 Borries et al. 1999; Palombit 2000; Weingrill 2000), cooperative defence (see Van Schaik
74 and Janson 2000; Pradhan and van Schaik 2008; Teichroeb et al. 2012), changes in group
75 sex-ratio (see Van Schaik and Kappeler 1997; Van Schaik and Janson 2000; Pradhan and van
76 Schaik 2008; Teichroeb et al. 2012), or multiple male mating to confuse paternity (see Van
77 Schaik and Janson 2000; Wolff and Macdonald 2004). It is also of note that female
78 infanticide is also common in rodents (Vom Saal et al. 1995; Yi et al. 1997).

79

80 The threat of infanticide has also been shown to have effects beyond the killing of the
81 offspring in mice. Parkes and Bruce (1961) reported that recently inseminated female mice
82 (*Mus musculus*) terminate their pregnancies following exposure to the urinary scent of an
83 unfamiliar male, and return to oestrus. This phenomenon, now known as the 'Bruce effect',
84 has been experimentally confirmed in numerous laboratory studies in house mice (Bruce
85 1960; Chipman and Fox 1966) and across a number of murine and microtine rodent species
86 (Clulow and Clarke 1968; Clulow and Langford 1971; Mallory and Clulow 1977). It has also
87 recently been observed in the wild in the primate *Theropithecus gelada* (Roberts et al. 2012).

88

89 The Bruce effect only occurs in mice if the female is exposed in early pregnancy, up to 4-5
90 days post mating (Bruce 1961), before embryo implantation. We have shown that if female
91 mice are exposed to an unfamiliar male in late pregnancy (approximately day 14), offspring
92 are of comparable birth weight, but growth is slower over lactation, resulting in lighter
93 weaning weights (Gale et al. 2013) compared with controls. Following weaning however,
94 the experimental offspring grew faster and caught up to the control offspring by the onset of
95 adulthood. Reduced weaning weight is possibly due to alteration of maternal behaviour
96 during lactation. As unfamiliar males typically kill pups sired by another male (Vom Saal
97 1985), the females may strategically reduce their investment in lactation to prevent wasting it

98 on 'doomed' offspring. If so, females are capable of utilising subtle tactics to optimise their
99 investment in reproduction in relation to their perceived chances of offspring survival. In
100 commensal habitats mice generally live in groups consisting of a number of breeding
101 females, potentially some non-breeding females and subordinate males, and a dominant male
102 who aggressively defends his territory against unfamiliar males (Hurst 1990). As territory
103 turnover does frequently occur (Oakeshott 1974; Bronson 1979), these counteradaptations to
104 the infanticide threat the unfamiliar presents are of great advantage to these females.
105 However, whether reduced offspring growth was a consequence of a strategic alteration of
106 investment by mothers in that study is unknown. Other possible causes of the offspring's
107 reduced weaning weight could include stress triggered by the unfamiliar male (stress has
108 previously been found to result in litters with a reduced weaning weight following lactation
109 (Barlow et al. 1978; Kinsley and Svare 1988; Meek et al. 2000)), or an epigenetic
110 modification in offspring gene expression.

111

112 Accelerating growth requires an increase in metabolic activity that can damage the organism
113 (Morgan et al. 2000). One such cost of accelerated growth, oxidative stress, has been
114 documented in zebra finches (Alonso - Alvarez et al. 2007) and damsel flies (De Block and
115 Stoks 2008). Oxidative stress results from an imbalance between the production of harmful
116 reactive oxygen species (ROS) and an organism's ability to mitigate and detoxify the
117 damaging effects (Monaghan et al. 2009). Failure to moderate this balance can result in
118 oxidative damage to key biological molecules such as DNA, proteins and lipids (Monaghan
119 et al. 2009) and can limit investment in other life history stages (Costantini 2008).

120

121 Modifications in maternal investment have been found to have further effects other than just
122 a reduced weaning weight. Mouse pups show great developmental plasticity, with various
123 aspects of their early environment affecting their life-history trajectories and having lasting
124 effects on adult phenotypes. Through the influence of maternal effects, mothers may alter the
125 phenotype of their offspring and affect their physiological, sexual and behavioural responses
126 as adults (Rossiter 1996; Sheldon and West 2004).

127

128 The aims of this investigation were to test if the degree of maternal investment during
129 lactation is affected by the threat of infanticide, and whether any such effects on maternal
130 investment impose further costs to the offspring later in life. We predicted that:

- 131 1) When females are faced with the threat of infanticide they will strategically alter
132 investment in their offspring by changing their behaviour during lactation.
- 133 2) Offspring that accelerate their growth rate following a reduced weaning weight will suffer
134 oxidative costs as a consequence of the rapid growth.
- 135 3) Offspring that have a reduced weaning weight will suffer costs to components of
136 reproductive effort such as scent-marking rates and composition for males and future
137 reproduction for females.

138

139 **Material and methods**

140 This study aimed to further investigate the cause of the reduced weaning weight we
141 previously observed in offspring whose mothers had been exposed to a novel male and also
142 test for further effects to the offspring's fitness. To do this we replicated the experimental
143 manipulation of our previous experiment (Gale et al. 2013) exposing females to either an
144 unfamiliar male or the paternal male late in their pregnancy on three occasions over the last
145 eight days of gestation. Building on from this previous experiment we added additional
146 measures to attempt to elucidate the cause of the reduced weaning weight observed in the
147 offspring and attempt to detect any costs of the altered growth rate. We added a third
148 unmated control treatment, in which females were housed with another female instead of a
149 male. The unmated treatment acted as a control for female weight and our marker of stress
150 (glucocorticoid metabolites). We then conducted scan sampling during lactation where we
151 examined female nursing behaviour, collected fecal samples to measure female stress, and
152 measured offspring oxidative stress levels at adulthood to test for oxidative damage
153 associated with compensatory growth. We also tested aspects of reproductive fitness in the
154 offspring including scent-marking rates and major urinary proteins in the males and we
155 mated the female offspring to assess their pup birth and growth rates.

156

157 **Animal Housing**

158 Experimental mice were all sixth-generation captive-bred house mice (*Mus musculus*)
159 originally derived from a population of wild mice acquired from a chicken farm in the
160 Northwest of Sydney, Australia (as in Gale et al., 2013). Females were weaned at 28 days of
161 age and were housed with their female siblings until the beginning of the experiment. Males
162 were also weaned at 28 days of age but were housed individually. Mice were maintained on
163 a 12:12 hour reversed light cycle. A dim red light was used for experimental procedures,

164 which were all undertaken in the dark phase as mice are nocturnal. Each mouse was housed
165 in a 315 x 180 x 125mm cage lined with corncob bedding and provided with tissues and
166 shredded newspaper for bedding. Vella Rat and Mouse Pellets and water were provided *ad*
167 *libitum*.

168

169 **Experimental protocol**

170 To investigate the consequences of a novel male's presence to pregnant females (post-
171 implantation of the embryo) we compared two controls (mated and unmated) with the
172 experimental 'novel male' treatment. The unmated control was exposed only to the presence
173 and odour of another unfamiliar female. The mated control or 'familiar male' treatment
174 females were mated and then exposed to the scent and presence of the same mate. Our third
175 treatment was the novel male treatment that was experimentally manipulated so that late
176 gestation the mated females were exposed to the scent and presence of a different male who
177 was unrelated to the mate. Because exposure to this new male was late in pregnancy, well
178 beyond the time frame when a Bruce effect is known to occur in mice, females carried
179 pregnancies through to parturition, therefore allowing us to examine whether females alter
180 their reproductive investment in their offspring and whether this has any long-lasting effects
181 on the offspring (as previously used in Gale et al. (2013)).

182

183 To begin each treatment replicate, a male (or another female for the unmated control)
184 unrelated to the focal female (all between 80-120 days of age) was placed into the focal
185 females cage for a two-week mating period. Males were subsequently removed, and the
186 females were transferred to a clean cage. Female weights were recorded every three days, a
187 weight gain of 3g indicated pregnancy (mean days \pm s.e. from reaching the 3g weight
188 threshold to giving birth for females exposed to the male they mated with: 6.9 ± 1.7 ; females
189 exposed to an unfamiliar male: 7.1 ± 1.9). As the gestation period of *Mus musculus* is 21
190 days (Jones et al., 2009), when the exposures took place the females were close to parturition
191 and the embryos had undergone substantial development. When the 3g-weight gain threshold
192 was reached, females were sequentially allocated into a treatment, to either exposure from
193 the paternal male (n=25) or a novel male (n=24). The unmated control females had already
194 been assigned to be exposed to another female (n=25). As the unmated control females did
195 not gain weight concurrent with pregnancy, one unmated control female was randomly
196 assigned to begin their exposure each day that a female from a different treatment hit the
197 weight gain threshold

198

199 The mice (males or females) added to the females' cages were separated by a metal divider
200 with nine small holes (5 mm diameter) which only allowed limited contact. The exposures
201 took place for three hours a day for three consecutive days. On each day a handful of the
202 respective male or female's used bedding (approximately 15-20g) was placed into the
203 female's cage. Following exposure, the female (prior to parturition) was placed into a clean
204 cage. There were no significant differences between the treatments in the number of days
205 between the removal of the breeding male and the beginning of the male exposure (mean
206 days \pm s.e. for females exposed to the male they mated with: 2.16 ± 1.1 ; an unfamiliar male:
207 2.08 ± 1.2). There was also no difference in the time between the exposure and the
208 subsequent birth (mean days \pm s.e. for females exposed to the male they mated with: $3.2 \pm$
209 1.9 ; an unfamiliar male: 3.7 ± 1.5).

210

211 To test for further costs associated with reduced investment during lactation, offspring were
212 tested for either effects to their reproductive fitness or for oxidative damage. At 13 weeks of
213 age half of the male and female offspring from both treatments were randomly assigned to a
214 reproductive fitness or oxidative stress group. The females in the reproductive fitness group
215 were mated to an unrelated male and the size, weight and pup mortality was recorded. The
216 males in the reproductive fitness group were tested for scent-marking rates and concentration
217 of major urinary proteins.

218

219 **Maternal Investment and offspring weights**

220 The body weights of the females were recorded every three days throughout the experiment.
221 Offspring were weighed at birth and every three days during lactation (to the nearest 0.1g) as
222 a quantification of maternal investment (Ortiz et al. 1984; Ross 1988; Pontier et al. 1989).
223 The offspring were weighed collectively as a litter until weaning (4 weeks old), after which
224 they were separated from their dam, and an individual weight was recorded. Another
225 individual weight was collected for each of the offspring at adulthood (13 weeks old).

226

227 **Maternal behaviour**

228 Maternal behaviour was evaluated by observing each of the females using a scan –sampling
229 technique twice a day, every day for the duration of their lactation (four weeks).
230 Observations were conducted during the dark phase using only a head torch with red light.

231 The first observation occurred in the second or third hour after the change from the light to
232 dark period and the second between four hours and one hour before the change from the dark
233 to light period. Each of the females was randomly assigned an observation order at the
234 beginning of each observation period and observations begun ten minutes after the red light
235 was turned on to allow them to habituate.

236
237 A single observer sequentially recorded each female's behaviour for a total of eight
238 observations with a five minute gap between each of the eight recordings. As this was done
239 morning and evening each day it totaled 16 scans a day. Behaviours recorded included; in
240 nest, nursing, grooming, licking pups, eating/ drinking, nest building, resting or active (e.g
241 see (Benus and Rondigs 1996; Koteja et al. 1999; Palanza et al. 2002). The unmated control
242 females were not included in any of the scan sampling of maternal behaviour.

243

244 **Glucocorticoid metabolites**

245 To determine whether the novel male elicits a stress response from the female, fecal samples
246 were obtained from the females at three points through the experiment and tested for
247 glucocorticoids metabolites (using methods as described by Palme and Möstl (1997).

248 Following a stressful event glucocorticoids are released into circulation (Sapolsky et al.
249 2000) and are hence used as an indicator of the stress response. The first sample was
250 taken at the beginning of the experiment (three weeks prior to being paired for mating), the
251 second on the second day of male exposure, and the third on day seven of lactation. The
252 unmated females were randomly allocated for fecal sample collection in order to correspond
253 with the timings of the females in the other treatments.

254 For collection of fecal samples, mice were placed in a large empty cage (565 x 387 x
255 203mm) made from H.D. polyethylene with a wire roof, for a maximum of 45 minutes. The
256 cages were placed topside down over another corresponding cage and fecal samples were
257 collected from the bottom cage. Immediately after collection, fecal samples were frozen at -
258 80 °C.

259

260 Fecal samples were homogenized and an aliquot of typically 0.05g faeces (Palme et al. 2013)
261 was extracted with 1ml of 80% methanol for 30min on a vortex. When there was insufficient
262 sample the protocol was adjusted accordingly (e.g., 0.25 g faeces in 2.5 mL methanol).

263 Samples were placed in a spinner overnight and then the supernatant was diluted (1:1000)
264 with assay buffer (Trizma, pH 7.5). Samples were then analysed in a double-antibody 5a-

265 pregnane- 3b,11b,21-triol-20-one enzyme immunoassay (EIA) which has been validated for
266 use in mice to assess concentration of glucocorticoid metabolites as described by (Touma et
267 al. 2003; Touma et al. 2004).

268

269 **Oxidative Stress**

270 At 13 weeks of age, half of the female and male offspring from the novel male and familiar
271 male treatment groups were culled humanely by cervical dislocation, and the liver, kidney,
272 heart and gastrocnemius muscle were quickly removed, snap-frozen in liquid nitrogen and
273 stored at -80°C . To assess oxidative stress in the mice two biomarkers of oxidation including
274 protein thiol content and aconitase enzyme activity were analysed in each of the tissues
275 (Gibson et al. 2015). Both of these biomarkers correlate negatively with oxidative stress.

276

277 Protein thiol content was measured by methods described by (Di Monte et al. 1984) but
278 modified for use on a 96 well plate reader (Vasilaki et al. 2006). Protein thiols are essential
279 for stability of and optimum function of proteins, but are highly susceptible to oxidation
280 (Halliwell and Gutteridge 1999), and therefore good markers of oxidative stress.

281

282 Aconitase is an enzyme of the tricarboxylic acid cycle that is highly susceptible to
283 deactivation by radical oxygen species (specifically superoxide) and therefore used as a
284 marker to indicate levels of reactive oxygen species and concomitantly oxidative stress
285 (Hausladen and Fridovich 1994; Gardner et al. 1995; Hausladen and Fridovich 1996;
286 Gardner 1997). As aconitase is located in part in the mitochondria (Wiegand and Remington
287 1986; Gardner et al. 1995), mitochondrial density was also assessed using citrate synthase
288 activity. Citrate synthase is an enzyme commonly used as an indicator of the content of intact
289 mitochondria (Holloszy et al. 1970) and was measured in homogenates according to
290 (Pichaud et al. 2008).

291

292 **Offspring reproductive fitness**

293 Dominant adults are known to scent mark more regularly than subordinates (Drickamer
294 1995) to communicate their territorial and sexual status (Bronson 1979; Hurst et al. 2001)
295 and competitive ability (Rich and Hurst 1998). These chemical scent marks are of high
296 importance to male fitness as they directly influence the attractiveness of a male to a female
297 (Rich and Hurst 1998). The main involatile scent component of male mouse urine is major

298 urinary proteins (MUPs) that bind volatile components of the urine and slowly release them
299 from the scent marks (Hurst et al. 1998). Scent marks may not prevent intruders invading the
300 territory, but they do allow males to use long-lasting signals of identity and dominance over
301 a territory to alert competitors and potential mates (Hurst et al. 1998; Hurst and Beynon
302 2004). Therefore, both the rates and the composition of scent-marks can substantially
303 influence male reproductive fitness.

304

305 Scent-marking rates

306 Scent-marking rates of all of the individually housed male offspring were assessed at 13
307 weeks of age. Scent-marking rates were measured by placing the individual males into an
308 empty (315 x 180 x 125mm) cage lined with Benchkote for one hour a day, for three
309 consecutive days. The scent marks were measured by the number of spatially separate marks
310 observed under UV light and the average number of marks for each male over the three trials
311 was used for analysis.

312

313 Major Urinary Protein Concentration

314 A urine sample was collected from each of the male offspring at 13 weeks of age. Males
315 were confined in a large empty cage (565 x 387 x 203mm) made from H.D. polyethylene
316 with a wire roof, for maximum of 180 minutes. The cages were placed topside down over
317 another corresponding cage to allow the mouse urine to pool in the bottom cage. Urine was
318 then pipetted into an Eppendorf tube and frozen at -20 °C. The concentrations of major
319 urinary proteins were established using Coomassie plus® protein assay reagent kit from
320 Perbio Science UK Ltd (Cramlington, Northumberland, UK) as described by Cheetham et al.
321 (2009). We also measured urinary creatinine (Beynon and Hurst 2004) using the method of
322 Cheetham et al. (2009) to correct for the urinary dilution.

323

324 **Statistics**

325 All statistical analyses were performed using SPSS software package version 2.1 (IBM Corp,
326 Armonk, NY, USA). The analyses were done with dam ID and experimental block fitted as
327 random factors to account for non-independence of individuals originating from the same
328 litter and time differences of each group of experimental mice unless otherwise described.
329 For the oxidative stress data we also fitted plate number as a random factor to control for

330 between plate variability. Scent marking frequency was transformed to $\log(x+1)$ to account
331 for measures of zero deposits and normalise the data. Significance was determined at $p \leq 0.05$.

332 **Results**

333 **Offspring weight**

334 In the novel male treatment, the females gave birth to pups that were of similar weight and
335 litter size as the familiar male treatment females (i.e. exposed to the paternal male). Out of
336 25 females mated in the familiar male treatment, 23 mothers gave birth within the time frame
337 with a mean litter size of 4.217 ± 0.77 and out of 24 mothers in the novel male treatment, 21
338 gave birth within the timeframe with a mean litter size of 4.14 ± 0.89 . The mean birth weight
339 of novel male treatment pups (1.619 ± 0.432 g) was not significantly different from that of the
340 familiar male treatment (1.628 ± 0.433 g; ANOVA: weight: $F_{1,182} = 0.022$, $p = 0.882$; litter
341 size: $F_{1,42} = 0.070$, $p = 0.792$). Two females in the control treatment and one female in the
342 experimental treatment group destroyed their litters within three days of giving birth. Some
343 pup mortality of unknown cause was observed over lactation; however, this is common in
344 captive breeding mice (Weber et al. 2013). The pup mortality was not significantly different
345 between the treatments (ANOVA: $F_{1,7} = 0.778$, $p = 0.407$) with six mothers of the novel male
346 treatment committing maternal infanticide (total 10 pups lost) and three mothers of the
347 familiar male (total seven pups lost).

348

349 Although the novel male treatment females gave birth to pups of a similar weight and litter
350 size as the familiar male treatment females, novel male treatment litters grew more slowly
351 over lactation. To analyse the differences in offspring growth rate we used repeated measures
352 ANOVA reporting within-subjects effects. Mauchly's Test of Sphericity indicated that the
353 assumption of sphericity had been violated ($\chi^2(2) = 34.210$, $p = <0.001$), therefore degrees of
354 freedom were corrected using Greenhouse-Geisser adjusted degrees of freedom. There was a
355 difference in growth rate between the treatments but there was no effect of sex (treatment:
356 $F_{1.7,310} = 7.49$, $p = 0.001$; interaction between treatment and time: $F_{1.7,310} = 5668$, $p = <0.001$;
357 sex x treatment: $F_{1.6,306} = 1.3$, $p = 0.270$). This resulted in novel male treatment pups being
358 significantly smaller (7.19 ± 0.61 g), at weaning (4 weeks old) (ANOVA: $F_{1,182} = 27.11$,
359 $p = <0.001$) than the familiar male treatment pups (7.99 ± 1.48 g) (see Figure 1). This difference
360 did not persist into adulthood however, as both males and females of the novel male
361 treatment exhibited catch-up growth following weaning and there were no weight differences
362 at adulthood (13 weeks old) (ANOVA: $F_{1,184} = 0.760$, $p = 0.384$) (see Figure 2). There were

363 also no significant interactions of treatment and sex in weight at weaning (sex x treatment
364 $F_{1,180} = 2.14, p=0.145$) or at adulthood (sex x treatment $F_{1,180}=0.044, p=0.868$)).

365

366

Insert Figure 1 and 2

367

368 **Maternal behaviour**

369 There was no significant difference in the amount of time the mothers spent in nest,
370 grooming, licking pups, eating/ drinking, nest building, resting or active (Table 1). The only
371 behaviour showing significant differences between treatments was nursing, with females
372 exposed to familiar males spent almost twice the time nursing (Table 1).

373

374

Insert Table 1

375

376 **Glucocorticoid metabolites in mothers**

377 Three fecal samples were taken from each female: The first sample was taken before the
378 experiment began to determine the normal level of fecal glucocorticoid metabolites (FGMs)
379 for each female. The second collection was taken on day two of the exposures to a male and
380 the third coincided with mid lactation in the mated treatments. Some of the females did not
381 provide a sample within the timeframe they were in the collection cages (45min) so the
382 sample sizes for the glucocorticoid metabolites are novel male $n=20$, familiar male $n=21$ and
383 unmated control $n=18$. Using a repeated measures ANOVA we found that FGM's did
384 change significantly over time (effect of time: $F_{2,112} = 131, p < 0.001$; interaction between
385 time and housing companion: $F_{4,112} = 29.5, p < 0.001$). In the novel male and familiar male
386 treatments the FGM's were higher in the second collection and dropped back down closer to
387 their normal levels of FGM's at the third collection. There was no difference in the FGMs in
388 the females before the experiment began (ANOVA: $F_{2,56}=0.094, p=0.911$), but there was
389 following exposure to their partners (ANOVA: $F_{2,56}=43.72, p < 0.001$) and at the end of the
390 experiment which coincided with mid lactation for the reproducing treatments ($F_{2,56}=4.6,$
391 $p=0.014$)(See Figure 3). After removing the unmated control from the analysis we found that
392 the effect of time was still significant ($F_{2,78} = 143, p < 0.001$) but the interaction between
393 time and housing companion was not ($F_{1,78}=1.89, p=0.158$). While there was no difference
394 between the FGMs between the novel male treatment and the familiar male treatment, there
395 was a difference between the mating treatments and females who were not mated. Females
396 from both mated treatments experienced a rise in FGM's levels in the second collection
397 following exposures to a male, however, by the third collection at mid lactation their FGM's

398 had returned closer to their normal level (collection one). The unmated control females FGM
399 levels did not show any pronounced variability over the three collections.

400

401

Insert Figure 3

402

403 **Oxidative stress in offspring**

404 Both markers of oxidative stress showed consistent evidence of oxidative damage with the
405 novel male treatment (n=47) in two (liver and heart) out of four organs tested having lower
406 protein thiol concentrations and aconitase enzyme activity (indicating higher levels of
407 oxidative damage) than the familiar male treatment (n=38). There was an effect of sex with
408 novel male treatment offspring males having lower protein thiol concentrations than those of
409 the familiar male treatment males in the heart but there were no other sex differences in any
410 of the analyses for oxidative damage (see Table 2 and Figure 4 and 5).

411

412 The same trend was observed when testing the aconitase enzyme activity to citrate synthase
413 activity (mitochondrial density) ratio. The liver and heart in the novel male treatment were
414 also found to have a significantly lower aconitase enzyme activity to citrate synthase activity
415 ratio indicating higher levels of reactive oxygen species (ROS) production in these organs.
416 This decrease in aconitase enzyme activity was not observed in either the kidneys or the
417 gastrocnemius and there were no effects of sex (See Table 2 and Figure 4 and 5).

418

419

Insert Table 2 and Figure 4

420

421 **Male and female offspring fitness components**

422 For the male offspring fitness components sample sizes were both n=17 for the treatments.
423 There was no difference in the concentration of major urinary proteins among the male
424 offspring produced by the novel male treatment females compared to the control offspring
425 (ANOVA: $F_{1,30} = 0.76$, $p = 0.543$). The mean protein (mg/mg creatinine) \pm s.e for the novel
426 male treatment was 18.44 ± 3.60 , and the familiar male treatment males were 20.73 ± 2.74 .
427 Experimental block also did not have an effect (ANOVA: $F_{1,30} = 5.8$, $p = 0.249$). The mean \pm
428 s.e scent-marking frequency for the novel male treatment offspring was 28.28 ± 10.54 and the
429 familiar male treatment males was 36.96 ± 10.49 . A repeated measures ANOVA found that
430 there was no difference in the frequency of scent-marks between the treatments ($F_{2,64} = 3.00$,
431 $p = 0.57$). For the female offspring produced that were bred at adulthood there was no
432 difference in the pup weights that the female offspring gave birth to (ANOVA: $F_{1,40} = 0.35$,

433 $p=0.556$) or of their pups weaning weights at the end of lactation ($F_{1,40}=1.002$, $p=0.323$) and
434 a repeated measures ANOVA showed there was no difference in growth rate between the
435 treatments (Sex x Treatment $F_{1,41}=1.29$ $p=0.262$). The mean birth weights between the
436 treatments were 1.26 ± 0.04 g and 1.355 ± 0.13 g for the novel male and familiar male
437 treatment females respectively and the mean weaning weights were 12.75 ± 2.09 g and $12.27 \pm$
438 0.13 g respectively. Experimental block also had no effect (ANOVA: $F_{1,40}=0.69$, $p=0.837$).

439

440 **Discussion**

441 Our results support the prediction that the 'Late Bruce Effect', in which female mice exposed
442 to a novel male late in pregnancy weaned at lower weights (Gale et al. 2013), may be due to
443 a strategic reduction in maternal investment in lactation. Notably, females exposed to a novel
444 male's scent spent less time nursing pups than females exposed to the scent of their mate.
445 Our results support the prediction that females adjust post-partum investment in relation to
446 pups' perceived chances of survival, potentially benefitting maternal lifetime fitness at the
447 expense of the current litter of offspring (Marshall and Uller 2007).

448

449 This finding suggests that patterns of post-partum investment can be adjusted by mothers in
450 relation to the risk of infanticide, in ways that mirror the strategic, spontaneous abortion of
451 pups under the Bruce effect (Bruce 1960, 1961; Hrdy 1979; Schwagmeyer 1979; Storey
452 1986), as well as evidence that a mechanism to abort fetuses strategically at the later stages
453 of pregnancy has evolved in other species including the gelada (Roberts et al. 2012) and
454 prairie vole (Clulow and Clarke 1968; Clulow and Langford 1971). As mice are only able to
455 abort in early pregnancy, altering their investment post-partum provides an opportunity to
456 respond to the threat of infanticide.

457

458 Other studies have shown that female mice strategically moderate investment during
459 lactation and that they may alter investment relative to the reproductive value of their
460 offspring. Mashoodh et al., (2012) found that females increased their investment during
461 lactation when mated with a male housed in lifelong socially enriched conditions compared
462 with females mated with males housed in impoverished conditions. This suggests that
463 females may invest more in offspring on the basis of paternal condition. König and Markl
464 (1987) showed that despite frequent suckling attempts, offspring do not get more milk than
465 corresponds to the maternal optimum during lactation, suggesting mothers have control over
466 reproductive allocation during lactation.

467

468 Another potential cause of the reduced weaning weight could be a stress response. In
469 rodents, maternal stress has been previously found to cause depressed growth in offspring
470 (Barlow et al. 1978; Kinsley and Svare 1988; Meek et al. 2000). Previous research on the
471 effects of prenatal stress on offspring is frequently contradictory. The nature, timing and
472 length of the stress inflicted varies from study to study, as do the results. One study, for
473 example that used crowding as a stressor found no difference in time spent nursing between
474 stressed and non-stressed dams (Moore and Power 1986), while another study that used
475 novelty stress found that stressed dams spent significantly more time nursing compared to
476 non-stressed dams (Muir et al. 1985). It is also important to note there is evidence that
477 maternal stress leads to faster (Dantzer et al. 2013) and increased growth of offspring at birth
478 and weaning (Szuran et al. 1991).

479

480 We found while females all had similar stress levels before the experiment began, females
481 that were mated and then exposed to either a novel or their mated male had significantly
482 higher FGMs than the females that were not mated in the experiment. Females from the
483 mated treatments both displayed a rise in their FGMs following the exposures, suggesting
484 that the females were equally stressed by encountering the paternal male as they were a
485 novel male. Higher levels of FGM could also result from exposure to a male, as our
486 experiment used another female as an unmated control we are unable to distinguish this
487 result properly. Only one other study we could find used a novel male conspecific as a
488 source of stress. Lerch et al. (2016) stressed pregnant or lactating females using unfamiliar
489 male faeces and examined maternal and offspring behaviour to investigate whether early
490 adverse experiences elevate the risk of developing psychiatric disorders. However, unlike
491 our study that used a familiar male as a control, they compared pregnant or lactating females
492 exposed to unfamiliar male faeces with a control group that didn't receive any faeces. The
493 scarcity of research that uses another conspecific as a stressor is surprising as this could be
494 expected to be a more environmentally relevant challenge than other manipulations like
495 forced immobilization challenges, for example. While the over production of maternal
496 glucocorticoids can be harmful (Korgun et al. 2012) levels are known to increase during
497 pregnancy in mice (Barlow et al. 1974; Dalle et al. 1978) as they are essential for fetal
498 development (Korgun et al. 2012). Post -partum, glucocorticoids also have important roles in
499 milk secretion and lactogenesis (Chida et al. 2011). The higher FGM levels that we found in

500 females from both of the mated, compared to the unmated treatment, may, therefore, be just
501 a normal consequence of pregnancy.

502

503 As in our previous study (Gale et al. 2013), the offspring of the females that were exposed to
504 the novel male were smaller at the end of lactation, but they caught up in size by maturity.
505 The catch-up growth exhibited by the offspring coincides with the time after lactation when
506 offspring begin to feed themselves on solid food. Offspring can acquire the resources to
507 accelerate their growth rate themselves. For compensation or catch-up growth to occur, the
508 benefits must outweigh the costs of not accelerating growth. While there may be a positive
509 association between size and fitness (Roff, 1992) and accelerated growth may increase
510 overall reproductive success, many studies have found that compensatory growth inflicts
511 various costs over different time-scales (reviewed in Metcalfe and Monaghan (2001)). Costs
512 that have been documented in rats (*Rattus norvegicus*) include deficiencies in protein
513 maintenance (Samuels and Baracos 1995), telomere abrasion rate (Jennings et al. 1999)
514 insulin regulation (Ozanne and Hales 1999), adult obesity (Waterland and Garza 1999), and,
515 perhaps most importantly, reduced lifespan (Jennings et al. 1999; Rollo 2002). While
516 compensatory growth may reduce lifespan it could still be adaptive if it increases overall
517 reproductive success (Metcalfe and Monaghan 2001) or short-term survival chances (Arendt
518 1997). In mice, reproductive allocation in adulthood is influenced by size, and so
519 compensatory growth may allow individuals to attain a normal reproductive rate in
520 adulthood, at least early in life. Female offspring that exhibited catch-up growth produced
521 litters of the same size and weight as those female offspring from the control group; male
522 offspring showed equivalent scent marking abilities in our assays. Thus, catch up growth
523 seems to allow offspring to attain a similar reproductive output early in life that is
524 comparable with the steadier growth in the familiar male group, although costs may be paid
525 for this in terms of late life reproduction or lifespan.

526

527 The existence of compensatory and catch up growth shows that organisms do not grow at
528 their maximal rate, but rather at a rate influenced by, and potentially optimized to, their
529 circumstances. Hector and Nakagawa (2012) distinguish these two terms by defining
530 compensatory growth as a faster than usual growth rate and catch-up growth an attainment of
531 control size. Mangel & Munch (2005) propose that growth leads to an accumulation of
532 damage at the cellular level that is expressed at the level of the organism and is an important
533 cost of compensatory growth. We tested for damage on a cellular level in form of oxidative

534 damage and oxidative stress. We found evidence for oxidative damage in the livers and
535 hearts of offspring from the novel male exposure treatment, highlighting that an olfactory
536 change in a pregnant mother's environment can elicit a variety of maternal and offspring
537 responses, ultimately influencing offspring physiological condition in adulthood. This may
538 be a consequence of catch-growth, but could also be a consequence of odour exposure itself,
539 or changes in maternal allocation in response to this. In future studies it may be of interest to
540 limit an offspring's ability to show compensatory growth (through a nutritional or genetic
541 manipulation) and test whether oxidative damage in offspring still occurs in adulthood.

542

543 Wild mice are highly territorial and turnover of the dominant male is a common occurrence
544 in wild populations (Oakeshott 1974; Bronson 1979). For the females in the territory that
545 means that they will be exposed to novel males which present the threat of infanticide
546 (Ebensperger 1998). Our experiment was designed with the rationale to mimic this turnover
547 by exposing the females to a novel male to see the effects it would have on reproductive
548 allocation and offspring fitness. Our results suggest that the very prospect of male territorial
549 turnover can have physiological consequences for the offspring and potentially alter their life
550 history. We also suggest that females are capable of strategically modulating their
551 investment relative to their current contextual conditions, which may offer significant fitness
552 benefits in the wild where offspring survival is much more variable and infanticide is a
553 common threat.

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568 **Acknowledgements**

569 We would like to thank Heather Try for all of her help with animal maintenance and Rebecca
570 Hobbs at the Wildlife Reproductive Centre (WRC) at Taronga Western Plains Zoo for
571 providing training on how to conduct an assay (using methods as described by Palme &
572 Möstl, 1997) for glucocorticoids metabolites in feces.

573

574 **Competing interests**

575 There are no competing interests to declare.

576

577 **Author contributions**

578 TG, MG and RB conceived the ideas and designed methodology;

579 TG collected the data;

580 TG, MG and RB analysed the data;

581 TG led the writing of the manuscript.

582 All authors, TG, MG and RB contributed critically to the drafts and gave final approval for
583 publication.

584

585 **Funding**

586 This study was funded by an Australian Research Council (ARC) Discovery Grant and
587 Fellowship [grant number: DP150100676] awarded to R.C. Brooks.

588 **Data accessibility**

589 We have archived our data on Dryad. A data set for this experiment is available from the
590 Dryad Digital Repository: doi:10.5061/dryad.762k24f

591

592

593

594 **References**

595

596 Agrell, J., J. O. Wolff, and H. Ylönen. 1998. Counter-strategies to infanticide in mammals:
597 costs and consequences. *Oikos*:507-517.

598 Alonso - Alvarez, C., S. Bertrand, B. Faivre, and G. Sorci. 2007. Increased susceptibility
599 to oxidative damage as a cost of accelerated somatic growth in zebra finches.
600 *Functional Ecology* 21:873-879.

601 Arendt, J. D. 1997. Adaptive intrinsic growth rates: an integration across taxa. *The*
602 *quarterly review of biology* 72:149-177.

603 Barlow, S., A. Knight, and F. Sullivan. 1978. Delay in postnatal growth and development
604 of offspring produced by maternal restraint stress during pregnancy in the rat.
605 *Teratology* 18:211-218.

606 Barlow, S. M., P. Morrison, and F. Sullivan. 1974. Plasma corticosterone levels during
607 pregnancy in the mouse: the relative contributions of the adrenal glands and
608 foeto-placental units. *Journal of Endocrinology* 60:473-483.

609 Benus, R. and M. Rondigs. 1996. Patterns of maternal effort in mouse lines
610 bidirectionally selected for aggression. *Animal Behaviour* 51:67-75.

611 Beynon, R. J. and J. L. Hurst. 2004. Urinary proteins and the modulation of chemical
612 scents in mice and rats. *Peptides* 25:1553-1563.

613 Borries, C., K. Launhardt, C. Epplen, J. T. Epplen, and P. Winkler. 1999. Males as infant
614 protectors in Hanuman langurs (*Presbytis entellus*) living in multimale groups—
615 defence pattern, paternity and sexual behaviour. *Behavioral Ecology and*
616 *Sociobiology* 46:350-356.

617 Bronson, F. 1979. The reproductive ecology of the house mouse. *The Quarterly review*
618 *of biology* 54:265-299.

619 Bronson, F. H. 1989. *Mammalian reproductive biology*. University of Chicago Press.

620 Bruce, H. M. 1960. A block to pregnancy in the mouse caused by proximity of strange
621 males. *Journal of Reproduction and Fertility* 1:96-103.

622 Bruce, H. M. 1961. Time relations in the pregnancy-block induced in mice by strange
623 males. *Journal of Reproduction and Fertility* 2:138-142.

624 Cheetham, S. A., A. L. Smith, S. D. Armstrong, R. J. Beynon, and J. L. Hurst. 2009. Limited
625 variation in the major urinary proteins of laboratory mice. *Physiology &*
626 *Behavior* 96:253-261.

627 Chida, D., K. Miyoshi, T. Sato, T. Yoda, T. Kikusui, and Y. Iwakura. 2011. The Role of
628 Glucocorticoids in Pregnancy, Parturition, Lactation, and Nurturing in
629 Melanocortin Receptor 2-Deficient Mice. *Endocrinology* 152:1652-1660.

630 Chipman, R. and K. Fox. 1966. Oestrous synchronization and pregnancy blocking in wild
631 house mice (*Mus musculus*). *Journal of reproduction and fertility* 12:233-236.

- 632 Clulow, F. and J. Clarke. 1968. Pregnancy-block in *Microtus agrestis* and induced
633 ovulator.
- 634 Clulow, F. and P. Langford. 1971. Pregnancy-block in the meadow vole, *Microtus*
635 *pennsylvanicus*. *Journal of reproduction and fertility* 24:275-277.
- 636 Clutton-Brock, T. and A. C. Vincent. 1991. Sexual selection and the potential
637 reproductive rates of males and females. *Nature* 351:58-60.
- 638 Costantini, D. 2008. Oxidative stress in ecology and evolution: lessons from avian
639 studies. *Ecology letters* 11:1238-1251.
- 640 Dalle, M., J. Giry, M. Gay, and P. Delost. 1978. Perinatal changes in plasma and adrenal
641 corticosterone and aldosterone concentrations in the mouse. *Journal of*
642 *Endocrinology* 76:303-309.
- 643 Dantzer, B., A. E. Newman, R. Boonstra, R. Palme, S. Boutin, M. M. Humphries, and A. G.
644 McAdam. 2013. Density triggers maternal hormones that increase adaptive
645 offspring growth in a wild mammal. *Science* 340:1215-1217.
- 646 De Block, M. and R. Stoks. 2008. Compensatory growth and oxidative stress in a
647 damselfly. *Proceedings of the Royal Society B: Biological Sciences* 275:781-785.
- 648 Di Monte, D., D. Ross, G. Bellomo, L. Eklöw, and S. Orrenius. 1984. Alterations in
649 intracellular thiol homeostasis during the metabolism of menadione by isolated
650 rat hepatocytes. *Archives of Biochemistry and Biophysics* 235:334-342.
- 651 Drickamer, L. C. 1995. Rates of urine excretion by house mouse (*Mus domesticus*):
652 differences by age, sex, social status, and reproductive condition. *Journal of*
653 *chemical ecology* 21:1481-1493.
- 654 Ebensperger, L. A. 1998. Do female rodents use promiscuity to prevent male
655 infanticide? *Ethology Ecology & Evolution* 10:129-141.
- 656 Ebensperger, L. A. and D. T. Blumstein. 2007. Nonparental infanticide. *Rodent societies:*
657 *an ecological and evolutionary perspective*:267-279.
- 658 Gale, T., A. B. Gibson, R. C. Brooks, and M. Garratt. 2013. Exposure to a novel male
659 during late pregnancy influences subsequent growth of offspring during
660 lactation. *Journal of Evolutionary Biology* 26:2057-2062.
- 661 Gardner, P. R. 1997. Superoxide-Driven Aconitase FE-S Center Cycling. *Bioscience*
662 *Reports* 17:33-42.
- 663 Gardner, P. R., I. Raineri, L. B. Epstein, and C. W. White. 1995. Superoxide Radical and
664 Iron Modulate Aconitase Activity in Mammalian Cells. *Journal of Biological*
665 *Chemistry* 270:13399-13405.

666 Gibson, A. B., M. Garratt, and R. C. Brooks. 2015. Experimental evidence that litter size
667 imposes an oxidative challenge to offspring. *The Journal of Experimental Biology*
668 218:3911-3918.

669 Halliwell, B. and J. Gutteridge. 1999. Detection of free radicals and other reactive
670 species: trapping and fingerprinting. Pp. 351-425. *Free radicals in biology and*
671 *medicine*. Oxford University Press Oxford.

672 Hausladen, A. and I. Fridovich. 1994. Superoxide and peroxynitrite inactivate
673 aconitases, but nitric oxide does not. *Journal of Biological Chemistry* 269:29405-
674 29408.

675 Hausladen, A. and I. Fridovich. 1996. [4] Measuring nitric oxide and superoxide: Rate
676 constants for aconitase reactivity. Pp. 37-41. *Methods in Enzymology*. Academic
677 Press.

678 Hector, K. L. and S. Nakagawa. 2012. Quantitative analysis of compensatory and catch -
679 up growth in diverse taxa. *Journal of Animal Ecology* 81:583-593.

680 Holloszy, J., L. Oscai, I. Don, and P. Mole. 1970. Mitochondrial citric acid cycle and
681 related enzymes: adaptive response to exercise. *Biochemical and biophysical*
682 *research communications* 40:1368-1373.

683 Hrdy, S. B. 1979. Infanticide among animals: a review, classification, and examination of
684 the implications for the reproductive strategies of females. *Ethology and*
685 *Sociobiology* 1:13-40.

686 Hurst, J. L. 1990. Urine marking in populations of wild house mice *Mus domesticus*
687 *rutty*. I. Communication between males. *Animal Behaviour* 40:209-222.

688 Hurst, J. L. and R. J. Beynon. 2004. Scent wars: the chemobiology of competitive
689 signalling in mice. *Bioessays* 26:1288-1298.

690 Hurst, J. L., C. E. Payne, C. M. Nevison, A. D. Marie, R. E. Humphries, D. H. Robertson, A.
691 Cavaggioni, and R. J. Beynon. 2001. Individual recognition in mice mediated by
692 major urinary proteins. *Nature* 414:631-634.

693 Hurst, J. L., D. H. Robertson, U. Tolladay, and R. J. Beynon. 1998. Proteins in urine scent
694 marks of male house mice extend the longevity of olfactory signals. *Animal*
695 *behaviour* 55:1289-1297.

696 Jennings, B., S. Ozanne, M. Dorling, and C. Hales. 1999. Early growth determines
697 longevity in male rats and may be related to telomere shortening in the kidney.
698 *FEBS letters* 448:4-8.

- 699 Kinsley, C. and B. Svare. 1988. Prenatal stress alters maternal aggression in mice.
700 *Physiology & behavior* 42:7-13.
- 701 Kirkwood, T. B. and S. N. Austad. 2000. Why do we age? *Nature* 408:233-238.
- 702 König, B. 1985. Die Brutpflegestrategie der Mäusemutter (*Mus musculus*): Eltern-
703 Nachkommen-Konflikt, Toleranz gegen fremde Junge und Reaktion auf
704 Nahrungsmittelmangel.
- 705 König, B. and H. Markl. 1987. Maternal care in house mice. *Behavioral Ecology and*
706 *Sociobiology* 20:1-9.
- 707 König, B., J. Riester, and H. Markl. 1988. Maternal care in house mice (*Mus musculus*): II.
708 The energy cost of lactation as a function of litter size. *Journal of Zoology*
709 216:195-210.
- 710 Korgun, E. T., A. Ozmen, G. Unek, and I. Mendilcioglu. 2012. The effects of
711 glucocorticoids on fetal and placental development. *development* 21:26.
- 712 Koteja, P., T. Garland, J. K. Sax, J. G. Swallow, and P. A. Carter. 1999. Behaviour of house
713 mice artificially selected for high levels of voluntary wheel running. *Animal*
714 *behaviour* 58:1307-1318.
- 715 Lerch, S., C. Dormann, C. Brandwein, P. Gass, and S. Chourbaji. 2016. The scent of stress:
716 environmental challenge in the peripartum environment of mice affects
717 emotional behaviours of the adult offspring in a sex-specific manner. *Laboratory*
718 *Animals* 50:167-178.
- 719 Mallory, F. and F. Clulow. 1977. Evidence of pregnancy failure in the wild meadow vole,
720 *Microtus pennsylvanicus*. *Canadian journal of zoology* 55:1-17.
- 721 Marshall, D. and T. Uller. 2007. When is a maternal effect adaptive? *Oikos* 116:1957-
722 1963.
- 723 Meek, L. R., K. M. Burda, and E. Paster. 2000. Effects of prenatal stress on development
724 in mice: maturation and learning. *Physiology & behavior* 71:543-549.
- 725 Metcalfe, N. B. and P. Monaghan. 2001. Compensation for a bad start: grow now, pay
726 later? *Trends in Ecology & Evolution* 16:254-260.
- 727 Monaghan, P., N. B. Metcalfe, and R. Torres. 2009. Oxidative stress as a mediator of life
728 history trade - offs: mechanisms, measurements and interpretation. *Ecology*
729 *letters* 12:75-92.
- 730 Moore, C. L. and K. L. Power. 1986. Prenatal stress affects mother-infant interaction in
731 Norway rats. *Developmental psychobiology* 19:235-245.

- 732 Morgan, I., I. McCarthy, and N. Metcalfe. 2000. Life - history strategies and protein
733 metabolism in overwintering juvenile Atlantic salmon: growth is enhanced in
734 early migrants through lower protein turnover. *Journal of Fish Biology* 56:637-
735 647.
- 736 Muir, J. L., H. P. Pfister, and A. Ivinskis. 1985. Effects of prepartum stress and
737 postpartum enrichment on mother-infant interaction and offspring problem-
738 solving ability in *Rattus norvegicus*. *J. Comp. Psychol.* 99:468.
- 739 Oakeshott, J. G. 1974. Social dominance, aggressiveness and mating success among male
740 house mice (*Mus musculus*). *Oecologia* 15:143-158.
- 741 Ortiz, C. L., B. J. L. Boeuf, and D. P. Costa. 1984. Milk Intake of Elephant Seal Pups: An
742 Index of Parental Investment. *The American Naturalist* 124:416-422.
- 743 Ozanne, S. E. and C. N. Hales. 1999. The long-term consequences of intra-uterine protein
744 malnutrition for glucose metabolism. *Proceedings of the Nutrition Society*
745 58:615-620.
- 746 Palanza, P. L., K. L. Howdeshell, S. Parmigiani, and F. S. vom Saal. 2002. Exposure to a
747 low dose of bisphenol A during fetal life or in adulthood alters maternal
748 behavior in mice. *Environmental health perspectives* 110:415.
- 749 Palme, R. and E. Möstl. 1997. Measurement of cortisol metabolites in faeces of sheep as
750 a parameter of cortisol concentration in blood. *Zeitschrift fuer Saeugetierkunde*
751 (Germany).
- 752 Palme, R., C. Touma, N. Arias, M. Dominchin, and M. Lepschy. 2013. Steroid extraction:
753 get the best out of faecal samples. *Wien Tierarztl Monatsschr* 100:238-246.
- 754 Palombit, R. 2000. Infanticide and the evolution of male-female bonds in animals.
755 Infanticide by males and its implications. Cambridge University Press,
756 Cambridge:239-268.
- 757 Palombit, R. A., R. M. Seyfarth, and D. L. Cheney. 1997. The adaptive value of
758 'friendships' to female baboons: experimental and observational evidence.
759 *Animal Behaviour* 54:599-614.
- 760 Parkes, A. and H. M. Bruce. 1961. Olfactory stimuli in mammalian reproduction. *Science*
761 134:1049-1054.
- 762 Pichaud, N., J. Pellerin, M. Fournier, S. Gauthier-Clerc, P. Rioux, and É. Pelletier. 2008.
763 Oxidative stress and immunologic responses following a dietary exposure to
764 PAHs in *Mya arenaria*. *Chemistry Central Journal* 2:1.

- 765 Pontier, D., J. M. Gaillard, D. Allaine, J. Trouvilliez, I. Gordon, and P. Duncan. 1989.
766 Postnatal-growth rate and adult body-weight in mammals - a new approach.
767 *Oecologia* 80:390-394.
- 768 Pradhan, G. and C. van Schaik. 2008. Infanticide-driven intersexual conflict over
769 matings in primates and its effects on social organization. *Behaviour* 145:251-
770 275.
- 771 Reznick, D. 1985. Costs of reproduction: an evaluation of the empirical evidence.
772 *Oikos*:257-267.
- 773 Rich, T. J. and J. L. Hurst. 1998. Scent marks as reliable signals of the competitive ability
774 of mates. *Animal Behaviour* 56:727-735.
- 775 Roberts, E. K., A. Lu, T. J. Bergman, and J. C. Beehner. 2012. A Bruce effect in wild
776 geladas. *Science* 335:1222-1225.
- 777 Rollo, C. D. 2002. Growth negatively impacts the life span of mammals. *Evolution &*
778 *development* 4:55-61.
- 779 Ross, C. 1988. The intrinsic rate of natural increase and reproductive effort in primates.
780 *Journal of Zoology* 214:199-219.
- 781 Rossiter, M. 1996. Incidence and consequences of inherited environmental effects.
782 *Annual review of ecology and systematics* 27:451-476.
- 783 Samuels, S. E. and V. E. Baracos. 1995. Tissue protein turnover is altered during catch-
784 up growth following *Escherichia coli* infection in weanling rats. *The Journal of*
785 *nutrition* 125:520.
- 786 Sapolsky, R. M., L. M. Romero, and A. U. Munck. 2000. How do glucocorticoids influence
787 stress responses? Integrating permissive, suppressive, stimulatory, and
788 preparative actions. *Endocrine reviews* 21:55-89.
- 789 Schwagmeyer, P. 1979. The Bruce effect: an evaluation of male/female advantages. *The*
790 *American Naturalist* 114:932-938.
- 791 Sheldon, B. C. and S. A. West. 2004. Maternal dominance, maternal condition, and
792 offspring sex ratio in ungulate mammals. *The American Naturalist* 163:40-54.
- 793 Smuts, B. B. 1985. Sex and friendship in baboons. Transaction Publishers.
- 794 Speakman, J. R. 2008. The physiological costs of reproduction in small mammals.
795 *Philosophical Transactions of the Royal Society B: Biological Sciences* 363:375-
796 398.
- 797 Stearns, S. C. 1992. *The evolution of life histories*. Oxford University Press Oxford.

- 798 Storey, A. E. 1986. Influence of sires on male-induced pregnancy disruptions in meadow
799 voles (*Microtus pennsylvanicus*) differs with stage of pregnancy. *J. Comp.*
800 *Psychol.* 100:15-20.
- 801 Szuran, T., E. Zimmerman, V. Pliska, H. P. Pfister, and H. Welzl. 1991. Prenatal stress
802 effects on exploratory activity and stress-induced analgesia in rats.
803 *Developmental Psychobiology* 24:361-372.
- 804 Teichroeb, J. A. and P. Sicotte. 2008. Infanticide in ursine colobus monkeys (*Colobus*
805 *vellerosus*) in Ghana: new cases and a test of the existing hypotheses. *Behaviour*
806 145:727-755.
- 807 Teichroeb, J. A., E. C. Wikberg, I. Bădescu, L. J. Macdonald, and P. Sicotte. 2012.
808 Infanticide risk and male quality influence optimal group composition for
809 *Colobus vellerosus*. *Behavioral Ecology*:ars128.
- 810 Touma, C., R. Palme, and N. Sachser. 2004. Analyzing corticosterone metabolites in fecal
811 samples of mice: a noninvasive technique to monitor stress hormones.
812 *Hormones and Behavior* 45:10-22.
- 813 Touma, C., N. Sachser, E. Möstl, and R. Palme. 2003. Effects of sex and time of day on
814 metabolism and excretion of corticosterone in urine and feces of mice. *General*
815 *and comparative endocrinology* 130:267-278.
- 816 Van Schaik, C. P. and C. H. Janson. 2000. Social counterstrategies against infanticide by
817 males in primates and other mammals. In: Kappeler PM, editor. *Primate males*,
818 Cambridge University Press 34-54.
- 819 Van Schaik, C. P. and P. M. Kappeler. 1997. Infanticide risk and the evolution of male-
820 female association in primates. *Proceedings of the Royal Society of London B:*
821 *Biological Sciences* 264:1687-1694.
- 822 Vasilaki, A., A. Mansouri, H. Van Remmen, J. H. Van Der Meulen, L. Larkin, A. G.
823 Richardson, A. McArdle, J. A. Faulkner, and M. J. Jackson. 2006. Free radical
824 generation by skeletal muscle of adult and old mice: effect of contractile activity.
825 *Aging Cell* 5:109-117.
- 826 Vom Saal, F. S. 1985. Time-contingent change in infanticide and parental behavior
827 induced by ejaculation in male mice. *Physiology & behavior* 34:7-15.
- 828 Vom Saal, F. S., P. Franks, M. Boechler, P. Palanza, and S. Parmigiani. 1995. Nest defense
829 and survival of offspring in highly aggressive wild Canadian female house mice.
830 *Physiology & Behavior* 58:669-678.

831 Waterland, R. A. and C. Garza. 1999. Potential mechanisms of metabolic imprinting that
 832 lead to chronic disease. *The American Journal of Clinical Nutrition* 69:179-197.

833 Weber, E. M., B. Algers, J. Hultgren, and I. A. S. Olsson. 2013. Pup mortality in laboratory
 834 mice – infanticide or not? *Acta Veterinaria Scandinavica* 55:83.

835 Weingrill, T. 2000. Infanticide and the value of male-female relationships in mountain
 836 chacma baboons. *Behaviour* 137:337-359.

837 Wiegand, G. and S. J. Remington. 1986. Citrate synthase - structure, control, and
 838 mechanism. *Annual Review of Biophysics and Biophysical Chemistry* 15:97-117.

839 Wolff, J. O. and D. W. Macdonald. 2004. Promiscuous females protect their offspring.
 840 *Trends in Ecology & Evolution* 19:127-134.

841 Wolff, J. O. and P. W. Sherman. 2008. *Rodent societies: an ecological and evolutionary*
 842 *perspective*. University of Chicago Press.

843 Yl, xf, H. nen, E. Koskela, and T. Mappes. 1997. Infanticide in the bank vole
 844 (*Clethrionomys glareolus*): occurrence and the effect of familiarity on female
 845 infanticide. *Annales Zoologici Fennici* 34:259-266.

846 Zhao, Q., C. Borries, and W. Pan. 2011. Male takeover, infanticide, and female
 847 countertactics in white-headed leaf monkeys (*Trachypithecus leucocephalus*).
 848 *Behavioral Ecology and Sociobiology* 65:1535-1547.

849
850
851
852
853
854
855
856
857
858
859
860
861

862 **Table 1:** Maternal behaviour of Novel male treatment and Familiar male treatment females
 863 over lactation

Comparison of the number of times maternal behaviours were recorded during lactation

Mated Control	Novel Male
---------------	------------

	F	d.f	P	Mean	SE	Mean	SE
In nest	3.77	1,41	0.059	114.34	4.15	103.47	3.54
Nursing	11.56	1,41	0.002	15.52	1.34	8.4	1.53
Licking Pups	1.79	1,41	0.188	1.82	0.63	0.80	0.34
Nest Building	0.12	1,41	0.730	0.65	0.27	0.80	0.31
Eating/ drinking	0.14	1,41	0.707	5.82	0.62	5.52	0.52
Grooming	0.14	1,41	0.705	9.73	0.68	10.14	0.82
Active	0.78	1,41	0.382	131.34	6.21	139.76	7.45
Resting	1.15	1,41	0.288	168.73	5.00	179	7.96

864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884

Table 2: Markers of oxidative stress in offspring of both treatments at adulthood. Showing protein thiol concentration ($\mu\text{mol g}^{-1}$ protein) and aconitase enzyme activity/citrate synthase (units/mg protein). Models were also fitted with plate number and block as a random factor to control for variation between assay plates.

Oxidative stress results for protein thiol concentration and aconitase enzyme activity

	Liver						Kidney					
	Protein thiol concentration			Aconitase enzyme activity			Protein thiol concentration			Aconitase enzyme activity		
	F	d.f	P	F	d.f	P	F	d.f	P	F	d.f	P
Treatment	60.65	1,80	<0.001	14.1	1,81	<0.001	1.44	1,80	0.234	1.47	1,80	0.228
Sex	0.130	1,80	0.719	0.20	1,81	0.656	0.01	1,80	0.997	0.32	1,80	0.569
Sex* treatment	1.43	1,80	0.234	1.59	1,81	0.210	0.61	1,80	0.436	1.85	1,80	0.177

	Heart						Gastrocnemius					
	Protein thiol concentration			Aconitase enzyme activity			Protein thiol concentration			Aconitase enzyme activity		
	F	d.f	P	F	d.f	P	F	d.f	P	F	d.f	P
Treatment	10.34	1,80	<0.001	20.42	1,81	<0.001	0.17	1,80	0.681	1.76	1,80	0.188
Sex	8.49	1,80	0.005	0.98	1,81	0.755	2.88	1,80	0.093	0.78	1,80	0.379
Sex* treatment	1.80	1,80	0.183	0.41	1,81	0.523	0.001	1,80	0.971	2.11	1,80	0.150

889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906

Author Manuscript

Mean litter Mass (g)

Author Manuscript

907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931

Figure 1: Average litter growth rate over lactation (4 weeks). Data are displayed as means \pm s.e.m.

932

933

Weight (g)

934

935

936 **Figure 2:** Comparison of individual offspring birth, weaning and adult weight between the

937 two treatments. Data are displayed as means \pm 3 s.e.m.

938

939

940

941

942

943

944

945

946

947

Mean G metabolites
(ng/g faeces)

Manuscript
Author Manuscript

948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971

Figure 3: Concentration of fecal glucocorticoid metabolites in females taken before the experiment started, on the second after exposure to a male/female and then mid lactation in the mated treatments. Females were either exposed to a novel male (novel male treatment) their mated male (familiar male) or another female (unmated control). Data are displayed as means \pm s.e.m.

Mean protein thiol concentration per organ
($\mu\text{mol g}^{-1}$ protein)

Author Manuscript

973

974 **Figure 4:** Mean protein thiol concentrations. Data is presented as estimated marginal means
975 ± 1 s.e.m. for each measure from general linear mixed models for each tissue sample.

976

977

978

979

980

981

982

983

984

985

986

987

988

989

990

991

Mean aconitase activity per organ
(units/mg protein)

Author Manuscript

992

993

994

995

Figure 5: Ratio of mean aconitase enzyme activity to citrate synthase. Data is presented as estimated marginal means \pm 1 s.e.m. for each measure from general linear mixed models for each tissue sample.