



Clinical outcomes and cost effectiveness of computer-guided versus conventional implant-retained hybrid prostheses: A long-term retrospective analysis of treatment protocols

Andrea Ravidà¹ | Shayan Barootchi¹ | Mustafa Tattan¹ | Muhammad H. A. Saleh¹ |
Jordi Gargallo-Albiol^{1,2} | Hom-Lay Wang¹

¹Graduate Periodontics, Department of Periodontics and Oral Medicine, University of Michigan School of Dentistry, Ann Arbor, MI

²International Master in Oral Surgery, Oral and Maxillo-facial Surgery Department, International University of Catalonia, Barcelona, Spain

Correspondence

Hom-Lay Wang, DDS, MSD, PhD, Department of Periodontics and Oral Medicine, University of Michigan School of Dentistry, 1011 North University Avenue, Ann Arbor, MI 48109-1078.

Email: homlay@umich.edu

Abstract

Background: Computer-guided systems were developed to facilitate implant placement at optimal positions in relation to the future prosthesis. However, the time, cost and, technique sensitivity involved with computer-guided surgery impedes its routine practice. The aim of this study is to evaluate survival rates and complications associated with computer-guided versus conventional implant placement in implant-retained hybrid prostheses. Furthermore, long-term economic efficiency of this approach was assessed.

Methods: Patients were stratified according to implant placement protocol into a test group, using computer-guided placement, and a control group, using traditional placement. Calibrated radiographs were used to measure bone loss around implants. Furthermore, the costs of the initial treatment and prosthetic complications, if any, were standardized and analyzed.

Results: Forty-five patients (149 implants in the test group and 111 implants in the control group) with a minimum follow-up of 5 years, and a mean follow-up of 9.6 years, were included in the study. While no significant difference was found between both groups in terms of biologic and technical complications, lower incidence of implant loss was observed in the test group ($P < 0.001$). A statistically significant difference in favor of the non-guided implant placement group was found for the initial cost ($P < 0.05$) but not for the prosthetic complications and total cost ($P > 0.05$).

Conclusions: Computer-guided implant placement for an implant-supported hybrid prosthesis is a valid, reliable alternative to the traditional approach for implant placement and immediate loading. Computer-guided implant placement showed higher implant survival rates and comparable long-term cost to non-guided implant placement.

KEYWORDS

computer-assisted surgery, cone-beam computed tomography, dental implants, implant-supported dental prosthesis, prostheses and implants



1 | INTRODUCTION

Dental implants have transformed the clinical perspective and rehabilitative approach to treating completely and partially edentulous patients.¹ In fact, the presence of implants is considered the most essential modifier of such patient therapy in the past 35 years.²

A key determinant of clinical implant success is accurate implant positioning and the avoidance of damage to the adjacent anatomical structures.³ Implementation of this pivotal factor has always been contingent upon operator skill and experience, in addition to other biologic and site-dependent factors.⁴ Hence, computer-guided systems were developed to facilitate implant placement in an optimal planned position and to retain the future prosthesis in an optimal biologic position.⁵ Such high precision is expected to decrease biologic and prosthetic complications, especially in more complex cases.⁶ Implants placed via computer-guided implant placement (CGIP) are reportedly within 1 mm and around 5° of deviation from the originally planned implant position.³ This is further supported by a recent systematic review which examined > 1,400 CGIP and showed mean global inaccuracies of 1.1 mm at entry point, 1.4 mm at the apex, and a 3.9° angular deviation.⁷

Quite helpfully, CGIP protocols use computed tomography (CT) scans for virtual identification and placement of implants in the exact positions and angulations avoiding vital anatomical structures.⁸ This treatment modality may eliminate the need for bone grafting, and even raising a flap, if appropriate bone dimension and morphology exists.^{9,10} This might, in turn, improve patient acceptance to the recommended treatment and reduce post-surgical morbidity.^{8,11}

Despite high documented survival rates for implants placed using CGIP (approximately 95% at a 7-year follow up period), a higher rate of prosthetic and biologic complications has also been reported.¹² It is crucial to keep in mind that precision of implant position remains subject to the guide accuracy and adherence of the clinician to the proposed surgical protocol.⁸ Although these high-precision technologies are used in the fabrication of such guides, several studies have shown some linear and angular deviations between the planned and placed implants.^{13–16} This may raise the question as to whether or not the use of a computer-guided protocol by an inexperienced operator expands the probability of technique-dependent complications. Conversely, Van de Wiele et al. reported that guided systems could facilitate and expedite ideal implant placement for clinicians with limited experience.¹⁷

The additional preparation time, greater cost, and technique sensitivity associated with CGIP are the chief obstacles to its routine practice.^{18,19} The results of a recent systematic review²⁰ questioned the economic benefits of using CGIP. Thus, another question that needs to be addressed is the

cost-effectiveness of CGIP across short- and long-time periods when compared with traditional implant placement. Finally, a commonly overlooked element of investigating the efficiency of CGIP, is whether or not the accuracy of these protocols results in decreasing the incidence of long-term postoperative complications, when compared to traditional non-computer-guided protocols. Thus, the aim of this study was to evaluate the long-term survival rate, complication rate, and cost of CGIP compared to traditional protocol. A null hypothesis was formulated, with the authors anticipating no influence using CGIP on the identified outcomes.

2 | MATERIALS AND METHODS

The present study was conducted according to the principles embodied in the Helsinki Declaration of 1975, as revised in 2000 for biomedical research involving human subjects, and was approved by the Institutional Review Board for Human Studies, School of Dentistry, University of Michigan, Ann Arbor, MI, (HUM00114382) to be conducted at the Department of Periodontology within the same institution.

This retrospective investigation enrolled all patients treated with implant-retained hybrids between January 1990 and September 2017 at the School of Dentistry, University of Michigan, Ann Arbor, MI. All papers and digital charts of edentulous patients treated with implant-retained hybrid prostheses were carefully scanned and analyzed by two authors (AR, SB). During each stage, all disagreements were resolved through discussion with a third reviewer (JG).

2.1 | Inclusion criteria

Inclusion criteria were 1) edentulous patients treated with implant-retained fixed hybrid prostheses and a documented follow-up of ≥ 5 years after implant placement; 2) cases where all implant fixtures associated with the prosthesis were placed within the same surgical procedure; and 3) patients who received an implant-retained fixed hybrid prosthesis, returning for regular maintenance, at the University of Michigan School of Dentistry.

2.2 | Exclusion criteria

Exclusion criteria were 1) edentulous patients treated with a removable overdenture or ceramic fixed dental prosthesis; 2) patients with ambiguous or incomplete charts; 3) patients with a < 5-year follow-up; 4) patients treated or maintained in centers outside the University of Michigan School of Dentistry; and 5) patients with inaccessible files due to bad debt or decease.




FIGURE 1 A) Prosthetically driven virtual positioning of implants and fixating screws using the software. B) Hybrid prosthesis (maxilla) in situ after delivery

2.3 | Data collection and classification

Within the review period, 222 patients were screened, their data subsequently evaluated against the aforementioned inclusion and exclusion criteria. In total, 45 patients were included in the study, while 175 were excluded for the following reasons: 1) 51 implant-retained hybrids with a < 5-year follow-up, 2) 49 implant-retained overdentures, 3) 32 implant-retained fixed bridges, 4) 32 inaccessible files, 5) 4 files with missing or incomplete data, 6) 3 destroyed files, 7) 3 deceased patients, and 8) 1 removable partial denture.

Later, the selected cases were separated into two groups: computer-guided implant placement (CGIP) as the test group (26 patients) and non-CGIP as the control group (19 patients).

Patient information, such as age (at the day of implant placement), sex, presence of a smoking habit (≥ 1 cigarette/day), diabetes (verified by checking full medical records) and history of periodontal disease were obtained. History of periodontal disease, determined by reviewing the periodontal chart, was defined as the presence of at least four sites with clinical attachment loss (AL) ≥ 3 mm and a past history of scaling and root planing.^{21,22} Additional data including time of implant placement (immediate, early, or delayed), time of implant loading (immediate, early, or delayed), number of implants and their positions, implant configuration (brand, length, and diameter), and whether or not bone augmentation or a flap procedure were performed.

The following prosthodontic/peri-implant complications and subsequent management were recorded at follow-up appointments: 1) fractured/chipped/replaced prosthetic tooth, 2) fractured prosthesis, 3) fractured bar, and 4) loosened abutment screw.

All complications have been classified into the following:

1. Biologic complications: peri-implant mucositis, peri-implantitis, implant failure and hyperplasias, prosthesis-induced ulcers of fibrous connective tissue, fistula formation, pain, or infection.

2. Early or delayed prosthetic complications: early prosthetic complications were defined as those occurring within 1 year of prosthetic loading, whereas late prosthetic complications were defined as those that occur 1 year following prosthetic loading.

3. Minor, moderate, or major prosthetic complications: minor complications are those managed within 24 hours of presentation, moderate are those managed between 2 to 7 days, while major complications required > 7 days to manage.

2.4 | Computer-guided implant placement (CGIP) group

The CGIP was planned according to manufacturer instruction. Digital three-dimensional (3D) diagnosis and treatment planning using software defines implant positions and sizes from an anatomical, surgical, and prosthetic perspective by combining the 3D future tooth setup according to the patient's anatomy. Anatomical conditions had to permit the placement of at least four implants in the positions ideal for full-arch prosthetic rehabilitation to be achieved. Treatment planning involved cone-beam computed tomography (CBCT)* or CT scans of both the patient and the prosthetically-driven radiographic guide according to the double-scan protocol: an initial scan of the patient wearing the radiographic guide prepared following tooth set-up, and a second scan of the template alone. Next, both scans were superimposed using the dedicated software to establish optimal implant positioning. The planning data was then sent to the manufacturer where a surgical template with hollow metallic sleeves was designed and later produced for implant placement according to the software-identified positions (Figure 1A). When immediate loading was necessary, full acrylic resin screw-retained provisional prostheses were prefabricated based on the surgical guide and the model obtained from the surgical templates were placed intraorally and fixed with ≥ 3 anchor pins. After

* 3DX Accutomo FPD; J Morita Mfg, Kyoto, Japan



correct placement and stabilization of the template, a flapless implant surgery was performed according to the software manufacturer's protocol, and fully guided drilling preceding implant insertion followed. Some patients were restored with a provisional fixed, immediately loaded prosthesis, while others went for early or delayed loading depending on primary stability.

2.5 | Non-CGIP group

Implant rehabilitation was planned on 2D (panoramic XR) or 3D (CBCT or CT) diagnostic imaging. Consequently, surgical guides were constructed from the diagnostic tooth set-up and cast model analysis using a light-polymerized composite material, where drill blanks placed in the prosthodontically driven implant position were set to assist the free-hand (non-CGIP).

A central, crestal arch incision was made on the alveolar ridge and a full thickness flap was elevated. When necessary, a distal vertical incision was performed. Subsequently, the drilling sequence proceeded according to manufacturer's instruction. A variety of implant systems*†‡§ were used in this group. Guided bone regeneration (GBR) was performed simultaneously, when necessary, using allograft particulate bone¶ and an absorbable collagen membrane# to repair bone defects and augment horizontal bone volume.

2.6 | Peri-implantitis and implant failure

To classify peri-implantitis, the definitions proposed by the 8th European Workshop on Periodontology in 2011 were adopted,²³ where peri-implantitis was defined as clinical inflammation together with radiographic marginal bone loss > 2 mm. Peri-implant marginal bone loss was measured at baseline (following the expected period of remodeling) and final follow-up via calibrated periapical and panoramic radiographs using software.^{||24} Two individual, calibrated examiners (JG and SB) performed the calculations separately using the designated software. Where significant differences were found, a third reviewer (AR) reassessed the radiographs to arrive at a final resolution. Peri-implantitis was evaluated per patient, then per implant individually. The incidence of peri-implantitis was recorded using a binary score for each implant (0 for a healthy implant, 1 for a diseased implant)

thus calculating the percentage of diseased implants. Similar dichotomous values were assigned to patients based on the presentation of peri-implantitis around any implants (0 for a patient with all healthy implants, 1 for a patient with radiographic signs of ≥ 1 diseased implant). Implant failure was defined as a removed, lost, mobile, or fractured implant and calculated for each implant individually and then each patient, with the same standards used previously for peri-implantitis.²⁵

2.7 | Prosthesis design

Only implant-retained fixed hybrid prostheses were included in this article. A titanium or gold bar was used to anchor the acrylic base, with a set of acrylic teeth in place (Figure 1B).

2.8 | Cost

The analysis of cost in this study was patient-focused, intended to identify all the necessary costs of diagnostic, therapeutic, and follow-up procedures. The primary objective of this analysis was to achieve a more comprehensive understanding of cost-effectiveness associated with both approaches, and their complications, discussed in this paper.

The average cost of clinical procedures across the 5- to 25-year follow-up period was determined beforehand and used as a method of standardization among the study sample. The costs were obtained and categorized into the following: 1) initial cost: implant + prosthesis placement fees; 2) cost of complication management: prosthetic + implant complication management fees; and 3) total cost: initial cost + cost of complication management.

The cost of all treatments related to initial placement and management procedures was predetermined based on an average of their individual costs every year since 1994, at the University of Michigan, School of Dentistry. This was performed to prevent the regular rate of inflation along the 5- to 23-year period interfering with the standardization and reliability of the cost analysis. After a price list was formulated based on these averages, all procedures pertaining to each patient file were scanned and recorded by one study investigator (MT). Wherever doubt arose, an expert in the matter (HLW) was referred to. With these records, the cost of treatment and management performed on each patient was noted and computed into the aforementioned categories of cost.

The purpose of analysis was to simulate a clinical setting where a patient is not pardoned for payments, just as a means of having a fair and elaborate comparison between the two treatment approaches. Therefore, whether or not the patient had actually paid for the provided treatments, actual cost was presumed within the particular patient's cost of treatment.

Within the initial cost, every treatment fee, such as preliminary consultation appointments, use of radiographic and/or

* NobelReplace, Tapered Groovy implant, Nobel Biocare, Gothenburg, Sweden

† Branemark Mark III® and IV®, Nobel Biocare, Gothenburg, Sweden

‡ Zimmer, Palm Beach Gardens, FL

§ Biohorizons IPH, Birmingham, AL

¶ Puros, Zimmer, Palm Beach Gardens, FL

Bio-Gide®, Geistlich Pharma AG, Wolhusen, Switzerland

|| ImageJ, US National Institutes of Health, Bethesda, MD




FIGURE 2 Distribution of patients in test and control groups according to the number of implants supporting the hybrid prosthesis

laboratory diagnostic aids, laboratory fees and preparations, and the entire cost of surgery, were included. Complication management cost included any fee related to follow-up maintenance, as well as management of any biologic or prosthetic complication pertaining to any of the components.

The average cost of each procedure was calculated as follows:

$$\text{Cost}_1 + \text{Cost}_2 + \text{Cost}_3 + \text{Cost}_4 \dots / n$$

where:

Cost_x = Procedure Cost at a Given Year

n = Total number of Cost_x events per procedure

2.9 | Statistical analyses

The demographic profile and clinical characteristics of the included sample were analyzed using: 1) descriptive statistics: mean, standard deviation, median; 2) Chi2 homogeneity tests (Chi2); 3) Fisher's exact test (Fis); and 4) Mann-Whitney (MW).

The association between prosthetic complications across both study groups was analyzed using: 1) descriptive analyses: number of cases (%) and mean \pm SD; 2) a simple binary logistic regression model for each type of complication: estimation of unadjusted odds ratio (OR); and 3) an MW test for homogeneity test of distributions in continuous variables.

The probability of peri-implantitis and implant failure in both groups was assessed using a generalized estimation equation (GEE): estimation of OR adjusted by sex, age, and follow-up time.

Cost analysis was performed using a general linear model: estimation of coefficients adjusted by number of implants and follow-up time (years).

3 | RESULTS

3.1 | Descriptive analyses

A total of 45 patients, 24 males (53.3%) and 21 females (46.7%), with a mean age of 58.9 ± 13.1 years (22 to 83 years), who received full-arch implant-retained hybrids during the past 27 years at the University of Michigan School of Dentistry, were selected.

Twenty-six (149 implants) of the total cases were treated with CGIP (test group), while the remaining 19 (111 implants) were treated traditionally (control group).

A total of 260 implants were included: 26 patients (80%) received five or six implants (40% each), five patients (11.1%) received eight implants and four patients (8.8%) received seven or four implants (4.4% each) (Figure 2). The average follow-up period was 116.0 ± 45.9 months (9.66 ± 3.82 years), where half the sample was monitored across a minimum of 9 years.

3.2 | Demographic profile and clinical characteristics

No statistically significant difference was observed with age ($P = 0.061$), although the test group did demonstrate a greater frequency of increased age, representing a mean age of 62.5 ± 10.7 , as opposed to the 53.9 ± 14.7 of the control group. Although not statistically significant ($P = 0.069$), the follow-up period was markedly longer in the control

TABLE 1 The characteristics and demographics of patients in control and test groups

	Parameters	Group		P value (test)
		Non-guided	Guided	
Patients	n (patients)	19	26	
	Follow up (months)	128.7 ± 62.3	101.0 ± 27.1	0.069 (MW)
	Age (years)	53.9 ± 14.7	62.55 ± 10.7	0.061 (MW)
	Male	6 (31.6)	18 (69.2)	0.012 (Chi²)
	Smokers	3 (15.8)	2 (7.7)	0.636 (Fis)
	Diabetes	3 (15.8)	2 (7.7)	0.636 (Fis)
	History of periodontitis	5 (26.3)	4 (15.4)	0.365 (Chi ²)
	Maxillary rehabilitations	8 (42.1)	15 (57.7)	0.302 (Chi ²)
	Immediate implants	2 (10.5)	2 (7.7)	1.000 (Fis)
	Immediate loading	4 (21.1)	17 (65.4)	0.003 (Chi²)
	Open Flap	19 (100)	2 (7.69)	<0.001 (Chi²)
	Bone Regeneration	11 (57.9)	2 (5.2)	0.001 (Chi²)
	Screwed prosthesis	17 (89.5)	26 (100)	0.176 (Chi ²)
	Implant failure (Patient)	4 (21.1)	5 (19.2)	0.880 (Chi ²)
Implants	n (implants)	111	149	
	Implant failure	22 (19.8)	5 (3.3)	<0.001 (Chi²)

Bold indicates statistically significant associations; Chi2 homogeneity tests (Chi2); Fisher's exact test (Fis); Mann-Whitney tests (MW).

group, with an approximately > 2-year difference. In order to avoid this difference interfering with the results, considering that it is plausible for a longer follow-up to be associated with more complications, this variable was controlled and adjusted for during statistical analysis. Clinical parameters such as implant loading (immediate versus delayed), the presence of a flap versus lack thereof and bone regeneration as a consequence of surgical planning also demonstrated significant differences between the two groups ($P < 0.01$); where guided surgery was normally associated with flapless surgery, immediate loading, and no bone grafting procedures. Contrarily, differences between smoking, diabetes, periodontitis, and both arches were not statistically significant

($P > 0.05$). The characteristics and demographics of the included patients are summarized in Table 1.

3.3 | Prosthetic complications

For each of the investigated parameters, no statistically significant difference was found between test and control ($P > 0.05$).

In both groups, tooth replacement was the most common problem, affecting 55.6% of the total sample. Denture removal, due to bulk fracture, was the second most detected complication (35.6%), followed by partial acrylic fracture (24.4%) (Figure 3).

Regarding the time of complication occurrence, 24.4% and 68.9% of patients presented with early and delayed complications, respectively. However, no statistically significant difference was found between test and control groups ($P > 0.05$); Table 2 depicts the incidence of prosthetic complications in both groups.

3.4 | Biologic complications

3.4.1 | Peri-implantitis

A generalized estimation equation (GEE), adjusted according to sex, age, and follow-up time, demonstrated a lower incidence of peri-implantitis within the CGIP group compared to the non-CGIP group; both per patient (34.6% vs. 52.6%) and per implant (13.4% vs. 24.3%). However, this was not statistically significant ($P = 0.230$; $P = 0.714$).

In addition, an observed trend, short of statistical significance ($P = 0.085$), depicted a lower probability of




FIGURE 3 Incidence of each of the prosthetic complications in either group, and prevalence of prosthetic complications in the total cohort

TABLE 2 The incidence and different types of prosthetic complications in test and control groups

Complications	Group		OR (95% CI)	P value
	Non-guided	Guided		
n (patients)	19	26		
Tooth replacement	11 (57.9)	14 (53.8)	0.84 (0.26-2.79)	0.787
Abutment breakage	1 (5.3)	2 (7.7)	1.50 (0.13-17.9)	0.748
Acrylic fracture	5 (26.3)	6 (23.1)	0.84 (0.21-3.30)	0.803
Dislodgment of entire prosthesis	1 (5.3)	1 (3.8)	0.72 (0.04-12.3)	0.820
Fracture of bar	2 (10.5)	2 (7.7)	0.71 (0.09-5.54)	0.742
Screw loosening	1 (5.3)	1 (3.8)	0.72 (0.04-12.3)	0.820
Denture removal	8 (42.1)	8 (30.8)	0.61 (0.18-2.09)	0.434
Early complications	5 (26.3)	6 (23.1)	0.84 (0.21-3.30)	0.803
Delayed complications	13 (68.4)	18 (69.2)	1.04 (0.29-3.72)	0.954
Times without prosthesis (resolved within 1-day)	2.4 ± 3.6	3.2 ± 3.6		0.598 (MW)
Times without prosthesis (resolved within 1-week)	0.4 ± 0.8	0.2 ± 0.5		0.516 (MW)
Times without prosthesis (resolved after 1-week)	0.8 ± 1.3	0.5 ± 0.8		0.791 (MW)

Mann-Whitney Tests (MW).

peri-implantitis with increasing age (OR = 0.95). Specifically, every additional year can be associated with a 5% reduction in risk of peri-implantitis.

3.4.2 | Prosthesis-induced biologic complications

A small number of prosthesis-induced biologic complications were observed within both groups. A clinical observation of an ulcer in one patient and an epulis fissuratum in another were recorded in the non-CGIP group. Three separate cases of ulcerations and a single presentation of soreness were documented within the CGIP group.

3.4.3 | Implant failure

A statistically significant difference in implant survival rate was found between the control (80.2%) and test group (96.7%) ($P < 0.001$). This was not the case when the same analysis was performed per patient ($P > 0.05$). The data showed that the 22 implant failures within the non-CGIP group were linked to only four patients (21.1%), whereas, the five documented failures within the CGIP group were associated with five (19.2%) individual patients (Table 1).

3.5 | Cost analyses

The analysis concluded that, when differences in both implant number and follow-up period were adjusted, neither mean total cost of CGIP versus non-CGIP nor the cost of the associated complications was significant ($P = 0.573$) and ($P = 0.860$), respectively (Figure 4). However, a comparison of both procedures' initial cost, considering the same number of implants, displayed statistical significance ($P < 0.05$), where CGIP surgery costed an additional \$659.10.




FIGURE 4 Projection of costs expended through the overall follow-up time, based on the number of implants used to support the framework

4 | DISCUSSION

Our results confirm that CGIP for implant-retained hybrid prostheses is an effective treatment option for experienced and inexperienced clinicians alike. The predicted null hypothesis for implant survival rate was rejected, since it was higher in CGIP, however, it was verified for the incidence of peri-implantitis, with no differences between both groups. When the cost of managing all complications throughout the follow-up period was considered, no difference was found between both groups; though, initially, the cost was higher in the test group.



Observational studies are able to create credible evidence of intervention effects through tracking large cohorts. This brings the benefits of generalizability, potential for real-world comparisons of treatment efficacy and long-term outcomes.²⁶ Using up-to-date methodological and statistical strategies can, when appropriately applied on long-term follow-up cohorts with sufficient data, improve result reliability.²⁷ In the past 2 decades, the paradigm of a prosthetically-driven implant surgery has been subject to fundamental evolution in practice. Proper implant positioning has obvious advantages, such as long-term stability of peri-implant hard and soft tissues, enhanced oral hygiene procedures, the potential for achieving optimal occlusion, and more favorable esthetic outcomes.²⁸⁻³⁰ Not limited to that, several groups advocated tailored, site-specific planning for implant placement, negating the need for augmentation, where several clinical studies reported excellent results for no-augmentation techniques, developed to restore edentulous patients with fixed prostheses.³¹⁻³³ Recently, the growing need for patient rehabilitation with implant-retained fixed prostheses was what pushed the industry towards applying present-day technology pursuing well-accepted, highly predictable, less invasive, and less technique-sensitive protocols for edentulous patient rehabilitation.¹⁸ The evident success of computer-guided systems means, for patients, that the entire procedure from surgery to final prosthetic restoration can be accomplished with reduced postoperative morbidity and overall treatment time.^{34,35}

In the current study, implant survival rate in the CGIP group was statistically more significant than the non-CGIP group ($P < 0.001$) with a 96.7% survival rate, concurring with previous studies, reporting 97.8%¹⁸ and 97.6%³⁶ survival rates at approximately 3 years of follow-up. This proves the external validity of our results, while a robust advantage of our study is the longer range of follow-up, with an average of 9.6 years. It should be highlighted that in our study, a significant difference, in survival, was only observed on an implant level ($P \leq 0.001$), but not on a patient level ($P = 0.88$).

A well-known concern with implant-retained hybrid prostheses is prosthetic complications, most commonly acrylic denture fracture. Similar studies reported a slightly higher rate of acrylic fracture 34.7%,³⁷ 30.3%,³⁸ and 30%,³⁹ as our in study 23.1% of cases had acrylic fractures. However, the most common technical complication encountered in the current study was prosthetic tooth loss and replacement with a noteworthy rate of 53.8%. Bulk fracture of acrylic teeth is a common observation in implant-retained acrylic prostheses, and recent studies suggested that teeth fabricated with improved materials are expected to perform better long-term.⁴⁰ In the current study, different types of acrylic teeth provided by the same manufacturer were used.*

To avoid bias and decrease confusion, we elected to utilize the definition of peri-implantitis proposed by the 8th European Workshop on Periodontology in 2011.²³ According to this definition, the incidence of peri-implantitis in our test group was 13.4%. We are not aware of comparable studies which have used the definition we have adopted, rather, most studies proposed their own definitions for peri-implantitis,³⁷ which generally renders incomparable and erratic results. We, therefore, assume that peri-implantitis incidence has been under-reported in similar studies, where Malo et al. reported 8.7% of cases to have peri-implantitis after one year,³⁷ and Puig et al. reporting an incidence of only 5.6%.⁴¹

One of the virtues of having a long-term follow-up, is the ability to compare the costs of resolving prosthetic and technical complications along a relatively long period. We are also not aware of similar studies that compared costs of managing late complications. It is well known that a computer-guided approach is more expensive than conventional implant placement due to software utilization, denture duplication, scanning patient's denture with CBCT, surgical template fabrication, extra laboratory fees and planning time.⁴² However, our study revealed a rather intriguing finding. Though CGIP approach did generate a greater initial cost, no significant difference in mean total costs was found, when managing short and long-term complications in both groups was taken into account.

In the CGIP group, all surgical templates were produced by the same manufacturer[†] and the same implant system was utilized, which limits the validity of our results to this particular manufacturer. It would be of interest to investigate whether other implant systems would suffer less incidence of peri-implantitis, for instance. Another limitation in our results, is that it include immediately loaded implants, which commonly have different survival rates and marginal bone levels than single or partial tooth rehabilitations.⁶

The very nature of this observational study, did not allow for the accuracy of used guides to be tested. However, though a guide was used, in two cases an open flap approach was mandatory, to correct fenestrations which occurred during implant placement, these sites were subsequently augmented with bone grafts and protected with barrier membranes.

Finally, 80% of the included cases received five or six implants, with the remaining 20% restored with four, seven, or eight implants. This presents a difference to other studies that have investigated a particular configuration like the "All-on-4" or "All-on-6,"^{32,41} again, limiting the validity of our results to that particular number of implants utilized for rehabilitation.

* Ivoclar Vivadent, Schaan, Liechtenstein

[†] NobelReplace, Tapered Groovy implant, Nobel Biocare, Gothenburg, Sweden



Observations from this study strongly suggest that the treatment team must be aware of the various steps involved in prosthesis fabrication, in order to yield an easier troubleshooting process during and after the surgery, if deemed necessary. Finally, upon case planning, it is critical to realize that every step will hold a lifetime consequence for restoration reliability.

5 | CONCLUSION

Our clinical results confirm that computer-guided implant placement for implant-supported hybrids is a valid, reliable alternative to the traditional approach of implant placement and immediate loading. Implants placed via guided surgery demonstrated higher survival rates and comparable long-term cost when compared with non-guided implant placement. No difference in technical complications was observed between the two groups. More consideration should be given to the routine use of computer-guided surgery in the treatment of edentulous cases.

ACKNOWLEDGMENTS

The authors declare no conflicts of interest within this study. The authors do not have any financial interests, neither directly nor indirectly, in the products or information listed in the paper.

REFERENCES

- Goodacre CJ, Naylor WP. Single implant and crown versus fixed partial denture: a cost-benefit, patient-centred analysis. *Eur J Oral Implant.* 2016; 9(Suppl 1):S59–68.
- Tarnow DP. Increasing Prevalence of Peri-implantitis: how Will We Manage. *J Dent Res.* 2016;95:7–8.
- Nickenig HJ, Eitner S. Reliability of implant placement after virtual planning of implant positions using cone beam CT data and surgical (guide) templates. *J Craniomaxillofac Surg.* 2007;35:207–211.
- Esposito M, Hirsch JM, Lekholm U, Thomsen P. Biological factors contributing to failures of osseointegrated oral implants. (II). Etiopathogenesis. *Eur J Oral Sci.* 1998;106:721–764.
- Vercruyssen M, Fortin T, Widmann G, Jacobs R, Quirynen M. Different techniques of static/dynamic guided implant surgery: modalities and indications. *Periodontol 2000.* 2014;66:214–227.
- Van Assche N, Vercruyssen M, Coucke W, Teughels W, Jacobs R, Quirynen M. Accuracy of computer-aided implant placement. *Clin Oral Implants Res.* 2012;23(Suppl 6):112–123.
- Tahmaseb A, Wismeijer D, Coucke W, Derksen W. Computer technology applications in surgical implant dentistry: a systematic review. *Int J Oral Maxillofac Implants.* 2014;29(Suppl):25–42.
- D'Haese J, Ackhurst J, Wismeijer D, De Bruyn H, Tahmaseb A. Current state of the art of computer-guided implant surgery. *Periodontol 2000.* 2017;73:121–133.
- van Steenberghe D, Glauser R, Blomback U, et al. A computed tomographic scan-derived customized surgical template and fixed prosthesis for flapless surgery and immediate loading of implants in fully edentulous maxillae: a prospective multicenter study. *Clin Implant Dent Relat Res.* 2005;7(Suppl 1):S111–120.
- Krekmanov L. Placement of posterior mandibular and maxillary implants in patients with severe bone deficiency: a clinical report of procedure. *Int J Oral Maxillofac Implants.* 2000;15:722–730.
- Vasak C, Kohal RJ, Lettner S, Rohner D, Zechner W. Clinical and radiological evaluation of a template-guided (NobelGuide) treatment concept. *Clin Oral Implants Res.* 2014;25:116–123.
- Lopes A, Malo P, de Araujo Nobre M, Sanchez-Fernandez E, Gravito I. The NobelGuide(R) All-on-4(R) Treatment Concept for Rehabilitation of Edentulous Jaws: a Retrospective Report on the 7-Years Clinical and 5-Years Radiographic Outcomes. *Clin Implant Dent Relat Res.* 2017;19:233–244.
- Van Assche N, van Steenberghe D, Guerrero ME, et al. Accuracy of implant placement based on pre-surgical planning of three-dimensional cone-beam images: a pilot study. *J Clin Periodontol.* 2007;34:816–821.
- Pettersson A, Kero T, Gillot L, et al. Accuracy of CAD/CAM-guided surgical template implant surgery on human cadavers: part I. *J Prosthet Dent.* 2010;103:334–342.
- Pettersson A, Kero T, Soderberg R, Nasstrom K. Accuracy of virtually planned and CAD/CAM-guided implant surgery on plastic models. *J Prosthet Dent.* 2014;112:1472–1478.
- Cushen SE, Turkyilmaz I. Impact of operator experience on the accuracy of implant placement with stereolithographic surgical templates: an in vitro study. *J Prosthet Dent.* 2013;109:248–254.
- Van de Wiele G, Teughels W, Vercruyssen M, Coucke W, Temmerman A, Quirynen M. The accuracy of guided surgery via mucosa-supported stereolithographic surgical templates in the hands of surgeons with little experience. *Clin Oral Implants Res.* 2015;26:1489–1494.
- Meloni SM, De Riu G, Pisano M, Cattina G, Tullio A. Implant treatment software planning and guided flapless surgery with immediate provisional prosthesis delivery in the fully edentulous maxilla. A retrospective analysis of 15 consecutively treated patients. *Eur J Oral Implant.* 2010;3: 245–251.
- Merli M, Bernardelli F, Esposito M. Computer-guided flapless placement of immediately loaded dental implants in the edentulous maxilla: a pilot prospective case series. *Eur J Oral Implant.* 2008;1:61–69.
- Laleman I, Bernard L, Vercruyssen M, Jacobs R, Bornstein MM, Quirynen M. Guided Implant Surgery in the Edentulous Maxilla: a Systematic Review. *Int J Oral Maxillofac Implants.* 2016;31(Suppl):s103–117.
- Armitage GC. Periodontal diagnoses and classification of periodontal diseases. *Periodontol 2000.* 2004;34:9–21.
- Pihlstrom BL, Michalowicz BS, Johnson NW. Periodontal diseases. *Lancet.* 2005;366:1809–1820.
- Tonetti M, Palmer R, Working Group 2 of the VEWoP. Clinical research in implant dentistry: study design, reporting and outcome measurements: consensus report of Working Group 2 of the VIII European Workshop on Periodontology. *J Clin Periodontol.* 2012;39(Suppl 12):73–80.



24. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods*. 2012;9:671–675.
25. Chrcanovic BR, Albrektsson T, Wennerberg A. Reasons for failures of oral implants. *J Oral Rehabil*. 2014;41:443–476.
26. Kalincik T, Butzkueven H. Observational data: understanding the real MS world. *Mult Scler*. 2016;22:1642–1648.
27. Trojano M, Tintore M, Montalban X, et al. Treatment decisions in multiple sclerosis – insights from real-world observational studies. *Nat Rev Neurol*. 2017;13:105–118.
28. Buser D, Chen ST, Weber HP, Belser UC. Early implant placement following single-tooth extraction in the esthetic zone: biologic rationale and surgical procedures. *Int J Periodontics Restorative Dent*. 2008;28:441–451.
29. Monje A, Galindo-Moreno P, Tozum TF, Suarez-Lopez del Amo F, Wang HL. Into the Paradigm of Local Factors as Contributors for Peri-implant Disease: short Communication. *Int J Oral Maxillofac Implants*. 2016;31:288–292.
30. Buser D, Halbritter S, Hart C, et al. Early implant placement with simultaneous guided bone regeneration following single-tooth extraction in the esthetic zone: 12-month results of a prospective study with 20 consecutive patients. *J Periodontol*. 2009;80:152–162.
31. Popper HA, Popper MJ, Popper JP. Teeth in a day. The Branemark Novum system. *N Y State Dent J*. 2003;69:24–27.
32. Malo P, Rangert B, Nobre M. All-on-Four™ immediate-function concept with Branemark System implants for completely edentulous mandibles: a retrospective clinical study. *Clin Implant Dent Relat Res*. 2003;5(Suppl 1):2–9.
33. Krekmanov L, Kahn M, Rangert B, Lindstrom H. Tilting of posterior mandibular and maxillary implants for improved prosthesis support. *Int J Oral Maxillofac Implants*. 2000;15:405–414.
34. Cannizzaro G, Leone M, Consolo U, Ferri V, Esposito M. Immediate functional loading of implants placed with flapless surgery versus conventional implants in partially edentulous patients: a 3-year randomized controlled clinical trial. *Int J Oral Maxillofac Implants*. 2008;23:867–875.
35. Esposito M, Grusovin MG, Maghaireh H, Coulthard P, Worthington HV. Interventions for replacing missing teeth: management of soft tissues for dental implants. *Cochrane Database Syst Rev*. 2007;CD006697.
36. Balshi SF, Wolfinger GJ, Balshi TJ. Guided implant placement and immediate prosthesis delivery using traditional Branemark System abutments: a pilot study of 23 patients. *Implant Dent*. 2008;17:128–135.
37. Malo P, de Araujo Nobre M, Lopes A. The use of computer-guided flapless implant surgery and four implants placed in immediate function to support a fixed denture: preliminary results after a mean follow-up period of thirteen months. *J Prosthet Dent*. 2007;97:S26–34.
38. Gillot L, Noharet R, Cannas B. Guided surgery and presurgical prosthesis: preliminary results of 33 fully edentulous maxillae treated in accordance with the NobelGuide protocol. *Clin Implant Dent Relat Res*. 2010;12(Suppl 1):e104–113.
39. Marra R, Acocella A, Rispoli A, Sacco R, Ganz SD, Blasi A. Full-mouth rehabilitation with immediate loading of implants inserted with computer-guided flap-less surgery: a 3-year multicenter clinical evaluation with oral health impact profile. *Implant Dent*. 2013;22:444–452.
40. Kappert PF, Kelly JR. Cyclic fatigue testing of denture teeth for bulk fracture. *Dent Mater*. 2013;29:1012–1019.
41. Pomares C. A retrospective study of edentulous patients rehabilitated according to the ‘all-on-four’ or the ‘all-on-six’ immediate function concept using flapless computer-guided implant surgery. *Eur J Oral Implantol*. 2010;3:155–163.
42. Vercruyssen M, Hultin M, Van Assche N, Svensson K, Naert I, Quirynen M. Guided surgery: accuracy and efficacy. *Periodontol 2000*. 2014;66:228–246.

How to cite this article: Ravidà A, Barootchi S, Tattan M, Saleh MHA, Gargallo-Albiol J, Wang H-L. Clinical outcomes and cost effectiveness of computer-guided versus conventional implant-retained hybrid prostheses: A long-term retrospective analysis of treatment protocols. *J Periodontol*. 2018;89:1015–1024. <https://doi.org/10.1002/JPER.18-0015>