

# Health Economics

## The Effect of Organized Breast Cancer Screening on Mammography Use: Evidence from France

Journal:	<i>Health Economics</i>
Manuscript ID	HEC-17-0599.R2
Wiley - Manuscript type:	Research Article
Keywords:	mammography, cancer, preventive health, disparities, France

SCHOLARONE™  
Manuscripts

Review

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1002/hec.3813](https://doi.org/10.1002/hec.3813)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

# The Effect of Organized Breast Cancer Screening on Mammography Use: Evidence from France

Thomas C. Buchmueller<sup>1</sup> and Léontine Goldzahl\*<sup>2</sup>

<sup>1</sup>Ross School of Business, University of Michigan and NBER

<sup>2</sup>Manchester Centre for Health Economics, School of Health Sciences, University of  
Manchester and EDHEC Business School

July 6, 2018

## Abstract

In 2004, France introduced a national program of organized breast cancer screening. The national program built on pre-existing local programs in some, but not all, *départements*. Using data from multiple waves of a nationally representative biennial survey of the French population, we estimate the effect of organized screening on the percentage of women obtaining a mammogram. The analysis uses difference-in-differences methods to exploit the fact that the program was targeted at women in a specific age group: 50 to 74 years old. We find that organized screening significantly raised mammography rates among women in the target age range. Just above the lower age threshold, the percentage of women reporting that they had a mammogram in the past two years increased by over 10 percentage points after the national program went into effect. Mammography rates increased even more among women in their sixties. Estimated effects are particularly large for women with less education and lower incomes, suggesting that France's organized screening program has reduced socioeconomic disparities in access to mammography.

---

\*For the second author, this research has been supported by a post-doctoral grant co-financed by the chair "Santé" at the Fondation du risque and the project ANR 11-LABX-0019. The authors wish to thank the IRDES who provided the ESPS data used in this paper.

# 1 Introduction

Breast cancer is the most frequently diagnosed cancer and the leading cause of death from cancer among women worldwide, accounting for one-quarter of all new cancer cases and roughly 15 percent of cancer deaths (Torre et al., 2015). In 2015, roughly 154,400 women in the 28 countries of the European Union and 46,400 women in the U.S. died of breast cancer (Ferlay et al. (2015)). If detected early, breast cancer is highly treatable, with very high rates of survival. Survival rates depend importantly on the stage at which breast cancer is detected. Five-year relative survival rate for breast cancer are 99 percent for localized cancers, 85 percent for regional cancers and 26 percent for metastatic cancers (Siegel, Miller and Jemal, 2016). Early screening increases the likelihood to detect a cancer at a more local stage, thereby improving survival.

The most common method of early detection is mammography, low dose X-ray imaging of the breasts used to identify abnormalities. In light of evidence from clinical trials indicating that screening mammography reduces mortality by detecting tumors at an earlier stage (Duffy and Paci, 2012), expert organizations, including the World Health Organization's International Agency for Research on Cancer and the American Cancer Society, recommend regular, biennial screening mammograms starting at age 50 (Perry et al., 2008). Nearly every European country has established a national breast cancer screening program that make mammograms available free of charge for women in the recommended age range (Altobelli and Lattanzi, 2014).

Despite the central role that organized screening programs play in national cancer prevention strategies, there is surprising little research evidence on their impact. A basic question is whether organized programs are effective in increasing the number of women who receive mammograms. We examine this question using data from France. The first organized programs in the country were introduced in the late 1980s and early 1990s at the level of the *département*.<sup>1</sup> In 2004, France established a national program that built on and strengthened the early local programs in several ways. In particular, the target age range was expanded slightly and the recommended frequency for mammography screening was increased to once every two years from once every three years. By 2009, France's organized breast cancer screening program was the largest in the European Union in terms of the number of mammogram performed each year (Séradour, 2010). We evaluate the impact of both the national program and the earlier local programs using data from a nationally representative survey of the French population spanning the period 2000 to 2010 and difference-in-differences models that exploit variation in exposure to organized screening related to age, geography and time.

Numerous studies from multiple countries have documented significant disparities related to socioeco-

---

<sup>1</sup>The *département* is the fundamental administrative and political jurisdiction in France. There are 95 *départements* in "metropolitan France", with an average land area of 5,666 square kilometers (3.5 times the median land area of US counties). In terms of population, in 2015 *départements* varied in size from 76,000 to 2.6 million, with an average population of 670,000.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

economic status in the use of a wide range of preventive health services, including cancer screening (Schueler, Chu and Smith-Bindman, 2008; Pruitt et al., 2009; Jusot, Or and Sirven, 2012; Carrieri and Wubker, 2013; Devaux, 2015). An important objective of population-based screening programs is to reduce such disparities in utilization, thereby reducing disparities in cancer detection and survival. We consider whether France's organized screening program has reduced disparities in screening by testing for heterogeneous treatment effects related to education and income.

Our results indicate that the early local programs increased the percentage of women who had a mammogram in the past two years by roughly 6 percentage points, a 14 percent effect relative to the rate in areas without an organized program. We estimate that the effect of the national program that went into effect in 2004 was nearly twice as large. Although the percentage of women reporting a recent mammogram increased for all ages within the target range, the change was greatest among women in their sixties. This result combined with the fact that we find a weaker effect of the program on the probability of *ever* having had a mammogram, suggests that the program has increased not only initiation to mammogram use, but also the regularity with which women obtain screening. Because we find that France's organized screening program had a larger effect on mammography rates for women with lower socioeconomic status, gradients with respect to education and income have declined since the national program has been in place.

## 2 Background and Previous Literature

### 2.1 Breast Cancer Screening

Breast cancer screening typically includes a clinical breast exam, a physical examination of the breast by a physician or nurse, and a mammogram, which is an X-ray of the breast tissue that provides detailed images of the breast from 2 angles (frontal and profile). The mammogram is performed and analyzed by a radiologist. Screening mammograms are given to asymptomatic women to look for suspicious markers. Diagnostic mammograms are typically given to women who have had a previous abnormal screening mammogram, have a family history of breast cancer, or have certain symptoms, such as the presence of lumps.

The accuracy of a mammogram depends on the density of the breast tissue, which tends to decline with age. For women over age 50, the sensitivity of the test (the probability of detecting an existing cancer) ranges between 66 and 90 percent and specificity (the probability that the test is negative for someone without the disease) reaches 95 percent (Smith, 2003). Thus, mammography is seen as an effective means of detecting breast tumor for women older than 50 years old. It is considered less appropriate for younger women whose

1  
2  
3  
4  
5 breast tissue tends to be denser.

6  
7 Several studies based on randomized control trials have concluded that mammography screening leads to a  
8 significant reduction in breast cancer mortality (Duffy and Paci, 2012). These results led to recommendations  
9 by medical societies and public agencies that women receive regular mammograms. Reviewing evidence from  
10 a number of micro-simulation studies, Cutler (2008) concludes that nearly half of the reduction in breast  
11 cancer mortality in the U.S. between 1990 and 2004 can be attributed to increased screening. Recently,  
12 however, there has been some debate about how the benefits of mammography related to reduced mortality  
13 should be weighed against the cost associated with false positive results and overdiagnosis possibly leading  
14 to overtreatment (Gøtzsche and Nielsen, 2009). In 2009, the U.S. Preventive Services Task Force (USPSTF)  
15 generated some controversy when it revised its guidelines to recommend that women begin to receive regular  
16 mammograms starting at age 50, rather than 40 as it had previously recommended. The USPSTF also revised  
17 its recommendation regarding the frequency of screening mammograms from annual to biennial. In 2012, a  
18 U.K. expert panel conducted a meta-analysis of the randomized trial evidence and concluded that screening  
19 mammograms reduced the relative risk of mortality by 20 percent, while acknowledging the problem of  
20 overdiagnosis (Marmot et al., 2012). The panel's conclusions supported the NHS policy of organized breast  
21 cancer screening beginning at age 50.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

## 33 **2.2 The Effect of Public Policies on Screening**

34  
35 Most countries have policies aimed at increasing the number of women who receive a regular mammogram.  
36 In the U.S., with its fragmented system of health care financing, different strategies have been targeted at  
37 women with different types of insurance coverage. Between the late 1980s and early 2000s, nearly every  
38 state enacted laws requiring private health insurance plans to include screening mammograms as a covered  
39 benefit. Recent research indicates that these benefit mandates significantly increased the percentage of  
40 women obtaining a mammogram, with especially large effects occurring when plans were prohibited from  
41 charging cost-sharing for the service (Bitler and Carpenter, 2016a).<sup>2</sup> The requirement that private insurance  
42 plans provide first-dollar coverage for screening mammograms became national policy as a result of the  
43 Affordable Care Act. A different U.S. policy, the National Breast and Cervical Cancer Early Detection  
44 Program (NBCCEDP), aims to increase cancer screening among uninsured low-income women. Although  
45 it is a Federal program, the NBCCEDP was rolled out incrementally by different states throughout the  
46  
47  
48  
49  
50  
51  
52

53  
54 <sup>2</sup>A related literature from the U.S. examines the effect of insurance coverage on mammography use. Busch and Duchovny  
55 (2005) and Finkelstein et al. (2012) find a positive effect of Medicaid. Several recent studies examine the effect of Massachusetts'  
56 2006 health reform on mammography use and obtain mixed results (Kolstad and Kowalski, 2012; Keating et al., 2013; Sabik  
57 and Bradley, 2016).  
58  
59  
60

1  
2  
3  
4  
5 1990s. Research exploiting that implementation pattern finds that this program also significantly increased  
6 the percentage of women obtaining mammograms (Adams et al., 2003; Bitler and Carpenter, 2016b).  
7

8 In Europe, breast cancer prevention strategies have centered on organized screening programs (Altobelli  
9 and Lattanzi, 2014). The earliest programs, established at the local level, date to the 1980s. Finland,  
10 Luxembourg, the UK and Sweden expanded their programs to the national level in the early 1990s. In 2003,  
11 the European Commission formally recommended population-based screening for women between the ages  
12 of 50 and 69. Soon thereafter, nearly every European country had established a national organized breast  
13 cancer screening program. Although the exact details vary, these national programs share several common  
14 elements. The modal program targets 50 to 69 year-olds.<sup>3</sup> Every two (or a few countries three) years, women  
15 in the target age range receive a letter inviting them to receive a free mammogram. The standard protocol  
16 involves two images per breast and double reading of normal mammograms.  
17  
18

19  
20  
21  
22  
23 Two recent studies evaluate the effect of organized breast cancer screening programs on the utilization  
24 of mammography, using difference-in-differences research designs. Pletscher (2017) analyzes the case of  
25 Switzerland, where a program targeted at women between the ages of 50 and 69 was rolled out over several  
26 years at the level of the canton. This pattern of implementation provides variation related to geography, age  
27 and time. However, it is not clear that this variation is orthogonal to other factors affecting cancer screening.  
28 According to Pletscher (2017), cantons that did and did not implement screening programs differed in terms  
29 of breast cancer incidence, treatment patterns, the supply of radiologists and patient preferences. In addition,  
30 the survey data he uses has an important limitation: the only question related to mammography use asks  
31 if female respondents had *ever* had a mammogram. Thus, to the extent that organized screening affects  
32 not only initiation to mammography but also the regularity of screening, his analysis will understate the  
33 impact of the policy. Pletscher (2017) finds that organized screening is associated with a 4.6 percentage  
34 point increase in the probability of ever having had a mammogram.  
35  
36  
37  
38  
39  
40  
41  
42

43 Carrieri and Wuebker (2016) use data from the Survey of Health, Ageing and Retirement in Europe  
44 (SHARE) to estimate cross-sectional difference-in-differences models that compare women from 13 different  
45 countries who were exposed to organized screening to those who were not exposed because of their age or  
46 geographic location. Their dependent variable is an indicator for whether a woman had a mammogram in  
47 the past two years. One limitation of their research design is that the geographic variation is measured at  
48 the level of the NUTS-2 region, a statistical unit that does not map directly to political or administrative  
49  
50  
51  
52  
53

54 <sup>3</sup>In addition to France, the Netherlands extends the target age range to 74. In Ireland and Estonia, the upper age limit is  
55 64 and 65, respectively. In the Czech Republic, Hungary and Portugal, organized screening begins at age 45. Eligibility for the  
56 national program of Austria and Sweden start at 40 years old.  
57

1  
2  
3  
4  
5 divisions, and therefore does not necessarily correspond to the geographic area covered by an organized  
6 screening program.<sup>4</sup> For example, NUTS-2 regions in Switzerland are larger than cantons, the level at where  
7 the organized screening programs analyzed by Pletscher (2017) were implemented. Similarly, the NUTS-2  
8 regions for France encompass multiple *départements*. Thus, it is likely that some women in the treatment  
9 group were not actually exposed to an eligible screening program, while some women in their control group  
10 were. And, even more than in the case of Switzerland's cantons, differences in health care financing and  
11 delivery among European countries raise concerns about the comparability of this study's treatments and  
12 controls. Their main estimates imply that organized screening programs raise the probability that a women  
13 in the target age range has had a mammogram in the past 2 years by roughly 17 percentage points.  
14  
15  
16  
17  
18  
19

20 Numerous studies document significant socioeconomic disparities in breast cancer screening. Jusot, Or  
21 and Sirven (2012) and Carrieri and Wubker (2013) use SHARE data to examine the socioeconomic correlates  
22 of several types of preventive care. Both studies find that, controlling for other factors, screening rates  
23 increase significantly with education and income. Significant disparities in mammography use have also  
24 been documented in the U.S. (Sabatino et al., 2008; Lantz et al., 2006; McMorrow, Kenney and Goin, 2014)  
25 and Canada (Katz and Hofer, 1994; Katz, Zemencuk and Hofer, 2000). In a comparative study of 19 OECD  
26 countries, France ranked fourth in terms of income-related inequalities in mammography use (Devaux, 2015).  
27  
28  
29  
30

31 If cost is an important reason that lower income women make less use of preventive care, organized  
32 programs that make mammograms available free of charge should reduce disparities in utilization. Similarly,  
33 to the extent that disadvantaged women have low rates of breast cancer screening because they are not  
34 well-informed about the benefits of mammography, the information that is provided as part of an organized  
35 program can also reduce disparities. On the other hand, more educated, higher income women may be more  
36 efficient users of health inputs, and therefore may be more responsive to the reduced cost and informational  
37 intervention associated with an organized screening program. Thus, it is possible that an organized program  
38 could amplify rather than reduce disparities in mammography use related to socioeconomic status (Goldman  
39 and Lakdawalla, 2001).  
40  
41  
42  
43  
44  
45

46 The existing evidence on the effect of public policies on disparities in mammography utilization is limited  
47 and mixed. Adams, Breen and Joski (2007) find that although the NBCCEDP increased breast and cervical  
48 cancer screening in the U.S., the program did not reduce disparities in screening outcomes related to race  
49 or ethnicity. Similarly, in his analysis of organized screening in Switzerland, Pletscher (2017) tests for  
50 heterogeneous effects related to education and income and finds no clear pattern with respect to either. In  
51  
52  
53

54  
55 <sup>4</sup>NUTS stands for Nomenclature of Units for Territorial Statistics. The system was developed by Eurostat for developing,  
56 collecting and analyzing harmonized statistics at the sub-national level.  
57



1  
2  
3  
4  
5 contrast, Carrieri and Wuebker (2016) find that organized programs have a larger effect on women with the  
6 lowest levels of education compared to those with medium or high levels of education.  
7  
8

### 9 10 **2.3 Organized Breast Cancer Screening in France**

11  
12 As in other European countries, organized screening in France began in the 1980s at the local level. Between  
13 1989 and 1998, programs were launched in 27 *départements*; by 2000, there were 32 local programs. These  
14 *départements*, which are shown as the darkest ones on the map (Figure 1), are scattered throughout the  
15 country. This effectively random geographic dispersion of local programs and the fact that France's health  
16 care system is highly centralized, with uniform benefits and reimbursement policies nationwide, minimizes the  
17 concern that the distribution of these local programs was correlated with other factors that likely influenced  
18 the demand for or supply of mammography.  
19  
20  
21  
22

23  
24 The national program was announced in 2003 and by 2004 was operational throughout France. The  
25 current system is financed at the national level by the public health insurance funds and is overseen by the  
26 national cancer institute, though the operation of the program is still managed at the level of the *département*,  
27 typically by a non-profit organization.  
28  
29

30  
31 The protocol for the early programs involved a single image, which was read by two radiologists. The  
32 national program is based on a new protocol, which includes two images per breast and a clinical breast exam.  
33 Whereas the local programs sent invitation letters every three years to women between the ages of 50 and  
34 69, under the national program women between the ages of 50 and 74 receive an invitation letter every two  
35 years. This letter provides information on the benefits of mammography in general as well as the specific  
36 advantages of the organized program, such as the second reading, and a list of radiologists participating  
37 in the program. Participating radiologists must undergo a specific training, perform a minimum of 500  
38 mammograms per year and agree to have their equipment inspected by a national agency. The goal of these  
39 requirements is to assure a standard level of quality throughout the country. For women who accept the  
40 invitation to the organized program, there is no charge for the mammogram. Currently, the amount that the  
41 public insurance system pays to the radiologists for a mammogram (the "standard tariff") is 66.42 euros.  
42  
43  
44  
45  
46  
47

48  
49 Whether or not a woman receives a letter of invitation, she can screen "opportunistically" by obtaining  
50 a prescription from a physician (usually a gynecologist or a general practitioner). In that case she would  
51 face some out-of-pocket expenses. Women screening opportunistically would need to pay the standard tariff  
52 upfront, though 70 percent of it would be reimbursed by the public health insurance system and part or all  
53 of the remaining 30 percent would be reimbursed by private complementary insurance. Some radiologists are  
54  
55  
56  
57



1  
2  
3  
4  
5 allowed to charge more than the standard tariff. It is estimated that in 2008 the physician's fee exceeded the  
6 standard tariff for over 80 percent of the mammograms performed outside of the national program, with an  
7 average extra-billing amount of 6 euros (Haute Autorité de Santé, 2011). This extra-billing is not reimbursed  
8 by the public system, though part may be reimbursed by private complementary insurance, depending on  
9 the contract. Mammograms done outside the organized program are not necessarily read by two radiologists  
10 and need not be combined with a clinical breast exam.

11  
12  
13  
14  
15 Thus, there are multiple channels by which the organized program may increase screening. First, there  
16 may be an educational or informational effect: the letter informs women about the benefits of mammmog-  
17 raphy and provides information about where they can obtain a mammogram. Second, the program lowers  
18 the financial cost of screening. Third, the program may also increase utilization by affecting perceptions of  
19 the quality of the program relative to opportunistic screening.<sup>5</sup>

20  
21  
22  
23  
24 With the program's two-year invitation cycle, women in the target age range receive an invitation letter  
25 20-22 months after their last mammogram received through the program (if they took up the previous offer)  
26 or their last invitation letter (if they did not). In the absence of an organized program, there would be no such  
27 follow-up for women who had obtained a mammogram by getting a prescription from their doctor. Thus, an  
28 organized screening program may not only affect the likelihood that a woman ever receives a mammogram,  
29 but the regularity with which she does. A recent study by Goldzahl and Jusot (2017) is suggestive of such  
30 an effect. They find that women who participate in the organized program are more likely than women who  
31 obtained a mammogram outside the program to receive a second mammogram within two years.

### 3 Data and Methods

#### 3.1 Data: The ESPS

32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43 Our analysis is based on data from the *Enquete Santé et Protection Sociale* (ESPS), a representative  
44 population-based survey that has been conducted biennially since 1988 by the *Institute de Recherche et*  
45 *Documentation en Economie de la Santé* (IRDES). The ESPS is administered to a sample of French house-  
46 holds randomly drawn from public health insurance files. The survey provides information on both the  
47 household and its members collected using both interviews (telephone or face to face) and self-administered  
48  
49  
50  
51

---

52  
53  
54  
55  
56  
57  
58  
59  
60  
<sup>5</sup>A number of studies find a positive relationship between the volume of mammograms that a physician interprets and accuracy, both in terms of sensitivity and specificity (Esserman et al., 2002; Smith-Bindman et al., 2005). Other research finds that double reading of mammograms significantly increases cancer detection rates (Thurfjell, Lernevall and Taube, 1994; Kopans, 2000). A qualitative study of target-age women in France found that those who had participated in the organized program viewed the second reading as an indicator of quality (Kalecinski et al., 2015).

1  
2  
3  
4  
5 questionnaires.<sup>6</sup> We use data from the 2000, 2002, 2006, 2008 and 2010 surveys, which are the only years in  
6  
7 which the survey includes questions on mammography. The ESPS has a longitudinal component, whereby  
8  
9 half of the sample is interviewed every 4 years. Thus, some respondents were interviewed two or three times  
10  
11 in the five survey waves that we analyze. In our main analysis we treat the sample as a repeated cross-section.  
12  
13 As a sensitivity check, we also use the panel dimension to estimate models with individual fixed effects.

14  
15 The timing of the surveys is well-suited for our study for several reasons. First, we have two periods of  
16  
17 data before and three periods after the implementation of the national screening program in 2004. Second,  
18  
19 all but one of the local programs that pre-dated the national program were in place by 2000.<sup>7</sup> This facilitates  
20  
21 clear cross-sectional comparisons during the “pre” period and simplifies the implementation of the difference-  
22  
23 in-differences model to estimate the effect of the national program. Third, the fact that our first year of  
24  
25 “post” data is two years after the program was put in place nationally, means that nearly all women in  
26  
27 the target age group should have received a letter of invitation by the time they responded to the survey.  
28  
29 Fourth, the four-year gap between our last year of data pre-dating the national program and our first year  
30  
31 of “post” data means that our estimates should not be affected by short-run dynamic effects around the  
32  
33 implementation year. That is, our estimates should not be subject to an upward bias caused by women  
34  
35 delaying a mammogram they otherwise would have had in 2003 in order to benefit from the program or by  
36  
37 a downward bias from a “learning curve” in the initial year of the program.

38  
39 There are two timing issues that could potentially cause the effect of the local program to be different  
40  
41 in 2000 and 2002. The first is that 8 of the 32 early local programs were started in 1998 or later. So, while  
42  
43 the program was in place by 2000, some eligible women may not have received their invitation letter by the  
44  
45 time they completed the 2000 ESPS. Second, some *départements* with local programs began transitioning to  
46  
47 the new protocol in 2001 and 2002. Both of these factors could result in the program having a larger impact  
48  
49 in 2002 than in 2000. However, we find no evidence of such a difference (as shown in table A.1). Therefore,  
50  
51 we pool the data from 2000 and 2002, treating it as a single “pre” period relative to the introduction of the  
52  
53 national program.

54  
55 As in previous studies on mammography use (Sabik and Bradley, 2016; Pletscher, 2017; Carrieri and  
56  
57 Wuebker, 2016; Bitler and Carpenter, 2016*a,b*) our dependent variables are based on self-reports. In the first  
58  
59 two waves, the ESPS asks women whether they ever had a mammogram. Possible answers are Yes, in the  
60  
past two years ; Yes, between two and three years ago ; Yes, more than three years ago ; No, never. We

---

<sup>6</sup>More information on the ESPS, including questionnaires for every year, is available at <http://www.irdes.fr/recherche/enquetes/esps-enquete-sur-la-sante-et-la-protection-sociale/actualites.html>.

<sup>7</sup>The exception was the Essonne, a *département* in the greater Paris region, which established its program in January 2000.

1  
2  
3  
4  
5 create a binary variable that equals who screened in the past two years. In the next three waves, the ESPS  
6 asks women whether they have ever had a mammogram. Those answering yes are then asked to give the  
7 date of their last mammogram. Based on these responses, we create a binary variable that equals one for  
8 women who have had a mammogram in the past two years and zero for all others (including both women  
9 who have never had a mammogram and women whose last mammogram was more than two years earlier).  
10 Although the mammography question is asked of all female respondents over age 15, mammography rates  
11 for younger women are extremely low. Therefore, we limit our main analysis to women age 35 and older.  
12

13  
14  
15  
16  
17 Table 1 presents summary statistics for the full sample and for women living in *départements* with and  
18 without a local organized screening program respectively in the pre and post-period. As we describe below,  
19 our estimate of the effect of the local program relies on a cross-sectional comparison of women in *départements*  
20 with and without such programs. Thus, the validity of our research design relies on the assumption that  
21 women in *départements* without a local program are good controls for women in *départements* with a program.  
22 The summary statistics support this assumption. In the pre-period, the two subsamples are remarkably  
23 similar in terms of observable characteristics. The mean age is 54.4 years old in *départements* that had an  
24 organized screening program and 55.1 years old in those that did not. In each group, just under 5 percent  
25 of women report their health as poor, 17 percent of women report their health as fair, and 16 percent  
26 say they are in excellent health. Between 13 and 14 percent of women in each type of *département* are  
27 current smokers. The percentage with private complementary health insurance—which has been shown to  
28 be positively correlated with the utilization of outpatient care (Buchmueller et al., 2004)—is very similar  
29 (87.1 percent in *départements* with a local program, 88.2 percent in those without) as is the distribution of  
30 income. The one notable difference is that *départements* with local programs tend to be less rural. Despite  
31 this, there is not a meaningful difference in the supply of services, as measured by the density of radiologists  
32 per capita. The two groups were also quite similar in terms of the breast cancer mortality rate for the  
33 years 1980 to 1988: in the average *département* with a local program the rate was 17.3 deaths per 100,000  
34 population compared to 16.5 per 100,000 in the average *département* without a local program.<sup>8</sup>  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

### 48 3.2 Empirical Strategy

49  
50 The quasi-experimental variation in our data allows for several different estimates of the effect of organized  
51 breast cancer screening programs on mammography use. Let  $T_t^k$  be the screening rate for women in the  
52

53  
54 <sup>8</sup>We use an averaged mortality rate data from the year 1980 to 1988. This is before local programs implementation. The  
55 reason is that we wanted to control for potential pre-local program implementation mortality differences between *départements*  
56 in case those differences explain why some *départements* launched a local program or not. GP and radiologist density rates  
57 were collected for the years 1998 to 2008 which is contemporaneously with surveys. The data comes from the Eco-Santé website  
58

1  
2  
3  
4  
5 target age range (50 to 74), and  $C_t^k$  be the screening rate for women outside this age range, i.e., the “control”  
6  
7 group. The superscript  $k$  indexes two types of *départements*: those with ( $L$ ) and without ( $N$ ) local screening  
8  
9 programs. The subscript  $t$  indexes two time periods. The “pre” period ( $t=0$ ) consists of the years 2000 and  
10  
11 2002, before the national program was established; the “post” period ( $t=1$ ) consists of data from 2006 to  
12  
13 2010, when the national program was in place in all *départements*.

14 Using this notation, a simple cross-sectional estimate of the average effect of the pre-existing local pro-  
15  
16 grams can be obtained by comparing the screening rates for the target age group in the two types of  
17  
18 *départements*:

$$19 \quad \Delta_L = T_0^L - T_0^N. \quad (1)$$

20  
21  
22 As just noted, *départements* with and without local programs are quite similar in terms of observable  
23  
24 characteristics. However, unobserved factors remain a potential source of bias. One can imagine the bias  
25  
26 going in either direction. On one hand, the “first mover” *départements* may be those with good referral  
27  
28 networks and radiologists with a strong orientation toward population screening. On the other hand, some  
29  
30 *départements* may have established organized programs in response to low rates of opportunistic screening.  
31  
32 If these other factors have the same effect on women within and outside the target age range, this potential  
33  
34 bias can be eliminated by using a cross-sectional difference-in-differences estimator:

$$35 \quad \Delta\Delta_L = (T_0^L - T_0^N) - (C_0^L - C_0^N). \quad (2)$$

36  
37  
38 The effect of the national program can be estimated by calculating changes in screening rates for target  
39  
40 group women who were not exposed to the earlier local programs, using changes for control group women  
41  
42 in the same *départements* to account for the effect of other factors that may have caused screening rates to  
43  
44 change over time. This difference-in-difference estimate can be calculated as

$$45 \quad \Delta\Delta_N = (T_1^N - T_0^N) - (C_1^N - C_0^N). \quad (3)$$

46  
47  
48 Given that the national program put in place in 2004 is stronger in several respects than the pre-existing  
49  
50 local programs, it is of interest to test whether screening rates increased in those areas that had local  
51  
52 programs in place in the “pre” period. The difference-in-difference estimate  
53  
54

$$55 \quad \Delta\Delta_{LN} = (T_1^L - T_0^L) - (C_1^L - C_0^L) \quad (4)$$

captures the combined effect of those specific elements of the national program that differ from the local programs: the use of a 2-year rather than 3-year invitation cycle, the extension of program eligibility to age 74 rather than 69, two images per breast are taken instead of one, and the fact that women receive a clinical exam in addition to a mammogram.

Because we have no data from years before the local programs were established, we cannot directly estimate the full effect of the national program in these *départements*. However, we can construct an estimate of that effect by summing the cross-sectional estimate of the effect of the local program,  $\Delta_L$ , with our estimate of the effect of moving from the local program to the national program  $\Delta\Delta_{LN}$ . Note that comparing this measure with our direct estimate of the effect of the national program,  $\Delta_N$ , provides a useful test of the assumption that the two types of *départements* are similar, apart from when they implemented their programs. If this assumption holds, the two estimates of the effect of the national program should be similar.

These various estimates can be obtained from the following regression estimated on pooled data:

$$y = \alpha_1 Local + \alpha_2 Target + \alpha_3 Post + \beta_1 Local \times Target + \beta_2 Local \times Post + \beta_3 Target \times Post + \gamma Local \times Target \times Post + X'\theta + \epsilon \quad (5)$$

In this equation *Local* is an indicator variable that equals one for women who live in a *département* that had a local screening program before 2004, *Target* is an indicator for women between the ages of 50 and 74, the indicator *Post* equals one for years after 2004, and  $X$  is a vector of control variables. The various estimates are then:

Effect of local program:	$\Delta\Delta_L =$	$\beta_1$
Effect of national program:	$\Delta\Delta_N =$	$\beta_3$
Effect of transition from local to national program:	$\Delta\Delta_{LN} =$	$\beta_3 + \gamma.$

### 3.3 Testing for Parallel Trends

Our strategy for estimating the effect of the national program relies on the assumption that the post-2004 experience of women outside the target age range represents an appropriate counterfactual for women that were targeted by the program. A standard way to test that assumption is to examine whether outcomes for the two groups were trending similarly before the policy was put in place. The fact that we have only two

years of data from the pre-period limits our ability to analyze pre-trends. Nonetheless, it is informative to test for differences in changes in mammography utilization between 2000 and 2002. A finding that utilization among target age women was already increasing relative to the control group before the national program was in place would suggest that our difference-in-differences estimates may overstate the program's effect.

Appendix Table A.1 presents the change in mammography utilization between 2000 and 2002 for the four categories of women defined by target group status and *département* type. We see that for each category, there was no significant change in utilization over this period. The most important comparison is between treatments and controls in *départements* without a pre-existing program, since these are the observations upon which our main estimates of the national program are based. Within these *départements*, the screening rate fell by a statistically insignificant one percentage point for women in the target age range (p-value = 0.535) and increased by a statistically insignificant 0.3 percentage points for the control group (p-value = 0.881). The difference between these two estimates is also not significantly different from zero (p-value = 0.741). While estimates based on only two years of data from the "pre" period do not provide definitive evidence of parallel trends, these null results, along with several other findings discussed below, provide support for our empirical strategy.

## 4 Results

### 4.1 Unadjusted Difference-in-Differences Estimates

Table 2 presents the percentage of women who had a mammogram in the past two years tabulated by membership to the target age group of 50 to 74 years old, residence in a *département* with a local program, and the years before or after the implementation of the national program. These figures can be used to calculate the various estimates just described.

Panel A reports data from the "pre" period (2000 and 2002). The first row pertains to women who are outside the target age range and therefore not eligible for organized screening regardless of where they live. A comparison of mammography rates for this group in the two types of *départements* provides information on whether there are other factors affecting mammography that are correlated with the existence of a program. The fact that the difference is not statistically significant, suggests that other confounding factors are not a concern. As a result, when we compare mammography rates across *département* types for women in the target group, the simple difference,  $\Delta_L$ , and the difference-in-differences estimate of the program effect,  $\Delta\Delta_L$  are essentially identical. Both imply that the local programs raised mammography rates among women in

1  
2  
3  
4  
5 the target age range by roughly 6 percentage points, or by about 14 percent of the rate in *départements*  
6 without an organized screening program.  
7

8 Panel B presents screening rates after the national program was in effect. The comparisons across  
9 *département* types again suggest a strong similarity between those with and without an organized screening  
10 program in place before 2004. As in the earlier period, there is no statistically significant difference between  
11 the two types of *départements* among women outside of target age range. The results for women in the  
12 target age range indicate that the implementation of the national screening program eliminated the gap in  
13 screening rates between *départements* with and without local program. In the “post” period, 55.3 percent  
14 of women in *départements* with an already established local program and 56.5 of women who were not  
15 exposed to organized screening prior to 2004. The 1.2 percentage point difference between these two rates is  
16 not statistically significant. The fact that screening rates for target group women were essentially identical  
17 when the policy environment was the same provides additional support for interpreting the cross-sectional  
18 difference in the pre-period as representing the effect of the local programs.  
19  
20  
21  
22  
23  
24  
25

26 The effect of the national program can be estimated based on changes over time, which are reported in  
27 Panel C of the table. Again, it is informative to note the differences that are not statistically significant.  
28 In both types of *départements* there was no significant change for women outside the targeted age range,  
29 suggesting that there were not other contemporaneous factors, such as a general increase in breast cancer  
30 awareness or a change in clinical guidelines that might have caused screening to increase over time. In  
31 contrast, we see significant changes for women who were eligible. For women in the target group who  
32 were not previously exposed to a local organized screening program (column 1), the percentage with a  
33 mammogram in the past two years increased by 13.8 percentage points, a 32 percent effect relative to the  
34 baseline rate. Since there was no significant change for women outside the target group, the difference-in-  
35 differences estimate of the effect of the program is essentially identical. The fact that this estimated effect  
36 is substantially larger than the estimated effect of the local programs suggests that the ways in which the  
37 national program strengthened the local programs had a meaningful effect. This conclusion is reinforced  
38 by the finding that screening rates increased over time in *départements* where there was already a local  
39 organized screening program in place. The difference-in-differences estimate of the effect of this change is  
40 6.9 percentage points ( $\Delta\Delta_{LN} = 0.063 + 0.006 = 0.069$ ).  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57



## 4.2 Basic Regression Results

In Table 3 we report key results from linear probability regressions that generate the same set of estimates. In the upper panel of the table we report estimated coefficients and robust standard errors (clustered by *département*). In the lower part of the table, we present the three difference-in-differences estimates of interest. The model reported in column 1 includes no covariates and therefore corresponds directly to the results in Table 2. The specifications in column 2 adds the following covariates: age, age squared, age cubed, quintiles of household income, self-reported health status (5 categories), education (5 categories), current or former occupation (7 categories), complementary health insurance status (3 categories) and indicator variables for smoking, having at least one chronic health condition, living in a rural area and living in the greater Paris region. We also control for radiologist density, GP density and the rate of mortality from breast cancer by *département*.

For the most part, including covariates does not have a major impact on the coefficients on the variables of interest. One important exception is the coefficient on the indicator for being in the target group, which is large and statistically significant in column 1 (0.177,  $p=0.000$ ) and is a precisely estimated zero in the adjusted model ( $p=0.999$ ). This result provides support for our difference-in-differences approach as it indicates that in the absence of an organized program there is not a discrete change in the probability of having a recent mammogram associated with turning 50. In other words, our estimates of the effect of organized screening are not capturing the effect of other factors, such as age-based clinical guidelines.

The full regression results are reported in Appendix Table A.2. In addition to the positive effect of age, the estimated coefficients on the other control variables are consistent with results from previous studies on the determinants of mammography use.<sup>9</sup> Women with private complementary insurance are more likely to have a recent mammogram than women who either have no complementary coverage or are covered by means-tested public insurance. Current smokers are significantly less likely than non-smokers to have had a recent mammogram, a result that may reflect preferences regarding risk and preventive medical care. There appears to be an inverse U-shaped relationship with self-reported health. The probability of having a mammogram in the past two years is highest for women who report their health as good and lower for those who say it is fair, poor or excellent. The results for education and income indicate a positive gradient with respect to socioeconomic status, an issue to which we return below.

Turning to the coefficients that represent the effect of organized screening program, we see that the

---

<sup>9</sup>Schueler, Chu and Smith-Bindman (2008) provide a comprehensive review of studies using data from the U.S. For a recent analysis of the correlates of mammography use in France, see Sicsic and Franc (2014) and Goldzahl and Jusot (2017).

1  
2  
3  
4  
5 adjusted differences-in-differences estimates are qualitatively similar to those from the unadjusted model.  
6 According to the model reported in column 2, the local programs raised the percentage of target aged women  
7 having a mammogram in the past two years by 6.6 percentage points. Modifying those programs to conform  
8 with the new national protocols and standards raised the two-year mammography rate by an additional  
9 4.2 percentage points. Combining these two estimates implies that the national program had the effect of  
10 raising the mammography rate by 10.8 percentage points. This is only slightly smaller than the difference-  
11 in-differences estimate of the effect of the national program that is identified by changes in *départements*  
12 that did not previously have an organized program ( $\Delta\Delta_N=.122$ ).  
13  
14  
15  
16  
17

18 Columns 3 and 4 report results a different outcome: a binary variable that equals one if a woman has ever  
19 had a mammogram and zero otherwise. The results from the model with covariates (column 4) imply that  
20 both the local and national programs raised the probability of every having a mammogram by between 3 and  
21 4 percentage points. These estimates are similar to the effects that Pletscher (2017) finds for Switzerland.  
22 The comparison between the two outcomes is informative as to the margins on which organized screening  
23 programs operate. The fact that we find a significant effect on ever having a mammogram but larger effects  
24 on the probability of screening in the past two years suggests that these programs have important effects on  
25 both initiation and the regularity with which women screen.  
26  
27  
28  
29  
30  
31  
32

### 33 4.3 Alternative Samples

34  
35 Table 4 reports key results from models using alternative estimation samples. For ease of comparison, the  
36 covariate-adjusted results from column 2 of Table 3 are presented in the first row. In row 2 we exploit the  
37 limited longitudinal feature of the ESPS data by limiting the sample to respondents who appear more than  
38 once in the data and estimating a model with individual fixed effects. The estimated program effects from  
39 this model are comparable to our main estimates.  
40  
41  
42

43 The next three rows illustrate the impact of changing the age criteria for the estimation sample or the  
44 target group. Because very young and very old women may not be good controls for women in the target  
45 age range, in row 3 we limit the sample to women between the ages of 40 and 80. In row 4, the sample  
46 includes all women 35 and older with the exception of those who are 50 or 51 at the time of the survey.  
47 While these women were eligible for the program, some may have not yet received an invitation letter. These  
48 modifications also do not have a material impact on the estimates.  
49  
50  
51  
52

53 A particular feature of our research design is that the control group combines women who are too young  
54 to participate in the program and women who are too old. Women under the age of 50 are a “cleaner”  
55  
56  
57

control group as they should have never received a letter inviting them to have a mammogram through the program. In contrast, some women above the age of 74 will have participated in the program, especially in the later years. Additionally, to the extent that their recall is not perfect, some women who received a mammogram through the program, say, three or four years ago, may mistakenly say that they had one in the past two years. Thus, we would expect to find stronger effects of the program if we compare the target group only to women under age 50. Consistent with this expectation, our estimates of  $\Delta\Delta_L$  and  $\Delta\Delta_N$  are slightly larger than the corresponding estimates for the full sample, though again the difference is small.

#### 4.4 Accounting for Heterogeneous Effects by Age

Even though the covariate-adjusted models reported in Table 3 control flexibly for age, they restrict the impact of organized screening to be the same for all women in the target groups. In Figure 2 and Table 5 we allow the effect of the program to vary with age, stratifying by *département* type to account for baseline differences in policy. The figure uses two-year age bins; in the regression we interact the “post” variable with indicators for five-year age categories.

The results for *départements* without a pre-existing local program (panel A of the figure and column 1 of the table) provide the clearest evidence of the impact of the national program. When there was no organized screening program in place, mammography rates increased with age up to age 60 and declined thereafter. A comparison of the age profiles for the “pre” and “post” periods confirms that the implementation of the national program had no effect on mammography rates for women under the age of 50. This is visually apparent from the figure and can be seen by the fact that the coefficient on “post” (which represents the change over time for 35 to 40 year olds) and the interaction of that variable and indicators for the next two age categories (40-45, 45-49) are all insignificantly different from zero.

Mammography rates increased for all age groups within the 50 to 74 year-old target range, but the effect was not uniform. The percentage of 50 to 54 year-olds having a recent mammogram increased by just under 9 percentage points. After the national program was in place, mammography rates increased slightly with age up until around age 70 and then began to decline thereafter. Because mammography rates peaked at an earlier age during the “pre” period, the implied effect of the program is larger for women in their 60s and 70s than for women in their 50s. The screening rate increased by roughly 20 percentage points for women in their 60s and by 29 percentage points for women between 70 and 74.

There was also a statistically significant increase among women between the ages of 75 and 79, who are just outside the target age range. Given that the dependent variable equals one for women who had a

1  
2  
3  
4  
5 mammogram in the past two years, many of these women may have participated in the national screening  
6 program just before aging out of the target group. Indeed, a supplementary analysis suggests that this is the  
7 case. We ran separate regressions for 75–79 year-olds, allowing the effect to differ between 75–76 year-olds,  
8 who would have been eligible for the organized program during the two year look-back period, and 77–79  
9 year-olds, who would not have been. For the younger group, the probability of having a recent mammogram  
10 increased by 25 percentage points, which is comparable to the effect for 70–74-year olds. The increase was  
11 roughly half as large for 77–79 year-olds, though it was still statistically significant ( $p=0.005$ ). This effect  
12 for women just above the upper age threshold may reflect a true spillover effect. For example, after being  
13 exposed to the organized program women may continue to have a strong demand for screening even after  
14 they age out of the program. Alternatively, it could reflect measurement error: women may be understating  
15 the time since their last mammogram.

16  
17  
18  
19  
20  
21  
22  
23 In *départements* with pre-existing local programs, the largest changes were for women between the ages  
24 of 70 and 74, who are covered by the new protocol but were outside the old protocol's target age range. For  
25 women in their 50s and early 60s, the change over time is much smaller and is not statistically significant.  
26 This suggest that the expansion of the target range is a major reason for the observed difference between  
27 the national program and the earlier local programs.

#### 28 29 30 31 32 33 **4.5 Impacts on Disparities in Screening**

34  
35 Having established that France's organized breast cancer screening increased mammography use, we now turn  
36 to the question of whether it also reduced socioeconomic disparities in screening. Table 6 presents screening  
37 rates tabulated by education and income categories. Because we are interested in how screening changed  
38 among affected women after the national program was implemented, we focus on women in the target age  
39 range. To account for differences in baseline conditions, we present separate results by *départements* type.<sup>10</sup>

40  
41  
42  
43 Looking first at *départements* without pre-existing local programs, we see that before the national pro-  
44 gram was established there were strong gradients with respect to education and income. Women with a  
45 middle school education were 16 percentage points more likely to report having had a mammogram in the  
46 past 2 years than women with a primary school education or less. The rate was higher still for women with  
47 a high school education and those with university degrees. The ratio of the rate for the highest education  
48  
49  
50  
51

52  
53 <sup>10</sup>Some studies in the literature on health disparities focus on unadjusted differences among groups, while others condition  
54 on proxies for health need and preferences as well as, in some cases, other observable characteristics. Because we are interested  
55 in overall differences in mammography use among targeted women, we focus on unadjusted differences across education and  
56 income groups. For a useful discussion of methodological issues concerning the measurement and analysis of health-related  
57 disparities, see Cook, McGuire and Zaslavsky (2012).

1  
2  
3  
4  
5 category to the rate for the lowest category was 1.79. The percentage of women with a recent mammogram  
6 also increased monotonically with income, with a slightly larger ratio between the highest and lowest quin-  
7 tiles. After the national program was in place, mammography rates increased for all education and income  
8 categories, though the change was generally larger at the lower ends of each distribution. The changes were  
9 18.3 and 20.9 percentage points for the lowest education and income categories, compared to just over 6  
10 percentage points for the highest categories. As a result of this differential effect, the ratio between the top  
11 and bottom income quintiles fell from 1.99 to 1.35.  
12  
13  
14  
15

16  
17 In *départements* with pre-existing programs, the baseline gradients were less steep, which suggests that  
18 organized screening had already reduced socioeconomic disparities there. Here, we also see greater increases  
19 for women in lower education categories, though the differences were less pronounced than in *départements*  
20 without pre-existing programs. In the “post” period, the ratio of the mammography rates for the top and  
21 bottom education and income categories were similar in the two types of *départements*.  
22  
23  
24

25  
26 The 2006 and 2008 ESPS surveys asked not only whether a woman had a mammogram in recent years,  
27 but those who had a mammogram were asked whether or not they received it through the organized pro-  
28 gram. In the last column of Table 6 we present the percentage of women in each education or income  
29 category that participated in the program. For these calculations, the sample is limited to women who had  
30 a mammogram in the past 2 years. Because the overall utilization rate and the gradients are so similar  
31 across *département* types, we pool them. The results indicate a negative relationship between socioeconomic  
32 status and participation in the organized program. Seventy percent of women with a primary education or  
33 less who had a mammogram in the past two years received it through the program, compared to 53 percent  
34 of women with at least a university degree. Similarly, the percentage using the program ranged from 71.5  
35 percent in the lowest income quintile to 55.8 percent in the highest. Multiplying these percentages by the  
36 mammography rates for the post period, gives the percentage of women in each group participating in the  
37 program. This calculation indicates that participation was quite similar across groups and that the gradients  
38 that remain are driven by the fact that higher SES women are more likely to go outside the program to  
39 receive a mammogram.  
40  
41  
42  
43  
44  
45  
46  
47

48  
49 The finding that the organized program had a larger impact on lower SES women is consistent with results  
50 found by Carrieri and Wuebker (2016) and not with those of Pletscher (2017). One possible explanation is  
51 that Pletscher (2017) tests for effects on the probability of ever having a mammogram, which we know is less  
52 sensitive to the policy. When we replicate the analysis of Table 6 using this outcome, we see that screening  
53 increased most for women in the lowest education and income categories, causing the ratio of screening  
54  
55  
56  
57

1  
2  
3  
4  
5 rates for the highest and lowest categories to decline (results available upon request). Thus, it appears that  
6 France's organized screening program reduced socioeconomic disparities in mammography initiation.  
7  
8  
9

## 10 5 Discussion and Conclusion

11  
12 Our results provide strong evidence that French women who were eligible for a publicly funded organized  
13 screening program were significantly more likely to have had a recent mammogram than women who were  
14 not eligible. The fact that our estimates of the impact of the French program are similar to those found  
15 for other European countries (Carrieri and Wuebker, 2016; Pletscher, 2017) strengthens the conclusion that  
16 organized screening programs are successful in increasing the percentage of women obtaining a mammogram.  
17  
18

19 As with any difference-in-differences analysis, the interpretation of our results as causal effects depends  
20 importantly on untestable assumptions about the comparability of treatment and control groups. In our  
21 case, there are two critical comparisons, one that is cross-sectional and one related to changes over time.  
22  
23

24 Our cross-sectional estimates of the effect of the local programs that existed before 2004 relies on the  
25 assumption that women in *départements* without local programs are good controls for women in *départements*  
26 with such programs. Several results support this assumption. First, women in the two types of *départements*  
27 are quite similar in observable demographic, economic and health characteristics. Second, in the years 2000  
28 and 2002, differences in mammography use are observed *only* for women in the target age range. Outside this  
29 age group, we observe no significant difference in screening rates. Third, after 2004, when a uniform policy  
30 was in place throughout France, we see no difference in mammography utilization between *départements* that  
31 did and did not have local programs prior to 2004. Taken together, these factors suggest that the roughly  
32 6 percentage point difference in screening rates that we observe in that earlier period represents the causal  
33 effect of the local programs that existed in some *départements* but not others.  
34  
35

36 Our estimate of the effect of the national program is identified by comparing changes over time for women  
37 within and outside the target age range. Several factors support the validity of this identification strategy  
38 as well. We see no trend in mammography rates for women in either the target or control group between  
39 2000 and 2002. Moreover, after 2004 screening rates remained constant for the control group, suggesting the  
40 non-importance of secular trends in breast cancer awareness, technology diffusion or other factors affecting  
41 clinical practice. For the target group, however, there is a large and significant increase in screening after the  
42 national program was put in place. Our preferred estimate, based on a regression model that conditions on  
43 individual demographic, economic and health characteristics implies that the national program led to a 12  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5 percentage point increase in the screening rate, a 29 percent effect relative to the baseline mean. Regression  
6 models that allow for heterogeneous effects related to age provide further evidence that the increases in  
7 mammography utilization occurred only for women age 50 or older.  
8  
9

10 One important difference between our study and that earlier research is that we test for heterogeneous  
11 effects related to age. Interestingly, we find the largest effects for women in their mid-60s, who are in the  
12 middle of the target age range. This result combined with the fact that we find weaker effects on the  
13 probability on the probability of *ever* having a mammogram suggest that the most important effect of the  
14 program may not be on mammography initiation, but rather on the regularity of screening.  
15  
16  
17

18 By eliminating financial barriers and increasing awareness of the importance of screening, organized  
19 programs have the potential to reduce socioeconomic disparities in cancer screening. The implementation of  
20 the national program increased mammography rates for women of all education and income levels. However,  
21 because the increases were greater for women of lower socioeconomic status, the policy had the effect of  
22 reducing disparities in mammography utilization.  
23  
24  
25

26 While our results suggest that France's organized breast cancer screening program was successful at  
27 increasing mammography use, the ultimate goal of any such program it to reduce mortality through the timely  
28 detection of treatable tumors. Although the benefits of mammography have long been taken as given, the  
29 strongest empirical evidence comes from randomized clinical trials, not population-based screening programs.  
30 The extent to which the higher rates of mammography use generated by this program led to reductions in  
31 mortality is an empirical question that merits further research.  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57



## References

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- Adams, E Kathleen, Curtis S Florence, Kenneth E Thorpe, Edmund R Becker, and Peter J Joski. 2003. "Preventive care: female cancer screening, 1996–2000." *American journal of preventive medicine*, 25(4): 301–307.
- Adams, E Kathleen, Nancy Breen, and Peter J Joski. 2007. "Impact of the National Breast and Cervical Cancer Early Detection Program on mammography and Pap test utilization among white, Hispanic, and African American women: 1996–2000." *Cancer*, 109(S2): 348–358.
- Altobelli, E, and A Lattanzi. 2014. "Breast cancer in European Union: An update of screening programmes as of March 2014 (Review)." *International journal of oncology*, 45(5): 1785–1792.
- Bitler, Marianne P, and Christopher S Carpenter. 2016a. "Health insurance mandates, mammography, and breast cancer diagnoses." *American Economic Journal–Economic Policy*, forthcoming.
- Bitler, Marianne P, and Christopher S Carpenter. 2016b. "New Evidence on the Effects of the National Breast and Cervical Cancer Early Detection Program." Vanderbilt University.
- Buchmueller, Thomas C, Agnès Couffinhal, Michel Grignon, and Marc Perronnin. 2004. "Access to physician services: does supplemental insurance matter? Evidence from France." *Health economics*, 13(7): 669–687.
- Busch, Susan H, and Noelia Duchovny. 2005. "Family coverage expansions: impact on insurance coverage and health care utilization of parents." *Journal of health economics*, 24(5): 876–890.
- Carrieri, Vincenzo, and Ansgar Wubker. 2013. "Assessing inequalities in preventive care use in Europe." *Health policy*, 113(3): 247–257.
- Carrieri, Vincenzo, and Ansgar Wuebker. 2016. "Quasi-Experimental Evidence on the Effects of Health Information on Preventive Behaviour in Europe." *Oxford Bulletin of Economics and Statistics*, 78(6): 765–791.
- Cook, Benjamin Lê, Thomas G McGuire, and Alan M Zaslavsky. 2012. "Measuring racial/ethnic disparities in health care: methods and practical issues." *Health services research*, 47(3pt2): 1232–1254.
- Cutler, David M. 2008. "Are we finally winning the war on cancer?" *The Journal of Economic Perspectives*, 22(4): 3–26.

- 1  
2  
3  
4  
5 **Devaux, Marion.** 2015. "Income-related inequalities and inequities in health care services utilisation in 18  
6 selected OECD countries." *The European Journal of Health Economics*, 16(1): 21–33.
- 7  
8  
9 **Duffy, S, and E Paci.** 2012. "Bénéfices et risques du dépistage du cancer du sein par mammographie."  
10 *Bulletin épidémiologique hebdomadaire*, 35: 406–411.
- 11  
12  
13 **Esserman, Laura, Helen Cowley, Carey Eberle, Alastair Kirkpatrick, Sophia Chang, Kevin**  
14 **Berbaum, and Alastair Gale.** 2002. "Improving the accuracy of mammography: volume and outcome  
15 relationships." *Journal of the National Cancer Institute*, 94(5): 369–375.
- 16  
17  
18  
19 **Ferlay, Jacques, Isabelle Soerjomataram, Rajesh Dikshit, Sultan Eser, Colin Mathers, Marise**  
20 **Rebelo, Donald Maxwell Parkin, David Forman, and Freddie Bray.** 2015. "Cancer incidence and  
21 mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012." *International journal*  
22 *of cancer*, 136(5): E359–E386.
- 23  
24  
25  
26 **Finkelstein, Amy, Sarah Taubman, Bill Wright, Mira Bernstein, Jonathan Gruber, Joseph P**  
27 **Newhouse, Heidi Allen, and Katherine Baicker.** 2012. "The Oregon Health Insurance Experiment:  
28 Evidence from the First Year\*." *The Quarterly journal of economics*, 127(3): 1057–1106.
- 29  
30  
31  
32 **Goldman, Dana, and Darius Lakdawalla.** 2001. "Understanding health disparities across education  
33 groups." National Bureau of Economic Research.
- 34  
35  
36 **Goldzahl, Léontine, and Florence Jusot.** 2017. "Determinants of breast cancer screening regularity in  
37 France." *Revue française d'économie*, 31(4): 109–152.
- 38  
39  
40 **Gøtzsche, Peter C, and Margrethe Nielsen.** 2009. "Screening for breast cancer with mammography."  
41 *Cochrane Database Syst Rev*, 4(1).
- 42  
43  
44 **Haute Autorité de Santé.** 2011. "La participation au dépistage du cancer du sein des femmes de 50 à 74  
45 ans en France Situation actuelle et perspectives d'évolution."
- 46  
47  
48 **Jusot, Florence, Zeynep Or, and Nicolas Sirven.** 2012. "Variations in preventive care utilisation in  
49 Europe." *European Journal of Ageing*, 9(1): 15–25.
- 50  
51  
52 **Kalecinski, Julie, Véronique Régnier-Denois, Samiratou Ouédraogo, Tienhan Sandrine**  
53 **Dabakuyo-Yonli, Agnès Dumas, Patrick Arveux, and Franck Chauvin.** 2015. "Dépistage organisé  
54 ou individuel du cancer du sein? Attitudes et représentations des femmes." *Santé Publique*, 27(2): 213–220.
- 55  
56  
57

- 1  
2  
3  
4  
5 **Katz, Steven J, and Timothy P Hofer.** 1994. "Socioeconomic disparities in preventive care persist  
6 despite universal coverage: breast and cervical cancer screening in Ontario and the United States." *Jama*,  
7 272(7): 530–534.  
8  
9
- 10 **Katz, Steven J, Judith K Zemencuk, and Timothy P Hofer.** 2000. "Breast cancer screening in the  
11 United States and Canada, 1994: socioeconomic gradients persist." *American Journal of Public Health*,  
12 90(5): 799.  
13  
14
- 15 **Keating, Nancy L, Elena M Kouri, Yulei He, Dee W West, and Eric P Winer.** 2013. "Effect  
16 of Massachusetts health insurance reform on mammography use and breast cancer stage at diagnosis."  
17 *Cancer*, 119(2): 250–258.  
18  
19
- 20 **Kolstad, Jonathan T, and Amanda E Kowalski.** 2012. "The impact of health care reform on hospital  
21 and preventive care: evidence from Massachusetts." *Journal of Public Economics*, 96(11): 909–929.  
22  
23
- 24 **Kopans, Daniel B.** 2000. "Double reading." *Radiologic Clinics of North America*, 38(4): 719–724.  
25  
26
- 27 **Lantz, Paula M, Mahasin Mujahid, Kendra Schwartz, Nancy K Janz, Angela Fagerlin, Barbara  
28 Salem, Lihua Liu, Dennis Deapen, and Steven J Katz.** 2006. "The influence of race, ethnicity, and  
29 individual socioeconomic factors on breast cancer stage at diagnosis." *American Journal of Public Health*,  
30 96(12): 2173–2178.  
31  
32
- 33 **Marmot, MG, DG Altman, DA Cameron, JA Dewar, SG Thompson, M Wilcox, et al.** 2012. "The  
34 benefits and harms of breast cancer screening: an independent review." *The Lancet*, 380(9855): 1778–1786.  
35  
36
- 37 **McMorrow, Stacey, Genevieve M Kenney, and Dana Goin.** 2014. "Determinants of receipt of recom-  
38 mended preventive services: implications for the Affordable Care Act." *American journal of public health*,  
39 104(12): 2392–2399.  
40  
41
- 42 **Perry, N, M Broeders, C De Wolf, S Tornberg, R Holland, and L Von Karsa.** 2008. "European  
43 guidelines for quality assurance in breast cancer screening and diagnosis." *Annals of Oncology*, 19(4): 614–  
44 622.  
45  
46
- 47 **Pletscher, Mark.** 2017. "The effects of organized screening programs on the demand for mammography in  
48 Switzerland." *The European Journal of Health Economics*, 18(5): 649–665.  
49  
50

- 1  
2  
3  
4  
5 **Pruitt, Sandi L, Matthew J Shim, Patricia Dolan Mullen, Sally W Vernon, and Benjamin C**  
6 **Amick.** 2009. "Association of area socioeconomic status and breast, cervical, and colorectal cancer screen-  
7 ing: a systematic review." *Cancer Epidemiology and Prevention Biomarkers*, 18(10): 2579–2599.  
8  
9  
10 **Sabatino, Susan A, Ralph J Coates, Robert J Uhler, Nancy Breen, Florence Tangka, and**  
11 **Kate M Shaw.** 2008. "Disparities in mammography use among US women aged 40–64 years, by race,  
12 ethnicity, income, and health insurance status, 1993 and 2005." *Medical care*, 46(7): 692–700.  
13  
14  
15 **Sabik, Lindsay M, and Cathy J Bradley.** 2016. "The Impact of Near-Universal Insurance Coverage on  
16 Breast and Cervical Cancer Screening: Evidence from Massachusetts." *Health economics*, 25(4): 391–407.  
17  
18  
19 **Schueler, Kristin M, Philip W Chu, and Rebecca Smith-Bindman.** 2008. "Factors associated with  
20 mammography utilization: a systematic quantitative review of the literature." *Journal of women's health*,  
21 17(9): 1477–1498.  
22  
23  
24  
25 **Séradour, B.** 2010. "Breast cancer screening in France: an overview in 2009." *La Revue du praticien*,  
26 60(2): 191–199.  
27  
28  
29 **Sicsic, Jonathan, and Carine Franc.** 2014. "Obstacles to the uptake of breast, cervical, and colorectal  
30 cancer screenings: what remains to be achieved by French national programmes?" *BMC health services*  
31 *research*, 14(1): 465.  
32  
33  
34  
35 **Siegel, Rebecca L, Kimberly D Miller, and Ahmedin Jemal.** 2016. "Cancer statistics, 2016." *CA: a*  
36 *cancer journal for clinicians*, 66(1): 7–30.  
37  
38  
39 **Smith-Bindman, Rebecca, Philip Chu, Diana L Miglioretti, Chris Quale, Robert D Rosenberg,**  
40 **Gary Cutter, Berta Geller, Peter Bacchetti, Edward A Sickles, and Karla Kerlikowske.** 2005.  
41 "Physician predictors of mammographic accuracy." *Journal of the National Cancer Institute*, 97(5): 358–  
42 367.  
43  
44  
45  
46 **Smith, Robert A.** 2003. "Iarc handbooks of cancer prevention, volume 7: Breast cancer screening." *Breast*  
47 *Cancer Res*, 5(4): 216–7.  
48  
49  
50  
51 **Thurfjell, Erik L, K Anders Lernevall, and AA Taube.** 1994. "Benefit of independent double reading  
52 in a population-based mammography screening program." *Radiology*, 191(1): 241–244.  
53  
54  
55  
56  
57

1  
2  
3  
4  
5 **Torre, Lindsey A, Freddie Bray, Rebecca L Siegel, Jacques Ferlay, Joannie Lortet-Tieulent, and**  
6 **Ahmedin Jemal.** 2015. "Global cancer statistics, 2012." *CA: a cancer journal for clinicians*, 65(2): 87–  
7 108.  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57

For Peer Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Figure 1: Implementation of the local and national programs <sup>a</sup>

<sup>a</sup>The grey areas represent the départements that adopted a local program before 2004.




Figure 2: The Percentage of Women with a Mammogram in the Past 2 Years by Age



Table 1: Summary Statistics: Full ESPS Sample, 2000-2010

	All		Depts without a local program		Depts with a local program		P-value of the diff.
Mammogram in the past 2 yrs	0.378	0.487	0.372	0.483	0.385	0.487	0.032
Ever had a mammogram	0.531	0.499	0.526	0.499	0.537	0.499	0.006
Poor health	0.046	0.194	0.045	0.207	0.048	0.213	0.218
Fair health	0.168	0.362	0.167	0.373	0.169	0.375	0.537
Good health	0.310	0.462	0.308	0.462	0.313	0.464	0.456
Excellent health	0.160	0.382	0.163	0.369	0.157	0.364	0.161
Health unknown	0.297	0.454	0.298	0.457	0.295	0.456	0.519
Private compl. insurance	0.877	0.328	0.882	0.323	0.871	0.335	0.005
Public compl. insurance	0.054	0.226	0.051	0.219	0.058	0.235	0.003
No compl. insurance	0.008	0.091	0.009	0.097	0.010	0.097	0.979
Unknown	0.009	0.097	0.008	0.091	0.008	0.091	0.866
Has long-term condition	0.157	0.363	0.156	0.363	0.157	0.364	0.963
Smoker	0.137	0.344	0.140	0.347	0.134	0.341	0.155
Education: Primary or less	0.303	0.459	0.301	0.459	0.305	0.461	0.391
Education: Middle school	0.271	0.444	0.267	0.442	0.276	0.447	0.083
Education: High school	0.192	0.394	0.196	0.397	0.188	0.391	0.111
Education: University	0.194	0.396	0.200	0.400	0.188	0.391	0.012
Education: Other	0.040	0.195	0.037	0.189	0.043	0.202	0.018
Occupation: Farmer	0.047	0.212	0.057	0.231	0.035	0.185	0.000
Occupation: Craftsman	0.048	0.214	0.053	0.223	0.043	0.202	0.000
Occupation: Executive	0.082	0.274	0.083	0.276	0.080	0.271	0.350
Occupation: Intermediate occup.	0.176	0.381	0.181	0.385	0.170	0.376	0.024
Occupation: Employees	0.431	0.495	0.416	0.493	0.449	0.497	0.000
Occupation: Workers	0.163	0.369	0.162	0.368	0.165	0.371	0.544
Occupation: Other/Unknown	0.053	0.224	0.049	0.215	0.058	0.234	0.001
Income: 1st quintile	0.106	0.308	0.108	0.311	0.104	0.305	0.229
Income: 2nd quintile	0.123	0.329	0.125	0.331	0.121	0.327	0.313
Income: 3rd quintile	0.132	0.338	0.134	0.341	0.129	0.335	0.156
Income: 4th quintile	0.171	0.376	0.167	0.373	0.176	0.381	0.027
Income: 5th quintile	0.182	0.386	0.179	0.384	0.185	0.389	0.168
Income: Refused to answer	0.286	0.452	0.287	0.452	0.285	0.451	0.689
Rural	0.293	0.455	0.347	0.476	0.227	0.419	0.000
Paris	0.124	0.330	0.113	0.317	0.137	0.344	0.000
Age	54.779	13.600	55.095	13.805	54.395	13.337	0.000
Radiologist density	9.053	2.362	8.900	2.450	9.239	2.238	0.000
GP density	85.202	11.480	85.374	12.190	84.990	10.560	0.004
Mortality rate 1980-88	16.96	2.192	17.30	2.099	16.54	2.23	0.000
Year: 2000	0.179	0.384	0.180	0.384	0.178	0.383	0.670
Year: 2002	0.184	0.387	0.184	0.388	0.184	0.387	0.846
Year: 2006	0.204	0.403	0.207	0.405	0.202	0.401	0.299
Year: 2008	0.213	0.410	0.215	0.411	0.211	0.408	0.372
Year: 2010	0.219	0.414	0.214	0.410	0.226	0.418	0.013
Observations	29,296		16,059		13,237		

Table 2: Unadjusted Difference-in-Differences Estimates

		Département has		
		a Pre-Existing Program?		
		No	Yes	Difference
A. Pre-National Program				
(2000 and 2002)		[1]	[2]	[2] - [1]
	Outside of Target Age Range	0.250	0.254	0.004
		(0.008)	(0.009)	(0.012)
	In Target Age Range	0.427	0.490	0.063***
		(0.009)	(0.011)	(0.014)
B. Post-National Program				
(2006 to 2010)				
	Outside of Target Age Range	0.235	0.248	0.012
		(0.006)	(0.008)	(0.009)
	In Target Age Range	0.565	0.553	-0.012
		(0.007)	(0.008)	(0.010)
C. Change over Time				
(B - A)				
	Outside of Target Age Range	-0.014	-0.006	
		(0.010)	(0.011)	
	In Target Age Range	0.138***	0.063***	
		(0.012)	(0.013)	

Table 3: Basic regression results

	Mammogram Last 2 Years?		Mammogram Ever?	
	(1)	(2)	(3)	(4)
Local ( $\alpha_1$ )	0.004 (0.015)	-0.011 (0.012)	0.007 (0.015)	-0.006 (0.012)
Target ( $\alpha_2$ )	0.177*** (0.013)	-0.002 (0.012)	0.186*** (0.015)	-0.043*** (0.013)
Post ( $\alpha_3$ )	-0.014 (0.009)	-0.026*** (0.009)	0.064*** (0.010)	0.035*** (0.008)
Local x Target ( $\beta_1$ )	0.059*** (0.021)	0.066*** (0.018)	0.031 (0.022)	0.037** (0.017)
Post x Local ( $\beta_2$ )	0.009 (0.013)	0.012 (0.014)	0.012 (0.014)	0.017 (0.013)
Post x Target ( $\beta_3$ )	0.153*** (0.015)	0.122*** (0.014)	0.065*** (0.017)	0.037*** (0.012)
Post x Local x Target ( $\gamma$ )	-0.084*** (0.019)	-0.080*** (0.017)	-0.061** (0.023)	-0.058*** (0.018)
$\Delta\Delta L$ ( $\beta_1$ )	0.059*** (0.021)	0.066*** (0.018)	0.031 (0.022)	0.037** (0.017)
$\Delta\Delta N$ ( $\beta_3$ )	0.153*** (0.015)	0.122*** (0.014)	0.065*** (0.017)	0.037*** (0.012)
$\Delta\Delta LN$ ( $\beta_3 + \gamma$ )	0.069*** (0.015)	0.042*** (0.011)	0.004 (0.016)	-0.021 (0.013)
N	29,296	29,296	29,296	29,296
Covariates?	No	Yes	No	Yes

Notes: The models in columns 2, 4 include the following covariates: age, age squared, age cubed, the number of radiologists and GP per capita in the département; indicators for smokers, individuals with long-standing chronic conditions, living in a rural area, living in the greater Paris region; and categorical variables for self-reported health, complementary health insurance coverage, education, current or former occupation, income quintile and breast cancer mortality rate. See Appendix Table A.2 for results corresponding to the model in column 2. \*\*\*  $p < .01$ ; \*\*  $p < .05$ ; \*  $p < .10$

Table 4: Regression results of robustness check specifications

	$\Delta\Delta L$	$\Delta\Delta N$	$\Delta\Delta LN$
1. Full Sample (N = 29,296)	0.066*** (0.018)	0.122*** (0.014)	0.042*** (0.011)
2. Panel analysis (N = 17,960)	0.054*** (0.018)	0.128*** (0.018)	0.062*** (0.014)
3. Ages 40 to 80 (N = 24,305)	0.064*** (0.020)	0.116*** (0.014)	0.041** (0.012)
4. Drop 50 & 51 Year-olds (N = 27,586)	0.067*** (0.019)	0.125*** (0.015)	0.049*** (0.012)
5. Control group 49 and younger (N=26,207)	0.079*** (0.019)	0.137*** (0.014)	0.039** (0.011)

Notes: Covariates are included in each specification. \*\*\* $p < .01$ ; \*\* $p < .05$ ; \* $p < .10$

Table 5: Regression results by age

	Dépts w/o Local Program		Dépts with Local Program	
	Coefficient	Std. Error	Coefficient	Std. Error
Age (ref: 35-39)				
Age 40 to 44	0.203***	(0.022)	0.200***	(0.020)
Age 45 to 49	0.308***	(0.020)	0.279***	(0.025)
Age 50 to 54	0.345***	(0.023)	0.386***	(0.020)
Age 55 to 59	0.408***	(0.023)	0.405***	(0.028)
Age 60 to 64	0.320***	(0.025)	0.424***	(0.030)
Age 65 to 69	0.272***	(0.025)	0.401***	(0.030)
Age 70 to 74	0.174***	(0.021)	0.257***	(0.041)
Age 75 to 79	-0.007	(0.019)	0.110***	(0.038)
Age 80 to 84	-0.072***	(0.019)	-0.029	(0.027)
Age 85+	-0.064***	(0.021)	-0.095***	(0.017)
Post	-0.015	(0.015)	-0.001	(0.015)
Post x 40 to 44	-0.024	(0.027)	-0.029	(0.023)
Post x 45 to 49	-0.031	(0.026)	0.002	(0.031)
Post x 50 to 54	0.088***	(0.026)	0.027	(0.023)
Post x 55 to 59	0.057**	(0.027)	0.042	(0.031)
Post x 60 to 64	0.192***	(0.030)	0.034	(0.040)
Post x 65 to 69	0.204***	(0.035)	0.079*	(0.040)
Post x 70 to 74	0.289***	(0.035)	0.219***	(0.041)
Post x 75 to 79	0.205***	(0.031)	0.081*	(0.045)
Post x 80 to 84	0.069**	(0.027)	0.041	(0.035)
Post x 85+	-0.009	(0.024)	0.038	(0.026)
Constant	0.113***	(0.011)	0.107***	(0.016)
N	16,059		13,237	

\*\*\*  $p < .01$ ; \*\*  $p < .05$ ; \*  $p < .10$

Table 6: Percentage of Women Receiving a Mammogram in Past 2 Years by income and education categories

	Dépt with no Local Program			Dépts with Local Program			% Screened via Organized Program
A. Differences by Education	Pre-2004 (N = 2,562)	Post-2004 (N = 4,806)	Change	Pre-2004 (N = 2,072)	Post-2004 (N = 3,994)	Change	Post-2004 (N = 3464)
Primary or less	31.8%	50.1%	18.3% ***	42.8%	50.3%	7.5%***	70.3%
Middle School	47.8%	57.7%	9.9% ***	50.0%	58.4%	8.4%**	59.6%
High School	53.8%	63.6%	9.8 % **	56.7%	59.3%	2.6%	61.7%
University or higher	56.8%	63.1%	6.3 % **	63.1%	62.2%	-0.9%	53.0%
University/Primary or less	1.79	1.26		1.47	1.24		
B. Differences by Income	Pre-2004 (N = 1,856)	Post-2004 (N = 3,535)	Change	Pre-2004 (N = 1,492)	Post-2004 (N = 2,954)	Change	Post-2004 (N = 3464)
1st Quintile	33.6%	54.5%	20.9% ***	50.2%	55.2%	4.9%	71.5%
2nd Quintile	42.1%	61.4%	19.3%***	54.5%	63.1%	8.6%**	66.0%
3rd Quintile	53.7%	66.9%	13.2% ***	60.3%	62.9%	2.6%	67.0%
4th Quintile	61.6%	67.9%	6.3% **	66.2%	68.9%	2.8%	61.3%
5th Quintile	66.7%	73.3%	6.5%**	66.3%	67.9%	1.7%	55.8%
5th Quintile/1st Quintile	1.99	1.35		1.32	1.23		

Notes: We assess if the change is statistically significant using t-tests. Screened via Organized program is only available for the years 2006 and 2008.

\*\*\* $p < .01$ ; \*\* $p < .05$ ; \* $p < .10$

## A Appendix

Table A.1: Parallel trends

	2000	2002	Difference
	(1)	(2)	(1)-(2)
<b>Local program</b>			
Outside of target age range	0.24	0.267	-0.027
	(0.012)	(0.012)	(0.017)
In target age range	0.487	0.493	-0.005
	(0.015)	(0.015)	(0.021)
<b>No local program</b>			
Outside of target age range	0.251	0.248	0.003
	(0.011)	(0.011)	(0.016)
In target age range	0.421	0.432	-0.010
	(0.014)	(0.013)	(0.019)



Table A.2: Basic regression results with covariates

	Coefficient	Std. Error
Local	-0.011	(0.012)
Target	-0.002	(0.012)
Post	-0.027***	(0.009)
Local x Target	0.066***	(0.018)
Post x Local	0.012	(0.014)
Post x Target	0.122***	(0.014)
Post x Local x Target	-0.080***	(0.017)
Health status (ref: good)		
Poor	-0.085***	(0.013)
Fair	-0.015*	(0.008)
Excellent	-0.030***	(0.008)
Unknown	-0.470***	(0.010)
Complementary health insurance (ref: private)		
Unknown	-0.044***	(0.017)
None	-0.088***	(0.010)
Public	-0.069***	(0.011)
Chronic disease (ref: no)		
Has a chronic disease	0.016**	(0.007)
Smoker	-0.043***	(0.008)
Education (ref: primary or lower)		
Junior High school	0.032***	(0.007)
High school	0.037***	(0.007)
University	0.040***	(0.009)
Other	0.002	(0.011)
Occupation (ref: executive)		
Farmer	-0.036***	(0.013)
Shopkeeper, craftsmen	-0.025	(0.017)
Intermediary profession	0.009	(0.011)
Employee	-0.015	(0.011)
Worker	-0.042***	(0.014)
Not working/other	-0.078***	(0.017)
Income (ref: 1st quintile)		
2nd quintile	0.023**	(0.011)
3rd quintile	0.041***	(0.012)
4th quintile	0.060***	(0.010)
5th quintile	0.082***	(0.010)
Unknown	0.007	(0.009)
Rural	-0.010	(0.007)
Parisian region	0.013	(0.009)
Age	0.167***	(0.009)
Age squared	-0.002***	(0.000)
Aged cubed	0.000***	(0.000)
Radiologist density	0.007***	(0.001)
GP density	0.000	(0.000)
Mortality rate of breast cancer	-0.001	(0.002)
Constant	-3.418***	(0.171)
Observations	29,296	
R-squared	0.340	

Notes: \*\*\* $p < .01$ ; \*\* $p < .05$ ; \* $p < .10$

Table A.3: Percentage of Women Receiving a Mammogram ever by income and education categories

	Dépt with no Local Program			Dépts with Local Program		
<u>A. Differences by Education</u>	<b>Pre-2004</b> (N = 2,562)	<b>Post-2004</b> (N = 4,806)	<b>Change</b>	<b>Pre-2004</b> (N = 2,072)	<b>Post-2004</b> (N = 3,994)	<b>Change</b>
Primary or less	46.25%	63.90%	17.65***	56.11%	64.16%	8.05***
Middle School	63.21%	70.98%	7.77***	61.72%	70.66%	8.94***
High School	65.88%	75.49%	9.61***	64.60%	72.77%	8.17**
University or higher	66.91%	74.22%	7.31**	72.48%	72.31%	-0.17
University/Primary or less	1.45	1.16		1.29	1.13	
<u>B. Differences by Income</u>	<b>Pre-2004</b> (N = 1,856)	<b>Post-2004</b> (N = 3,535)	<b>Change</b>	<b>Pre-2004</b> (N = 1,492)	<b>Post-2004</b> (N = 2,954)	<b>Change</b>
1st Quintile	52.20%	77.46%	25.26***	65.33%	74.46%	9.13**
2nd Quintile	60.69%	77.18%	16.49***	67.77%	79.92%	12.15***
3rd Quintile	73.03%	79.56%	6.53**	79.04%	77.72%	-1.32
4th Quintile	77.42%	81.74%	4.32*	77.46%	81.45%	3.99
5th Quintile	78.69%	84.64%	5.95**	78.86%	81.39%	2.53
5th Quintile/1st Quintile	1.51	1.09		1.21	1.09	

Notes: We assess if the change is statistically significant using t-tests.

\*\*\* $p < .01$ ; \*\* $p < .05$ ; \* $p < .10$