




Developmental changes in resting-state functional networks among individuals with and without internalizing psychopathologies

Katie L. Burkhouse¹  | Jonathan P. Stange¹  | Rachel H. Jacobs¹ | Runa Bhaumik¹ | Katie L. Bessette^{1,2} | Amy T. Peters^{1,2} | Natania A. Crane¹ | Kayla A. Kreutzer¹ | Kate Fitzgerald³ | Christopher S. Monk⁴ | Robert C. Welsh⁵ | K. Luan Phan^{1,2,6,7} | Scott A. Langenecker^{1,2,5} 

¹Department of Psychiatry, University of Illinois–Chicago, Chicago, USA

²Department of Psychology, University of Illinois–Chicago, Chicago, USA

³Department of Psychiatry, University of Michigan, Ann Arbor, USA

⁴Department of Psychology, University of Michigan, Ann Arbor, USA

⁵Department of Psychiatry, The University of Utah Medical Center, Salt Lake City, USA

⁶Mental Health Service Line, Jesse Brown VA Medical Center, Chicago, USA

⁷Department of Anatomy and Cell Biology & the Graduate Program in Neuroscience, University of Illinois–Chicago, Chicago, USA

Correspondence

Katie L. Burkhouse, Department of Psychiatry, University of Illinois–Chicago, Chicago, IL 60608. Email: kburkho@uic.edu.

Funding information

National Institute of Mental Health, Grant/Award Numbers: MH086517, MH091811, MH101487, MH112769, MH113793; Mind and Life Foundation; Klingenstein Third Generation Foundation

Background: Three well-established intrinsic connectivity networks (ICNs) involved in cognitive-affective processing include the cognitive control network (CCN), default mode network (DMN), and salience and emotional network (SEN). Despite recent advances in understanding developmental changes in these ICNs, the majority of research has focused on single seeds or networks in isolation with limited age ranges. Additionally, although internalizing psychopathologies (IPs), such as anxiety and depression, are often characterized by maladaptive cognitive-affective processing styles, it is not clear how IP history influences age-related changes in brain networks.

Method: The current study aimed to characterize the normative development of the CCN, DMN, and SEN across a large age-span (7–29 year olds) of typically developing (TD) individuals ($n = 97$). We also explore how age may impact differences in network connectivity between TD individuals and patients with IPs ($n = 136$).

Results: Among TD individuals, DMN and CCN connectivity strengthened with age, whereas connectivity between the SEN and ventromedial prefrontal cortex weakened across development. When exploring group (IP vs. TD) differences, the IP group was characterized by greater connectivity between the CCN and cerebellum and between the SEN and caudate from childhood to early adulthood, relative to TD individuals. In addition, patients with IPs, versus TD individuals, exhibited reduced connectivity between the SEN and medial frontal gyrus from adolescence to adulthood.

Conclusions: The current findings shed light on differential age-related changes in brain network patterns among psychiatrically free, TD individuals and those with internalizing disorders, and may provide plausible targets for novel mechanism-based treatments that differ based on developmental stage.

KEYWORDS

age, anxiety, depression, development, internalizing psychopathologies, intrinsic connectivity networks (ICN)

1 | INTRODUCTION

Substantial evidence suggests that the brain is organized into many functional networks that flexibly interact to support various types of information processing (Menon, 2011; Seeley et al., 2007; Yeo et al., 2011). To capture a representation of these networks, researchers

have examined functional connectivity patterns, measured via resting-state functional magnetic resonance imaging (rs-fMRI), which represent the temporal correlation of brain regions (Biswal et al., 2010) and capture spontaneous low frequency fluctuations in the blood oxygen-level dependent signal (Fox & Raichle, 2007). In recent years, increased attention has been placed on examining age-related changes in these

intrinsic connectivity networks (ICNs) to capture typical maturational shifts in brain development, and also to characterize the biological basis of psychiatric disorders in which normal developmental processes are disrupted (Kaiser & Pizzagalli, 2015).

Three ICNs that have been well established and serve distinct roles in cognitive and emotional processing include the default mode network (DMN), salience and emotional network (SEN), and cognitive control network (CCN) (Fox & Raichle, 2007; Menon, 2011; Seeley et al., 2007). Briefly, the DMN is hypothesized to support self-referential or introspective mental processes and includes regions that are typically active during rest and inactive during cognitively demanding tasks (e.g., Raichle et al., 2001). The SEN on the other hand, encompasses limbic and salience regions that are active during the detection and processing of external or internal emotional and novel stimuli (Seeley et al., 2007). Finally, the CCN, also often referred to as the central executive network, is a frontoparietal system that is thought to be instrumental in problem-solving and executive functioning (Menon, 2011).

Given the importance of these three networks for key aspects of information processing implicated in internalizing disorders, increased attention has been placed in recent years to better understand age-related changes in the ICNs. Consistent with the notion of enhanced cognitive and mental processing with age, there is evidence that connectivity among regions in the DMN increases in strength from childhood to late adolescence (Fair et al., 2008; Power, Fair, Schlaggar, & Petersen, 2010; Rubia, 2013). Additional research has found evidence for increased connectivity among regions within the CCN among adults relative to children (Barber, Caffo, Pekar, & Mostofsky, 2013; Jolles, van Buchem, Crone, & Rombouts, 2010). These previous studies are consistent with functional magnetic resonance imaging (fMRI) studies highlighting increased activation in frontal and parietal regions from childhood to adulthood during cognitive control tasks, suggesting that efficient top-down control processes may not be fully developed until adulthood (i.e., Luna et al., 2001). Regarding SEN development, there is evidence for an increase in connectivity between the nucleus accumbens and subcortical regions among a sample of 8–25 year olds (van Duijvenvoorde, Achterberg, Braams, Peters, & Crone, 2016). However, connectivity between the nucleus accumbens and ventromedial prefrontal cortex (vmPFC), a region shown to support down regulation of key regions within the SEN (i.e., amygdala, striatum) (Etkin, Egner, & Kalisch, 2011), declined with age among these individuals. This is consistent with a separate study showing positive amygdala with the medial prefrontal cortex (mPFC) connectivity during childhood, versus the opposite pattern in adolescence and adulthood (Gee et al., 2013), perhaps implicating enhanced regulatory abilities with development.

Although these previous studies shed light on typical maturational processes of ICN patterns, the majority of past research has focused on single seeds or networks in isolation with limited age ranges. In contrast, a recent study did examine age-related changes in several resting-state networks from childhood to late-adolescence. Specifically, developmental changes in connectivity among six resting-state networks linked to cognitive and emotional processes were assessed among healthy youth ranging in age from 7 to 18 (Solé-Padullés et al., 2016). In this study, within CCN and DMN intrinsic

connectivity increased from childhood to adolescence. Additionally, SEN was synchronized with more anterior regions with increasing age, such as the inferior and superior frontal cortices and the left insula. However, connectivity between the SEN and mPFC declined from childhood to adolescence, replicating previous studies (Gee et al., 2013; van Duijvenvoorde et al., 2016).

The current study sought to replicate and extend these previous findings in two important ways. First, we aimed to characterize the development of intrinsic connectivity within and between three networks (DMN, CCN, and SEN) among a larger age-span (i.e., 7–29 year olds) of psychiatrically free, typically developing (TD) individuals by utilizing rs-fMRI. The inclusion of a wider age range when exploring age-related changes in network connectivity is essential given evidence that maturation of the corpus callosum occurs well into early adulthood (i.e., late 20s) (Keshavan et al., 2002; Pujol, Vendrell, Junqué, Martí-Vilalta, & Capdevila, 1993). Consistent with previous studies (Barber et al., 2013; Fair et al., 2008; Jolles et al., 2010; Power et al., 2010; Rubia, 2013; Solé-Padullés et al., 2016), we predicted that connectivity within the CCN and DMN would be positively associated with age among TD individuals. For the SEN, we predicted that there would be enhanced connectivity with subcortical regions (i.e., hippocampus, amygdala) with age (van Duijvenvoorde et al., 2016); however, consistent with previous studies (Solé-Padullés et al., 2016) and the notion of enhanced regulatory abilities with age, we predicted that connectivity between the SEN and vmPFC would decline with age.

A second extension of the current study was to explore whether individuals with a history of internalizing psychopathologies (IPs) exhibit parallel patterns of network connectivity with development relative to the TD sample. Specifically, IPs, such as anxiety and depression, are highly comorbid across the lifespan (e.g., Kessler et al., 2005; Moffitt et al., 2007) and derive from common mechanisms. Indeed, there is substantial evidence demonstrating that depressive and anxiety disorders are characterized by deviant patterns of connectivity within DMN, SEN, and CCN regions relative to TD individuals in specific developmental and adult age spans (for reviews, see Hulvershorn, Cullen, & Anand, 2011; Kim et al., 2011; MacNamara, DiGangi, & Phan, 2016; Mulders, van Eijndhoven, Schene, Beckmann, & Tendolkar, 2015). However, few studies to date have examined network changes between individuals with IPs and psychiatrically free individuals across the lifespan, particularly during childhood development when neural plasticity is most active. A better understanding of age-related changes in ICN patterns among individuals with IPs is essential to increase our understanding of these disorders across the lifespan and to develop plausible targets for early identification and novel mechanism-based treatments that may differ based on stage of development. Moreover, exploring the specificity of these network patterns early on in development (i.e., during childhood and adolescence) may provide insight into biological markers that can be targeted to prevent the development and persistence of anxiety and depression in youth.

Thus, this extension of the current study was to examine how developmental stage (i.e., age) may impact differences in network connectivity between TD individuals and patients with IPs. Consistent with previous adolescent and adult studies (for reviews, see Kaiser, Andrews-Hanna, Wager, & Pizzagalli, 2015; MacNamara et al., 2015;

Mulders et al., 2015), we predicted that relative to the TD group, individuals with a history of anxiety and/or depression would exhibit greater connectivity within the DMN from childhood to adulthood. Investigations of CCN and SEN connectivity across the lifespan have been more mixed among individuals with anxiety and depression (Kaiser et al., 2015; Mulders et al., 2015), with evidence for both decreased (Bluhm et al., 2009; Cullen et al., 2009; Sylvester et al., 2012; Veer et al., 2010) and increased (Davey, Harrison, Yücel, & Allen, 2012; Sheline, Price, Yan, & Mintun, 2010; Sylvester et al., 2012) connectivity within these networks among adolescent and adult populations. We have previously shown that within the CCN, adolescents (ages 12–17) with a history of depression exhibit increased connectivity (Peters, Burkhouse, Feldhaus, Langenecker, & Jacobs, 2016), whereas the opposite pattern is found among remitted depressed young adults ranging in age from 18 to 23 (Stange et al., 2017). Moreover, when examining cross-network connectivity patterns, remitted depressed adolescents exhibited increased connectivity between the SEN and DMN (Peters et al., 2016), while young adults in remission from MDD exhibited stable and reliable hyperconnectivities of the DMN and SEN with the CCN (Bessette et al., 2018; Jacobs et al., 2014). Thus, taken together, these previous studies suggest that developmental stage may impact the pattern of network connectivity among individuals with a history of depressive and anxiety disorders.

In the current study, we sought to extend these previous findings by studying a wider age range to comprise the full developmental span from childhood to adulthood and by including a larger sample of patients with internalizing disorders to explore how period of development may impact ICN patterns among individuals with IPs and no history of psychopathology. The inclusion of a larger age range is essential to characterize the biological basis of these disorders given that anxiety and depressive disorders are highly prevalent and cause significant disability across the lifespan, and knowing that cortex maturation occurs into early adulthood (Keshavan et al., 2002; Pujol et al., 1993). Given the mixed findings reported to date across a number of different age and developmental windows, we did not have *a priori* hypotheses regarding age-by-disorder interactions.

2 | METHOD

2.1 | Participants

The reported participant sample consisted of 233 children, adolescents, and adults recruited through the University of Michigan (UM) and University of Illinois at Chicago (UIC)¹. Participants were enrolled in one of five research studies at UIC or UM. For each of these studies, participants were recruited using flyers and multiple postings on the internet. All participants completed an assessment battery of self-report and diagnostic measures, followed by an MRI scan. Youth diagnoses were obtained using the Schedule of Affective Disorders and Schizophrenia for School-Age Children (Kaufman et al., 1997). For adults, diagnoses were determined using either the Structured Clinical Interview for DSM-IV (SCID; First, Spitzer, & Gibbon, 2002) or the Diagnostic Interview for Genetic Studies (Nurnberger et al., 1994). To

be included in the TD group, participants could not meet criteria for a current or past DSM-IV Axis-I psychological disorder. To be included in the IP patient group, participants had to meet criteria for a current or past DSM-IV depressive or anxiety disorder. Exclusionary criteria for the IP group included substance abuse or dependence within the past 6 months, history of bipolar or schizophrenia disorders, intellectual disability, and pervasive development disorders. The reported sample included 97 TD individuals and 136 patients with IPs between 7 and 29 years of age. Participant demographic and clinical characteristics are presented in Table 1. The current study was approved by the University of Michigan (UM) and the University of Illinois at Chicago (UIC) Institutional Review Boards and all adults and youth provided written informed consent and assent, respectively.

2.2 | Symptom measures

Given that the current study consisted of data across five different studies, participants did not complete a uniformed measure of anxiety or depressive symptoms. To assess anxiety symptoms, participants either completed the Beck Anxiety Inventory (BAI; Beck & Steer, 1990) or the Multidimensional Anxiety Scale for Children (MASC; March, Parker, Sullivan, Stallings, & Conners, 1997). To assess depressive symptoms, participants either completed the Beck Depression Inventory (BDI; Beck, Steer, & Brown, 1996), Children's Depression Inventory (CDI; Kovacs, 2004), or the Reynold's Adolescent Depression Scale (RADs; Reynolds, 2004). As such, standardized anxiety and depressive scores were created for all participants by calculating age and where possible, sex-corrected Z scores utilizing published normative data from community samples for each of the measures (see Supp. Table 1 for additional details).

2.3 | rs-fMRI acquisition

At UM, an eyes-open resting-state scan was acquired over 8 min on a 3.0 T GE Signa scanner (Milwaukee, WI) using T2*-weighted single-shot reverse spiral sequence with the following parameters: 90° flip angle, field-of-view 20 cm, matrix size = 64 × 64, slice thickness = 4 mm, 30 ms echo time, 29 sequentially acquired slices. Eyes-open, resting scans at UIC were collected over 8 min on a 3.0 T GE Discovery scanner (Milwaukee, WI) using parallel imaging with ASSET and T2* gradient-echo axial EPI with the following parameters: 90° flip angle, field-of-view 22 cm, matrix size = 64 × 64, slice thickness = 3 mm, 22.2 ms echo time, 44 interleaved slices. At both sites, high-resolution anatomic T1 scans were obtained for spatial normalization; motion was minimized with foam pads, a visual tracking line (UIC only) and/or cross (UIC and UM) on the display, and by conveying the importance of staying still to participants, with TRs of 2000 ms and a total of 240 TRs.

2.4 | rs-fMRI preprocessing

Time series data were detrended and mean-centered. Physiologic correction was performed by regressing out white matter and cerebral spinal fluid signals (Behzadi, Restom, Liau, & Liu, 2007). Motion parameters were regressed out (Jo et al., 2013). Based upon recent literature

TABLE 1 Demographics and Clinical Characteristics of the Sample

	TD (n = 97)	IP (n = 136)	Statistic (χ^2 or t-value)
Demographics			
Age (M, SD)	19.26 (4.56)	18.64 (4.47)	-1.03
Sex (% female)	58.8%	65.4%	0.38
Race/ethnicity			
Caucasian	58.7%	59.6%	0.18
American Indian/Alaskan Native	0%	0.7%	0.89
African American	8.2%	11.0%	0.64
Asian	18.6%	5.1%	10.99*
Other/biracial	11.3%	21.3%	4.04
Hispanic	10.4%	19.8%	3.83
Diagnoses			
Current anxiety disorder	0.0%	72.1%	-
Current MDD	0.0%	29.4%	-
Past MDD	0.0%	47.8%	-
Comorbid anxiety/depression history	0.0%	49.3%	-
Clinical characteristics (M, SD)			
Depressive symptoms (Z Score)	-1.02 (0.41)	0.45 (1.43)	9.82**
Anxiety symptoms (Z Score)	-0.51 (0.71)	0.46 (1.06)	7.87**
Study characteristics			
Study site (% UIC)	73.2%	71.3%	-0.75

Note. TD, typically developing; IP, internalizing psychopathology; MDD, major depressive disorder.

* $p < 0.01$.

** $p < 0.001$.

(Jo et al., 2013; Power et al., 2012), motion volumes were identified based on any TR to TR movement exceeding 1.5° , or 3 consecutive TRS exceeding the same in any of direction pitch, roll, or yaw (Power et al., 2012). In addition, we examined normality plots of the average standard deviation of movement values in the pitch, roll, and yaw planes for outliers. Participants with movement values in any plane greater than .05 mm were excluded. IP and TD groups did not differ in regards to these movement values (i.e., pitch, roll, and yaw planes).

Slice timing was completed with SPM8 (version R4667) (<http://www.fil.ion.ucl.ac.uk/spm/doc/>) and motion detection algorithms were applied using FSL (version 5.0.0.2) (<http://fsl.fmrib.ox.ac.uk/fsl/fslwiki/>). Co-registration of structural images to functional images was followed with spatial normalization of the co-registered T1-SPGR to the Montreal Neurological Institute (MNI) 152 brain template. The resulting normalization matrix was then applied to the slice-time-corrected time-series data. These normalized T2* time-series data were spatially smoothed with a 5 mm Gaussian kernel and spatially resampled to T2* images with isotropic voxels, 2 mm on each side.

To define the mask for the CCN, we selected key nodes based upon Yeo et al. (2011). We included bilateral dorsolateral prefrontal cortex nodes within the middle frontal gyrus (± 45 29 32), inferior parietal lobule (± 52 -50 49), and dorsal anterior cingulate (± 5 22 47) to use as seeds in the connectivity analysis (Yeo et al., 2011). For each of these three bilateral seeds, we ran cross-correlation time series analyses as described above (and similar to Stange et al., 2017). Similar procedures were used for three bilateral seeds for the SEN [amygdala

(± 23 -5 -19; Jacobs et al., 2014), ventral striatum inferior (± 9 9 -8; Di Martino et al., 2008), anterior insula (± 33 12 -6; Deen et al., 2010)] and for the DMN [posterior cingulate (± 5 -49 25), dorsal medial cingulate (± 7 46 -2), temporoparietal junction (± 49 -63 45)] (Yeo et al., 2011).

2.5 | Analytical plan

Using the three masks (CCN, DMN, and SEN) derived from the available download (Yeo et al., 2011), second-level models in SPM8 tested for voxels that were significantly related to age (as a continuous factor) within the masks among the TD group. In these analyses, we controlled for study site, sex, and movement parameters (i.e., average standard deviation of movement values in each of the pitch, roll, and yaw planes). The threshold of significance reported for the fMRI analyses was $p < 0.001$ and $k \geq 150$ [3dClustsim with whole brain corrected p -value of 0.001 per analysis (December 16, 2015, updated release; 10,000 iterations; http://afni.nimh.nih.gov/pub/dist/doc/program_help/3dClustSim.html; Eklund, Nichols, & Knutsson, 2016)]. Using whole-brain correction thresholds, despite focused hypotheses that utilized only a subset of voxels (DMN, SEN, and CCN masks, for clarity in interpretation) represents a conservative approach (family-wise WB error, $p < 0.003$). Data were extracted from the suprathreshold ROIs that were associated with age.

Next, we explored Group \times Age interactions in SPM8. Specifically, an identical set of analyses to the TD model described above was conducted with the addition of the Group (IP, TD) \times Age

(continuous) interaction. Data were extracted from significant ROIs for the Age \times Group interaction contrasts: positive contrast = IP group exhibits stronger connectivity with age, relative to the TD group; negative contrast = IP group exhibits weaker connectivity with age, relative to the TD group. To follow-up on significant interactions, we utilized the Johnson–Neyman technique in SPSS (Hayes & Rockwood, 2017), which outputs a value of the moderator (Age), at which the association between the independent variable (IP vs. TD) and outcome crosses the threshold for statistical significance ($p < 0.01$).

3 | RESULTS

3.1 | Participant characteristics

As shown in Table 1, the TD and IP groups did not differ in regards to age, sex, ethnicity, and study site. However, a greater number of TD participants identified as Asian American, relative to the IP group. As expected, the IP group reported a greater number of anxiety and depressive symptoms, relative to the TD group.

3.2 | Age-related changes in network connectivity among TD individuals

Table 2 displays the findings of the influence of age on connectivity within the three networks among TD participants. As shown in Figures 1 and 2, second-level analyses in SPM8 revealed a positive linear effect of age on connectivity between the CCN (average of connectivity between three bilateral seeds of CCN mask) and the following regions: inferior frontal gyrus (IFG), middle frontal gyrus (MFG), rostral anterior cingulate cortex (ACC), right ACC, inferior parietal lobe (IPL), and putamen. Age was also positively correlated with connectivity between the DMN (average of connectivity between three bilateral seeds of DMN mask) and dorsal anterior cingulate and middle prefrontal cortex. Finally, results revealed a negative

TABLE 2 Location of Clusters of Connectivity that were Significantly related to Age among the Typically Developing (TD) Participants, for each of the Three Networks

Lobe	Gyrus	MNI Coordinates					
		BA	x	y	z	Z	k
Age increasing							
CCN mask							
Frontal	Inferior	45	54	34	12	5.14	233
Frontal	Middle	9	38	10	36	4.37	222
Limbic	Cingulate	32	0	44	4	4.78	360
Limbic	Cingulate	32	-2	24	34	4.23	165
Parietal	Inferior	40	-40	-52	32	4.05	209
Subcortical	Putamen	-	18	14	-4	4.87	273
DMN mask							
Frontal	Cingulate	32	6	20	38	4.82	456
Frontal	Middle	9	44	22	22	4.67	997
Age decreasing							
SEN mask							
Frontal	Ventromedial	9	2	54	4	-4.78	462

Note. CCN, cognitive control network; DMN, default mode network; SEN, salience emotion network; k, number of voxels; MNI, Montreal Neurological Institute.

correlation between age and SEN (average of connectivity between three bilateral seeds of SEN mask) connectivity with the vmPFC.

3.3 | Group differences in network connectivity

Group \times Age interactions were examined in SPM8 (Table 3, Figure 3, Suppl. Figure 1). For the CCN model, results revealed significant Group \times Age interactions with the cerebellum. Follow-up analyses utilizing the Johnson–Neyman technique revealed that TD individuals, relative to patients, exhibited greater connectivity




FIGURE 1 Top panel: The seed by cluster effects for the typically developing comparison subjects in panels A (left lateral), B (left medial), C (right putamen), and D (right inferior and middle frontal). Blue: Increasing connectivity with default mode network (DMN) seeds with age. Red: Decreasing connectivity with age to salience emotional network (SEN) seeds. Green: Increasing connectivity with age for cognitive control network (CCN) seeds. Bottom panels (E–H): illustrate the three primary network masks, DMN (cyan), SEN (violet), and CCN (yellow) in the same laterality of panels to match panels A–D for easy comparison of within and across network connectivity comparisons




FIGURE 2 Scatter plots of the relation between age (continuous) and mean Z-corrected connectivity between (a) cognitive control network (CCN) with inferior frontal gyrus (IFG); (b) CCN with middle frontal gyrus (MFG); (c) CCN with right cingulate; (d) CCN with rostral anterior cingulate cortex (rACC); (e) CCN with inferior parietal lobule (IPL); (f) CCN with putamen; (g) default mode network (DMN) with dorsal ACC (dACC); (h) DMN with MFG; (i) salience emotional network (SEN) with ventromedial prefrontal cortex (vmPFC). The colors represent the age ranges for illustrative purposes, and the linear error bar is the 95% confidence interval for the mean

between the CCN and cerebellum between the ages of 7 and 20. For the SEN model, significant Group \times Age interactions were observed for the caudate, superior temporal gyrus, and the medial frontal gyrus. Post hoc analyses revealed that IP patients exhibited greater connectivity between the SEN and caudate between ages 7 and 24, relative to the TD group. Conversely, the IP, relative to the TD, group exhibited reduced connectivity between the SEN and medial frontal gyrus beginning at age 12. Additionally, patients exhibited reduced

connectivity between the SEN and the superior temporal gyrus later in development (i.e., ages 21–29), relative to TD individuals.

3.4 | Potential confounds

Findings were similar when excluding the 21 participants from the patient group taking a psychotropic medication at the time of the scan. We also conducted the above analyses utilizing data from UIC

TABLE 3 Location of Significant Clusters of Connectivity for the Group (IP vs. TD) × Age Interaction, for each of the Three Networks

Cluster Location	MNI Coordinates					
	BA	x	y	z	Z	k
CCN mask – positive						
Cerebellum/culmen/ midbrain/pons	-	12	-36	-24	3.91	273
SEN mask – negative						
Caudate	-	-10	18	6	4.41	153
Superior temporal gyrus	22	-66	-44	16	3.92	169
Medial frontal gyrus	6	8	-30	60	3.91	641

Note: IP, internalizing psychopathology; TD, typically developing; positive, regions in which the IP group exhibits stronger connectivity with age, relative to the TD group; negative, regions in which the IP group exhibits weaker connectivity with age, relative to the TD group; CCN, cognitive control network; DMN, default mode network; SEN, salience emotion network; k, number of voxels; MNI, Montreal Neurological Institute.

participants only (i.e., excluding UM participants) to rule out site differences. The findings remained unchanged.

4 | DISCUSSION

The current study sought to examine age-related changes of three commonly studied ICNs (i.e., CCN, DMN, and SEN) across a broad age-span (i.e., 7–29 year olds) of psychiatrically healthy individuals. We also explored regions of network connectivity that differed between psychiatrically free, TD individuals and patients with IPs (i.e., anxiety and depression) across development. Several notable findings emerged from the current study. First, when focusing analyses on findings with psychiatrically free individuals, we found age-related increases in CCN

connectivity with several regions involved in higher level cognitive and mental processing (i.e., IFG, MFG, rostral ACC, right ACC, IPL, and putamen). Age was also positively correlated with connectivity between the DMN and several regions comprising the CCN (i.e., MFG, rostral ACC). These findings replicate previous studies (Barber et al., 2013; Fair et al., 2008; Jolles et al., 2010; Power et al., 2010; Rubia, 2013; Solé-Padullés et al., 2016) by demonstrating potential maturation of the DMN and CCN at rest throughout development among a larger and broader sample of individuals with no prior history of a psychological disorder.

When focusing on SEN development among healthy individuals, the current study found no evidence for increased connectivity with limbic regions with age when utilizing a salience emotional network. This contrasted with another recent study, which reported increased connectivity between the nucleus accumbens and other subcortical regions with age (van Duijvenvoorde et al., 2016). As we elected to only study regions within the network that had common patterns of connectivity change, our results are not directly comparable to seed-based analyses of the nucleus accumbens. Our finding revealing age-related decreases in connectivity between the SEN and vmPFC is consistent with previous studies suggesting that connectivity between limbic regions and the mPFC decreases with age during rest and face processing tasks (Gee et al., 2013; Solé-Padullés et al., 2016; van Duijvenvoorde et al., 2016). This might be reflective of age-specific increases in emotion regulatory capacity.

In a distinct set of analyses, we also explored how period of development may impact ICN patterns that differ between TD participants and patients with IPs. Although previous studies have examined differences in patterns of connectivity between TD individuals and patients with IPs, no studies to date have examined how these effects may differ across development utilizing a broad age-span of individuals ranging




FIGURE 3 Scatter plot depicting group (TD vs. IP) differences in network connectivity at high and low (± 1 SD) age for illustrative purposes. (A) Cognitive control network (CCN) to cerebellum: groups differ statistically from age 7 to 20; (B) salience emotional network (SEN) to caudate: groups differ statistically from age 7 to 24; (C) SEN to superior temporal gyrus: groups differ statistically from age 21 to 29; and (D) SEN to medial frontal gyrus: groups differ from age 12 to 29

from childhood to early adulthood. Focusing first on the CCN, patients with IPs were characterized by increased connectivity between the CCN and cerebellum from childhood to early adulthood (ages 7–20), relative to the TD group. Studies have provided increasing evidence for the cerebellum's involvement in cognitive function and emotional reactions, and subsequently, IP pathophysiology (Schmahmann, 2010; Stoodley, 2012). Notably, other studies have demonstrated that depressed adults exhibit decreased connectivity between the CCN and cerebellum (Liu et al., 2012; Ma, Zeng, Shen, Liu, & Hu, 2013; Zeng et al., 2012), relative to TD individuals. Our findings contribute to the literature by highlighting that these group differences are most pronounced in young children, potentially suggesting a plausible brain-based target for early identification of IPs.

Next, when examining group differences in SEN connectivity across age, results revealed several interesting findings. First, patients, compared to TD individuals, exhibited increased connectivity between the SEN and caudate, a region of the striatum involved in the automated control of cognitive and emotional processes, between ages 7 and 24. Interestingly, studies to date provide evidence of cortical-striatal dysfunction in a variety of mood and anxiety disorders (for a review, see Marchand, 2010), and it has been posited that a cortical-striatal-thalamic circuit is involved in the pathophysiology of depression (Drevets, 2001). Results also revealed that patients, relative to TD individuals, exhibited reduced connectivity between the SEN and superior temporal gyrus later in development (i.e., ages 20–29). This finding is consistent with a separate study demonstrating that adults with depression exhibit a wide-spread reduction in intrinsic connectivity of the amygdala with several regions involved in emotion processing and regulation, including the superior temporal gyrus (Ramasubbu et al., 2014). Notably, this group difference was not observed among younger individuals; therefore, characteristics of the disease may be contributing to the observed connectivity pattern, as there is evidence for the role of number of episodes and length of illness on changes in neural structures (e.g., Frodl et al., 2003; MacQueen et al., 2003; Sheline, Sanghavi, Mintun, & Gado, 1999). The current study did not comprehensively assess disease characteristics for all participants; therefore, future studies are needed to assess specific factors contributing to these ICN patterns across development.

Replicating previous studies, we also found evidence for reduced connectivity between the SEN and medial frontal gyrus, which plays a pivotal role in the modulation and inhibition of excessive limbic activity (Etkin et al., 2011). Interestingly, the observed group difference began during the adolescent period (i.e., age 12), which is when rates of depression escalate, particularly among females (Hankin et al., 1998). This finding corresponds to previous research showing reduced connectivity between key regions within the SEN and mPFC among individuals with IPs (Gee et al., 2013; Kim et al., 2011; Pannekoek et al., 2014). Notably, there is increasing evidence in support of the theory that early adolescence and puberty-specific changes contribute to increased emotional and stress reactivity and responsiveness (Dahl & Gunnar, 2009; Spear, 2009), which have been linked to IPs across the lifespan (see Aldao, Nolen-Hoeksema, & Schweizer, 2010 for a meta-analysis). Although speculative, increased frontolimbic connectivity

may be one potential mechanism implicated in the development of depression during the adolescent period. Future, longitudinal designs are needed to test this possibility.

Finally, contrary to our original hypothesis, we found limited evidence for group differences in connectivity within the DMN across developmental groups. This finding differs from some previous studies, which have provided support for DMN hyperconnectivity among adults with current depression (Mulders et al., 2015). However, studies examining DMN connectivity among anxious populations have been more mixed with evidence for increased (Liao et al., 2011), decreased (Sylvester et al., 2012), and intact (Pannekoek et al., 2013) DMN connectivity patterns. Future studies might specifically stratify recruitment of major depression, anxiety disorders, and comorbid groups to test additional hypotheses, as well as including serial assessments to evaluate the effects of illness. Our findings do suggest intact DMN connectivity patterns during the resting state among individuals with highly overlapping anxiety and depressive disorders across a large age-span.

The current findings may provide plausible targets to identify risk for internalizing disorders. They could also be used as targets for novel mechanism-based treatments that may differ based on developmental stage. For instance, there is evidence that attention bias modification training and cognitive remediation can alter connectivity within the CCN among patients with IPs (Beevers, Clasen, Enock, & Schnyer, 2015; Lanius et al., 2015). Alternatively, neurofeedback training utilizing electroencephalography (EEG) has been shown to assist in the regulation of major brain networks such as the SEN (Lanius, Frewen, Tursich, Jetly, & McKinnon, 2015). Lastly, repetitive transcranial stimulation (Fischer, Keller, & Etkin, 2016), transcranial direct current stimulation, and novel pharmacotherapeutic treatments (Watts et al., 2013) developed specifically to target intrinsic network functioning may prove to be beneficial for patients with IPs.

There were several limitations to the current study that should be addressed. First, in order to fully understand the development of these three ICNs, a longitudinal investigation is needed to examine nonlinear age effects, changes within individuals, and how different characteristics of IPs (i.e., episodes, length) influence ICN patterns over time. Second, given the role of pubertal status and sex hormones on connectivity patterns (Klapwijk et al., 2013), it will be important for future studies to examine their influence on patterns of network maturation. Next, because of the sample size and high documented comorbidity among individuals with IPs, we were unable to examine group differences based on specific diagnosis. Although the inclusion of a highly comorbid sample of individuals with IPs is highly generalizable to the community, it will be important for future larger studies to examine how these ICNs may differ for specific IPs across development. Fourth, despite the strength of rs-fMRI for identifying individual differences in ICNs, future studies involving task-based fMRI (e.g., those requiring cognitive control, emotion regulation, and internal thought) might provide additional information about age-related changes and differences in the functioning of these networks in other contexts (e.g., Spreng, 2012).

An additional limitation of the current study was the collection of data across five different studies, which had different goals ranging

from trait assessments in the remitted state to pretreatment assessments in the active state. Despite being able to demonstrate that the current results did not differ across sites and scanners (UIC vs. UM), we were unable to compare findings across the five studies due to inadequate statistical power and nonoverlapping age ranges. Although study site was adjusted for in all analyses, independent replication is needed before strong conclusions can be generated. Additionally, given the use of data across several research studies, we were unable to have uniformed symptom or functioning measures that spanned across development, beyond the broad categorical definition provided. To explore symptom relations in the current study, standardized z-scores were created based on published normative data. However, an important direction for future studies will be to examine whether differences in ICN patterns that span across development relate to broader aspects of behavior and functioning.

Despite these limitations, the current study addresses a gap in the literature by examining how stage of development influences differences in network connectivity patterns between psychiatrically free, TD individuals and those with a history of IPs. Future larger studies are needed to better understand illness/disease factors (e.g., number of episodes, chronicity, family history of IPs) contributing to the observed differences in ICN patterns among individuals with IPs across the different age groups. If replicated, these findings have the potential to inform novel brain-based therapeutics that may differ based on developmental stage.

ACKNOWLEDGMENTS

Subsets of these samples have been reported in prior work, including from MH091811 awarded to Langenecker (Bessette et al., 2018; Bhaumik et al., 2017; DelDonno et al., 2017; Jacobs et al., 2014, 2016; Jenkins et al., 2017; Rao et al., 2016; Stange et al., 2017), MH086517 awarded to Monk and Phan (Hamm et al., 2014; Wu et al., 2016), UIC CCTS UL1TR000050, Mind and Life Foundation, and Klingenstein Third Generation Foundation Fellowship awarded to Jacobs (Jacobs et al., 2016; Peters et al., 2016). Data is also present from MH101487 (awarded to Langenecker). Burkhouse is supported by K23-MH113793, Stange is supported by K23-MH112769, and Bessette is supported by T32-MH067631.

ENDNOTE

¹ Twenty-six individuals (TD = 16, IP = 10) were excluded based on excessive movement (criteria described in "Method" section) during the resting scan.

ORCID

Katie L. Burkhouse  <https://orcid.org/0000-0002-3232-526X>

Jonathan P. Stange  <https://orcid.org/0000-0003-3718-5945>

Scott A. Langenecker  <https://orcid.org/0000-0002-1241-1662>

REFERENCES

- Aldao, A., Nolen-Hoeksema, S., & Schweizer, S. (2010). Emotion-regulation strategies across psychopathology: A meta-analytic review. *Clinical Psychology Review, 30*(2), 217–237.

- Barber, A. D., Caffo, B. S., Pekar, J. J., & Mostofsky, S. H. (2013). Developmental changes in within-and between-network connectivity between late childhood and adulthood. *Neuropsychologia, 51*(1), 156–167.
- Beck, A. T., & Steer, R. A. (1990). *Manual for the Beck anxiety inventory*. San Antonio, TX: Psychological Corporation.
- Beck, A. T., Steer, R. A., & Brown, G. K. (1996). Beck depression inventory-II. *San Antonio, 78*(2), 490–498.
- Beevers, C. G., Clasen, P. C., Enock, P. M., & Schnyer, D. M. (2015). Attention bias modification for major depressive disorder: Effects on attention bias, resting state connectivity, and symptom change. *Journal of Abnormal Psychology, 124*(3), 463–475.
- Behzadi, Y., Restom, K., Liu, J., & Liu, T. T. (2007). A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. *Neuroimage, 37*(1), 90–101.
- Bessette, K. L., Jenkins, L. M., Skerrett, K. A., Gowins, J. R., Deldonno, S. R., Zubieta, J. K., ... Langenecker, S. A. (2018). Reliability, convergent validity and time invariance of default mode network deviations in early adult major depressive disorder. *Frontiers in Psychiatry, 9*, <https://doi.org/10.3389/fpsy.2018.00244>
- Bhaumik, R., Jenkins, L. M., Gowins, J. R., Jacobs, R. H., Barba, A., Bhaumik, D. K., & Langenecker, S. A. (2017). Multivariate pattern analysis strategies in detection of remitted major depressive disorder using resting state functional connectivity. *NeuroImage: Clinical, 16*, 390–398.
- Biswal, B. B., Mennes, M., Zuo, X. N., Gohel, S., Kelly, C., Smith, S. M., ... Dagonowski, A. M. (2010). Toward discovery science of human brain function. *Proceedings of the National Academy of Sciences, 107*(10), 4734–4739.
- Bluhm, R., Williamson, P., Lanius, R., Théberge, J., Densmore, M., Bartha, R., ... Osuch, E. (2009). Resting state default-mode network connectivity in early depression using a seed region-of-interest analysis: Decreased connectivity with caudate nucleus. *Psychiatry and Clinical Neurosciences, 63*(6), 754–761.
- Craighead, W. E., Smucker, M. R., Craighead, L. W., & Ilardi, S. S. (1998). Factor analysis of the children's depression inventory in a community sample. *Psychological Assessment, 10*(2), 156–165.
- Crane, N. A., Jenkins, L. M., Bhaumik, R., Dion, C., Gowins, J. R., Mickey, B. J., ... Langenecker, S. A. (2017). Multidimensional prediction of treatment response to antidepressants with cognitive control and functional MRI. *Brain, 140*(2), 472–486.
- Cullen, K. R., Gee, D. G., Klimes-Dougan, B., Gabbay, V., Hulvershorn, L., Mueller, B. A., ... Lim, K. O. (2009). A preliminary study of functional connectivity in comorbid adolescent depression. *Neuroscience Letters, 460*(3), 227–231.
- Dahl, R. E., & Gunnar, M. R. (2009). Heightened stress responsiveness and emotional reactivity during pubertal maturation: Implications for psychopathology. *Development and Psychopathology, 21*(1), 1–6.
- Dalley, M. B., Bolocofsky, D. N., Alcorn, M. B., & Baker, C. (1992). Depressive symptomatology, attributional style, dysfunctional attitude, and social competency in adolescents with and without learning disabilities. *School Psychology Review, 21*(3), 444–458.
- Davey, C. G., Harrison, B. J., Yücel, M., & Allen, N. B. (2012). Regionally specific alterations in functional connectivity of the anterior cingulate cortex in major depressive disorder. *Psychological Medicine, 42*(10), 2071–2081.
- DelDonno, S. R., Jenkins, L. M., Crane, N. A., Nusslock, R., Ryan, K. A., Shankman, S. A., ... & Langenecker, S. A. (2017). Affective traits and history of depression are related to ventral striatum connectivity. *Journal of affective disorders, 221*, 72–80.
- Deen, B., Pitskel, N. B., & Pelphrey, K. A. (2010). Three systems of insular functional connectivity identified with cluster analysis. *Cerebral cortex, 21*(7), 1498–1506.

- Di Martino, A., Scheres, A., Margulies, D. S., Kelly, A. M. C., Uddin, L. Q., Shehzad, Z., ... & Milham, M. P. (2008). Functional connectivity of human striatum: a resting state fMRI study. *Cerebral cortex*, *18*(12), 2735–2747.
- Dozois, D. J., Dobson, K. S., & Ahnberg, J. L. (1998). A psychometric evaluation of the Beck Depression Inventory-II. *Psychological Assessment*, *10*(2), 83–89.
- Drevets, W. C. (2001). Neuroimaging and neuropathological studies of depression: Implications for the cognitive–emotional features of mood disorders. *Current Opinion in Neurobiology*, *11*(2), 240–249.
- Eklund, A., Nichols, T. E., & Knutsson, H. (2016). Cluster failure: Why fMRI inferences for spatial extent have inflated false-positive rates. *Proceedings of the National Academy of Sciences*, *113*(28), 7900–7905.
- Etkin, A., Egner, T., & Kalisch, R. (2011). Emotional processing in anterior cingulate and medial prefrontal cortex. *Trends in Cognitive Sciences*, *15*(2), 85–93.
- Fair, D. A., Cohen, A. L., Dosenbach, N. U., Church, J. A., Miezin, F. M., Barch, D. M., ... Schlaggar, B. L. (2008). The maturing architecture of the brain's default network. *Proceedings of the National Academy of Sciences*, *105*(10), 4028–4032.
- First, M. B., Spitzer, R. L., & Gibbon, M. W. (2002). *Structured clinical interview for DSM-IV-TR axis I disorders, research version, patient edition (SCID-I/P)*. New York: Biometrics Research.
- Fischer, A. S., Keller, C. J., & Etkin, A. (2016). The clinical applicability of functional connectivity in depression: Pathways toward more targeted intervention. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, *1*(3), 262–270.
- Fox, M. D., & Raichle, M. E. (2007). Spontaneous fluctuations in brain activity observed with functional magnetic resonance imaging. *Nature Reviews Neuroscience*, *8*(9), 700–711.
- Frodil, T., Meisenzahl, E. M., Zetzsche, T., Born, C., Jäger, M., Groll, C., ... Möller, H. J. (2003). Larger amygdala volumes in first depressive episode as compared to recurrent major depression and healthy control subjects. *Biological Psychiatry*, *53*(4), 338–344.
- Gee, D. G., Humphreys, K. L., Flannery, J., Goff, B., Telzer, E. H., Shapiro, M., ... Tottenham, N. (2013). A developmental shift from positive to negative connectivity in human amygdala–prefrontal circuitry. *Journal of Neuroscience*, *33*(10), 4584–4593.
- Gillis, M. M., Haaga, D. A., & Ford, G. T. (1995). Normative values for the Beck Anxiety Inventory, Fear Questionnaire, Penn State Worry Questionnaire, and Social Phobia and Anxiety Inventory. *Psychological Assessment*, *7*(4), 450–455.
- Hamm, L. L., Jacobs, R. H., Johnson, M. W., Fitzgerald, D. A., Fitzgerald, K. D., Langenecker, S. A., ... & Phan, K. L. (2014). Aberrant amygdala functional connectivity at rest in pediatric anxiety disorders. *Biology of mood & anxiety disorders*, *4*(1), 15.
- Hankin, B. L., Abramson, L. Y., Moffitt, T. E., Silva, P. A., McGee, R., & Angell, K. E. (1998). Development of depression from preadolescence to young adulthood: Emerging gender differences in a 10-year longitudinal study. *Journal of Abnormal Psychology*, *107*(1), 128–140.
- Hammar, A., & Ardal, G. (2009). Cognitive functioning in major depression—A summary. *Frontiers in Human Neuroscience*, *3*(26), 1–7.
- Hayes, A. F., & Rockwood, N. J. (2017). Regression-based statistical mediation and moderation analysis in clinical research: Observations, recommendations, and implementation. *Behaviour Research and Therapy*, *98*, 39–57.
- Hulvershorn, L. A., Cullen, K., & Anand, A. (2011). Toward dysfunctional connectivity: A review of neuroimaging findings in pediatric major depressive disorder. *Brain Imaging and Behavior*, *5*(4), 307–328.
- Jacobs, R. H., Jenkins, L. M., Gabriel, L. B., Barba, A., Ryan, K. A., Weisenbach, S. L., ... Gotlib, I. H. (2014). Increased coupling of intrinsic networks in remitted depressed youth predicts rumination and cognitive control. *PLoS One*, *9*(8), e104366.
- Jacobs, R. H., Watkins, E. R., Peters, A. T., Feldhaus, C. G., Barba, A., Carbray, J., & Langenecker, S. A. (2016). Targeting ruminative thinking in adolescents at risk for depressive relapse: Rumination-focused cognitive behavior therapy in a pilot randomized controlled trial with resting state fMRI. *PLoS One*, *11*(11), e0163952.
- Jenkins, L. M., Stange, J. P., Barba, A., DelDonno, S. R., Kling, L. R., Briceño, E. M., ... & Langenecker, S. A. (2017). Integrated cross-network connectivity of amygdala, insula, and subgenual cingulate associated with facial emotion perception in healthy controls and remitted major depressive disorder. *Cognitive, Affective, & Behavioral Neuroscience*, *17*(6), 1242–1254.
- Jo, H. J., Gotts, S. J., Reynolds, R. C., Bandettini, P. A., Martin, A., Cox, R. W., & Saad, Z. S. (2013). Effective preprocessing procedures virtually eliminate distance-dependent motion artifacts in resting state fMRI. *Journal of Applied Mathematics*.
- Jolles, D. D., van Buchem, M. A., Crone, E. A., & Rombouts, S. A. (2010). A comprehensive study of whole-brain functional connectivity in children and young adults. *Cerebral Cortex*, *21*(2), 385–391.
- Kaiser, R. H., Andrews-Hanna, J. R., Wager, T. D., & Pizzagalli, D. A. (2015). Large-scale network dysfunction in major depressive disorder: A meta-analysis of resting-state functional connectivity. *JAMA Psychiatry*, *72*(6), 603–611.
- Kaiser, R. H., & Pizzagalli, D. A. (2015). Dysfunctional connectivity in the depressed adolescent brain. *Biological Psychiatry*, *78*(9), 594–595.
- Kaufman, J., Birmaher, B., Brent, D., Rao, U. M. A., Flynn, C., Moreci, P., ... Ryan, N. (1997). Schedule for affective disorders and schizophrenia for school-age children-present and lifetime version (K-SADS-PL): Initial reliability and validity data. *Journal of the American Academy of Child & Adolescent Psychiatry*, *36*(7), 980–988.
- Keshavan, M. S., Diwadkar, V. A., DeBellis, M., Dick, E., Kotwal, R., Rosenberg, D. R., ... Pettegrew, J. W. (2002). Development of the corpus callosum in childhood, adolescence and early adulthood. *Life Sciences*, *70*(16), 1909–1922.
- Kessler, R. C., Berglund, P., Demler, O., Jin, R., Merikangas, K. R., & Walters, E. E. (2005). Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*, *62*(6), 593–602.
- Kim, M. J., Loucks, R. A., Palmer, A. L., Brown, A. C., Solomon, K. M., Marchante, A. N., & Whalen, P. J. (2011). The structural and functional connectivity of the amygdala: From normal emotion to pathological anxiety. *Behavioural Brain Research*, *223*(2), 403–410.
- Klapwijk, E. T., Goddings, A. L., Heyes, S. B., Bird, G., Viner, R. M., & Blake-More, S. J. (2013). Increased functional connectivity with puberty in the mentalising network involved in social emotion processing. *Hormones and Behavior*, *64*(2), 314–322.
- Kovacs, M. (2004). *Children's depression inventory (CDI)*. Toronto: Multi-Health Systems Inc.
- Kujawa, A., MacNamara, A., Fitzgerald, K. D., Monk, C. S., & Phan, K. L. (2015). Enhanced neural reactivity to threatening faces in anxious youth: Evidence from event-related potentials. *Journal of Abnormal Child Psychology*, *43*(8), 1493–1501.
- Lanius, R. A., Frewen, P. A., Tursich, M., Jetly, R., & McKinnon, M. C. (2015). Restoring large-scale brain networks in PTSD and related disorders: A proposal for neuroscientifically-informed treatment interventions. *European Journal of Psychotraumatology*, *6*(1), 1–12.
- Liao, W., Xu, Q., Mantini, D., Ding, J., Machado-de-Sousa, J. P., Hallak, J. E., ... Crippa, J. A. S. (2011). Altered gray matter morphometry and

- resting-state functional and structural connectivity in social anxiety disorder. *Brain Research*, 1388, 167–177.
- Liu, L., Zeng, L. L., Li, Y., Ma, Q., Li, B., Shen, H., & Hu, D. (2012). Altered cerebellar functional connectivity with intrinsic connectivity networks in adults with major depressive disorder. *PLoS One*, 7(6), e39516.
- Luna, B., Thulborn, K. R., Munoz, D. P., Merriam, E. P., Garver, K. E., Minshew, N. J., ... Sweeney, J. A. (2001). Maturation of widely distributed brain function subserves cognitive development. *Neuroimage*, 13(5), 786–793.
- Ma, Q., Zeng, L. L., Shen, H., Liu, L., & Hu, D. (2013). Altered cerebellar-cerebral resting-state functional connectivity reliably identifies major depressive disorder. *Brain Research*, 1495, 86–94.
- March, J. S., Parker, J. D., Sullivan, K., Stallings, P., & Conners, C. K. (1997). The Multidimensional Anxiety Scale for Children (MASC): Factor structure, reliability, and validity. *Journal of the American Academy of Child & Adolescent Psychiatry*, 36(4), 554–565.
- Marchand, W. R. (2010). Cortico-basal ganglia circuitry: A review of key research and implications for functional connectivity studies of mood and anxiety disorders. *Brain Structure and Function*, 215(2), 73–96.
- MacNamara, A., DiGangi, J., & Phan, K. L. (2016). Aberrant spontaneous and task-dependent functional connections in the anxious brain. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, 1(3), 278–287.
- MacQueen, G. M., Campbell, S., McEwen, B. S., Macdonald, K., Amano, S., Joffe, R. T., ... Young, L. T. (2003). Course of illness, hippocampal function, and hippocampal volume in major depression. *Proceedings of the National Academy of Sciences*, 100(3), 1387–1392.
- Menon, V. (2011). Large-scale brain networks and psychopathology: A unifying triple network model. *Trends in Cognitive Sciences*, 15(10), 483–506.
- Moffitt, T. E., Harrington, H., Caspi, A., Kim-Cohen, J., Goldberg, D., Gregory, A. M., & Poulton, R. (2007). Depression and generalized anxiety disorder: Cumulative and sequential comorbidity in a birth cohort followed prospectively to age 32 years. *Archives of General Psychiatry*, 64(6), 651–660.
- Mulders, P. C., van Eijndhoven, P. F., Schene, A. H., Beckmann, C. F., & Tendolkar, I. (2015). Resting-state functional connectivity in major depressive disorder: A review. *Neuroscience & Biobehavioral Reviews*, 56, 330–344.
- Muris, P., Merckelbach, H., Ollendick, T., King, N., & Bogie, N. (2002). Three traditional and three new childhood anxiety questionnaires: Their reliability and validity in a normal adolescent sample. *Behaviour Research and Therapy*, 40(7), 753–772.
- Nurnberger, J. I., Blehar, M. C., Kaufmann, C. A., York-Cooler, C., Simpson, S. G., Harkavy-Friedman, J., ... Reich, T. (1994). Diagnostic interview for genetic studies: Rationale, unique features, and training. *Archives of General Psychiatry*, 51(11), 849–859.
- Pannekoek, J. N., Werff, S. J. A., Meens, P. H., Bulk, B. G., Jolles, D. D., Veer, I. M., ... Vermeiren, R. R. (2014). Aberrant resting-state functional connectivity in limbic and salience networks in treatment-naïve clinically depressed adolescents. *Journal of Child Psychology and Psychiatry*, 55(12), 1317–1327.
- Pannekoek, J. N., Veer, I. M., van Tol, M. J., van der Werff, S. J., Demenescu, L. R., Aleman, A., ... van der Wee, N. J. (2013). Resting-state functional connectivity abnormalities in limbic and salience networks in social anxiety disorder without comorbidity. *European Neuropsychopharmacology*, 23(3), 186–195.
- Peters, A. T., Burkhouse, K., Feldhaus, C. C., Langenecker, S. A., & Jacobs, R. H. (2016). Aberrant resting-state functional connectivity in limbic and cognitive control networks relates to depressive rumination and mindfulness: A pilot study among adolescents with a history of depression. *Journal of Affective Disorders*, 200, 178–181.
- Perrin, J. S., Merz, S., Bennett, D. M., Currie, J., Steele, D. J., Reid, I. C., & Schwarzbauer, C. (2012). Electroconvulsive therapy reduces frontal cortical connectivity in severe depressive disorder. *Proceedings of the National Academy of Sciences*, 109(14), 5464–5468.
- Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2012). Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. *Neuroimage*, 59(3), 2142–2154.
- Power, J. D., Fair, D. A., Schlaggar, B. L., & Petersen, S. E. (2010). The development of human functional brain networks. *Neuron*, 67(5), 735–748.
- Pujol, J., Vendrell, P., Junqué, C., Martí-Vilalta, J. L., & Capdevila, A. (1993). When does human brain development end? Evidence of corpus callosum growth up to adulthood. *Annals of Neurology*, 34(1), 71–75.
- Raichle, M. E., MacLeod, A. M., Snyder, A. Z., Powers, W. J., Gusnard, D. A., & Shulman, G. L. (2001). A default mode of brain function. *Proceedings of the National Academy of Sciences*, 98(2), 676–682.
- Rao, J. A., Jenkins, L. M., Hymen, E., Feigon, M., Weisenbach, S. L., Zubieta, J. K., & Langenecker, S. A. (2016). Differential resting state connectivity patterns and impaired semantically cued list learning test performance in early course remitted major depressive disorder. *Journal of the International Neuropsychological Society*, 22(2), 225–239.
- Ramasubbu, R., Konduru, N., Cortese, F., Bray, S., Gaxiola, I., & Goodyear, B. (2014). Reduced intrinsic connectivity of amygdala in adults with major depressive disorder. *Frontiers in Psychiatry*, 5, <https://doi.org/10.3389/fpsy.2014.00017>
- Reynolds, W. M. (2004). Reynolds adolescent depression scale. *Comprehensive handbook of psychological assessment*, 2, 224–236.
- Rubia, K. (2013). Functional brain imaging across development. *European Child & Adolescent Psychiatry*, 22(12), 719–731.
- Schmahmann, J. D. (2010). The role of the cerebellum in cognition and emotion: Personal reflections since 1982 on the dysmetria of thought hypothesis, and its historical evolution from theory to therapy. *Neuropsychology Review*, 20(3), 236–260.
- Seeley, W. W., Menon, V., Schatzberg, A. F., Keller, J., Glover, G. H., Kenna, H., ... Greicius, M. D. (2007). Dissociable intrinsic connectivity networks for salience processing and executive control. *Journal of Neuroscience*, 27(9), 2349–2356.
- Sheline, Y. I., Price, J. L., Yan, Z., & Mintun, M. A. (2010). Resting-state functional MRI in depression unmasks increased connectivity between networks via the dorsal nexus. *Proceedings of the National Academy of Sciences*, 107(24), 11020–11025.
- Sheline, Y. I., Sanghavi, M., Mintun, M. A., & Gado, M. H. (1999). Depression duration but not age predicts hippocampal volume loss in medically healthy women with recurrent major depression. *Journal of Neuroscience*, 19(12), 5034–5043.
- Solé-Padullés, C., Castro-Fornieles, J., de la Serna, E., Calvo, R., Baeza, I., Moya, J., ... Sugranyes, G. (2016). Intrinsic connectivity networks from childhood to late adolescence: Effects of age and sex. *Developmental Cognitive Neuroscience*, 17, 35–44.
- Spear, L. P. (2009). Heightened stress responsivity and emotional reactivity during pubertal maturation: Implications for psychopathology. *Development and Psychopathology*, 21(1), 87–97.
- Spreng, R. N. (2012). The fallacy of a "task-negative" network. *Frontiers in Psychology*, 3, <https://doi.org/10.3389/fpsy.2012.00145>
- Stange, J. P., Bessette, K. L., Jenkins, L. M., Peters, A. T., Feldhaus, C., Crane, N. A., ... Langenecker, S. A. (2017). Attenuated intrinsic connectivity within cognitive control network among individuals with remitted

- depression: Temporal stability and association with negative cognitive styles. *Human Brain Mapping*, 38(6), 2939–2954.
- Stoodley, C. J. (2012). The cerebellum and cognition: Evidence from functional imaging studies. *The Cerebellum*, 11(2), 352–365.
- Sylvester, C. M., Corbetta, M., Raichle, M. E., Rodebaugh, T. L., Schlaggar, B. L., Sheline, Y. I., ... Lenze, E. J. (2012). Functional network dysfunction in anxiety and anxiety disorders. *Trends in Neurosciences*, 35(9), 527–535.
- van Duijvenvoorde, A. C. K., Achterberg, M., Braams, B. R., Peters, S., & Crone, E. A. (2016). Testing a dual-systems model of adolescent brain development using resting-state connectivity analyses. *Neuroimage*, 124, 409–420.
- Veer, I. M., Beckmann, C. F., Van Tol, M. J., Ferrarini, L., Milles, J., Veltman, D. J., ... Rombouts, S. A. (2010). Whole brain resting-state analysis reveals decreased functional connectivity in major depression. *Frontiers in Systems Neuroscience*, 41, 1–10.
- Watts, B. V., Schnurr, P. P., Mayo, L., Young-Xu, Y., Weeks, W. B., & Friedman, M. J. (2013). Meta-analysis of the efficacy of treatments for posttraumatic stress disorder. *Journal of Clinical Psychiatry*, 74(6), 541–550.
- Wu, M., Kujawa, A., Lu, L. H., Fitzgerald, D. A., Klumpp, H., Fitzgerald, K. D., ... & Phan, K. L. (2016). Age-related changes in amygdala-frontal connectivity during emotional face processing from childhood into young adulthood. *Human brain mapping*, 37(5), 1684–1695.
- Yeo, B. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M., ... Fischl, B. (2011). The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *Journal of Neurophysiology*, 106(3), 1125–1165.
- Zeng, L. L., Shen, H., Liu, L., Wang, L., Li, B., Fang, P., ... Hu, D. (2012). Identifying major depression using whole-brain functional connectivity: A multivariate pattern analysis. *Brain*, 135(5), 1498–1507.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Burkhouse KL, Stange JP, Jacobs RH, et al. Developmental changes in resting-state functional networks among individuals with and without internalizing psychopathologies. *Depress Anxiety*. 2019;36:141–152. <https://doi.org/10.1002/da.22864>