

Table S1. Stepwise multivariate logistic regression analysis for independent risk factors for thrombosis

Age		β-coefficient	OR (95%CI)	p-value
	<25	0.006	1.00 (0.99-1.02)	0.85
	25-35	-0.16	0.85 (0.47-4.53)	0.29
	>35	0.12	1.13(0.61-2.10)	0.17
Diagnosis of concomitant autoimmune disease		0.06	1.07(0.57-1.98)	0.19
Cardiovascular risk factors (Any)		0.96	2.63(1.18-5.83)	0.01
	<i>Hypertension</i>	0.13	1.15(0.55-2.22)	0.18
	<i>Diabetes on medication</i>	0.07	1.05(0.58-1.99)	0.57
	<i>Hyperlipidemia on medication</i>	0.005	1.03(0.91-1.04)	0.78
	<i>Obesity (BMI > 30)</i>	0.18	1.17(0.65-2.63)	0.12
	<i>Smoking (ever)</i>	0.21	1.15(0.55-2.22)	0.18
aPL (any)		0.23	1.25 (1.78-3.14)	0.11
	Multiple positivity	1.03	2.81 (1.48-5.22)	0.003
Type of pregnancy morbidity				
	<i>≥ Three (pre)-embryonic loss</i>	0.008	1.01 (0.98-1.32)	0.87
	<i>Fetal Loss</i>	0.007	1.00 (0.97-1.05)	0.86
	<i>Premature Birth < 34 week</i>	0.006	1.00 (0.99-1.03)	0.91