

# Space Weather

## MEETING REPORT

10.1029/2018SW002127

**Special Section:**  
Space Weather Capabilities  
Assessment

### Key Points:

- A working team has been established to develop a process for validation of auroral precipitation and electrodynamic models
- Validation of auroral electrodynamic parameters requires generation of ground truth data sets for selected events
- Current observations and data assimilation techniques continue to improve the accuracy of global auroral electrodynamic specification

### Correspondence to:

R. Robinson,  
robert.m.robinson@nasa.gov

### Citation:

Robinson, R., Zhang, Y., Garcia-Sage, K., Fang, X., Verkhoglyadova, O. P., Ngwira, C., et al. (2019). Space weather modeling capabilities assessment: Auroral precipitation and high-latitude ionospheric electrodynamic. *Space Weather*, 17, 212–215. <https://doi.org/10.1029/2018SW002127>

Received 24 NOV 2018






Accepted 11 JAN 2019

Accepted article online 15 JAN 2019

Published online 5 FEB 2019

©2019. American Geophysical Union.  
All Rights Reserved.

## Space Weather Modeling Capabilities Assessment: Auroral Precipitation and High-Latitude Ionospheric Electrodynamics

Robert Robinson<sup>1</sup> , Yongliang Zhang<sup>2</sup> , Katherine Garcia-Sage<sup>1</sup> , Xiaohua Fang<sup>3</sup> , Olga P. Verkhoglyadova<sup>4</sup> , Chigomezzyo Ngwira<sup>1</sup> , Suzy Bingham<sup>5</sup>, Burcu Kosar<sup>1</sup> , Yihua Zheng<sup>6</sup> , Stephen Kaeppler<sup>7</sup>, Michael Liemohn<sup>8</sup> , James B. Weygand<sup>9</sup> , Geoffrey Crowley<sup>10</sup> , Viacheslav Merkin<sup>2</sup> , Ryan McGranaghan<sup>4</sup> , and Anthony J. Mannucci<sup>4</sup> 

<sup>1</sup>Department of Physics, The Catholic University of America, Washington, DC, USA, <sup>2</sup>Johns Hopkins University Applied Physics Laboratory, Laurel, MD, USA, <sup>3</sup>University of Colorado Laboratory for Atmospheric and Space Physics, <sup>4</sup>Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA, <sup>5</sup>UK Met Office, Exeter, UK, <sup>6</sup>NASA Goddard Space Flight Center, Greenbelt, MD, USA, <sup>7</sup>Department of Physics and Astronomy, Clemson University, Clemson, SC, USA, <sup>8</sup>Climate and Space Sciences and Engineering, University of Michigan, Ann Arbor, MI, USA, <sup>9</sup>Earth, Planetary, and Space Sciences, University of California, Los Angeles, CA, USA, <sup>10</sup>ASTRA, Greenbelt, MD, USA

**Abstract** As part of its International Capabilities Assessment effort, the Community Coordinated Modeling Center initiated several working teams, one of which is focused on the validation of models and methods for determining auroral electrodynamic parameters, including particle precipitation, conductivities, electric fields, neutral density and winds, currents, Joule heating, auroral boundaries, and ion outflow. Auroral electrodynamic properties are needed as input to space weather models, to test and validate the accuracy of physical models, and to provide needed information for space weather customers and researchers. The working team developed a process for validating auroral electrodynamic quantities that begins with the selection of a set of events, followed by construction of ground truth databases using all available data and assimilative data analysis techniques. Using optimized, predefined metrics, the ground truth data for selected events can be used to assess model performance and improvement over time. The availability of global observations and sophisticated data assimilation techniques provides the means to create accurate ground truth databases routinely and accurately.

### 1. Meeting Report

As part of its International Capabilities Assessment effort (see <https://ccmc.gsfc.nasa.gov/assessment/forum-topics.php>), the Community Coordinated Modeling Center (CCMC) initiated several working teams, one of which is focused on auroral precipitation and high-latitude ionospheric electrodynamic model validation. The goal of the Auroral Precipitation and High Latitude Electrodynamic (AuroraPHILE) working team is to establish quantitative means to measure the accuracy and reliability of modeled properties of the auroral ionosphere, including particle precipitation, conductivities, electric fields, neutral density and winds, currents, Joule heating, auroral boundaries, and ion outflow. The working team's objective is to establish a set of properties that describe the state of auroral particle precipitation and electrodynamic and then quantify the accuracy and reliability currently achievable using a combination of data and models. Working team discussions were held during the International CCMC-Living With a Star Working Meeting: Assessing Space Weather Understanding and Applications, 3–7 April 2017, in Cape Canaveral, Florida ([https://ccmc.gsfc.nasa.gov/CCMC-LWS\\_Meeting/](https://ccmc.gsfc.nasa.gov/CCMC-LWS_Meeting/)) and in teleconferences before and after the meeting.

Properties of the auroral ionosphere are critical for improving resilience to impacts of space weather events. Auroral electrodynamic properties are needed as input to space weather models, to test and validate the accuracy of physical models, and to provide needed information for space weather customers and researchers. The aurora is a manifestation of energy input to the upper atmosphere that heats the thermosphere, resulting in increased satellite drag. Auroral precipitation modifies the ionospheric electrical conductivity, needed to specify and predict the currents causing ground-based magnetic perturbations that threaten the electric power grid. Through ionization and convection, the aurora modifies the ionospheric electron density,

Auroral Parameter:	Energetic Particle Fluxes	Auroral Electrical Conductances	Ionospheric Electric Fields	Ionospheric Currents	Field-aligned Currents	Joule Heating	Auroral Boundaries	Ionospheric Electron Density	Neutral Density and Winds	Ion Outflow	Poynting Flux
<b>Space Weather Application</b>											
Satellite Drag	S	S	S	S	S	S	S	S	P	S	S
HF Radio Propagation	S	S	S	S	S	S	S	P	S	S	S
Transionospheric Radio Propagation	S	S	S	S	S	S	S	P	S	S	S
Navigation	S	S	S	S	S	S	S	P	S	S	S
Satellite Operations	P	S	S	S	S	S	S	S	S	S	S
Human Spaceflight	P	S	S	S	S	S	S	S	S	S	S
Commercial Aviation--Radiation	P	S	S	S	S	S	S	S	S	S	S
Commercial Aviation--Comm and Nav	S	S	S	S	S	S	S	P	S	S	S
Electric Power	S	S	S	P	S	S	S	S	S	S	S

**Figure 1.** Auroral parameters important for mitigating impacts to space weather applications.


resulting in disturbances and disruptions to transionospheric radiowave transmissions needed for navigation and communication. Auroral electrodynamic parameters are also needed as input to and validation of many different types of space weather models. Finally, an accurate specification of auroral properties is important for assessing surface charging effects on space assets traversing through the auroral zones.

Figure 1 lists the space weather applications for which auroral parameters are important. For each application, the orange highlight indicates the primary (*P*) auroral property that must be modeled or observed to mitigate the associated space weather effects. The yellow highlights indicate secondary (*S*) properties that either indirectly impact the application or are needed as input to accurately model and predict the impact. Given the overall importance of auroral properties to mitigating space weather effects on applications, it is essential to quantitatively assess the accuracy with which those properties can be observed and modeled.

The AuroraPHILE working team began by compiling comprehensive lists of models, data, and data sources (both ground based and space based) available over the past 25 years that can be used to test and validate knowledge of the auroral ionosphere and the capability to both specify and forecast high-latitude ionospheric properties. These lists are accessible on the CCMC web site (<https://ccmc.gsfc.nasa.gov/assessment/topics/iono-joule.php>). Based on a careful consideration of the available models and data and their associated uncertainties, the working team developed a methodology for assessing the accuracy with which auroral precipitation and high-latitude electrodynamic quantities can be specified and forecast. For a pre-selected group of events, all available data would be used to determine the most accurate values of auroral electrodynamic parameters. We refer to this as the *ground truth* data set, although the values may be determined by a combination of direct measurements, data assimilation, and other models needed to fill in gaps and extend observations. Once the optimum ground truth data set has been determined, any model can be tested with respect to its accuracy in replicating *reality*. Thus, all models will be evaluated against the same standard and for the same events. New models, or upgrades to existing models, would be tested against the same events so that improvements can be unambiguously tracked over time.

In considering the set of events to include in the ground truth database, the AuroraPHILE working team noted the importance of including a broad range of geomagnetic conditions. Overlap with events selected by other working teams will help facilitate the assembly of observations for ground truth data sets. For example, the AuroraPHILE working team overlaps with the Geospace Environment Modeling Challenge working group and other CCMC working teams developing metrics for ionospheric parameters and geomagnetic indices (see, e.g., Liemohn et al., 2018; Welling et al., 2018).

Because auroral properties span a broad range of temporal and spatial scales and are highly variable in space and time, assembling the necessary measurements to create ground truth databases is a major



**Figure 2.** Functional relationship between two dimensional (in rectangles) and global (in circles) auroral electrodynamic parameters. HPI is the Hemispheric Power Index (Fuller-Rowell & Evans, 1987).

challenge. The current availability of ground-based and space-based measurements of auroral properties, coupled with the improving sophistication of assimilative models, has made specification of high-latitude electrodynamic parameters more accurate and more global than ever before. Figure 2 shows how auroral electrodynamic parameters are functionally connected. Two-dimensional parameters are in rectangular boxes and global quantities are indicated by circles. With the availability of global field-aligned currents from the Iridium satellite constellation (Anderson et al., 2000) and Active Magnetosphere and Planetary Response Experiment (Anderson et al., 2014), one very important piece of the puzzle is now in place. Conductances are also critical to the calculation of electrodynamic parameters as they are used to compute electric fields, currents, Joule heating, and precipitating particle energy flux.

Accurate identification of auroral boundaries is important for many space weather applications. Both poleward and equatorward boundaries are often necessary, and boundaries may differ depending on the process or phenomenon that is most important to the application. Boundary identification algorithms have used optical observations from the

IMAGE (Imager for Magnetopause-to-Aurora Global Exploration) satellite (Longden et al., 2010) and from the Global Ultraviolet Imager on the Thermosphere Ionosphere Mesosphere Energetics and Dynamics satellite (Christensen et al., 2003; Zhang et al., 2010), and from DMSP (Defense Meteorological Satellite Program) measurements of precipitating particles (Kilcommons et al., 2017; Ngwira et al., 2013; Redmon et al., 2010). A novel approach to auroral boundary identification is the Aurorasaurus project database (Case et al., 2016; MacDonald et al., 2015), which offers a collection of geotagged and time-stamped signals of auroral visibility collected from citizen scientists.

The AuroraPHILE working team identified a number of different observations that can be used with sophisticated assimilative mapping programs to fill in gaps, constrain measurements, and minimize inconsistencies (e.g., Cosgrove et al., 2009, 2014). In some cases, quantities such as Joule heating cannot be measured directly but are calculated with certain assumptions from other validated measurements (Thayer, 1998; Verkhoglyadova et al., 2016, 2017). When the ground truth data are model dependent, all the model assumptions must be thoroughly documented for future review and possible revision. Another important aspect of constructing a ground truth database is specifying the errors and uncertainties in the results. These errors and uncertainties can arise not only from the measurements but also from the models used to derive physical quantities from the observations.

The working team discussed approaches to quantitatively assess model results using a carefully selected collection of metrics. For any given auroral electrodynamic parameter, there may be several metrics by which to compare model output and ground truth data. Metrics can be user dependent (e.g., operational versus scientific metrics). In one case the timing of an event may be more important than the amplitude of the parameter. For some applications, the ability to capture small-scale or highly time-varying features will be more important than capturing the large-scale changes taking place. To constrain the number of metrics for auroral electrodynamic parameters, some compromises are inevitable. A good metric is one that will reflect overall improvement in model capability for all or most applications. Additionally, metrics are most useful when they not only assess the validity and accuracy of models but also provide information about the source of model strengths or weaknesses. Quantifying model accuracy in the presence of rapidly time-evolving patterns can lead to different results depending on the resolution of the model and the data (see, e.g., Merkin et al., 2007, 2013). Different metrics should be used for two-dimensional (or three-dimensional) images as opposed to time series data. An example is a multidimensional correlation coefficient, including both spatial and temporal variables. The metric multidimensional root-mean-square error is another approach. Other approaches include calculation of the median absolute deviation (see, e.g., McGranaghan et al., 2016). Alternatively, image recognition software used for other applications may be appropriate for comparing model results with global measurements (e.g., Wiltberger et al., 2017). Specific metrics for forecast evaluation have been described by Murphy (1991) and Kubo et al. (2017).

The AuroraPHILE working team recommended next steps to implement the planned model validation activities. Essential to the process is the construction of the ground truth data sets for selected events. The working team discussions highlighted the improved capabilities currently available for global and continuous specification of auroral electrodynamic parameters. Although far from ideal, the AuroraPHILE working team concluded that accuracies are sufficient for model validation and testing, and for monitoring the improvement in models over time.

#### Acknowledgments

This work was supported at Goddard Space Flight Center by NASA Cooperative Agreement NNG11PL10A and the Community Coordinated Modeling Center. Portions of this research were carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with NASA. R. M. M. was supported by the NASA Living With a Star Jack Eddy Postdoctoral Fellowship Program, administered by the University Corporation for Atmospheric Research and coordinated through the Cooperative Programs for the Advancement of Earth System Science (CPAESS). Garcia-Sage's effort was supported by the NASA Heliophysics Internal Scientist Funding Model (HISFM18-0006). M. W. L.'s effort was supported by NSF grant 1663770. No new data, models, or model output were created in preparing this meeting report. Models, data, and data sources considered by the AuroraPHILE Working Team are available through the CCMC web site at <https://ccmc.gsfc.nasa.gov/assessment/topics/iono-joule.php>.

#### References

- Anderson, B. J., Korth, H., Waters, C. L., Green, D. L., Merkin, V. G., Barnes, R. J., & Dyrud, L. P. (2014). Development of large-scale Birkeland currents determined from the active magnetosphere and planetary electrodynamics response experiment. *Geophysical Research Letters*, *41*, 3017–3025. <https://doi.org/10.1002/2014GL059941>
- Anderson, B. J., Takahashi, K., & Toth, B. A. (2000). Sensing global Birkeland currents with iridium engineering magnetometer data. *Geophysical Research Letters*, *27*(24), 4045–4048. <https://doi.org/10.1029/2000GL000094>
- Case, N. A., MacDonald, E. A., & Viereck, R. (2016). Using citizen science reports to define the equatorial extent of auroral visibility. *Space Weather*, *14*, 198–209. <https://doi.org/10.1002/2015SW001320>
- Christensen, A. B., Christensen, A. B., Paxton, L. J., Avery, S., Craven, J., Crowley, G., et al. (2003). Initial observations with the Global Ultraviolet Imager (GUVI) in the NASA TIMED satellite mission. *Journal of Geophysical Research*, *108*(A12), 1451. <https://doi.org/10.1029/2003JA009918>
- Cosgrove, R. B., Bahcivan, H., Chen, S., Strangeway, R. J., Ortega, J., Alhassan, M., et al. (2014). Empirical model of Poynting flux derived from FAST data and a cusp signature. *Journal of Geophysical Research: Space Physics*, *119*, 411–430. <https://doi.org/10.1002/2013JA019105>
- Cosgrove, R. B., Lu, G., Bahcivan, H., Matsuo, T., Heinselman, C. J., & McCready, M. A. (2009). Comparison of AMIE-modeled and Sondrestrom-measured Joule heating: A study in model resolution and electric field–conductivity correlation. *Journal of Geophysical Research*, *114*, A04316. <https://doi.org/10.1029/2008JA013508>
- Fuller-Rowell, T. J., & Evans, D. S. (1987). Height-integrated Pedersen and Hall conductivity patterns inferred from the TIROS-NOAA satellite data. *Journal of Geophysical Research*, *92*, 7606–7618.
- Kilcommons, L., Redmon, R. J., & Knipp, D. J. (2017). A new DMSP magnetometer dataset and estimates of field aligned currents in dynamic auroral boundary coordinates. *Journal of Geophysical Research: Space Physics*, *122*, 9068–9079. <https://doi.org/10.1002/2016JA023342>
- Kubo, Y., Den, M., & Ishii, M. (2017). Verification of operational solar flare forecast: Case of Regional Warning Center Japan. *Space Weather and in Space Climate*, *7*, A20. <https://doi.org/10.1051/swsc/2017018>
- Liemohn, M. W., McCollough, J. P., Jordanova, V. K., Ngwira, C. M., Morley, S. K., Cid, C., et al. (2018). Model evaluation guidelines for geomagnetic index prediction. *Space Weather*, *16*. <https://doi.org/10.1029/2018SW002067>, 2079–2102.
- Longden, N., Chisham, G., Freeman, M. P., Abel, G. A., & Sotirelis, T. (2010). Estimating the location of the open-closed magnetic field line boundary from auroral images. *Annales de Geophysique*, *28*(9), 1659–1678. <https://doi.org/10.5194/angeo-28-1659-2010>
- MacDonald, E. A., Case, N. A., Clayton, J. H., Hall, M. K., Heavner, M., Lalone, N., et al. (2015). Aurorasaurus: A citizen science platform for viewing and reporting the aurora. *Space Weather*, *13*, 548–559. <https://doi.org/10.1002/2015SW001214>
- McGranaghan, R., Knipp, D. J., Matsuo, T., & Cousins, E. (2016). Optimal interpolation analysis of high-latitude ionospheric Hall and Pedersen conductivities: Application to assimilative ionospheric electrodynamics reconstruction. *Journal of Geophysical Research: Space Physics*, *121*, 4898–4923. <https://doi.org/10.1002/2016JA022486>
- Merkin, V. G., Anderson, B. J., Lyon, J. G., Korth, H., Wiltberger, M., & Motoba, T. (2013). Global evolution of Birkeland currents on 10 min timescales: MHD simulations and observations. *Journal of Geophysical Research: Space Physics*, *118*, 4977–4997. <https://doi.org/10.1002/jgra.50466>
- Merkin, V. G., Lyon, J. G., Anderson, B. J., Korth, H., Goodrich, C. C., & Papadopoulos, K. (2007). A global MHD simulation of an event with a quasi-steady northward IMF component. *Annales Geophysicae*, *25*(6), 1345–1358.
- Murphy, A. H. (1991). Forecast verification: Its complexity and dimensionality. *Monthly Weather Review*, *119*, 1590–1601. [https://doi.org/10.1175/1520-0493\(1991\)119<1590:FVICAD>2.0.CO;2](https://doi.org/10.1175/1520-0493(1991)119<1590:FVICAD>2.0.CO;2)
- Ngwira, C. M., Pulkkinen, A., Wilder, F. D., & Crowley, G. (2013). Extended study of extreme geoelectric field event scenarios for geomagnetically induced current applications. *Space Weather*, *11*, 121–131. <https://doi.org/10.1002/swe.20021>
- Redmon, R. J., Peterson, W. K., Andersson, L., Kihn, E. A., Denig, W. F., Hairston, M., & Coley, R. (2010). Vertical thermal O+ flows at 850 km in dynamic auroral boundary coordinates. *Journal of Geophysical Research*, *115*, A00J08. <https://doi.org/10.1029/2010JA015589>
- Thayer, J. P. (1998). Height-resolved joule heating rates in the high-latitude E region and the influence of neutral winds. *Journal of Geophysical Research*, *103*(A1), 471–487. <https://doi.org/10.1029/97JA02536>
- Verkhoglyadova, O. P., Meng, X., Mannucci, A. J., Mlynczak, M. G., Hunt, L. A., & Lu, G. (2017). Ionosphere-thermosphere energy budgets for the ICME storms of March 2013 and 2015 estimated with GITM and observational proxies. *Space Weather*, *15*, 1102–1124. <https://doi.org/10.1002/2017SW001650>
- Verkhoglyadova, O. P., Meng, X., Mannucci, A. J., Tsurutani, B. T., Hunt, L. A., Mlynczak, M. G., et al. (2016). Estimation of energy budget of ionosphere-thermosphere system during two CIR-HSS events: Observations and modeling. *Journal of Space Weather and Space Climate*, *6*, A20. <https://doi.org/10.1051/swsc/2016013>
- Welling, D. T., Ngwira, C. M., Opgenoorth, H., Haiducek, J. D., Savani, N. P., Morley, S. K., Cid, C., et al. (2018). Recommendations for next-generation ground magnetic perturbation validation. *Space Weather*, *16*, 1912–1920. <https://doi.org/10.1029/2018SW002064>
- Wiltberger, M., Rigler, E. J., Merkin, V., & Lyon, J. G. (2017). Structure of high latitude currents in magnetosphere-ionosphere models. *Space Science Reviews*, *206*, 575. <https://doi.org/10.1007/s11214-016-0271-2>
- Zhang, Y., Paxton, L., Bilitza, D., & Doe, R. (2010). Near real-time assimilation in IRI of auroral peak E-region density and equatorward boundary. *Advances in Space Research*, *46*(8), 1055–1063.