

JGR Space Physics

COMMENTARY

10.1029/2018JA026282

Key Points:

- Advancements on first-principle ring current models, new empirical models on IM fields/waves/plasma and application of innovative techniques
- Advanced knowledge of IM characteristics, e.g., compositions, fields, coupling with ionosphere/tail region, and wave particle interactions
- Challenges remain in numerical representation of IM and its linkage with other related areas; validation needed across various IM models

Correspondence to:

Y. Yu,
yiqunyu17@gmail.com

Citation:

Yu, Y., Liemohn, M. W., Jordanova, V. K., Lemon, C., & Zhang, J. (2019). Recent advancements and remaining challenges associated with inner magnetosphere cross-energy/population interactions (IMCEPI). *Journal of Geophysical Research: Space Physics*, 124, 886–897. <https://doi.org/10.1029/2018JA026282>

Received 12 NOV 2018

Accepted 17 JAN 2019

Accepted article online 31 JAN 2019

Published online 23 FEB 2019

©2019. American Geophysical Union.
All Rights Reserved.

Recent Advancements and Remaining Challenges Associated With Inner Magnetosphere Cross-Energy/Population Interactions (IMCEPI)

Yiqun Yu¹ , Mike W. Liemohn² , Vania K. Jordanova³ , Colby Lemon⁴ , and Jichun Zhang⁵ 

¹School of Space and Environment, Beihang University, Beijing, China, ²Department of Climate and Space Sciences and Engineering, University of Michigan, Ann Arbor, MI, USA, ³Space Science and Application, Los Alamos National Laboratory, Los Alamos, NM, USA, ⁴The Aerospace Corporation, El Segundo, CA, USA, ⁵Previously at Space Science Center and Department of Physics, University of New Hampshire, Durham, NH, USA

Abstract The geospace inner magnetosphere, within about 10 Earth radii, contains various plasma populations with energy from a few electron volts to megaelectron volts and plays important roles in regulating the energy density of the magnetosphere, the magnetic field configuration, and wave dynamics. As an integrated part of the magnetosphere, the inner magnetosphere region also ties to other regions and can change the global geospace circulation. Therefore, understanding both internal and external cross-energy/population interactions can help further our knowledge of the inner magnetosphere dynamics and nonlinear feedback processes. In view of this, in the past 5 years (2014–2018), the Geospace Environment Modeling (GEM) Focus Group “inner magnetosphere cross-energy/population interactions (IMCEPI)” has gathered and boosted community-wide interactions among observation, simulation, and modeling studies. This commentary reports some major accomplishments of the interactive inner magnetosphere community that were advanced by the IMCEPI Focus Group discussions and layouts remaining challenges that need to be carried on.

1. Introduction

The GEM Inner magnetosphere cross-energy/population interactions (IMCEPI) Focus Group (FG) was termed from 2014 to 2018, aiming to improve physical knowledge and modeling of the inner magnetosphere, particularly the ring current interactions and feedback with other populations in the magnetosphere and ionosphere. There are three main plasma populations in the inner magnetosphere, which reside in an overlapped region around the Earth ($3 < L < 7$) and are categorized based on their distinct features. The plasmasphere is populated with cold ions (< 1 eV) with density of hundreds of to thousands of per cubic centimeters. The ring current mainly consists of ions with energy from hundreds of electron volts to hundreds of kiloelectron volts with density of few per cubic centimeters. The radiation belts are mostly dominant by megaelectron volt electrons with tenuous density ($\ll 1$ cm⁻³). Figure 1 shows the coupling relations between different populations via many processes. Although the plasmasphere and radiation belts do not contribute significantly to the inner magnetosphere current systems (Ganushkina, Liemohn, et al., 2015), they still play a very important role in the dynamics of the inner magnetosphere. The plasmasphere influences the plasma wave environment that controls pitch angle scattering rates and subsequent precipitation losses of ring current and radiation belt particles (e.g., Albert, 2004; Jordanova et al., 2001). In addition to directly altering the phase space density in the inner magnetosphere, the precipitation also enhances ionospheric conductivity and outflow rate, which modifies the inner magnetospheric electric field and composition and in turn influences the magnetic field via the modified current systems (Ebihara et al., 2004; Liemohn et al., 2005). All of these require the proper interactions among different populations and regimes in inner magnetosphere models.

The broad scientific goals of IMCEPI FG were, through implementation of physics in existing models, to contribute to the physics-based understanding of (1) the mechanisms responsible for the ring current growth and decay, (2) the interactions with particles in other regions, and (3) the nonlinear feedback mechanisms. The main deliverable was to obtain more comprehensive, self-consistent physics-based circulation models, the ultimate goal of the NSF/GEM program. The numerical blueprint is demonstrated in Figure 2, a framework for modeling the coupled inner magnetosphere system, which puts forward the guiding picture for models.




Figure 1. The schematics of inner magnetosphere coupling physics, showing the connection between the electric/magnetic fields with particles, together with other effects (adopted from Liemohn, 2006).

In this commentary, we describe recent achievements in the inner magnetosphere community toward the above goals and discuss near-future challenges remaining in the field. Note that roughly a dozen FGs are active at any one time within the GEM program, including several focusing on other aspects of inner magnetospheric physics. The brief review below is not meant to be comprehensive but rather highlighting advancements in the specific scientific goals of the IMCEPI FG mentioned above.

2. Advancements on IMCEPI Topics During the IMCEPI Years (2014–2018)

In the past 5 years, the physics-based inner magnetosphere models become more mature, more self-consistent with more physics included. Different models are interconnected to represent the sophisticated,




Figure 2. The schematics of the modeling framework for the inner magnetosphere coupling processes. The center is the primary inner magnetosphere model for plasmasphere (PS), ring current (RC), and radiation belt (RB). The surrounding model components represent processes being coupled with the inner magnetosphere populations.

nonlinearly coupled geospace system, allowing for a better understanding of the internal interactions. Compared to earlier models (e.g., Fok et al., 1993, 2001; Ilie et al., 2012; Jordanova et al., 1994, 2010; Lemon et al., 2004; Liemohn et al., 2001; Toffoletto et al., 2003), the inner magnetosphere models are now capable of resolving particle dynamics across a broader range of energy or regions, covering thermal-energy plasmasphere, warm ring current particles, and energetic radiation belt populations (Fok et al., 2014; Ganushkina, Amariutei, et al., 2015; Huba & Sazykin, 2014; Huba et al., 2017; Jordanova et al., 2014, 2016; Krall et al., 2017); they are more self-consistently linked with the ionosphere system by taking into account more physics-based ionosphere-thermosphere processes (Raeder et al., 2016; Wiltberger et al., 2017; Xi et al., 2016; Yu et al., 2018); they can be driven by various tail dynamics using different approaches such as injecting particles within prescribed electromagnetic fields (e.g., Brito et al., 2017; Ganushkina et al., 2014; Jordanova et al., 2018) or by earthward propagating bubbles (e.g., Cramer et al., 2017; Yang et al., 2015, 2016). They also include more realistic representation of the influence of plasma waves by including more types of waves or using newly derived pitch angle/energy/cross-energy diffusion coefficients or loss rates based on tremendously increased data base in space, leading to significant improvement in the modeling of the energization/decay of inner magnetosphere populations (e.g., Aryan et al., 2017; Jordanova et al., 2016; Kang et al., 2016; Ma et al., 2018; Tu et al., 2014) and ionospheric precipitation/conductance (Chen, Lemon, Guild, et al., 2015; Chen, Lemon, Orlova, et al., 2015; Perlongo et al., 2017; Yu et al., 2016). Following earlier efforts in combining kinetic models with global MHD models (De Zeeuw et al., 2004; Ebihara & Tanaka, 2013; Glocer et al., 2009, 2013; Pembroke et al., 2012), more ring current models have been equipped with such capability during the past few years by coupling with global MHD models (e.g., Cramer et al., 2017; Raeder et al., 2016; Welling et al., 2018; Yu et al., 2017, 2014; Y. Yu et al., 2015). The above advanced models are largely capable of reproducing various particle dynamics within the global magnetosphere and providing important feedback processes on particle populations, the electric/magnetic fields, and dynamics in other geospace regions.

While the numerical representation of the geospace system is being greatly improved, better characterization of physics in the near-Earth environment has also been consistently obtained. For example, the wave dynamics and their impact on plasma dynamics have been analyzed in great detail (e.g., Blum et al., 2015, 2016, 2017; Fu et al., 2017; Kim et al., 2016; W. Li et al., 2014; J. Li, Ni, et al., 2016; L. Y. Li, Yu, et al., 2016, 2017; L. Li, Zhou, et al., 2017; Ma, Li, et al., 2016; Murphy et al., 2015; Shi et al., 2018; Shprits, Drozdov, et al., 2016; Turner et al., 2014, 2017; Usanova et al., 2016; Wang, Pan, et al., 2017; Wang, Zhai, et al., 2017; J. Yu et al. 2015; J.-C. Zhang et al., 2014; X.-J. Zhang et al., 2016; Zhao et al., 2014; Zhou et al., 2015). Furthermore, the diffusion rates of inner magnetosphere particles have been upgraded from simple and empirical rates (e.g., Albert, 1999; Schulz, 1998) to more comprehensive, pitch angle and energy-resolved, and even event-specific diffusion rates based on a larger plasma wave data set. These new rates account for a variety of responsible causes, including electromagnetic ion cyclotron (EMIC) waves (e.g., Cao et al., 2016; Kang et al., 2015; Kersten et al., 2014; Ni et al., 2015; Usanova et al., 2014), whistler mode waves (e.g., Agapitov et al., 2014, 2018; Glauert et al., 2014; Horne et al., 2013; Ma et al., 2017; Mourenas et al., 2014; Orlova et al., 2014; Orlova & Shprits, 2015; Ripoll et al., 2017), field line curvature scattering (Ji & Shen, 2014), magnetosonic waves (Shprits, 2016), and radial diffusion (e.g., Ali et al., 2016; Liu et al., 2016). In addition, electron scattering and acceleration by broad electrostatic turbulence around plasma injection regions (e.g., Ma, Mourenas, et al., 2016; Mozer et al., 2015) and electron losses due to Alfvén waves (e.g., Chaston et al., 2018; Malaspina et al., 2015) are found to play important roles in the inner magnetosphere dynamics.

Compelling knowledge is also achieved in the area of inner magnetosphere-ionosphere coupling. For example, it was found that small-scale electrojet turbulence plays a critical role in changing the electrodynamics in the ionosphere and the plasma transport in the inner magnetosphere (Wiltberger et al., 2017). Field-aligned electric potential drop, which had been long neglected in global models, was incorporated for the first time into the global MHD model and was found to impact remarkably on the tail reconnection as well as the plasma transport in supplying sources to the inner magnetosphere (Xi et al., 2016). Observational evidence of field-aligned currents (both regions 1 and 2) connecting the ionosphere with the magnetosphere was discovered with the aid of joint operation of Cluster and Swarm missions (Dunlop et al., 2015). Systematic physical insights were also gained on the origin of diffuse auroras (Ni et al., 2016; X.-J. Zhang et al., 2015), and observational evidence was found for the chorus wave-associated precipitation being the

driver of pulsating aurora (Kasahara et al., 2018). Moreover, during precipitation, the affiliated production of secondary superthermal electrons reflected between hemispheres was found to participate in ionosphere-magnetosphere energy redistribution (Khazanov et al., 2014, 2015, 2017), enhancing ionospheric conductance (Khazanov et al., 2018). Other ionospheric phenomena were also investigated to search for the magnetospheric drivers. For instance, subauroral arcs in the premidnight during substorms were found to be connected to localized ring current pressure gradients in the R2 source region (Motoba et al., 2015). The subauroral proton aurora was suggested to link to the flow bursts moving from the tail to the inner magnetosphere (Nishimura et al., 2014). In addition, a narrow luminous structure across the night sky in the subauroral region, called the Strong Thermal Emission Velocity Enhancement or “STEVE,” recently caught the attention of the scientific community. However, Gallardo-Lacourt et al. (2018) found that STEVE is unrelated to magnetospheric particle precipitation and is likely to be generated by ionosphere-thermosphere interactions.

The in situ and ground-based data sets have tremendously grown in the past few years due to active spacecraft missions such as Cluster, Van Allen Probes, and Arase, providing substantial data samples for understanding the inner magnetosphere environment. The ample data sets help derive long-term trends and evolutions of the physical processes within the inner magnetosphere in response to different driving conditions, by looking into various wave properties (e.g., occurrence rate, amplitudes, spatial sizes, obliquity, and propagation; Artemyev et al., 2016; Aryan et al., 2014, 2016; Fu et al., 2014; Kersten et al., 2014; W. Li et al., 2015; W. Li et al., 2016; Malaspina et al., 2016, 2017; Meredith et al., 2014; Nemec et al., 2016; Ni et al., 2017; Saikin et al., 2015; Santolík, Macúsová, et al., 2014; Spasojevic et al., 2015; Yue et al., 2017; Zhima et al., 2014, 2015), inner magnetosphere plasma compositions (Claudepierre et al., 2016; Fernandes et al., 2017; Kistler & Mouikis, 2016; Kistler, Mouikis, Spence, et al., 2016; Sarno-Smith et al., 2015; Yue et al., 2018), ion mass density along closed magnetic field lines as well as mass loading in response to geomagnetic activity levels (Sandhu et al., 2016, 2017), plasma sheet composition outside the inner magnetosphere (e.g., Denton et al., 2017; Nosé, 2016), the spacecraft surface charging environment (Sarno-Smith et al., 2016), plasmopause model (He et al., 2017; Liu et al., 2015; X.-X. Zhang et al., 2017), electric field model (Califf et al., 2014), and global magnetospheric field model (Tsyganenko & Andreeva, 2017), which upgraded from a series of previous models (e.g., Tsyganenko, 1989, 1996, 2002; Tsyganenko & Sitnov, 2007).

With many kinds of inner magnetosphere models coexisting in the community, their application values are of particular interest for operational purposes. “Challenges” on realistically representing geospace have been conducted in order to show the capability and limitation of individual models. Therefore, a few models have participated in the challenges to determine different capabilities, such as reproducing global indices (i.e., Dst and Kp indices; Glocer et al., 2016; Rastätter et al., 2013), ground-based magnetic perturbations ΔB or dB/dt (Pulkkinen et al., 2013; Rastätter et al., 2014; Welling et al., 2017), radiation belt electron dynamics (Ma et al., 2018), and spacecraft surface charging environment (Yu et al., 2019). These challenge studies have shown to the community how these models compare to each other in capturing features of the system and in what circumstances they are applicable.

3. Future Challenges

Although remarkable progress has been made in modeling and understanding the inner magnetosphere during the IMCEPI years, a number of challenges still stand ahead. Modeling of the coupling of the inner magnetosphere to other regions remains an important topic, especially the coupling to the magnetotail and ionosphere. How the injected plasma is supplied to the inner magnetosphere and how the inner magnetosphere populations couple with the ionosphere-thermosphere via a more physics-based approach challenges the current techniques because these regions involve different physics and any single theoretical method cannot satisfy these complex cross-scale interactions.

In order to better represent the electromagnetic drivers for the inner magnetospheric plasma, it is necessary to capture the mutually consistent electric field and magnetospheric configuration. As inductive electric field is closely associated with time-varying magnetospheric configuration, its effects on the transport and energization of particles cannot be neglected (e.g., Ilie et al., 2017; Zaharia et al., 2008). On the other hand, the convective electric field involves complex physical and chemical processes in the ionosphere-thermosphere system. To represent a more accurate magnetosphere-ionosphere system, one of the next

steps is to extend the inner magnetosphere modeling capability to include the ionosphere-thermosphere system. This requirement poses significant challenges for the modeling community and will require different techniques and collaboration across The Coupling, Energetics, and Dynamics of Atmospheric Regions and GEM communities.

The bridge linking the inner magnetosphere with the ionosphere-thermosphere, to a large extent, pertains to wave-particle diffusion processes in the magnetosphere, because the plasma waves drive particle precipitation down to the upper atmosphere. The wave dynamics, such as wave excitation due to anisotropic plasma distributions, wave propagation, and wave diffusion processes, is essential to the particle acceleration and loss. However, it is challenging to incorporate the microscale wave dynamics self-consistently and efficiently into the macroscale models without simplifications. The first attempt was made by Jordanova et al. (1997, 2001), who were able to simulate the EMIC waves generation and resulting ion precipitation self-consistently with the evolving ring current ion dynamics using quasi-linear theory. These results were further expanded to calculate self-consistently the EMIC wave amplitude based on first principles by Khazanov et al. (2003) and Gamayunov et al. (2009, 2014) and based on hybrid simulations by Bortnik et al. (2011) and Fu et al. (2016). Clearly, more studies of self-consistent modeling of the interactions between various types of waves and plasma are further needed. In addition, it is also challenging to move beyond the extensively applied quasi-linear theory to the nonlinear wave-particle interactions (Albert et al., 2013) because quasi-linear theory fails with large wave amplitude (e.g., Santolik, Kletzing, et al., 2014; Tao et al., 2012) and the nonlinearity is particularly important in wave generation (e.g., Demekhov et al., 2017; Omura et al., 2013). Therefore, more work is needed to investigate how to substitute quasi-linear theory with nonlinear theory for the inner magnetosphere study under unusual conditions.

Another challenge is to advance the plasmasphere modeling. In contrast to the ring current and radiation belt models that have been persistently improved in our community, the modeling of the low-energy plasmasphere regime seems left behind. While advancements have been recently achieved in the 3-D ionosphere-plasmasphere model SAMI3, which is self-consistently coupled to the ring current model RCM (e.g., Huba & Sazykin, 2014; Huba et al., 2017), many existing plasmasphere models incorporated in the inner magnetosphere models are decades old, and still empirical based. Advancing the modeling of plasmasphere as part of the inner magnetosphere and its effects on other collocated populations and wave dynamics is thus needed.

We also need awareness of the complicated inner magnetosphere composition and their impact on the evolution of the ring current, especially oxygen and nitrogen ions. In the past years, oxygen ions have received extensive attention and investigation (Kistler, 2016; Welling et al., 2015; B. Zhang et al., 2016), but nitrogen ions, which behave similarly to oxygen, have not been well explored. It is not yet clear if the nitrogen ions contribute the same impact on the inner magnetosphere dynamics (Ilie & Liemohn, 2016). In addition, it was also suggested that the inner magnetosphere current systems be partitioned into several categories, including banana current, eastward ring current, and partial ring current (e.g., Ganushkina, Liemohn, et al., 2015; Liemohn et al., 2013). As controversies in determining these current densities exist, Liemohn et al. (2016) have elucidated the steps toward the resolution to tackle the issue, which requires dedicated community-wide effort. Furthermore, the magneto-tail currents can also contribute to the inner magnetospheric configuration (e.g., Artemyev et al., 2018; Lühr et al., 2016). It is therefore necessary to determine the individual contributions of these current systems.

Challenges further lie in how to take full advantage of the rapidly growing massive satellite measurements and maturing physics-based models by using effective tools, such as machine learning and data assimilation. While these techniques have been widely and successfully applied in the ionosphere, they are quite limited in the inner magnetosphere field (there are only a few research groups dedicating to this area, e.g., Bortnik et al., 2016; Chu et al., 2017; Godinez et al., 2016; Kellerman et al., 2014; Shprits et al., 2013; Zhelavskaya et al., 2017). How to extend their application to other geospace regions and dynamics to achieve a full 3-D representation of the inner magnetosphere is one challenging objective.

Nanosatellites have recently emerged as an economical and valuable platform for understanding near-Earth space. For example, the Colorado Student Space Weather Experiment successfully demonstrated the feasibility of studying inner magnetosphere at much lower cost (Li et al., 2013). One of the next goals would be to explore the space environment more economically by utilizing such CubeSats spacecraft.

Finally, a long-lasting challenge is associated with model validation. Since different models utilize different settings, parameters, and numerical schemes, and different models may show different levels of capabilities for different storm events, it is challenging to distinguish the key factors that control the model performance in capturing the inner magnetosphere environment. Standardizing assessment metrics will help, such as those put forward for geomagnetic indices by Liemohn et al. (2018). More challenge studies reproducing important geospace features are still needed to further look into the limitation of the models and to help improve them.

4. Summary

The Earth's inner magnetosphere plays a key role in governing global magnetospheric dynamics and its rich internal cross-population interactions, and coupling to other geospace regions has long been a fundamental scientific focus. With tremendous progress being made in the past few years in fulfilling more realistic physical models to better represent and understand the inner magnetosphere, it is noteworthy that challenges still remain toward fully comprehending the plasma and field dynamics and to eventually realize the science-to-operation application purpose. We need continued community-wide collaboration and investigation.

Acknowledgments

Acknowledgment This work is supported by NSFC Grant 41574156. The authors would like to thank the NSF GEM program and committee members for supporting our Focus Group IMCEPI during 2014–2018. The authors also wish to thank George Khazanov and Yihua Zheng for helpful conversations. No data are used in this article.

References

- Agapitov, O. V., Artemyev, A. V., Mourenas, D., Kasahara, Y., & Krasnoselskikh, V. (2014). Inner belt and slot region electron lifetimes and energization rates based on AKEBONO statistics of whistler waves. *Journal of Geophysical Research: Space Physics*, *119*, 2876–2893. <https://doi.org/10.1002/2014JA019886>
- Agapitov, O. V., Mourenas, D., Artemyev, A. V., Mozer, F. S., Hospodarsky, G., Bonnell, J., & Krasnoselskikh, V. (2018). Synthetic empirical chorus wave model from combined Van Allen Probes and Cluster statistics. *Journal of Geophysical Research: Space Physics*, *123*, 297–314. <https://doi.org/10.1002/2017JA024843>
- Albert, J. M. (1999). Analysis of quasi-linear diffusion coefficients. *Journal of Geophysical Research*, *104*(A2), 2429–2441. <https://doi.org/10.1029/1998JA900113>
- Albert, J. M. (2004). Using quasi-linear diffusion to model acceleration and loss from wave-particle interactions. *Space Weather*, *2*, S09S03. <https://doi.org/10.1029/2004SW000069>
- Albert, J. M., Tao, X., & Bortnik, J. (2013). Aspects of nonlinear wave-particle interactions. In D. Summers, I. R. Mann, D. N. Baker, & M. Schulz (Eds.), *Dynamics of the Earth's radiation belts and inner magnetosphere* (pp. 255–264). Washington, DC: American Geophysical Union. <https://doi.org/10.1029/2012GM001324>
- Ali, A. F., Malaspina, D. M., Elkington, S. R., Jaynes, A. N., Chan, A. A., Wygant, J., & Kletzing, C. A. (2016). Electric and magnetic radial diffusion coefficients using the Van Allen probes data. *Journal of Geophysical Research: Space Physics*, *121*, 9586–9607. <https://doi.org/10.1002/2016JA023002>
- Artemyev, A., Agapitov, O., Mourenas, D., Krasnoselskikh, V., Shastun, V., & Mozer, F. (2016). Oblique whistler mode waves in the Earth's inner magnetosphere: Energy distribution, origins, and role in radiation belt dynamics. *Space Science Reviews*, *200*, 1–95. <https://doi.org/10.1007/s11214-016-0252-5>
- Artemyev, A. V., Zhang, X.-J., Angelopoulos, V., Runov, A., Spence, H. E., & Larsen, B. A. (2018). Plasma anisotropies and currents in the near-Earth plasma sheet and inner magnetosphere. *Journal of Geophysical Research: Space Physics*, *123*, 5625–5639. <https://doi.org/10.1029/2018JA025232>
- Aryan, H., Sibeck, D., Balikhin, M., Agapitov, O., & Kletzing, C. (2016). Observation of chorus waves by the Van Allen Probes: Dependence on solar wind parameters and scale size. *Journal of Geophysical Research: Space Physics*, *121*, 7608–7621. <https://doi.org/10.1002/2016JA022775>
- Aryan, H., Sibeck, D. G., Kang, S.-B., Balikhin, M. A., Fok, M.-C., Agapitov, O., et al. (2017). CIMI simulations with newly developed multiparameter chorus and plasmaspheric hiss wave models. *Journal of Geophysical Research: Space Physics*, *122*, 9344–9357. <https://doi.org/10.1002/2017JA024159>
- Aryan, H., Yearby, K., Balikhin, M., Agapitov, O., Krasnoselskikh, V., & Boynton, R. (2014). Statistical study of chorus wave distributions in the inner magnetosphere using AE and solar wind parameters. *Journal of Geophysical Research: Space Physics*, *119*, 6131–6144. <https://doi.org/10.1002/2014JA019939>
- Blum, L. W., Agapitov, O., Bonnell, J. W., Kletzing, C., & Wygant, J. (2016). EMIC wave spatial and coherence scales as determined from multipoint Van Allen Probe measurements. *Geophysical Research Letters*, *43*, 4799–4807. <https://doi.org/10.1002/2016GL068799>
- Blum, L. W., Bonnell, J. W., Agapitov, O., Paulson, K., & Kletzing, C. (2017). EMIC wave scale size in the inner magnetosphere: Observations from the dual Van Allen Probes. *Geophysical Research Letters*, *44*, 1227–1233. <https://doi.org/10.1002/2016GL072316>
- Blum, L. W., Halford, A., Millan, R., Bonnell, J. W., Goldstein, J., Usanova, M., et al. (2015). Observations of coincident EMIC wave activity and duskside energetic electron precipitation on 18–19 January 2013. *Geophysical Research Letters*, *42*, 5727–5735. <https://doi.org/10.1002/2015GL065245>
- Bortnik, J., Li, W., Thorne, R. M., & Angelopoulos, V. (2016). A unified approach to inner magnetospheric state prediction. *Journal of Geophysical Research: Space Physics*, *121*, 2423–2430. <https://doi.org/10.1002/2015JA021733>
- Bortnik, J., Omid, N., Chen, L., Thorne, R. M., & Horne, R. B. (2011). Saturation characteristics of electromagnetic ion cyclotron waves. *Journal of Geophysical Research*, *116*, A09219. <https://doi.org/10.1029/2011JA016638>
- Brito, T. V., Woodroffe, J., Jordanova, V. K., Henderson, M., & Birn, J. (2017). Particle tracing modeling of ion fluxes at geosynchronous orbit. *Journal of Atmospheric and Solar-Terrestrial Physics*, *177*, 131–140. <https://doi.org/10.1016/j.jastp.2017.10.008>
- Califf, S., Li, X., Blum, L., Jaynes, A., Schiller, Q., Zhao, H., et al. (2014). THEMIS measurements of quasi-static electric fields in the inner magnetosphere. *Journal of Geophysical Research: Space Physics*, *119*, 9939–9951. <https://doi.org/10.1002/2014JA020360>

- Cao, X., Ni, B., Liang, J., Xiang, Z., Wang, Q., Shi, R., et al. (2016). Resonant scattering of central plasma sheet protons by multiband EMIC waves and resultant proton loss timescales. *Journal of Geophysical Research: Space Physics*, *121*, 1219–1232. <https://doi.org/10.1002/2015JA021933>
- Chaston, C. C., Bonnell, J. W., Halford, A. J., Reeves, G. D., Baker, D. N., Kletzing, C. A., & Wygant, J. R. (2018). Pitch angle scattering and loss of radiation belt electrons in broadband electromagnetic waves. *Geophysical Research Letters*, *45*, 9344–9352. <https://doi.org/10.1029/2018GL079527>
- Chen, M. W., Lemon, C. L., Guild, T. B., Keesee, A. M., Lui, A., Goldstein, J., et al. (2015). Effects of modeled ionospheric conductance and electron loss on self-consistent ring current simulations during the 5–7 April 2010 storm. *Journal of Geophysical Research: Space Physics*, *120*, 5355–5376. <https://doi.org/10.1002/2015JA021285>
- Chen, M. W., Lemon, C. L., Orlova, K., Shprits, Y., Hecht, J., & Walterscheid, R. L. (2015). Comparison of simulated and observed trapped and precipitating electron fluxes during a magnetic storm. *Geophysical Research Letters*, *42*, 8302–8311. <https://doi.org/10.1002/2015GL065737>
- Chu, X., Bortnik, J., Li, W., Ma, Q., Denton, R., Yue, C., et al. (2017). A neural network model of three-dimensional dynamic electron density in the inner magnetosphere. *Journal of Geophysical Research: Space Physics*, *122*, 9183–9197. <https://doi.org/10.1002/2017JA024464>
- Claudepierre, S. G., Chen, M. W., Roeder, J. L., & Fennell, J. F. (2016). An empirical model of ion plasma in the inner magnetosphere derived from CRRES/MICS measurements. *Journal of Geophysical Research: Space Physics*, *121*, 11,780–11,797. <https://doi.org/10.1002/2016JA023468>
- Cramer, W. D., Raeder, J., Toffoletto, F. R., Gilson, M., & Hu, B. (2017). Plasma sheet injections into the inner magnetosphere: Two-way coupled OpenGGCM-RCM model results. *Journal of Geophysical Research: Space Physics*, *122*, 5077–5091. <https://doi.org/10.1002/2017JA024104>
- De Zeeuw, D. L., Sazykin, S., Wolf, R. A., Gombosi, T. I., Ridley, A. J., & Toth, G. (2004). Coupling of a global MHD code and an inner magnetospheric model: Initial results. *Journal of Geophysical Research*, *109*(A12), A12219. <https://doi.org/10.1029/2003JA010366>
- Demekhov, A. G., Taubenschuss, U., & Santolik, O. (2017). Simulation of VLF chorus emissions in the magnetosphere and comparison with THEMIS spacecraft data. *Journal of Geophysical Research: Space Physics*, *122*, 166–184. <https://doi.org/10.1002/2016JA023057>
- Denton, M. H., Reeves, G. D., Larsen, B. A., Friedel, R. H. W., Thomsen, M. F., Fernandes, P. A., et al. (2017). On the origin of low-energy electrons in the inner magnetosphere: Fluxes and pitch-angle distributions. *Journal of Geophysical Research: Space Physics*, *122*, 1789–1802. <https://doi.org/10.1002/2016JA023648>
- Dunlop, M. W., Yang, J.-Y., Yang, Y.-Y., Xiong, C., Lühr, H., Bogdanova, Y. V., et al. (2015). Simultaneous field-aligned currents at Swarm and Cluster satellites. *Geophysical Research Letters*, *42*, 3683–3691. <https://doi.org/10.1002/2015GL063738>
- Ebihara, Y., Fok, M.-C., Wolf, R. A., Immel, T. J., & Moore, T. E. (2004). Influence of ionospheric conductivity on the ring current. *Journal of Geophysical Research*, *109*(A8), A08205. <https://doi.org/10.1029/2003JA010351>
- Ebihara, Y., & Tanaka, T. (2013). Fundamental properties of substorm time energetic electrons in the inner magnetosphere. *Journal of Geophysical Research: Space Physics*, *118*, 1589–1603. <https://doi.org/10.1002/jgra.50115>
- Fernandes, P. A., Larsen, B. A., Thomsen, M. F., Skoug, R. M., Reeves, G. D., Denton, M. H., et al. (2017). The plasma environment inside geostationary orbit: A Van Allen Probes HOPE survey. *Journal of Geophysical Research: Space Physics*, *122*, 9207–9227. <https://doi.org/10.1002/2017JA024160>
- Fok, M.-C., Buzulukova, N. Y., Chen, S.-H., Glocer, A., Nagai, T., Valek, P., & Perez, J. D. (2014). The comprehensive inner magnetosphere-ionosphere model. *Journal of Geophysical Research: Space Physics*, *119*, 7522–7540. <https://doi.org/10.1002/2014JA020239>
- Fok, M.-C., Kozyra, J. U., Nagy, A. F., Rasmussen, C. E., & Khazanov, G. V. (1993). A decay model of equatorial ring current and the associated aeronomical consequences. *Journal of Geophysical Research*, *98*(A11), 19,381–19,393. <https://doi.org/10.1029/93JA01848>
- Fok, M.-C., Wolf, R. A., Spiro, R. W., & Moore, T. E. (2001). Comprehensive computational model of Earth's ring current. *Journal of Geophysical Research*, *106*(A5), 8417–8424. <https://doi.org/10.1029/2000JA000235>
- Fu, H. S., Cao, J. B., Zhima, Z., Khotyaintsev, Y. V., Angelopoulos, V., Santolik, O., et al. (2014). First observation of rising-tone magnetosonic waves. *Geophysical Research Letters*, *41*, 7419–7426. <https://doi.org/10.1002/2014GL061867>
- Fu, X., Cowee, M. M., Jordanova, V. K., Gary, S. P., Reeves, G. D., & Winske, D. (2016). Predicting electromagnetic ion cyclotron wave amplitude from unstable ring current plasma conditions. *Journal of Geophysical Research: Space Physics*, *121*, 10,954–10,965. <https://doi.org/10.1002/2016JA023303>
- Fu, X., Gary, S. P., Reeves, G. D., Winske, D., & Woodroffe, J. R. (2017). Generation of highly oblique lower band chorus via nonlinear three-wave resonance. *Geophysical Research Letters*, *44*, 9532–9538. <https://doi.org/10.1002/2017GL074411>
- Gallardo-Lacourt, B., Liang, J., Nishimura, Y., & Donovan, E. (2018). On the origin of STEVE: Particle precipitation or ionospheric sky-glow? *Geophysical Research Letters*, *45*, 7968–7973. <https://doi.org/10.1029/2018GL078509>
- Gamayunov, K. V., Engebretson, M. J., Zhang, M., & Rassoul, H. K. (2014). Model of electromagnetic ion cyclotron waves in the inner magnetosphere. *Journal of Geophysical Research: Space Physics*, *119*, 7541–7565. <https://doi.org/10.1002/2014JA020032>
- Gamayunov, K. V., Khazanov, G. V., Liemohn, M. W., Fok, M.-C., & Ridley, A. J. (2009). Self-consistent model of magnetospheric electric field, ring current, plasmasphere, and electromagnetic ion cyclotron waves: Initial results. *Journal of Geophysical Research*, *114*(A3), A03221. <https://doi.org/10.1029/2008JA013597>
- Ganushkina, N. Y., Amariutei, O. A., Welling, D., & Heynderickx, D. (2015). Nowcast model for low-energy electrons in the inner magnetosphere. *Space Weather*, *13*, 16–34. <https://doi.org/10.1002/2014SW001098>
- Ganushkina, N. Y., Liemohn, M. W., Amariutei, O. A., & Pitchford, D. (2014). Low-energy electrons (5–50 keV) in the inner magnetosphere. *Journal of Geophysical Research: Space Physics*, *119*, 246–259. <https://doi.org/10.1002/2013JA019304>
- Ganushkina, N. Y., Liemohn, M. W., Dubyagin, S., Daglis, I. A., Dandouras, I., de Zeeuw, D. L., et al. (2015). Defining and resolving current systems in geospace. *Annales de Geophysique*, *33*, 1369–1402. <https://doi.org/10.5194/angeo-33-1369-2015>
- Glauert, S. A., Horne, R. B., & Meredith, N. P. (2014). Three-dimensional electron radiation belt simulations using the BAS Radiation Belt Model with new diffusion models for chorus, plasmaspheric hiss, and lightning-generated whistlers. *Journal of Geophysical Research: Space Physics*, *119*, 268–289. <https://doi.org/10.1002/2013JA019281>
- Glocer, A., Fok, M., Meng, X., Toth, G., Buzulukova, N., Chen, S., & Lin, K. (2013). CRCM + BATS-R-US two-way coupling. *Journal of Geophysical Research: Space Physics*, *118*, 1635–1650. <https://doi.org/10.1002/jgra.50221>
- Glocer, A., Rastätter, L., Kuznetsova, M., Pulkkinen, A., Singer, H. J., Balch, C., et al. (2016). Community-wide validation of geospace model local K index predictions to support model transition to operations. *Space Weather*, *14*, 469–480. <https://doi.org/10.1002/2016SW001387>

- Glocer, A., Toth, G., Fok, M., Gombosi, T., & Liemohn, M. (2009). Integration of the radiation belt environment model into the space weather modeling framework. *Journal of Atmospheric and Solar - Terrestrial Physics*, *71*(16), 1653–1663. <https://doi.org/10.1016/j.jastp.2009.01.003>
- Godinez, H. C., Yu, Y., Lawrence, E., Henderson, M. G., Larsen, B., & Jordanova, V. K. (2016). Ring current pressure estimation with RAM-SCB using data assimilation and Van Allen Probe flux data. *Geophysical Research Letters*, *43*, 11,948–11,956. <https://doi.org/10.1002/2016GL071646>
- He, F., Zhang, X.-X., Lin, R.-L., Fok, M.-C., Katus, R. M., Liemohn, M. W., et al. (2017). A new solar wind-driven global dynamic plasmamodel: 2. Model and validation. *Journal of Geophysical Research: Space Physics*, *122*, 7172–7187. <https://doi.org/10.1002/2017JA023913>
- Horne, R. B., Kersten, T., Glauert, S. A., Meredith, N. P., Boscher, D., Sicard-Piet, A., & Li, W. (2013). A new diffusion matrix for whistler mode chorus waves. *Journal of Geophysical Research: Space Physics*, *118*, 6302–6318. <https://doi.org/10.1002/jgra.50594>
- Huba, J. D., & Sazykin, S. (2014). Storm time ionosphere and plasmasphere structuring: SAMI3-RCM simulation of the 31 March 2001 geomagnetic storm. *Geophysical Research Letters*, *41*, 8208–8214. <https://doi.org/10.1002/2014GL062110>
- Huba, J. D., Sazykin, S., & Coster, A. (2017). SAMI3-RCM simulation of the 17 March 2015 geomagnetic storm. *Journal of Geophysical Research: Space Physics*, *122*, 1246–1257. <https://doi.org/10.1002/2016JA023341>
- Ilie, R., Daldorff, L. K. S., Liemohn, M. W., Toth, G., & Chan, A. A. (2017). Calculating the inductive electric field in the terrestrial magnetosphere. *Journal of Geophysical Research: Space Physics*, *122*, 5391–5403. <https://doi.org/10.1002/2017JA023877>
- Ilie, R., & Liemohn, M. W. (2016). The outflow of ionospheric nitrogen ions: A possible tracer for the altitude-dependent transport and energization processes of ionospheric plasma. *Journal of Geophysical Research: Space Physics*, *121*, 9250–9255. <https://doi.org/10.1002/2015JA022162>
- Ilie, R., Liemohn, M. W., Toth, G., & Skoug, R. M. (2012). Kinetic model of the inner magnetosphere with arbitrary magnetic field. *Journal of Geophysical Research*, *117*(A4), A04208. <https://doi.org/10.1029/2011JA017189>
- Ji, Y., & Shen, C. (2014). The loss rates of O⁺ in the inner magnetosphere caused by both magnetic field line curvature scattering and charge exchange reactions. *Physics of Plasmas*, *21*, 032903. <https://doi.org/10.1063/1.4868863>
- Jordanova, V. K., Delzanno, G. L., Henderson, M. G., Godinez, H. C., Jeffery, C. A., Lawrence, E. C., et al. (2018). Specification of the near-Earth space environment with SHIELDS. *Journal of Atmospheric and Solar-Terrestrial Physics*, *177*, 148–159. <https://doi.org/10.1016/j.jastp.2017.11.006>
- Jordanova, V. K., Farrugia, C. J., Thorne, R. M., Khazanov, G. V., Reeves, G. D., & Thomsen, M. F. (2001). Modeling ring current proton precipitation by electromagnetic ion cyclotron waves during the May 14–16, 1997, storm. *Journal of Geophysical Research*, *106*(A1), 7–22. <https://doi.org/10.1029/2000JA002008>
- Jordanova, V. K., Kozyra, J. U., Khazanov, G. V., Nagy, A. F., Rasmussen, C. E., & Fok, M.-C. (1994). A bounce-averaged kinetic model of the ring current ion population. *Geophysical Research Letters*, *21*(25), 2785–2788. <https://doi.org/10.1029/94GL02695>
- Jordanova, V. K., Kozyra, J. U., Nagy, A. F., & Khazanov, G. V. (1997). Kinetic model of the ring current-atmosphere interactions. *Journal of Geophysical Research*, *102*(A7), 14,279–14,291. <https://doi.org/10.1029/96JA03699>
- Jordanova, V. K., Tu, W., Chen, Y., Morley, S. K., Panaitescu, A. D., Reeves, G. D., & Kletzing, C. A. (2016). RAM-SCB simulations of electron transport and plasma wave scattering during the October 2012 “double-dip” storm. *Journal of Geophysical Research: Space Physics*, *121*, 8712–8727. <https://doi.org/10.1002/2016JA022470>
- Jordanova, V. K., Yu, Y., Niehof, J. T., Skoug, R. M., Reeves, G. D., Kletzing, C. A., et al. (2014). Simulations of inner magnetosphere dynamics with an expanded RAM-SCB model and comparisons with Van Allen Probes observations. *Geophysical Research Letters*, *41*, 2687–2694. <https://doi.org/10.1002/2014GL059533>
- Jordanova, V. K., Zaharia, S., & Welling, D. T. (2010). Comparative study of ring current development using empirical, dipolar, and self-consistent magnetic field simulations. *Journal of Geophysical Research*, *115*, A00J11. <https://doi.org/10.1029/2010JA015671>
- Kang, S.-B., Fok, M.-C., Glocer, A., Min, K.-W., Choi, C.-R., Choi, E., & Hwang, J. (2016). Simulation of a rapid dropout event for highly relativistic electrons with the RBE model. *Journal of Geophysical Research: Space Physics*, *121*, 4092–4102. <https://doi.org/10.1002/2015JA021966>
- Kang, S.-B., Min, K.-W., Fok, M.-C., Hwang, J., & Choi, C.-R. (2015). Estimation of pitch angle diffusion rates and precipitation timescales of electrons due to EMIC waves in a realistic field model. *Journal of Geophysical Research: Space Physics*, *120*, 8529–8546. <https://doi.org/10.1002/2014JA020644>
- Kasahara, S., Miyoshi, Y., Yokota, S., Mitani, T., Kasahara, Y., Matsuda, S., et al. (2018). Pulsating aurora from electron scattering by chorus waves. *Nature*, *554*, 337–340. <https://doi.org/10.1038/nature25505>
- Kellerman, A. C., Shprits, Y. Y., Kondrashov, D., Subbotin, D., Makarevich, R. A., Donovan, E., & Nagai, T. (2014). Three-dimensional data assimilation and reanalysis of radiation belt electrons: Observations of a four-zone structure using five spacecraft and the VERB code. *Journal of Geophysical Research: Space Physics*, *119*, 8764–8783. <https://doi.org/10.1002/2014JA020171>
- Kersten, T., Horne, R. B., Glauert, S. A., Meredith, N. P., Fraser, B. J., & Grew, R. S. (2014). Electron losses from the radiation belts caused by EMIC waves. *Journal of Geophysical Research: Space Physics*, *119*, 8820–8837. <https://doi.org/10.1002/2014JA020366>
- Khazanov, G. V., Gamayunov, K. V., & Jordanova, V. K. (2003). Self-consistent model of magnetospheric ring current and electromagnetic ion cyclotron waves: The 2–7 May 1998 storm. *Journal of Geophysical Research*, *108*(A12). <https://doi.org/10.1029/2003JA009856>
- Khazanov, G. V., Glocer, A., & Himwich, E. W. (2014). Magnetosphere-ionosphere energy interchange in the electron diffuse aurora. *Journal of Geophysical Research: Space Physics*, *119*, 171–184. <https://doi.org/10.1002/2013JA019325>
- Khazanov, G. V., Robinson, R. M., Zesta, E., Sibeck, D. G., Chu, M., & Grubbs, G. A. (2018). Impact of precipitating electrons and magnetosphere-ionosphere coupling processes on ionospheric conductance. *Space Weather*, *16*, 829–837. <https://doi.org/10.1029/2018SW001837>
- Khazanov, G. V., Sibeck, D. G., & Zesta, E. (2017). Major pathways to electron distribution function formation in regions of diffuse aurora. *Journal of Geophysical Research: Space Physics*, *122*, 4251–4265. <https://doi.org/10.1002/2017JA023956>
- Khazanov, G. V., Tripathi, A. K., Singhal, R. P., Himwich, E. W., Glocer, A., & Sibeck, D. G. (2015). Superthermal electron magnetosphere-ionosphere coupling in the diffuse aurora in the presence of ECH waves. *Journal of Geophysical Research: Space Physics*, *120*, 445–459. <https://doi.org/10.1002/2014JA020641>
- Kim, K.-H., Shiokawa, K., Mann, I. R., Park, J.-S., Kwon, H.-J., Hyun, K., et al. (2016). Longitudinal frequency variation of long-lasting EMIC Pc1-Pc2 waves localized in the inner magnetosphere. *Geophysical Research Letters*, *43*, 1039–1046. <https://doi.org/10.1002/2015GL067536>
- Kistler, L. M. (2016). The Impact of O⁺ on Magnetotail Dynamics. In C. R. Chappell, R. W. Schunk, P. M. Banks, J. L. Burch, & R. M. Thorne (Eds.), *Magnetosphere-ionosphere coupling in the solar system*, *Geophysical Monograph Series* (Chap. 6). Washington, DC: American Geophysical Union. <https://doi.org/10.1002/9781119066880.ch6>

- Kistler, L. M., & Mouikis, C. G. (2016). The inner magnetosphere ion composition and local time distribution over a solar cycle. *Journal of Geophysical Research: Space Physics*, *121*, 2009–2032. <https://doi.org/10.1002/2015JA021883>
- Kistler, L. M., Mouikis, C. G., Spence, H. E., Menz, A. M., Skoug, R. M., Funsten, H. O., et al. (2016). The source of O⁺ in the storm time ring current. *Journal of Geophysical Research: Space Physics*, *121*, 5333–5349. <https://doi.org/10.1002/2015JA022204>
- Krall, J., Huba, J. D., & Sazykin, S. (2017). Erosion of the plasmasphere during a storm. *Journal of Geophysical Research: Space Physics*, *122*, 9320–9328. <https://doi.org/10.1002/2017JA024450>
- Lemon, C., Wolf, R. A., Hill, T. W., Sazykin, S., Spiro, R. W., Toffoletto, F. R., et al. (2004). Magnetic storm ring current injection modeled with the Rice Convection Model and a self-consistent magnetic field. *Geophysical Research Letters*, *31*, L21801. <https://doi.org/10.1029/2004GL020914>
- Li, J., Ni, B., Ma, Q., Xie, L., Pu, Z., Fu, S., et al. (2016). Formation of energetic electron butterfly distributions by magnetosonic waves via Landau resonance. *Geophysical Research Letters*, *43*, 3009–3016. <https://doi.org/10.1002/2016GL067853>
- Li, L., Zhou, X.-Z., Zong, Q.-G., Rankin, R., Zou, H., Liu, Y., et al. (2017). Charged particle behavior in localized ultralow frequency waves: Theory and observations. *Geophysical Research Letters*, *44*, 5900–5908. <https://doi.org/10.1002/2017GL073392>
- Li, L. Y., Yu, J., Cao, J. B., Wang, Z. Q., Yu, Y. Q., Reeves, G. D., & Li, X. (2016). Effects of ULF waves on local and global energetic particles: Particle energy and species dependences. *Journal of Geophysical Research: Space Physics*, *121*, 11,007–11,020. <https://doi.org/10.1002/2016JA023149>
- Li, L. Y., Yu, J., Cao, J. B., Yang, J. Y., Li, X., Baker, D. N., et al. (2017). Roles of whistler mode waves and magnetosonic waves in changing the outer radiation belt and the slot region. *Journal of Geophysical Research: Space Physics*, *122*, 5431–5448. <https://doi.org/10.1002/2016JA023634>
- Li, W., Ma, Q., Thorne, R. M., Bortnik, J., Kletzing, C. A., Kurth, W. S., et al. (2015). Statistical properties of plasmaspheric hiss derived from Van Allen Probes data and their effects on radiation belt electron dynamics. *Journal of Geophysical Research: Space Physics*, *120*, 3393–3405. <https://doi.org/10.1002/2015JA021048>
- Li, W., Ni, B., Thorne, R. M., Bortnik, J., Nishimura, Y., Green, J. C., et al. (2014). Quantifying hiss-driven energetic electron precipitation: A detailed conjunction event analysis. *Geophysical Research Letters*, *41*, 1085–1092. <https://doi.org/10.1002/2013GL059132>
- Li, W., Santolik, O., Bortnik, J., Thorne, R. M., Kletzing, C. A., Kurth, W. S., & Hospodarsky, G. B. (2016). New chorus wave properties near the equator from Van Allen Probes wave observations. *Geophysical Research Letters*, *43*, 4725–4735. <https://doi.org/10.1002/2016GL068780>
- Li, X., Schiller, Q., Blum, L., Califf, S., Zhao, H., Tu, W., et al. (2013). First results from CSSWE CubeSat: Characteristics of relativistic electrons in the near-Earth environment during the October 2012 magnetic storms. *Journal of Geophysical Research: Space Physics*, *118*, 6489–6499. <https://doi.org/10.1002/2013JA019342>
- Liemohn, M. W. (2006). Introduction to special section on “Results of the National Science Foundation Geospace Environment Modeling Inner Magnetosphere/Storms Assessment Challenge”. *Journal of Geophysical Research*, *111*, A11S01. <https://doi.org/10.1029/2006JA011970>
- Liemohn, M. W., Ganushkina, N. Y., Ilie, R., & Welling, D. T. (2016). Challenges associated with near-Earth nightside current. *Journal of Geophysical Research: Space Physics*, *121*, 6763–6768. <https://doi.org/10.1002/2016JA022948>
- Liemohn, M. W., Ganushkina, N. Y., Katus, R. M., De Zeeuw, D. L., & Welling, D. T. (2013). The magnetospheric banana current. *Journal of Geophysical Research: Space Physics*, *118*, 1009–1021. <https://doi.org/10.1002/jgra.50153>
- Liemohn, M. W., Kozyra, J. U., Thomsen, M. F., Roeder, J. L., Lu, G., Borovsky, J. E., & Cayton, T. E. (2001). The dominant role of the asymmetric ring current in producing the stormtime Dst*. *Journal of Geophysical Research*, *106*(A6), 10,883–10,904. <https://doi.org/10.1029/2000JA000326>
- Liemohn, M. W., McCollough, J. P., Jordanova, V. K., Ngwira, C. M., Morley, S. K., Cid, C., et al. (2018). Model evaluation guidelines for geomagnetic index predictions. *Space Weather*, *16*, 2079–2102. <https://doi.org/10.1029/2018SW002067>
- Liemohn, M. W., Ridley, A. J., Brandt, P. C., Gallagher, D. L., Kozyra, J. U., Mitchell, D. G., et al. (2005). Parametric analysis of nightside conductance effects on inner magnetospheric dynamics for the 17 April 2002 storm. *Journal of Geophysical Research*, *110*, A12S22. <https://doi.org/10.1029/2005JA011109>
- Liu, W., Tu, W., Li, X., Sarris, T., Khotyaintsev, Y., Fu, H., et al. (2016). On the calculation of electric diffusion coefficient of radiation belt electrons with in situ electric field measurements by THEMIS. *Geophysical Research Letters*, *43*, 1023–1030. <https://doi.org/10.1002/2015GL067398>
- Liu, X., Liu, W., Cao, J. B., Fu, H. S., Yu, J., & Li, X. (2015). Dynamic plasmopause model based on THEMIS measurements. *Journal of Geophysical Research: Space Physics*, *120*, 10,543–10,556. <https://doi.org/10.1002/2015JA021801>
- Lühr, H., Xiong, C., Olsen, N., & Le, G. (2016). Near-earth magnetic field effects of large-scale magnetospheric currents. *Space Science Reviews*, *206*, 521–545. <https://doi.org/10.1007/s11214-016-0267-y>
- Ma, Q., Li, W., Bortnik, J., Thorne, R. M., Chu, X., Ozeke, L. G., et al. (2018). Quantitative evaluation of radial diffusion and local acceleration processes during GEM challenge events. *Journal of Geophysical Research: Space Physics*, *123*, 1938–1952. <https://doi.org/10.1002/2017JA025114>
- Ma, Q., Li, W., Thorne, R. M., Bortnik, J., Kletzing, C. A., Kurth, W. S., & Hospodarsky, G. B. (2016). Electron scattering by magnetosonic waves in the inner magnetosphere. *Journal of Geophysical Research: Space Physics*, *121*, 274–285. <https://doi.org/10.1002/2015JA021992>
- Ma, Q., Mourenas, D., Artemyev, A., Li, W., Thorne, R. M., & Bortnik, J. (2016). Strong enhancement of 10–100 keV electron fluxes by combined effects of chorus waves and time domain structures. *Geophysical Research Letters*, *43*, 4683–4690. <https://doi.org/10.1002/2016GL069125>
- Ma, Q., Mourenas, D., Li, W., Artemyev, A., & Thorne, R. M. (2017). VLF waves from ground-based transmitters observed by the Van Allen Probes: Statistical model and effects on plasmaspheric electrons. *Geophysical Research Letters*, *44*, 6483–6491. <https://doi.org/10.1002/2017GL073885>
- Malaspina, D. M., Claudepierre, S. G., Takahashi, K., Jaynes, A. N., Elkington, S. R., Ergun, R. E., et al. (2015). Kinetic Alfvén waves and particle response associated with a shock-induced, global ULF perturbation of the terrestrial magnetosphere. *Geophysical Research Letters*, *42*, 9203–9212. <https://doi.org/10.1002/2015GL065935>
- Malaspina, D. M., Jaynes, A. N., Boul, C., Bortnik, J., Thaller, S. A., Ergun, R. E., et al. (2016). The distribution of plasmaspheric hiss wave power with respect to plasmopause location. *Geophysical Research Letters*, *43*, 7878–7886. <https://doi.org/10.1002/2016GL069982>
- Malaspina, D. M., Jaynes, A. N., Hospodarsky, G., Bortnik, J., Ergun, R. E., & Wygant, J. (2017). Statistical properties of low-frequency plasmaspheric hiss. *Journal of Geophysical Research: Space Physics*, *122*, 8340–8352. <https://doi.org/10.1002/2017JA024328>

- Meredith, N. P., Horne, R. B., Kersten, T., Fraser, B. J., & Grew, R. S. (2014). Global morphology and spectral properties of EMIC waves derived from CRRES observations. *Journal of Geophysical Research: Space Physics*, *119*, 5328–5342. <https://doi.org/10.1002/2014JA020064>
- Motoba, T., Ohtani, S., Anderson, B. J., Korth, H., Mitchell, D., Lanzerotti, L. J., et al. (2015). On the formation and origin of substorm growth phase/onset auroral arcs inferred from conjugate space-ground observations. *Journal of Geophysical Research: Space Physics*, *120*, 8707–8722. <https://doi.org/10.1002/2015JA021676>
- Mourenas, D., Artemyev, A. V., Agapitov, O. V., & Krasnoselskikh, V. (2014). Consequences of geomagnetic activity on energization and loss of radiation belt electrons by oblique chorus waves. *Journal of Geophysical Research: Space Physics*, *119*, 2775–2796. <https://doi.org/10.1002/2013JA019674>
- Mozer, F. S., Agapitov, O. V., Artemyev, A., Drake, J. F., Krasnoselskikh, V., Lejosne, S., & Vasko, I. (2015). Time domain structures: What and where they are, what they do, and how they are made. *Geophysical Research Letters*, *42*, 3627–3638. <https://doi.org/10.1002/2015GL063946>
- Murphy, K. R., Mann, I. R., & Sibeck, D. G. (2015). On the dependence of stormtime ULF wave power on magnetopause location: Impacts for ULF wave radial diffusion. *Geophysical Research Letters*, *42*, 9676–9684. <https://doi.org/10.1002/2015GL066592>
- Nemec, F., Hospodarsky, G., Pickett, J. S., Santolik, O., Kurth, W. S., & Kletzing, C. (2016). Conjugate observations of quasiperiodic emissions by the Cluster, Van Allen Probes, and THEMIS spacecraft. *Journal of Geophysical Research: Space Physics*, *121*, 7647–7663. <https://doi.org/10.1002/2016JA022774>
- Ni, B., Cao, X., Zou, Z., Zhou, C., Gu, X., Bortnik, J., et al. (2015). Resonant scattering of outer zone relativistic electrons by multiband EMIC waves and resultant electron loss time scales. *Journal of Geophysical Research: Space Physics*, *120*, 7357–7373. <https://doi.org/10.1002/2015JA021466>
- Ni, B., Gu, X., Fu, S., Xiang, Z., & Lou, Y. (2017). A statistical survey of electrostatic electron cyclotron harmonic waves based on THEMIS FFF wave data. *Journal of Geophysical Research: Space Physics*, *122*, 3342–3353. <https://doi.org/10.1002/2016JA023433>
- Ni, B., Thorne, R. M., Zhang, X., Bortnik, J., Pu, Z., Xie, L., et al. (2016). Origins of the Earth's diffuse auroral precipitation. *Space Science Reviews*, *200*, 205–259. <https://doi.org/10.1007/s11214-016-0234-7>
- Nishimura, Y., Bortnik, J., Li, W., Lyons, L. R., Donovan, E. F., Angelopoulos, V., & Mende, S. B. (2014). Evolution of nightside subauroral proton aurora caused by transient plasma sheet flows. *Journal of Geophysical Research: Space Physics*, *119*, 5295–5304. <https://doi.org/10.1002/2014JA020029>
- Nosé, M. (2016). Long-term variations in the plasma sheet ion composition and substorm occurrence over 23 years. *Geoscience Letters*, *3*, 1–8. <https://doi.org/10.1186/s40562-015-0033-0>
- Omura, Y., Nunn, D., & Summers, D. (2013). Generation processes of whistler mode chorus emissions: Current status of nonlinear wave growth theory. In D. Summers, I. R. Mann, D. N. Baker, & M. Schulz (Eds.), *Dynamics of the Earth's radiation belts and inner magnetosphere* (Vol. 199, pp. 243–254). Washington, DC: American Geophysical Union. <https://doi.org/10.1029/2012GM001347>
- Orlova, K., & Shprits, Y. (2015). Model of lifetimes of the outer radiation belt electrons in a realistic magnetic field using realistic chorus wave parameters, 6199–6206. <https://doi.org/10.1002/2015JA021354>
- Orlova, K., Spasojevic, M., & Shprits, Y. (2014). Activity-dependent global model of electron loss inside the plasmasphere. *Geophysical Research Letters*, *41*, 6413–6419. <https://doi.org/10.1002/2014GL061184>
- Pembroke, A., Toffoletto, F., Sazykin, S., Wiltberger, M., Lyon, J., Merkin, V., & Schmitt, P. (2012). Initial results from a dynamic coupled magnetosphere-ionosphere-ring current model. *Journal of Geophysical Research*, *117*, A02211. <https://doi.org/10.1029/2011JA016979>
- Perlongo, N. J., Ridley, A. J., Liemohn, M. W., & Katus, R. M. (2017). The effect of ring current electron scattering rates on magnetosphere-ionosphere coupling. *Journal of Geophysical Research: Space Physics*, *122*, 4168–4189. <https://doi.org/10.1002/2016JA023679>
- Pulkkinen, A., Rastätter, L., Kuznetsova, M., Singer, H., Balch, C., Weimer, D., et al. (2013). Community-wide validation of geospace model ground magnetic field perturbation predictions to support model transition to operations. *Space Weather*, *11*, 369–385. <https://doi.org/10.1002/swe.20056>
- Raeder, J., Cramer, W. D., Jensen, J., Fuller-Rowell, T., Maruyama, N., Toffoletto, F., & Vo, H. (2016). Sub-auroral polarization streams: A complex interaction between the magnetosphere, ionosphere, and thermosphere. *Journal of Physics: Conference Series*, *767*, 012021. <https://doi.org/10.1088/1742-6596/767/1/012021>
- Rastätter, L., Kuznetsova, M. M., Gloer, A., Welling, D., Meng, X., Raeder, J., et al. (2013). Geospace environment modeling 2008–2009 challenge: Dst index. *Space Weather*, *11*, 187–205. <https://doi.org/10.1002/swe.20036>
- Rastätter, L., Tóth, G., Kuznetsova, M. M., & Pulkkinen, A. A. (2014). CalcDeltaB: An efficient postprocessing tool to calculate ground-level magnetic perturbations from global magnetosphere simulations. *Space Weather*, *12*, 553–565. <https://doi.org/10.1002/2014SW001083>
- Ripoll, J.-F., Santolik, O., Reeves, G. D., Kurth, W. S., Denton, M. H., Lorida, V., et al. (2017). Effects of whistler mode hiss waves in March 2013. *Journal of Geophysical Research: Space Physics*, *122*, 7433–7462. <https://doi.org/10.1002/2017JA024139>
- Saikin, A. A., Zhang, J.-C., Allen, R. C., Smith, C. W., Kistler, L. M., Spence, H. E., et al. (2015). The occurrence and wave properties of H⁺, He⁺, and O⁺-band EMIC waves observed by the Van Allen Probes. *Journal of Geophysical Research: Space Physics*, *120*, 7477–7492. <https://doi.org/10.1002/2015JA021358>
- Sandhu, J. K., Yeoman, T. K., Fear, R. C., & Dandouras, I. (2016). A statistical study of magnetospheric ion composition along the geomagnetic field using the Cluster spacecraft for L values between 5.9 and 9.5. *Journal of Geophysical Research: Space Physics*, *121*, 2194–2208. <https://doi.org/10.1002/2015JA022261>
- Sandhu, J. K., Yeoman, T. K., Rae, I. J., Fear, R. C., & Dandouras, I. S. (2017). The dependence of magnetospheric plasma mass loading on geomagnetic activity using Cluster. *Journal of Geophysical Research: Space Physics*, *122*, 9371–9395. <https://doi.org/10.1002/2017JA024171>
- Santolik, O., Kletzing, C. A., Kurth, W. S., Hospodarsky, G. B., & Bounds, S. R. (2014). Fine structure of large-amplitude chorus wave packets. *Geophysical Research Letters*, *41*, 293–299. <https://doi.org/10.1002/2013GL058889>
- Santolik, O., Macúsová, E., Kolmasová, I., Cornilleau-Wehrlin, N., & de Conchy, Y. (2014). Propagation of lower-band whistler-mode waves in the outer Van Allen belt: Systematic analysis of 11 years of multi-component data from the Cluster spacecraft. *Geophysical Research Letters*, *41*, 2729–2737. <https://doi.org/10.1002/2014GL059815>
- Sarno-Smith, L. K., Larsen, B. A., Skoug, R. M., Liemohn, M. W., Breneman, A., Wygant, J. R., & Thomsen, M. F. (2016). Spacecraft surface charging within geosynchronous orbit observed by the Van Allen Probes. *Space Weather*, *14*, 151–164. <https://doi.org/10.1002/2015SW001345>
- Sarno-Smith, L. K., Liemohn, M. W., Katus, R. M., Skoug, R. M., Larsen, B. A., Thomsen, M. F., et al. (2015). Postmidnight depletion of the high-energy tail of the quiet plasmasphere. *Journal of Geophysical Research: Space Physics*, *120*, 1646–1660. <https://doi.org/10.1002/2014JA020682>

- Schulz, M. (1998). Particle drift and loss rates under strong pitch angle diffusion in Dungey's model magnetosphere. *Journal of Geophysical Research*, 103(A1), 61–67. <https://doi.org/10.1029/97JA02042>
- Shi, R., Li, W., Ma, Q., Claudepierre, S. G., Kletzing, C. A., & William, S. (2018). Van Allen Probes observation of plasmaspheric hiss modulated by injected energetic electrons, (January), 1–34
- Shprits, Y., Kellerman, A., Kondrashov, D., & Subbotin, D. (2013). Application of a new data operator-splitting data assimilation technique to the 3-D VERB diffusion code and CRRES measurements. *Geophysical Research Letters*, 40, 4998–5002. <https://doi.org/10.1002/grl.50969>
- Shprits, Y. Y. (2016). Estimation of bounce resonant scattering by fast magnetosonic waves. *Geophysical Research Letters*, 43, 998–1006. <https://doi.org/10.1002/2015GL066796>
- Shprits, Y. Y., Drozdov, A. Y., Spasojevic, M., Kellerman, A. C., Usanova, M. E., Engebretson, M. J., et al. (2016). Wave-induced loss of ultra-relativistic electrons in the Van Allen radiation belts. *Nature Communications*, 7, 12883. <https://doi.org/10.1038/ncomms12883>
- Spasojevic, M., Shprits, Y. Y., & Orlova, K. (2015). Global empirical models of plasmaspheric hiss using Van Allen Probes. *Journal of Geophysical Research: Space Physics*, 120, 10,370–10,383. <https://doi.org/10.1002/2015JA021803>
- Tao, X., Bortnik, J., Albert, J. M., & Thorne, R. M. (2012). Comparison of bounce-averaged quasi-linear diffusion coefficients for parallel propagating whistler mode waves with test particle simulations. *Journal of Geophysical Research*, 117, A10205. <https://doi.org/10.1029/2012JA017931>
- Toffoletto, F., Sazykin, S., Spiro, R., & Wolf, R. (2003). Inner magnetospheric modeling with the Rice Convection Model. *Space Science Reviews*, 107(1/2), 175–196. <https://doi.org/10.1023/A:1025532008047>
- Tsyganenko, N. A. (1989). A magnetospheric magnetic field model with a warped tail current sheet. *Planetary and Space Science*, 37(1), 5–20. [https://doi.org/10.1016/0032-0633\(89\)90066-4](https://doi.org/10.1016/0032-0633(89)90066-4)
- Tsyganenko, N. A. (1996). Modeling the Earth's magnetospheric magnetic field, confined within a realistic magnetopause. *Journal of Geophysical Research*, 100, 5599–5612. <https://doi.org/10.1029/94JA03193>
- Tsyganenko, N. A. (2002). A model of the near magnetosphere with a dawn-dusk asymmetry: 1. Mathematical structure. *Journal of Geophysical Research*, 107(A8), 1179. <https://doi.org/10.1029/2001JA000219>
- Tsyganenko, N. A., & Andreeva, V. A. (2017). A hybrid approach to empirical magnetosphere modeling. *Journal of Geophysical Research: Space Physics*, 122, 8198–8213. <https://doi.org/10.1002/2017JA024359>
- Tsyganenko, N. A., & Sitnov, M. I. (2007). Magnetospheric configurations from a high-resolution data-based magnetic field model. *Journal of Geophysical Research*, 112, A06225. <https://doi.org/10.1029/2007JA012260>
- Tu, W., Cunningham, G. S., Chen, Y., Morley, S. K., Reeves, G. D., Blake, J. B., et al. (2014). Event-specific chorus wave and electron seed population models in DREAM3D using the Van Allen Probes. *Geophysical Research Letters*, 41, 1359–1366. <https://doi.org/10.1002/2013GL058819>
- Turner, D. L., Angelopoulos, V., Li, W., Bortnik, J., Ni, B., Ma, Q., et al. (2014). Competing source and loss mechanisms due to wave-particle interactions in Earth's outer radiation belt during the 30 September to 3 October 2012 geomagnetic storm. *Journal of Geophysical Research: Space Physics*, 119, 1960–1979. <https://doi.org/10.1002/2014JA019770>
- Turner, D. L., Lee, J. H., Claudepierre, S. G., Fennell, J. F., Blake, J. B., Jaynes, A. N., et al. (2017). Examining coherency scales, sub-structure, and propagation of whistler mode chorus elements with magnetospheric multiscale (MMS). *Journal of Geophysical Research: Space Physics*, 122, 11,201–11,226. <https://doi.org/10.1002/2017JA024474>
- Usanova, M. E., Drozdov, A., Orlova, K., Mann, I. R., Shprits, Y., Robertson, M. T., et al. (2014). Effect of EMIC waves on relativistic and ultrarelativistic electron populations: Ground-based and Van Allen Probes observations. *Geophysical Research Letters*, 41, 1375–1381. <https://doi.org/10.1002/2013GL059024>
- Usanova, M. E., Malaspina, D. M., Jaynes, A. N., Bruder, R. J., Mann, I. R., Wygant, J. R., & Ergun, R. E. (2016). Van Allen Probes observations of oxygen cyclotron harmonic waves in the inner magnetosphere. *Geophysical Research Letters*, 43, 8827–8834. <https://doi.org/10.1002/2016GL070233>
- Wang, Z. Q., Pan, Z., Zhai, H., Gao, Z., Sun, K., & Zhang, Y. (2017). The nonlinear interactions between O⁺ ions and oxygen band EMIC waves: The nonlinear interactions of O⁺ ions. *Journal of Geophysical Research: Space Physics*, 122, 7097–7109. <https://doi.org/10.1002/2017JA024113>
- Wang, Z., Zhai, H., & Gao, Z. (2017). The effects of hydrogen band EMIC waves on ring current H⁺ ions. *Geophysical Research Letters*, 44, 11,722–11,728. <https://doi.org/10.1002/2017GL075843>
- Welling, D. T., Anderson, B. J., Crowley, G., Pulkkinen, A. A., & Rastätter, L. (2017). Exploring predictive performance: A reanalysis of the geospace model transition challenge. *Space Weather*, 15, 192–203. <https://doi.org/10.1002/2016SW001505>
- Welling, D. T., Jordanova, V. K., Glocer, A., Toth, G., Liemohn, M. W., & Weimer, D. R. (2015). The two-way relationship between ionospheric outflow and the ring current. *Journal of Geophysical Research: Space Physics*, 120, 4338–4353. <https://doi.org/10.1002/2015JA021231>
- Welling, D. T., Toth, G., Jordanova, V. K., & Yu, Y. (2018). Integration of RAM-SCB into the Space Weather Modeling Framework. *Journal of Atmospheric and Solar-Terrestrial Physics*, 177, 160–168. <https://doi.org/10.1016/j.jastp.2018.01.007>
- Wiltberger, M., Merkin, V., Zhang, B., Toffoletto, F., Oppenheim, M., Wang, W., et al. (2017). Effects of electrojet turbulence on a magnetosphere-ionosphere simulation of a geomagnetic storm. *Journal of Geophysical Research: Space Physics*, 122, 5008–5027. <https://doi.org/10.1002/2016JA023700>
- Xi, S., Lotko, W., Zhang, B., Wiltberger, M., & Lyon, J. (2016). Effects of auroral potential drops on plasma sheet dynamics. *Journal of Geophysical Research: Space Physics*, 121, 11,129–11,144. <https://doi.org/10.1002/2016JA022856>
- Yang, J., Toffoletto, F. R., & Wolf, R. A. (2016). Comparison study of ring current simulations with and without bubble injections. *Journal of Geophysical Research: Space Physics*, 121, 374–379. <https://doi.org/10.1002/2015JA021901>
- Yang, J., Toffoletto, F. R., Wolf, R. A., & Sazykin, S. (2015). On the contribution of plasma sheet bubbles to the storm time ring current. *Journal of Geophysical Research: Space Physics*, 120, 7416–7432. <https://doi.org/10.1002/2015JA021398>
- Yu, J., Li, L. Y., Cao, J. B., Yuan, Z. G., Reeves, G. D., Baker, D. N., et al. (2015). Multiple loss processes of relativistic electrons outside the heart of outer radiation belt during a storm sudden commencement. *Journal of Geophysical Research: Space Physics*, 120, 10,275–10,288. <https://doi.org/10.1002/2015JA021460>
- Yu, Y., Cao, J., Fu, H., Lu, H., & Yao, Z. (2017). The effects of bursty bulk flows on global-scale current systems. *Journal of Geophysical Research: Space Physics*, 122, 6139–6149. <https://doi.org/10.1002/2017JA024168>
- Yu, Y., Jordanova, V., Welling, D., Larsen, B., Claudepierre, S. G., & Kletzing, C. (2014). The role of ring current particle injections: Global simulations and Van Allen Probes observations during 17March 2013 storm. *Geophysical Research Letters*, 41, 1126–1132. <https://doi.org/10.1002/2014GL059322>

- Yu, Y., Jordanova, V., Zou, S., Heelis, R., Ruohoniemi, M., & Wygant, J. (2015). Modeling subauroral polarization streams during the 17 March 2013 storm. *Journal of Geophysical Research: Space Physics*, *120*, 1738–1750. <https://doi.org/10.1002/2014JA020371>
- Yu, Y., Jordanova, V. K., McGranaghan, R. M., & Solomon, S. C. (2018). Self-consistent modeling of electron precipitation and responses in the ionosphere: Application to low-altitude energization during substorms. *Geophysical Research Letters*, *45*, 6371–6381. <https://doi.org/10.1029/2018GL078828>
- Yu, Y., Jordanova, V. K., Ridley, A. J., Albert, J. M., Horne, R. B., & Jeffery, C. A. (2016). A new ionospheric electron precipitation module coupled with RAM-SCB within the geospace general circulation model. *Journal of Geophysical Research: Space Physics*, *121*, 8554–8575. <https://doi.org/10.1002/2016JA022585>
- Yu, Y., Rastaetter, L., Jordanova, V. K., Zheng, Y., Engel, M., Fok, M.-C., & Kuznetsova, M. (2019). Initial results from the GEM challenge on the spacecraft surface charging environment. *Space Weather*, *17*. <https://doi.org/10.1029/2018SW002031>
- Yue, C., Bortnik, J., Li, W., Ma, Q., Gkioulidou, M., Reeves, G. D., Wang, C. P., et al. (2018). The composition of plasma inside geostationary orbit based on Van Allen Probes observations. *Journal of Geophysical Research: Space Physics*, *123*, 6478–6493. <https://doi.org/10.1029/2018JA025344>
- Yue, C., Chen, L., Bortnik, J., Ma, Q., Thorne, R. M., Angelopoulos, V., et al. (2017). The characteristic response of whistler mode waves to interplanetary shocks. *Journal of Geophysical Research: Space Physics*, *122*, 10,047–10,057. <https://doi.org/10.1002/2017JA024574>
- Zaharia, S., Jordanova, V. K., Thomsen, M. F., & Reeves, G. D. (2008). Self-consistent geomagnetic storm simulation: The role of the induced electric fields. *Journal of Atmospheric and Solar-Terrestrial Physics*, *70*, 511–518. <https://doi.org/10.1016/j.jastp.2007.08.067>
- Zhang, B., Brambles, O. J., Lotko, W., Ouellette, J. E., & Lyon, J. G. (2016). The role of ionospheric O⁺ outflow in the generation of earthward propagating plasmoids. *Journal of Geophysical Research: Space Physics*, *121*, 1425–1435. <https://doi.org/10.1002/2015JA021667>
- Zhang, J.-C., Saikin, A. A., Kistler, L. M., Smith, C. W., Spence, H. E., Mouikis, C. G., et al. (2014). Excitation of EMIC waves detected by the Van Allen Probes on 28 April 2013. *Geophysical Research Letters*, *41*, 4101–4108. <https://doi.org/10.1002/2014GL060621>
- Zhang, X.-J., Angelopoulos, V., Ni, B., & Thorne, R. M. (2015). Predominance of ECH wave contribution to diffuse aurora in Earth's outer magnetosphere. *Journal of Geophysical Research: Space Physics*, *120*, 295–309. <https://doi.org/10.1002/2014JA020455>
- Zhang, X.-J., Li, W., Thorne, R. M., Angelopoulos, V., Bortnik, J., Kletzing, C. A., et al. (2016). Statistical distribution of EMIC wave spectra: Observations from Van Allen Probes. *Geophysical Research Letters*, *43*, 12,348–12,355. <https://doi.org/10.1002/2016GL071158>
- Zhang, X.-X., He, F., Lin, R.-L., Fok, M.-C., Katus, R. M., Liemohn, M. W., et al. (2017). A new solar wind-driven global dynamic plasmapause model: 1. Database and statistics. *Journal of Geophysical Research: Space Physics*, *122*, 7153–7171. <https://doi.org/10.1002/2017JA023912>
- Zhao, H., Li, X., Blake, J. B., Fennell, J. F., Claudepierre, S. G., Baker, D. N., et al. (2014). Peculiar pitch angle distribution of relativistic electrons in the inner radiation belt and slot region. *Geophysical Research Letters*, *41*, 2250–2257. <https://doi.org/10.1002/2014GL059725>
- Zhelavskaya, I. S., Shprits, Y. Y., & Spasojevic, M. (2017). Empirical modeling of the plasmasphere dynamics using neural networks. *Journal of Geophysical Research: Space Physics*, *122*, 11,227–11,244. <https://doi.org/10.1002/2017JA024406>
- Zhima, Z., Chen, L., Fu, H., Cao, J., Horne, R. B., & Reeves, G. (2015). Observations of discrete magnetosonic waves off the magnetic equator. *Geophysical Research Letters*, *42*, 9694–9701. <https://doi.org/10.1002/2015GL066255>
- Zhima, Z., Cao, J. B., Liu, W. L., Fu, H. S., Wang, T. Y., Zhang, X. M., & Shen, X. H. (2014). Storm-time evolution of ELF/VLF waves observed by DEMETER satellite. *Journal of Geophysical Research: Space Physics*, *119*, 2612–2622. <https://doi.org/10.1002/2013JA019237>
- Zhou, X.-Z., Wang, Z.-H., Zong, Q.-G., Claudepierre, S. G., Mann, I. R., Kivelson, M. G., et al. (2015). Imprints of impulse-excited hydro-magnetic waves on electrons in the Van Allen radiation belts. *Geophysical Research Letters*, *42*, 6199–6204. <https://doi.org/10.1002/2015GL064988>