

Eplerenone versus placebo renal function stratified dose comparisons in the EMPHASIS-HF trial

João Pedro Ferreira^{1,2}; Paula Abreu³; John J. V. McMurray⁴; Dirk J. van Veldhuisen⁵; Karl Swedberg⁶; Stuart J. Pocock⁷; John Vincent³; Katharina Lins³; Patrick Rossignol¹; Bertram Pitt⁸; Faiez Zannad¹

¹ National Institute of Health and Medical Research (INSERM), Center for Clinical Multidisciplinary Research 1433, INSERM U1116, University of Lorraine, Regional University Hospital of Nancy, French Clinical Research Infrastructure Network (F-CRIN) Investigation Network Initiative-Cardiovascular and Renal Clinical Trialists (INI-CRCT), Nancy, France.

² Department of Physiology and Cardiothoracic Surgery, Cardiovascular Research and Development Unit, Faculty of Medicine, University of Porto, Porto, Portugal.

³ Pfizer Inc., New York, USA.

⁴ BHF Cardiovascular Research Centre, University of Glasgow, Glasgow, Scotland, UK.

⁵ Department of Cardiology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands.

⁶ University of Gothenburg, Gothenburg, Sweden.

⁷ Department of Medical Statistics, London School of Hygiene and Tropical Medicine, London, UK.

⁸ Department of Medicine, University of Michigan School of Medicine, Ann Arbor, Michigan.

Corresponding author: Pr Faiez Zannad

Centre d'Investigations Cliniques-INSERM CHU de Nancy

Institut Lorrain du Cœur et des Vaisseaux Louis Mathieu

4 rue du Morvan

54500 Vandoeuvre Lès Nancy

France

Tel : +33 3 83 15 73 20

Fax : +33 3 83 15 73 24

f.zannad@chru-nancy.fr

Word-count: 2152 (excluding abstract and references)

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1002/ejhf.1400](https://doi.org/10.1002/ejhf.1400)

Abstract

Background: Current heart failure (HF) guidelines recommend target eplerenone dose of 50mg/day. We have examined the effect of different eplerenone doses based on prespecified renal function stratification in the Eplerenone in Patients with Systolic Heart Failure and Mild Symptoms (EMPHASIS-HF) trial.

Methods: In EMPHASIS-HF, the target dose of eplerenone/placebo was stratified at randomisation according to estimated glomerular filtration rate (eGFR): 50mg/day if eGFR ≥ 50 ml/min/1.73m² and 25mg/day if eGFR 30-49ml/min/1.73m². Patients remained within these dose ranges during the trial (*per* stratification). The primary outcome was a composite of heart failure hospitalization (HFH) or cardiovascular mortality (CVM).

Results: Eplerenone was superior to placebo within each respective eGFR stratum: hazard ratio (HR) eplerenone vs. placebo in the eGFR ≥ 50 ml/min/1.73m² stratum=0.58 (0.45-0.74) and HR in the eGFR 30-49ml/min/1.73m² stratum=0.62 (0.49-0.78); p for interaction=0.89. Despite receiving lower eplerenone doses patients in the eGFR 30-49ml/min/1.73m² stratum had more often hyperkalemia, renal failure events, and drug discontinuation.

Conclusion: In EMPHASIS-HF the eplerenone effect was not influenced by the eGFR. Patients with impaired renal function experienced more adverse events despite receiving lower eplerenone doses. The current guidelines do not specify eplerenone dose recommendation according to renal function and should thus be adapted in the light of these data.

Key-words: eplerenone, heart failure, treatment dose, renal function, stratification.

ClinicalTrials.gov identifier: NCT00232180

Introduction

Current heart failure (HF) guidelines recommend up-titration of angiotensin converting enzyme inhibitor (ACEi), angiotensin receptor blocker (ARB), and beta-blocker (BB) doses to evidence-based targets based upon those used in pivotal clinical trials in heart failure with reduced ejection fraction (HFrEF). In ordinary practice, these doses are not attained in many patients despite the randomized evidence showing the benefit of higher ACEi/ARB and BB doses¹⁻⁶.

By contrast, there is no study comparing different doses of a mineralocorticoid receptor antagonists (MRA)⁷. For eplerenone, the current guidelines recommend a starting dose of 25 mg/day and a target dose of 50 mg/day regardless of renal function^{1,2}.

In the Eplerenone in Patients with Systolic Heart Failure and Mild Symptoms trial (EMPHASIS-HF), eplerenone reduced the risk of death and the risk of hospitalization, compared with placebo, in patients with heart failure and reduced left ventricular ejection fraction (LVEF) who were in New York Heart Association (NYHA) functional class II at the time of randomization⁸. Because of pharmacokinetic and safety considerations, patients were stratified at randomization to either a higher target dose (50 mg/day) of placebo/eplerenone or to a lower target dose (up to 25 mg/day), according to estimated glomerular filtration rate (eGFR) strata. We used this pre-specified dose-stratification to compare the efficacy and safety of low-dose eplerenone vs. low-dose placebo and high-dose eplerenone vs. high-dose placebo by renal function strata.

Methods

EMPHASIS-HF trial design

The design of EMPHASIS-HF is published⁸. In short, EMPHASIS-HF was a randomized, double-blind trial in which 2,737 patients in NYHA functional class II and with LVEF \geq 35% were

randomized to eplerenone or placebo, added to other recommended therapies. The primary outcome was a composite of death from cardiovascular causes (CVM) or hospitalization for heart failure (HFH). The median duration of follow-up was 21 months. The primary outcome occurred in 18.3% of patients in the eplerenone group, compared with 25.9% in the placebo group: HR (95%CI) =0.63 (0.54-0.74); $p < 0.001$.

Eplerenone dose attribution and adjustment

Patients were stratified to receive “high-dose” or “low-dose” study treatment according to eGFR as *per* stratification protocol. The main reason why a lower target dose of eplerenone was chosen in patients with an eGFR between 30 and 49 ml/min/1.73 m² was because in the Eplerenone Post-Acute Myocardial Infarction Heart Failure Efficacy and Survival Study (EPHESUS)⁹, where no prespecified dose allocation was performed, patients with eGFR <50 ml/min/1.73m² had higher incidence of serious hyperkalemia with eplerenone compared to placebo (10.1% vs. 5.9%; $p = 0.006$), whereas in patients with an eGFR ≥ 50 ml/min/1.73m² the corresponding hyperkalemia rates were much lower (4.6 % vs. 3.5%; $p = 0.04$). In order to avoid excessive side-effects in high-risk patients with impaired renal function, these received lower study drug doses by protocol prespecification.

In concordance, placebo/eplerenone was started at a dose of 25 mg/day and could be increased after 4 weeks up to 50 mg/day if the eGFR was ≥ 50 ml/min/1.73 m²; or started at 25 mg on alternate days, and increased to 25 mg/day if the eGFR was 30 to 49 ml/min/1.73 m². By protocol, eplerenone/placebo doses were maintained in these dose ranges with drug-dose adjustments allowed according to potassium levels, as follows: if the serum potassium level was 5.5 to 5.9 mmol/L the study drug dose would be decreased and if the serum potassium level was ≥ 6.0 mmol/L the study drug would be temporarily stopped. Potassium was to be re-measured within 72 hours after the dose reduction or study-drug withdrawal, and the study drug was to be restarted only if the level was below 5.0 mmol/L.

Statistical analysis

In descriptive analyses, continuous variables are expressed as mean \pm standard deviation (SD). Categorical variables are expressed as frequencies and proportions (%). Comparison of patients in the low-dose and high-dose strata and within each dose strata (placebo versus eplerenone) was performed using an independent samples t-test and a chi-square test for categorical variables. Normality assumptions were verified.

The primary outcome was a composite of HFH or CVM. Cox proportional hazard regression models were used to model long-term event rates both in univariable and multivariable analysis. Cox

proportional-hazards assumptions were assessed and no violations were found. The variables used to adjust outcomes were those used in a published risk model developed in EMPHASIS-HF¹⁰ *i.e.*, age, sex, systolic blood pressure (SBP), eGFR¹¹, diabetes, prior HFH, haemoglobin, prior myocardial infarction (MI)/coronary artery bypass grafting (CABG), body mass index (BMI).

All analyses were performed with SAS® software version 9.4 (SAS Institute Inc., Cary, N.C., USA).

Results

Characteristics of the study population

Within the respective eGFR stratum the randomization resulted in treatment groups that were well balanced in terms of their clinical characteristics, in accordance with the study overall. **Table 1.**

Comparison of eplerenone and placebo doses during the trial by eGFR strata

The mean eplerenone/placebo doses in the eGFR ≥ 50 ml/min/1.73m² stratum were of 25 mg/day at the study start, increased to ≈ 40 mg/day at week 4 and were maintained stable at ≈ 40 mg/day during the trial. **Table 2.** The mean eplerenone/placebo doses in the eGFR 30-49 ml/min/1.73m² stratum were inferior to 17 mg/day at the study start, increased up to 23 mg/day at week 4 and did not exceed 30 mg/day during the trial. **Table 2.**

Comparison of eplerenone with placebo by eGFR strata

The event rate reduction with eplerenone compared to placebo was similar within each eGFR stratum: HR eplerenone vs. placebo in the eGFR ≥ 50 ml/min/1.73m² stratum =0.58 (0.45-0.74) and HR eplerenone vs. placebo in the eGFR 30-49 ml/min/1.73m² stratum =0.62 (0.49-0.78); between strata p for interaction =0.89. **Table 3 & Figure 1.**

Adverse events

Hyperkalemia ($K^+ > 5.5$ mmol/L) was more frequent with eplerenone compared to placebo regardless of the eGFR stratum. However, hyperkalemia, renal failure and drug discontinuation were more frequent with low-dose eplerenone/placebo (*i.e.*, eGFR 30-49 ml/min/1.73m² stratum) compared with high-dose (*i.e.*, eGFR ≥ 50 ml/min/1.73m² stratum). For example, hyperkalemia was observed in 1% and 4% of patients randomized to placebo and eplerenone, respectively, in the eGFR ≥ 50 ml/min/1.73m² stratum, whereas in the eGFR 30-49 ml/min/1.73m² stratum these proportions increased to 7% with placebo and 13% with eplerenone ($p < 0.001$). **Table 4.**

Discussion

In EMPHASIS-HF the eplerenone effect was not influenced by the eGFR *i.e.* the treatment effect was similar regardless of the eGFR stratum. However, as *per* stratification, eplerenone/placebo doses were much lower in patients with eGFR below 50ml/min/1.73m²; and despite these lower doses, side effects were observed more often. Therefore, using high (up to 50 mg/day) eplerenone doses in patients with impaired renal function may greatly increase the rate of adverse events and drug discontinuation. The current guidelines do not specify eplerenone dose recommendation in HFrEF and should thus be adapted in order to reflect the protocol of the EMPHASIS-HF trial.

To date, no randomized trials exist directly comparing different doses of eplerenone (or any other aldosterone antagonist). There are, however, two large trials in which patients were prospectively randomized to a high or low dose of ACE inhibitor or ARB: The Assessment of Treatment with Lisinopril and Survival trial (ATLAS) and the Heart failure Endpoint evaluation with the Angiotensin II Antagonist Losartan trial (HEAAL). In ATLAS, 3,164 HF patients with an ejection fraction $\geq 30\%$ were randomized to double-blind treatment with either low doses (2.5 to 5.0 mg daily, n=1596) or high doses (32.5 to 35 mg daily, n=1568) of the ACE inhibitor lisinopril. Compared with the low-dose group, patients in the high-dose group had a significant 12% lower relative risk of death or hospitalization for any reason (p=0.002) and 24% fewer hospitalizations for HF (p=0.002). Drug discontinuation due to side-effects was similar between groups. In HEAAL, 3,846 HF patients with an ejection fraction $\geq 40\%$ and intolerance to ACE inhibitors were randomly assigned to low-dose (50 mg daily, n=1919) or high-dose (150 mg daily, n=1927) of the ARB losartan. Compared with the low-dose group, patients in the high-dose group had a significant 10% lower relative risk of death or hospitalization for HF (p=0.027) and 13% fewer hospitalizations for HF (p=0.025). Drug discontinuation due to side-effects was also similar between groups. These findings indicate that HF patients should not be maintained on low doses of an ACE inhibitor or ARB (unless these are the only doses that can be tolerated). In contrast, MRA dose comparisons have not been performed to date. The design of EMPHASIS-HF was different: by stratification two dose-levels were compared with placebo rather than directly low vs. high treatment dose. It should be pointed out, that unlike ATLAS and HEAAL, EMPHASIS-HF patients were randomized within two strata which were determined by renal function, hence high-dose vs. low-dose treatment cannot be compared. In EMPHASIS-HF low-dose was as effective as high-dose eplerenone when used in appropriate patients (*i.e.*, low-dose for patients with eGFR 30-49ml/min/1.73m² and high-dose for patients with eGFR ≥ 50 ml/min/1.73m²), supporting the use of eplerenone at doses around 25 mg/day in patients with eGFR 30-49ml/min/1.73m²

and around 50 mg/day in patients with eGFR ≤ 50 ml/min/1.73m², adapting for potassium levels when required.

Stratification is usually performed to ensure that strong outcome or treatment response predictors are balanced between randomization groups, but stratification is also the only situation in which balanced randomization is maintained in subgroups, since the randomization is performed within each stratum¹². Therefore, strata analyses are less permeable to bias caused by imbalances in treatment allocation and patients' characteristics, which inevitably hamper all analyses made on subgroups defined from non-randomized baseline characteristics. In the absence of a statistical interaction (*i.e.* similar between-strata hazard ratios, as observed herein) the treatment effect can be considered similar between both strata provided that the same strata-treatment doses are used in clinical practice. These findings, should thus change the current guidelines where no eGFR-specific eplerenone dose recommendation is provided^{1,2}; and many patients may be receiving inappropriate doses of eplerenone contributing to higher hyperkalemia rates and drug discontinuation¹³.

In summary, the present analysis of stratified randomized data from EMPHASIS-HF provides robust evidence that eplerenone is equally beneficial and should be used in clinical practice at the respective target doses of 50 mg/day in patients with eGFR ≤ 50 ml/min/1.73m² and 25 mg/day in patients with eGFR between 30 and 49 ml/min/1.73m².

Limitations

This is an analysis of prespecified strata. Hence, our findings are as robust as the main randomized clinical trial because no statistical interaction (*i.e.* treatment effect differences) was observed between strata.

Conclusion

In EMPHASIS-HF the eplerenone effect was not influenced by the eGFR. Patients with impaired renal function experienced more adverse events despite receiving lower eplerenone doses. The current guidelines do not specify eplerenone dose recommendation according to renal function and should thus be adapted in the light of these data.

Funding

The EMPHASIS-HF Study was sponsored by Pfizer.

Disclosures


J.J.V.M., D.J.vV., K.S., S.J.P., B.P., and F.Z. are members of the EMPHASIS-HF Writing Committee and report having received fees and travel support in the past from the study sponsor, Pfizer Inc., for participation in and travelling to meetings of the committee. P.A., J.V. and K.L. are currently employed by Pfizer and own stock in Pfizer Inc., the makers of eplerenone. K.S. has received research support from Pfizer, Amgen, Novartis, and Servier. S.J.P. reports receiving consulting fees from Servier, Amgen, AstraZeneca, and Novartis, and that his institution receives grants from Servier and AstraZeneca on his behalf. B.P. reports receiving fees for serving on the board of Novartis, consulting fees from Takeda, AstraZeneca, Boehringer Ingelheim, GE Healthcare, Relypsa, BG Medicine, Nile Therapeutics, Merck, Forest Laboratories, and Novartis, grant support from Forest Laboratories and Novartis, and stock options from Relypsa, BG Medicine, Nile Therapeutics, and Aurasenc, that his institution receives grant support from Forest Laboratories on his behalf and he and his institution receive grant support from Bayer. F.Z. reports receiving fees for serving on the board of Boston Scientific, consulting fees from Novartis, Takeda, AstraZeneca, Boehringer Ingelheim, GE Healthcare, Relypsa, Servier, Boston Scientific, Bayer, Johnson & Johnson, and Resmed, and speaker's fees from Pfizer and AstraZeneca, and that his institution receives grant support from BG Medicine and Roche Diagnostics on his behalf. J.P.F. has reported that he has no relationships relevant to the contents of this paper to disclose.

Bibliography

1. Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JG, Coats AJ, Falk V, Gonzalez-Juanatey JR, Harjola VP, Jankowska EA, Jessup M, Linde C, Nihoyannopoulos P, Parissis JT, Pieske B, Riley JP, Rosano GM, Ruilope LM, Ruschitzka F, Rutten FH and van der Meer P. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC). Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur J Heart Fail.* 2016.
2. Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE, Jr., Colvin MM, Drazner MH, Filippatos G, Fonarow GC, Givertz MM, Hollenberg SM, Lindenfeld J, Masoudi FA, McBride PE, Peterson PN, Stevenson LW and Westlake C. 2016 ACC/AHA/HFSA Focused Update on New Pharmacological Therapy for Heart Failure: An Update of the 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *J Am Coll Cardiol.* 2016.
3. Maggioni AP, Anker SD, Dahlstrom U, Filippatos G, Ponikowski P, Zannad F, Amir O, Chioncel O, Leiro MC, Drozd J, Erglis A, Fazlibegovic E, Fonseca C, Fruhwald F, Gatzov P, Goncalvesova E, Hassanein M, Hradec J, Kavoliuniene A, Lainscak M, Logeart D, Merkely B, Metra M, Persson H, Seferovic P, Temizhan A, Tousoulis D and Tavazzi L. Are hospitalized or ambulatory patients with

- heart failure treated in accordance with European Society of Cardiology guidelines? Evidence from 12,440 patients of the ESC Heart Failure Long-Term Registry. *Eur J Heart Fail.* 2013;15:1173-84.
4. Lam PH, Dooley DJ, Fonarow GC, Butler J, Bhatt DL, Filippatos GS, Deedwania P, Forman DE, White M, Fletcher RD, Arundel C, Blackman MR, Adamopoulos C, Kanonidis IE, Aban IB, Patel K, Aronow WS, Allman RM, Anker SD, Pitt B and Ahmed A. Similar clinical benefits from below-target and target dose enalapril in patients with heart failure in the SOLVD Treatment trial. *Eur J Heart Fail.* 2017.
 5. Khan MS, Fonarow GC, Ahmed A, Greene SJ, Vaduganathan M, Khan H, Marti C, Gheorghiade M and Butler J. Dose of Angiotensin-Converting Enzyme Inhibitors and Angiotensin Receptor Blockers and Outcomes in Heart Failure: A Meta-Analysis. *Circ Heart Fail.* 2017;10.
 6. Bristow MR, Gilbert EM, Abraham WT, Adams KF, Fowler MB, Hershberger RE, Kubo SH, Narahara KA, Ingersoll H, Krueger S, Young S and Shusterman N. Carvedilol produces dose-related improvements in left ventricular function and survival in subjects with chronic heart failure. MOCHA Investigators. *Circulation.* 1996;94:2807-16.
 7. Ferreira JP, Mentz RJ, Pizard A, Pitt B and Zannad F. Tailoring mineralocorticoid receptor antagonist therapy in heart failure patients: are we moving towards a personalized approach? *Eur J Heart Fail.* 2017.
 8. Zannad F, McMurray JJ, Krum H, van Veldhuisen DJ, Swedberg K, Shi H, Vincent J, Pocock SJ and Pitt B. Eplerenone in patients with systolic heart failure and mild symptoms. *N Engl J Med.* 2011;364:11-21.
 9. Pitt B, Williams G, Remme W, Martinez F, Lopez-Sendon J, Zannad F, Neaton J, Roniker B, Hurley S, Burns D, Bittman R and Kleiman J. The EPHEsus trial: eplerenone in patients with heart failure due to systolic dysfunction complicating acute myocardial infarction. Eplerenone Post-AMI Heart Failure Efficacy and Survival Study. *Cardiovasc Drugs Ther.* 2001;15:79-87.
 10. Collier TJ, Pocock SJ, McMurray JJ, Zannad F, Krum H, van Veldhuisen DJ, Swedberg K, Shi H, Vincent J and Pitt B. The impact of eplerenone at different levels of risk in patients with systolic heart failure and mild symptoms: insight from a novel risk score for prognosis derived from the EMPHASIS-HF trial. *Eur Heart J.* 2013;34:2823-9.
 11. Levey AS, Bosch JP, Lewis JB, Greene T, Rogers N and Roth D. A more accurate method to estimate glomerular filtration rate from serum creatinine: a new prediction equation. Modification of Diet in Renal Disease Study Group *Ann Intern Med* United States; 1999(130): 461-70.
 12. Kernan WN, Viscoli CM, Makuch RW, Brass LM and Horwitz RI. Stratified randomization for clinical trials. *J Clin Epidemiol.* 1999;52:19-26.
 13. Juurlink DN, Mamdani MM, Lee DS, Kopp A, Austin PC, Laupacis A and Redelmeier DA. Rates of hyperkalemia after publication of the Randomized Aldactone Evaluation Study. *N Engl J Med.* 2004;351:543-51.

Figure 1. Kaplan-Meier curves for the primary outcome of HFH or CVM of eplerenone vs. placebo within eGFR strata



Legend: HFH, hospitalization for heart failure; CVM, cardiovascular mortality; eGFR, estimated glomerular filtration rate.

Between eGFR strata P for interaction =0.89.

Table 1. Comparison of patients within each eGFR stratum at week 4: e50 ml/min/1.73m² or 30-49 ml/min/1.73m²

Patients` characteristics	eGFR stratum: e50 ml/min/1.73m ²		p-value	eGFR stratum: 30-49 ml/min/1.73m ²		p-value
	High-dose eplerenone N=742	High-dose placebo N=825		Low-dose eplerenone N=618	Low-dose placebo N=544	
<i>Demographics</i>						
Age (years)	67.7±7.3	67.7±7.4	NS	69.8±7.9	70.1±7.0	NS
Male gender, n (%)	579 (78%)	660 (80%)	NS	474 (77%)	409 (75%)	NS
BMI (Kg/m ²)	27.6±4.9	27.6±4.7	NS	27.4±5.0	27.4±5.0	NS
SBP (mmHg)	125±17	125±16	NS	123±18	122±17	NS
Heart rate (bpm)	74±16	73±15	NS	73±15	74±16	NS
<i>Laboratory</i>						
eGFR (ml/min/1.73m ²)	78±20	76±20	0.048	63±20	62±22	NS
Hemoglobin (g/dL)	13.9±1.5	13.9±1.6	NS	13.7±1.6	13.6±1.6	NS
Sodium (mmol/L)	140±4	140±4	NS	140±4	139±4	NS
Potassium (mmo/L)	4.2±0.4	4.3±0.4	<0.001	4.3±0.5	4.3±0.4	NS
<i>Ecocardiography</i>						
LVEF (%)	26±5	26±5	NS	26±5	26±5	NS
<i>Co-morbidities</i>						
Diabetes, n (%)	235 (32%)	225 (27%)	NS	222 (36%)	173 (32%)	NS
AFib, n (%)	221 (30%)	243 (30%)	NS	188 (30%)	189 (35%)	NS
Prior HFH, n (%)	365 (49%)	409 (50%)	NS	346 (56%)	313 (58%)	NS
Prior MI, n (%)	378 (51%)	414 (50%)	NS	352 (57%)	279 (51%)	NS
ICD/CRT, n (%)	133 (18%)	160 (19%)	NS	166 (27%)	150 (28%)	NS
<i>Medications</i>						
ACEi/ARB, n (%)	708 (95%)	784 (95%)	NS	589 (95%)	507 (93%)	NS
Beta-blocker, n (%)	661 (89%)	745 (90%)	NS	540 (87%)	468 (86%)	NS

Legend: BMI, body mass index; SBP; systolic blood pressure; eGFR, estimated glomerular filtration rate; LVEF, left ventricular ejection fraction; AFib, atrial fibrillation; MI, myocardial infarction; ICD/CRT; intracardiac defibrillator/cardiac resynchronization therapy; ACEi/ARB, angiotensin converting enzyme inhibitors/angiotensin receptor blocker.

Table 2. Mean eplerenone/placebo doses (in mg) within each eGFR stratum during the trial

eGFR strata	eGFR ≥ 50 ml/min/1.73m ² stratum		eGFR 30-49 ml/min/1.73m ² stratum	
	High-dose eplerenone	High-dose placebo	Low-dose eplerenone	Low-dose placebo
Study start	24.7±2.6	24.8±2.7	17.0±6.4	16.1±5.6
Week 4	38.9±13.2	40.4±12.7	23.4±9.6	23.3±9.8
Month 5	42.0±12.3	43.6±11.2	24.8±10.8	27.3±11.6
Month 12	42.3±12.4	43.9±11.1	26.8±12.3	30.2±12.6
Month 24	41.6±12.5	43.7±11.3	28.0±11.2	30.8±12.7
Study end	39.8±13.3	41.8±12.3	24.6±11.6	26.3±12.7

Table 3. Adjusted hazard ratio of within each eGFR strata

Study outcomes	eGFR e50 ml/min/1.73m ² stratum: High-dose eplerenone vs. placebo					eGFR 30-49 ml/min/1.73m ² stratum: Low-dose eplerenone vs. placebo					P for interaction
	Event (%) Eplerenone	Event (%) Placebo	ARD (%)	HR (95%CI)*	p-value	Event (%) Eplerenone	Event (%) Placebo	ARD (%)	HR (95%CI)*	p-value	
HFH/CVM	100 (13.5)	178 (21.6)	-8.1	0.58 (0.45-0.74)	<0.001	149 (24.1)	177 (32.5)	-8.4	0.62 (0.49-0.78)	<0.001	0.89
CVM	51 (6.9)	89 (10.8)	-3.9	0.61 (0.43-0.86)	0.004	96 (15.5)	95 (17.5)	-1.9	0.77 (0.58-1.04)	0.084	
HFH**	69 (9.3)	126 (15.3)	-6.0	0.56 (0.42-0.76)	<0.001	95 (15.4)	127 (23.3)	-8.0	0.55 (0.41-0.72)	<0.001	

*Model adjusted on age (75+ vs. <75 y), gender (male vs. female), systolic blood pressure (<130 vs. 130+ mmHg), heart rate (<80 vs. 80+ bpm), diabetes (yes vs. no), hemoglobin (<11 vs. 11-12.9 vs. 13+ g/dL), prior HFH (yes vs. no), eGFR (<60 vs. 60+ ml/min/1.73m²), prior MI/CABG (yes vs. no), BMI (<25 vs. 25+ Kg/m²). Note: unadjusted models provide similar results.



**Rate Ratio including repeated events.

Legend: HFH, hospitalization for heart failure; CVM, cardiovascular mortality; ARD, absolute risk difference; HR, hazard ratio.

Table 4. Investigator reported adverse events (AEs) by allocation dose

Adverse event (AE)	eGFR e 50 ml/min/1.73m ² stratum		p-value	eGFR 30-49 ml/min/1.73m ² stratum		p-value	p-value high vs. low eplerenone	p-value high vs. low placebo
	High-dose eplerenone	High-dose placebo		Low-dose eplerenone	Low-dose placebo			
Hyperkalemia	27 (3.6%)	11 (1.3%)	0.005	82 (13.3%)	39 (7.2%)	<0.001	<0.001	<0.001
Hypokalemia	5 (0.7%)	18 (2.2%)	0.018	11 (1.8%)	12 (2.2%)	0.68	0.064	1
Renal failure	9 (1.2%)	14 (1.7%)	0.53	29 (4.7%)	27 (5.0%)	0.89	<0.001	0.001
AE drug discontinuation	74 (10.0%)	108 (13.1%)	0.058	114 (18.4%)	114 (21.0%)	0.30	<0.001	<0.001

Author Manuscript



No. at Risk by Treatment Group

Eplerenone	618	369	222	86
Placebo	544	263	160	60

No. at Risk by Treatment Group

Eplerenone	742	555	340	146
Placebo	825	585	352	139

Fig_1.tif

Permission note:

“The authors do hereby declare that all illustrations and figures in the manuscript are entirely original and do not require reprint permission.”