

# Evaluation of vitamin D biosynthesis and pathway target genes reveals *UGT2A1/2* and *EGFR* polymorphisms associated with epithelial ovarian cancer in African American Women

RUNNING TITLE: Genetic association with variants in the vitamin D pathway

KEYWORDS: ovarian cancer, genetic association, vitamin D pathway, African ancestry risk

Delores J. Grant<sup>1</sup>, Ani Manichaikul<sup>2</sup>, Anthony J. Alberg<sup>3</sup>, Elisa V. Bandera<sup>4</sup>, Jill Barnholtz-Sloan<sup>5</sup>, Melissa Bondy<sup>6</sup>, Michele L. Cote<sup>7</sup>, Ellen Funkhouser<sup>8</sup>, Patricia G. Moorman<sup>9</sup>, Lauren C. Peres<sup>2</sup>, Edward S. Peters<sup>10</sup>, Ann G. Schwartz<sup>7</sup>, Paul D. Terry<sup>11</sup>, Xin-Qun Wang<sup>2</sup>, Temitope O. Keku<sup>12</sup>, Cathrine Hoyo<sup>13</sup>, Andrew Berchuck<sup>14</sup>, Dale P. Sandler<sup>15</sup>, Jack A. Taylor<sup>15</sup>, Katie M. O'Brien<sup>16</sup>, Digna R. Velez Edwards<sup>17</sup>, Todd L. Edwards<sup>18</sup>, Alicia Beeghly-Fadiel<sup>19</sup>, Nicolas Wentzensen<sup>20</sup>, Celeste Leigh Pearce<sup>21,22</sup>, Anna H. Wu<sup>22</sup>, Alice S. Whittemore<sup>23,24</sup>, Valerie McGuire<sup>23</sup>, Weiva Sieh<sup>25,26</sup>, Joseph H. Rothstein<sup>25,26</sup>, Francesmary Modugno<sup>27,28,29</sup>, Roberta Ness<sup>30</sup>, Kirsten Moysich<sup>31</sup>, Mary Anne Rossing<sup>32,33</sup>, Jennifer A. Doherty<sup>34</sup>, Thomas A. Sellers<sup>35</sup>, Jennifer B. Permuth-Way<sup>35</sup>, Alvaro N. Monteiro<sup>35</sup>, Douglas A. Levine<sup>36,37</sup>, Veronica Wendy Setiawan<sup>38</sup>, Christopher A. Haiman<sup>38</sup>, Loic LeMarchand<sup>39</sup>, Lynne R. Wilkens<sup>40</sup>, Beth Y. Karlan<sup>41</sup>, Usha Menon<sup>42</sup>, Susan Ramus<sup>43,44</sup>, Simon Gayther<sup>45,46</sup>, Aleksandra Gentry-Maharaj<sup>42</sup>, Kathryn L. Terry<sup>47,48</sup>, Daniel W. Cramer<sup>47,48</sup>, Ellen L. Goode<sup>49</sup>, Melissa C. Larson<sup>50</sup>, Scott H. Kaufmann<sup>51</sup>, Rikki Cannioto<sup>52</sup>, Kunle Odunsi<sup>53</sup>, John L. Etter<sup>31</sup>, Ruea-Yea Huang<sup>54</sup>, Marcus Q. Bernardini<sup>55</sup>, Alicia A. Tone<sup>55</sup>, Taymaa May<sup>55</sup>, Marc T. Goodman<sup>56,57</sup>, Pamela J. Thompson<sup>56</sup>, Michael E. Carney<sup>58</sup>, Shelley S. Tworoger<sup>59</sup>, Elizabeth M. Poole<sup>60</sup>, Diether Lambrechts<sup>61,62</sup>, Ignace Vergote<sup>63</sup>, Adriaan Vanderstichele<sup>63</sup>, Els Van Nieuwenhuysen<sup>63</sup>, Hoda Anton-Culver<sup>64</sup>, Argyrios Ziogas<sup>65</sup>, James D. Brenton<sup>66</sup>, Line Bjorge<sup>67,68</sup>, Helga B. Salvensen<sup>67,68</sup>, Lambertus A. Kiemeny<sup>69</sup>, Leon F. A. G. Massuger<sup>70</sup>, Tanja Pejovic<sup>71,72</sup>, Amanda Bruegl<sup>71,72</sup>, Melissa Moffitt<sup>71,72</sup>, Linda Cook<sup>73</sup>, Nhu D. Le<sup>74</sup>, Angela Brooks-Wilson<sup>75,76</sup>, Linda Kelemen<sup>77</sup>, Paul D. P. Pharoah<sup>78</sup>, Honglin Song<sup>79</sup>, Ian Campbell<sup>80,81</sup>, Diana Eccles<sup>82</sup>, Anna DeFazio<sup>83,84</sup>, Catherine J. Kennedy<sup>83,84</sup>, Joellen M. Schildkraut<sup>85</sup>

1. Department of Biological and Biomedical Sciences, Cancer Research Program, JLC-Biomedical/Biotechnology Research Institute, North Carolina Central University, Durham, North Carolina, USA
2. Center for Public Health Genomics, University of Virginia, Charlottesville, Virginia, USA

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/CAM4.1996](https://doi.org/10.1002/CAM4.1996)

This article is protected by copyright. All rights reserved

3. Department of Epidemiology and Biostatistics, Arnold School of Public Health, University of South Carolina, Columbia, SC, USA
4. Department of Population Science, Rutgers Cancer Institute of New Jersey, New Brunswick, New Jersey, USA
5. Case Comprehensive Cancer Center, Case Western Reserve University School of Medicine, Cleveland, Ohio, USA
6. Cancer Prevention and Population Sciences Program, Baylor College of Medicine, Houston, Texas, USA
7. Department of Oncology and the Karmanos Cancer Institute Population Studies and Disparities Research Program, Wayne State University School of Medicine, Detroit, Michigan, USA
8. Division of Preventive Medicine, University of Alabama at Birmingham, Birmingham, Alabama, USA
9. Department of Community and Family Medicine, Duke University Medical Center, Durham, North Carolina, USA
10. Epidemiology Program, Louisiana State University Health Sciences Center School of Public Health, New Orleans, Louisiana, USA
11. Department of Medicine, University of Tennessee Medical Center – Knoxville, Knoxville, Tennessee, USA
12. Departments of Medicine and Nutrition, Division of Gastroenterology and Hepatology, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA
13. Department of Biological Sciences, North Carolina State University, Raleigh, North Carolina, USA
14. Department of Obstetrics and Gynecology, Duke University Medical Center, Durham, North Carolina, USA
15. Epidemiology Branch, Division of Intramural Research, National Institute of Environmental Health Sciences, National Institutes of Health, Research Triangle Park, North Carolina, USA
16. Biostatistics and Computational Biology Branch, Division of Intramural Research, National Institute of Environmental Health Sciences, National Institutes of Health, Research Triangle Park, North Carolina, USA
17. Vanderbilt Epidemiology Center, Center for Human Genetics Research, Department of Obstetrics and Gynecology, Vanderbilt University Medical Center, Nashville, Tennessee, USA
18. Division of Epidemiology, Center for Human Genetics Research, Department of Medicine, Vanderbilt University Medical Center, Nashville, Tennessee, USA
19. Division of Epidemiology, Department of Medicine, Vanderbilt Epidemiology Center, Institute for Medicine and Public Health, Vanderbilt University Medical Center, Vanderbilt-Ingram Cancer Center, Nashville, Tennessee, USA
20. Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, Maryland, USA

21. Department of Epidemiology, University of Michigan School of Public Health, Ann Arbor, Michigan, USA
22. Department of Preventive Medicine, Keck School of Medicine, University of Southern California Norris Comprehensive Cancer Center, Los Angeles, California, USA
23. Department of Health Research and Policy, Stanford University School of Medicine, Stanford, California, USA
24. Department of Biomedical Data Science, Stanford University School of Medicine, Stanford, California, USA
25. Department of Population Health Science and Policy, Icahn School of Medicine at Mount Sinai, New York, New York, USA
26. Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, New York, USA
27. Division of Gynecologic Oncology, Department of Obstetrics, Gynecology and Reproductive Sciences, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA
28. Department of Epidemiology, University of Pittsburgh Graduate School of Public Health, Pittsburgh, Pennsylvania, USA
29. Ovarian Cancer Center of Excellence, Womens Cancer Research Program, Magee-Womens Research Institute and University of Pittsburgh Cancer Institute, Pittsburgh, Pennsylvania, USA
30. The University of Texas School of Public Health, Houston, Texas, USA
31. Department of Cancer Prevention and Control, Roswell Park Cancer Institute, Buffalo, NY
32. Program in Epidemiology, Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, Washington, USA
33. Department of Epidemiology, University of Washington, Seattle, Washington, USA
34. Huntsman Cancer Institute, Department of Population Health Sciences, University of Utah, Salt Lake City, Utah, USA
35. Department of Cancer Epidemiology, Moffitt Cancer Center, Tampa, Florida, USA
36. Gynecology Service, Department of Surgery, Memorial Sloan Kettering Cancer Center, New York, New York, USA
37. Gynecologic Oncology, Laura and Isaac Pearlmuter Cancer Center, New York University Langone Medical Center, New York, New York, USA
38. University of Southern California Norris Comprehensive Cancer Center, Los Angeles, California, USA
39. University of Hawaii Cancer Center, Honolulu, Hawaii, USA
40. Cancer Epidemiology Program, University of Hawaii Cancer Center, Hawaii, USA
41. Women's Cancer Program at the Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, California, USA

42. Department of Women's Cancer, Institute for Women's Health, University College London, London, UK
43. School of Women's and Children's Health, University of New South Wales, New South Wales, Australia
44. The Kinghorn Cancer Centre, Garvan Institute of Medical Research, Darlinghurst, New South Wales, Australia
45. Center for Cancer Prevention and Translational Genomics, Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, California, USA
46. Department of Biomedical Sciences, Cedars-Sinai Medical Center, Los Angeles, California, USA
47. Obstetrics and Gynecology Epidemiology Center, Brigham and Women's Hospital, Boston, Massachusetts, USA
48. Harvard T. H. Chan School of Public Health, Boston, Massachusetts, USA
49. Department of Health Science Research, Division of Epidemiology, Mayo Clinic, Rochester, Minnesota, USA
50. Department of Health Science Research, Division of Biomedical Statistics and Informatics, Mayo Clinic, Rochester, Minnesota, USA
51. Departments of Medicine and Pharmacology, Mayo Clinic, Rochester, Minnesota, USA
52. Cancer Pathology & Prevention, Division of Cancer Prevention and Population Sciences, Roswell Park Cancer Institute, Buffalo, New York, USA
53. Department of Gynecological Oncology, Roswell Park Cancer Institute, Buffalo, New York, USA
54. Center For Immunotherapy, Roswell Park Cancer Institute, Buffalo, New York, USA
55. Division of Gynecologic Oncology, Princess Margaret Hospital, University Health Network, Toronto, Ontario, Canada
56. Cancer Prevention and Control, Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, California, USA
57. Community and Population Health Research Institute, Department of Biomedical Sciences, Cedars-Sinai Medical Center, Los Angeles, California, USA
58. Department of Obstetrics and Gynecology, John A. Burns School of Medicine, University of Hawaii, Honolulu, Hawaii, USA
59. Channing Division of Network Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts, USA
60. Biostatistics, Sanofi Genzyme, Boston, Massachusetts, USA
61. Vesalius Research Center, VIB, Leuven, Belgium
62. Laboratory for Translational Genetics, Department of Oncology, University of Leuven, Belgium
63. Division of Gynecologic Oncology, Department of Obstetrics and Gynaecology and Leuven Cancer Institute, University Hospitals Leuven, Leuven, Belgium

64. Department of Epidemiology, Director of Genetic Epidemiology Research Institute, Center for Cancer Genetics Research & Prevention, School of Medicine, University of California Irvine, Irvine, California, USA
65. Department of Epidemiology, University of California Irvine, Irvine, California, USA
66. Cancer ResearchUK Cambridge Institute, University of Cambridge, Cambridge, UK
67. Department of Gynecology and Obstetrics, Haukeland University Hospital, Bergen, Norway
68. Centre for Cancer Biomarkers, Department of Clinical Science, University of Bergen, Bergen, Norway
69. Radboud University Medical Center, Radboud Institute for Health Sciences, Nijmegen, Netherlands
70. Radboud University Medical Center, Radboud Institute for Molecular Life sciences, Department of Gynaecology, Nijmegen, The Netherlands
71. Department of Obstetrics & Gynecology, Oregon Health & Science University, Portland, Oregon, USA
72. Knight Cancer Institute, Oregon Health & Science University, Portland, Oregon, USA
73. Division of Epidemiology and Biostatistics, Department of Internal Medicine, University of New Mexico, Albuquerque, New Mexico, USA
74. Cancer Control Research, British Columbia Cancer Agency, Vancouver, British Columbia, Canada
75. Canada's Michael Smith Genome Sciences Centre, British Columbia Cancer Agency, Vancouver, British Columbia, Canada
76. Department of Biomedical Physiology and Kinesiology, Simon Fraser University, Burnaby, British Columbia, Canada
77. Hollings Cancer Center and Department of Public Health Sciences, Medical University of South Carolina, Charleston, South Carolina, USA
78. Department of Oncology, Department of Public Health and Primary Care, University of Cambridge, Strangeways Research laboratory, Cambridge, UK
79. Department of Oncology, University of Cambridge, Strangeways Research Laboratory, Cambridge, UK
80. Cancer Genetics Laboratory, Research Division, Peter MacCallum Cancer Centre, East Melbourne, Victoria, Australia
81. Department of Pathology, University of Melbourne, Parkville, Victoria, Australia
82. Faculty of Medicine, University of Southampton, Southampton, UK
83. Centre for Cancer Research, The Westmead Institute for Medical Research, The University of Sydney, Sydney, New South Wales, Australia
84. Department of Gynaecological Oncology, Westmead Hospital, Sydney, New South Wales, Australia

## FINANCIAL SUPPORT

This work was supported by the National Institutes of Health (Genetic Associations and Mechanisms in Oncology (GAME-ON) (U19-CA148112); R01-CA114343 and R01-CA114343-S1 for the genotyping, bioinformatics, and biostatistical analysis for MAY, NCO, TBO, and TOR; P30-CA15083 for the Mayo Clinic Genotyping Shared Resource; R01-CA142081 for AACES; R01-CA112523 and R01-CA87538 for DOV; R01-CA58598, N01-CN-55424 and N01-PC-67001 for HAW; K07-CA080668, R01-CA95023, P50-CA159981 and MO1-RR000056 for HOP; R01-CA122443, P30-CA15083, and P50-CA136393 for MAY; CA54281, CA164973, and CA63464 for MEC; R01-CA76016 for NCO; R01-CA54419 and P50-CA105009 for NEC; UM1-CA186107, P01-CA87969, R01-CA49449, UM1-CA176726, and R01-CA67262 for NHS; R01CA160669-01A1 for OVA; P50-CA159981 and R01-CA126841 for RPC; Z01-ES044005 and Z01-ES049033 for SIS and the Intramural Research Program of the NIH, National Institute of Environmental Health Sciences for SIS; U01-CA71966 and U01-CA69417 for STA; R01-CA058860 for UCI; P01-CA17054, P30-CA14089, R01-CA61132, N01-PC67010, R03-CA113148, R03-CA115195, and N01-CN025403 for USC; the Intramural Research Program of the National Cancer Institute for PLC); Nationaal Kankerplan (BEL), a Vanderbilt CTSA grant from the National Institutes of Health/National Center for Advancing Translational Sciences (ULTR000445 for BVU); the National Institutes of Health Research Cambridge Biomedical Research Centre and Cancer Research UK (for RMH; C490/A10119 and C490/A10124 for SEA; and Cambridge Cancer Centre for CAM); the Ovarian Cancer Research Fund (DKE); the Department of Defense (DAMD17-02-1-0669 for HOP; DAMD17-02-1-0666 for NCO; W81XWH-10-1-02802 for NEC); the American Cancer Society Early Detection Professorship (SIOP-06-258-01-COUN) and the National Center for Advancing Translational Sciences (UL1TR000124) for LAX; the Mayo Foundation, the Minnesota Ovarian Cancer Alliance and the Katherine B. Andersen Foundation (for MAY); the Moffitt Cancer Center (P30-CA076292), Merck Pharmaceuticals, the state of Florida, Hillsborough County and the city of Tampa (for MOF); Helse Vest, the Norwegian Cancer Society, and the Research Council of Norway (for NOR); Radboud University Medical Centre (for NTH); the OHSU Foundation (for ORE); the Canadian Institutes of Health (MOP-86727 for OVA); Royal Marsden Hospital (for RMH); the UK National Institute for Health Research Biomedical Research Centres at the

University of Cambridge (SEA); the Lon V Smith Foundation (LVS-39420 for UCI); Princess Margaret Cancer Centre Foundation-Bridge for the Cure (for UHN); The Eve Appeal (The Oak Foundation) and the National Institute for Health Research University College London Hospitals Biomedical Research Centre (for UKO); the California Cancer Research Program (00-01389V-20170, 2110200 for USC); the National Health and Medical Research Council of Australia (310670 and 628903 for WMH); Cancer Institute NSW (12/RIG/1-17 and 15/RIG/1-16 for WMH); and North Carolina Central University/University of North Carolina at Chapel Hill, U54 Cooperative Agreement Grant, National Cancer Institute (1U54CA156735-01 for D.J. Grant, T.O. Keku, C. Hoyo, J.M. Schildkraut).

**CORRESPONDING AUTHOR:**

Delores J. Grant, Ph.D.

[dgrant@ncsu.edu](mailto:dgrant@ncsu.edu)

1801 Fayetteville Street

Department of Biological and Biomedical Sciences

North Carolina Central University

Durham, NC 27707

WORD COUNT: 2,885

NUMBER OF TABLES: 2

NUMBER OF SUPPLEMENTAL TABLES: 8

NUMBER OF FIGURES: 0

CONFLICT OF INTEREST: The authors have no conflicts of interest to disclose.

**NOVELTY & IMPACT STATEMENTS**

Women of African ancestry present with more advanced ovarian tumors and have lower five-year survival than Caucasian women. Progress in determining risk factors contributing to those differences in ovarian cancer development and mortality is hampered by the dearth of research. This study shows novel associations of genetic variants from *EGFR* and *UGT2A1/2* gene regions with ovarian cancer in African Americans that is suggestive of proliferation and tobacco metabolism pathway alterations that contribute to health disparities.

## ACKNOWLEDGMENTS

The **BEL** study would like to thank Gilian Peuteman, Thomas Van Brussel, Annick Van den Broeck and Joke De Roover for technical assistance. The **CAM** study was supported by Cancer Research UK, the University of Cambridge, and the National Institute for Health Research Cambridge Biomedical Research Centre. The **MOF** study would like to thank the Total Cancer Care™ Protocol and the Collaborative Data Services and Tissue Core Facilities at the H. Lee Moffitt Cancer Center & Research Institute, an NCI designated Comprehensive Cancer Center, Merck Pharmaceuticals and the state of Florida (MOF). The **NHS/NHSII** studies thank the following state cancer registries for their help: AL, AZ, AR, CA, CO, CT, DE, FL, GA, ID, IL, IN, IA, KY, LA, ME, MD, MA, MI, NE, NH, NJ, NY, NC, ND, OH, OK, OR, PA, RI, SC, TN, TX, VA, WA, and WY. The **SEA** study thanks the SEARCH team, Craig Luccarini, Caroline Baynes, Don Conroy. The **UKO** study thanks I. Jacobs, M. Widschwendter, E. Wozniak, A. Ryan, J. Ford and N. Balogun for their contribution to the study. The **WMH** study thanks the Gynaecological Oncology Biobank at Westmead, a member of the Australasian Biospecimen Network-Oncology group.



DR. DELORES GRANT (Orcid ID : 0000-0001-8981-8930)

DR. JILL BARNHOLTZ-SLOAN (Orcid ID : 0000-0001-6190-9304)

DR. KIRSTEN MOYSICH (Orcid ID : 0000-0002-4678-2058)

DR. JENNIFER PERMUTH (Orcid ID : 0000-0002-4726-9264)

DR. DANIEL W CRAMER (Orcid ID : 0000-0002-8024-3066)

Article type : Original Research

Evaluation of vitamin D biosynthesis and pathway target genes reveals *UGT2A1/2* and *EGFR* polymorphisms associated with epithelial ovarian cancer in African American Women

RUNNING TITLE: Genetic association with variants in the vitamin D pathway

KEYWORDS: ovarian cancer, genetic association, vitamin D pathway, African ancestry risk

WORD COUNT: 2,885

NUMBER OF TABLES: 2

NUMBER OF SUPPLEMENTAL TABLES: 8

NUMBER OF FIGURES: 0

CONFLICT OF INTEREST: The authors have no conflicts of interest to disclose.

## NOVELTY & IMPACT STATEMENTS

Women of African ancestry present with more advanced ovarian tumors and have lower five-year survival than Caucasian women. Progress in determining risk factors contributing to those differences in ovarian cancer development and mortality is hampered by the dearth of research. This study shows novel associations of genetic variants from *EGFR* and *UGT2A1/2* gene regions with ovarian cancer in African Americans that is suggestive of proliferation and tobacco metabolism pathway alterations that contribute to health disparities.

## ABSTRACT

An association between genetic variants in the vitamin D receptor (*VDR*) gene and epithelial ovarian cancer (EOC) was previously reported in women of African ancestry (AA). We sought to examine associations in *VDR* and additional genes from vitamin D biosynthesis and pathway targets (*EGFR*, *UGT1A*, *UGT2A1/2*, *UGT2B*, *CYP3A4/5*, *CYP2R1*, *CYP27B1*, *CYP24A1*, *CYP11A1*, and *GC*). Genotyping was performed using the custom-designed 533,631 SNP Illumina OncoArray with imputation to the 1,000 Genomes Phase 3 v5 reference set in 755 EOC cases, including 537 high-grade serous (HGSOC), and 1,235 controls. All subjects are of African ancestry (AA). Logistic regression was performed to estimate odds ratios (OR) and 95% confidence intervals (CI). We further evaluated statistical significance of selected SNPs using the Bayesian False Discovery Probability (BFDP). A significant association with EOC was identified in the *UGT2A1/2* region for the SNP rs10017134 (per allele OR = 1.4, 95% CI = (1.2, 1.7)  $p=1.2 \times 10^{-6}$ , BFDP=0.02); and an association with HGSOC was identified in the *EGFR* region for the SNP rs114972508 (per allele OR = 2.3, 95% CI = (1.6, 3.4)  $p=1.6 \times 10^{-5}$ , BFDP=0.29) and in the *UGT2A1/2* region again for rs1017134 (per allele OR = 1.4, 95% CI = (1.2, 1.7)  $p=2.3 \times 10^{-5}$ , BFDP=0.23). Genetic variants in the *EGFR* and *UGT2A1/2* may increase susceptibility of EOC in AA women. Future studies to validate these

findings are warranted. Alterations in *EGFR* and *UGT2A1/2* could perturb enzyme efficacy, proliferation in ovaries, impact and mark susceptibility to EOC.

## INTRODUCTION

Women of African ancestry (AA) have the lowest incidence of ovarian cancer worldwide, but they tend to present with more advanced tumors and have lower five-year survival (35%) compared to women of European descent (47%) in nearly every cancer subtype<sup>1,2</sup>. Compared to Caucasian women, there have been fewer published studies investigating the association between common risk factors, such as tubal ligation, use of hormonal contraceptives, obesity, body powder and dietary patterns, and ovarian cancer risk in AA<sup>1,3-9</sup>. Moreover, the investigation of genetic susceptibility to epithelial ovarian cancer (EOC) in AA has not been comprehensive. The limited assessment of genetic susceptibility among AA are in modest sized study populations of candidate genes including the repeat polymorphisms of the androgen receptor (*AR*), vitamin D receptor (*VDR*) and cellular transport genes, where an association with risk of ovarian cancer was observed<sup>10-12</sup>.

The vitamin D receptor mediates the regulation of a pleotropic cascade of physiological responses; including those involved in phase I and phase II detoxification and the epidermal growth factor receptor (*EGFR*) proliferation pathways in ovarian and other cancer cell lines; through *VDR*/DNA interactions and bioavailability of vitamin D<sup>13-17</sup>. A *VDR* variant, rs7305032, was associated with ovarian cancer in 125 cases and 155 controls of AA but other observations were limited because of small sample size<sup>11</sup>. Moreover, known genetic variations in the *VDR*/vitamin D biosynthesis and pathway target genes have been implicated in AA disease risk. Therefore an objective of this study was assess those variants in ovarian cancer in women of African ancestry in a large sample.

Using a candidate gene approach, SNPs were selected from genes involved in vitamin D biosynthesis and metabolism; and putative targets of *VDR* regulation. Genes of the vitamin D biosynthesis pathways included cytochrome P450s: *CYP2R1*,

*CYP27B1*, *CYP24A1*, *CYP11A1*, and group-specific component-Vitamin D-binding protein (*GC*) which collectively are responsible for the homeostatic control and bioavailability of vitamin D<sup>18-23</sup>. The candidate genes involved in vitamin D metabolic processes included *CYP3A4/5* and UDP-glucuronosyltransferase 1A (*UGT1A*) locus members responsible for glucuronidation and hydroxylation of the biologically active and circulatory forms of vitamin D. These genes are also inclusive of candidates regulated by vitamin D/VDR binding which included *CYP3A4/5*, *UGT1A* locus members, *EGFR* and UDP-glucuronosyltransferase 2 (*UGT2*) locus members; and associated, in part, with other cancers in AA individuals<sup>24-39</sup>. Thus, variants in *VDR* and additional genes from vitamin D biosynthesis and pathway targets are viable candidates to investigate the genetic underpinnings of ovarian cancer risk in women of African descent.

In this study, SNPs from 11 gene regions: *VDR*, *EGFR*, *UGT1A*, *UGT2A1/2*, *UGT2B*, *CYP3A4/5*, *CYP2R1*, *CYP27B1*, *CYP24A1*, *CYP11A1*, and *GC*, were genotyped, imputed then assessed for risk of EOC and high grade serous ovarian cancer (HGSOC) in cases and controls of AA from the African American Cancer Epidemiology Study (AACES)<sup>40</sup> and the Ovarian Cancer Association Consortium (OCAC)<sup>41</sup>.

## **MATERIALS AND METHODS**

### Study Populations

The Genetic Associations and Mechanisms in Oncology (GAME-ON) project comprised 63, mostly, case-control studies from four continents (North America, Europe, Asia and Australia). Only 32 studies contributed subjects of African Ancestry, including AACES and studies in OCAC, and were included in the current analysis (Supplemental Table 1). AACES, previously described elsewhere<sup>40</sup>, is a multi-center population-based case-control study of newly diagnosed invasive EOC in African American women that enrolled study subjects between 2010 and 2015. Established in 2005, OCAC is an international consortium focused on genetic association and pooled risk factor analyses. The current analyses included 1,990 samples: 1,235 controls and 755 invasive EOC cases who passed quality control filters, all of whom were AA. The majority of the EOC cases were HGSOC (n= 537, 71%), followed by 49 mucinous cases (7%), 28

endometrioid cases (4%), 23 clear cell cases (3%), 12 mixed histology (2%) and 53 other (7%). All subjects included in this analysis provided written informed consent as well as data and blood samples under ethically approved protocols.

### Genotyping, ancestry analysis and quality control

Genotyping of AA women from OCAC was completed using the custom-designed 533,631 SNP array, the Illumina OncoArray. Sample level quality control included restriction to females, filter on call rate > 95%, heterozygosity (either too big or too small), removal of ineligible samples, and relationship inference to check for unexpected first degree relatives. SNP level quality control included filter on call rate > 95%, and Hardy-Weinberg Equilibrium p-value >  $1 \times 10^{-5}$ . After applying these procedures, 471,780 SNPs remained.

Intercontinental ancestry was calculated for the OCAC and AACES samples using the software package FastPop<sup>42</sup> that was developed specifically for the OncoArray Consortium. Only the African ancestry samples defined as having >50% AA were used for the present analyses reported here. Seventy-seven cases and 120 controls were omitted due to African ancestry <50% and one gender mismatch. Principal components computed using FastPop were further used to adjust for population structure in our analyses.

### Genotype Imputation Analysis

Using the genotyped SNPs that passed quality control, haplotypes were phased using SHAPEIT v2 followed by imputation to the 1,000 Genomes Phase 3 v5 reference set using Minimac3.

### Gene Region and SNP Selection

Eleven gene regions were defined based on human genome build 37. SNPs within the selected regions were filtered on imputation quality score (minimac imputation R-squared) > 0.5 for imputed SNPs, or Hardy-Weinberg Equilibrium p-value >  $1.0 \times 10^{-5}$  for genotyped SNPs. Quantile-quantile plots on the EOC and HGSOC dataset (Manichaikul et al., unpublished) have lambdas of 1.01 each within normal range<sup>43</sup>.

The imputation quality scores for significant SNPs are provided. We further applied filter on effective heterozygosity count  $> 30$ . After applying filters, the following numbers of SNPs were remaining in each of the selected gene regions for EOC: 288 in *VDR*, 433 in *UGT2A1/2*, 6302 in *UGT2B*, 919 in *UGT1A*, 963 in *EGFR*, 17 in *CYP2R1*, 4 in *CYP27B1*, 113 in *CYP24A1*, 90 in *CYP11A1*, 411 in *CYP3A4/5* and 296 in *GC*. For selected regions for HGSOC analysis, the number of SNPs were: 234 in *VDR*, 413 in *UGT2A1/2*, 5674 in *UGT2B*, 833 in *UGT1A*, 824 in *EGFR*, 15 in *CYP2R1*, 4 in *CYP27B1*, 106 in *CYP24A1*, 82 in *CYP11A1*, 375 in *CYP3A4/5* and 282 in *GC*.

### Statistical Analysis

Genetic association testing was carried out with adjustment for two principal components (PCs) of ancestry using a logistic regression model that accounts for genotype uncertainty under a score test as implemented in SNPTEST v2.5.2 to estimate odds ratios (OR) and 95 % confidence intervals (CI). For each gene region, we applied a gene-specific Bonferroni-threshold for statistical significance defined as  $0.05 / \text{number of SNPs examined for that gene}$ . We further assessed the main results with an alternative to the Bonferroni threshold using the Bayesian False Discovery Probability (BFDP) which provides the posterior probability of a false discovery based on a given prior probability of non-null association at a given SNP<sup>44</sup>. For this study we specified a prior probability of association at each SNP under investigation based on the total number of SNPs within each candidate gene region as  $0.5 \times 1 / (N_{\text{SNP}} / 3)$  where  $N_{\text{SNP}}$  represents the number of SNPs in the given candidate gene region. We considered  $N_{\text{SNP}}/3$  to be an approximation of the effective number of independent SNPs within in each gene region, taking into account the fact that many SNPs will be correlated due to linkage disequilibrium. Accordingly, the specified prior indicates a 50% chance of true discovery within each gene region, with the prior probability of non-null association distributed randomly among all SNPs within the region. In order to avoid spurious positive associations, we applied a filter on effective heterozygosity count (HC)  $> 30$  in each of cases and controls. Here, HC is defined as  $N \times \text{MAF} \times (1 - \text{MAF})$  for each SNP,  $N$  represents the sample size (either the number of cases or the number of controls), and MAF represents the SNP minor allele frequency. Based on 755 EOC cases and 537

HGSOC cases, respectively, applying this filter equates to applying a SNP MAF filter of 4.2% and 6% in analysis of EOC and HGSOC, respectively. Statistical power calculations for AA study participants and Caucasians are included in Supplemental Tables 2 and 3.

## RESULTS

### ***VDR Pathway Gene Regions and risk of EOC***

SNPs from eleven gene regions (*CYP3A4/5*, *CYP2R1*, *CYP27B1*, *CYP24A1*, *CYP11A1*, *EGFR*, *GC*, *UGT1A*, *UGT2A1/2*, *UGT2B* and *VDR*) from VDR biosynthesis and pathway targets were assessed for association with EOC (Supplemental Table 4). The top associations are reported in table 1. Individuals carrying the major allele of SNP rs10017134 of the *UGT2A1/2* gene region had an increased odds of EOC when corrected for multiple comparisons (OR=1.4, 95% CI=(1.2, 1.7),  $p=1.2 \times 10^{-6}$ ). The BFDP for rs10017134 of 0.020 corresponds to 98% posterior probability of non-null association for this SNP. Significant associations with EOC were also observed for *UGT2A1/2* SNPs, rs2288741 and rs11939884. The variants are found in both *UGT2A1* and *UGT2A2* as the genes share common exons 2 through 6<sup>45</sup>. Supplemental Table 5 summarizes other notable ( $p < 0.01$ ) SNP associations with EOC in the OncoArray analysis.

### ***VDR Pathway Gene Regions and risk of HGSOC***

SNPs from the ten gene regions from VDR biosynthesis and pathway targets were assessed for association with HGSOC (Supplemental Table 4). The top associations are reported in table 2. Individuals carrying the minor allele of *EGFR* SNP rs114972508 had more than two-fold increased odds of HGSOC (OR=2.3, 95% CI=(1.2, 3.4),  $p=1.6 \times 10^{-5}$ ) (Table 2). The posterior BFDP is 29% for SNP rs114972508 corresponds to 71% posterior probability of non-null association. SNP rs10017134 of the *UGT2A1/2* gene region also showed association with HGSOC (OR=1.4, 95% CI=(1.2, 1.7),  $p=2.3 \times 10^{-5}$ ) (Table 2). The posterior BFDP is 22.8%. Supplemental

Table 6 summarizes other notable ( $p < 0.01$ ) SNP associations with HGSOE in the OncoArray analysis.

## DISCUSSION

Few studies have investigated the genetic susceptibility for ovarian cancer among women of African descent. The assessment of candidate SNPs from chromosomal regions that contain genes regulated by VDR activity provides some evidence of association with EOC risk. The notable findings from this analysis show, for the first time, that risk assessments of variants in the *UGT2A1/2* and *EGFR* gene regions are suggestive of associations with EOC and HGSOE. The results also demonstrate evidence of associations for other SNPs from the candidate gene regions with EOC and HGSOE. Although the candidate SNPs are located in intronic regions there is ample evidence that many gene regulatory regions are present in those regions including encoded microRNAs, alternate splice sites, and cis-regulatory modules and transcription factors binding sites<sup>46-48</sup>. In addition, recent studies have shown using targeted RNAseq analysis that there are numerous splice variants of the *UGT* genes<sup>49</sup>.

The *UGT2A1* and *2A2* genes are distinguished by unique first exons joined to common exons 2–6 and are located downstream of *UGT2B4* on chromosome 4<sup>45</sup>. *UGT2A* transcripts have been detected in several extrahepatic tissues such as the lung, trachea, larynx, intestine, pancreas and kidney<sup>50</sup>. *UGT2A1* is an extrahepatic enzyme that is expressed mainly in the nasal epithelium, catalyzing the glucuronidation of testosterone and epitestosterone at considerable rates and has similar kinetics as the *UGT2B* gene family members<sup>51</sup>. There are reports that this enzyme also has activity toward estrogen metabolites epiestradiol and  $\beta$ -estradiol<sup>52</sup>. *UGT2A1* has exhibited highest expression in the lung, followed by trachea, tonsil, larynx, colon, olfactory<sup>53</sup>. *UGT2A2* mRNA expression was reported in fetal and adult nasal mucosa tissues<sup>54</sup>. However, unlike *UGT2A1*, other expression analyses suggested that wild-type *UGT2A2* had the highest expression in the breast, followed by trachea, larynx, and kidney<sup>55</sup>.

Neither the *UGT2A1* gene, nor *UGT2A2* expression have been examined in ovarian tissue. However, VDR ChIPseq peak locations have been identified 430 kb downstream of the *UGT2A1/2* locus in experiments with THP-1 cells treated with



1 $\alpha$ ,25(OH)<sub>2</sub>D<sub>3</sub>, the biologically active form of the vitamin D hormone, suggestive of a regulatory role for vitamin D<sup>56</sup>. Splice variants found in *UGT2A1/2* that are highly conserved among both *UGT1A* and *UGT2* gene families have been implicated in altered glucuronidation activity against tobacco carcinogenesis<sup>49, 53, 55, 57</sup>. Two of the *UGT2A1* SNPs associated with EOC and HGSOE in this study are intron variants (rs10017134 and rs2288741) while the third (rs11939884) is a 3' UTR variant. It is probable that these variants alter enzyme function in target tissues including ovarian and/or alter risk in AA smokers. Of note, cigarette smoking has been found to be associated with the risk of mucinous EOC, but not HGSOE among Caucasian women<sup>58</sup>. Moreover, providing some plausibility for the mechanism of the observed SNP association, a recent report suggests that cigarette smoking may be associated with serous EOC among African American women although a dose-response relationship was not observed<sup>59</sup>. The association of genes from the *UGT* superfamily with ovarian cancer in AA is consistent with significant associations observed for Caucasian women for *UGT1A*<sup>12</sup>. However in that study, no association was observed for AA samples which had a MAF of 0.42 for the risk allele while Caucasians had MAF of 0.07. Some but not all MAFs for the relationships observed in this study differ by race so it is unlikely to explain racial differences in race.

The *EGFR* gene product has been a chemotherapeutic target for EOC since overexpression has been linked to poor prognosis in ovarian cancer patients<sup>60-62</sup>. The signaling pathway for EGFR is mediated by ligands including the epidermal growth factor in the regulation of cell proliferation, differentiation and apoptosis in normal cells. Research into the mechanisms of EGFR overexpression has focused on mutations and amplifications in the coding region of the gene containing the receptor tyrosine kinase domain<sup>63</sup>. However, few studies on SNP variants in this region have been linked to EOC or other ovarian cancer histologic subtypes<sup>61, 63</sup>. *EGFR* SNP rs114972508 is located in intron 1 of the *EGFR* gene. The location of the SNP is approximately 70 kb upstream of a *VDR* binding site also within *EGFR* intron 1 that has been shown experimentally to down regulate *EGFR* expression and proliferative function<sup>15</sup>. Perhaps changes in the intron sequences may impact *EGFR* function and subsequently be as critical to cellular homeostasis as the receptor tyrosine function that has been

extensively researched. Thus, *EGFR* SNPs could be abrogating vitamin D hormone regulation of ovarian cell proliferation and increasing susceptibility for the development of HGSOC in AA women.

Although we were unable to confirm the association between previously identified *VDR* variants and risk of EOC, a recent case-control study of women of European ancestry (10,065 cases, 21,654 controls) showed that SNPs associated with decreased circulating 25-hydroxyvitamin D were associated with ovarian cancer and HGSOC<sup>64</sup> while another study showed that AA women exposed to increased sunlight had a decreased risk for ovarian cancer<sup>8</sup>. These observations suggest that other mechanisms affecting vitamin D hormone activity independent of the *VDR* may be important in ovarian cancer etiology.

The main observations in the current study result from imputations of genotyped SNPs but independent of *VDR* variant association with EOC and HGSOC. The *VDR* SNPs previously observed to be associated with the risk of EOC<sup>11</sup>, including rs7975232 and rs7305032, were not associated with risk of EOC in the current study (Supplemental table 7). A look up of the significant study SNPs in archived OCAC data on Caucasians shows no significant associations for the *UGT2A1/2* SNPs. Data on the *EGFR* SNP was not available (Supplemental Table 8). Other *VDR* SNPs showed nominal (non-Bonferroni corrected) associations with EOC but not with HGSOC (Supplemental Table 7). Although the largest study to date of genetic association with EOC in AA, the modest sample size remains a limitation of the current study and therefore some of the nominal SNP associations may be a result of inadequate power. The analyses are underpowered for discovery analysis across the selected gene regions and important associations may have been missed, nonetheless, we still found significant associations with EOC and HGSOC. Several suggestive and nominal SNP associations (outside of Bonferroni significance) may provide some insight and consideration for future experimental studies to further explore the relevance of vitamin D biosynthesis and pathway target genes. Larger studies of AA are warranted to clarify these findings.

In summary, this study reports, for the first time, an association between *EGFR* and *UGT2A1/2* variants with ovarian cancer risk in AA women. These gene variants

could perturb cell proliferation and enzyme efficacy in ovaries and impact susceptibility to ovarian cancer by altering growth and intercellular hormone metabolism. Future studies are needed to validate the associations of the imputed SNPs and to determine their impact on cancer development. Currently, there are no published reports of population studies of *UGT2A1/2* polymorphisms in Europeans or other racially distinct groups in larger sample sizes than this AA study that would allow intricate gene-environment analysis. At this present time, there is only speculative evidence that *UGT2B* gene region variants may be associated with differences in nicotine metabolism across African American, Native Hawaiian, Caucasian, Latino, and Japanese American smokers<sup>65, 66</sup>. Analyses of the *UGT2A1/2* variants across populations may reveal differential risk to ovarian disease. In addition, expression and functional analysis in ovarian tissue needs to be accomplished to elucidate the impact on tissue homeostasis. In spite of the limitations of this study, these results provide new insight into proliferative and hormone target pathways that may represent important opportunities for the development of chemotherapeutic targets and intervention strategies.

Author Manuscript

## REFERENCES

1. Chornokur G, Amankwah EK, Schildkraut JM, Phelan CM. Global ovarian cancer health disparities. *Gynecologic oncology* 2013;**129**: 258-64.
2. Torre LA, Trabert B, DeSantis CE, Miller KD, Samimi G, Runowicz CD, Gaudet MM, Jemal A, Siegel RL. Ovarian cancer statistics, 2018. *CA: a cancer journal for clinicians* 2018.
3. Moorman PG, Palmieri RT, Akushevich L, Berchuck A, Schildkraut JM. Ovarian cancer risk factors in African-American and white women. *American journal of epidemiology* 2009;**170**: 598-606.
4. Schildkraut JM, Calvocoressi L, Wang F, Wrensch M, Bondy ML, Wiemels JL, Claus EB. Endogenous and exogenous hormone exposure and the risk of meningioma in men. *Journal of neurosurgery* 2014;**120**: 820-6.
5. Schildkraut JM, Abbott SE, Alberg AJ, Bandera EV, Barnholtz-Sloan JS, Bondy ML, Cote ML, Funkhouser E, Peres LC, Peters ES, Schwartz AG, Terry P, Crankshaw S, Camacho F, Wang F, Moorman PG. Association between Body Powder Use and Ovarian Cancer: the African American Cancer Epidemiology Study (AACES). *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology* 2016.

6. Bandera EV, Qin B, Moorman PG, Alberg AJ, Barnholtz-Sloan JS, Bondy M, Cote ML, Funkhouser E, Peters ES, Schwartz AG, Terry P, Schildkraut JM. Obesity, weight gain, and ovarian cancer risk in African American women. *International journal of cancer* 2016;**139**: 593-600.
7. Erondy CO, Alberg AJ, Bandera EV, Barnholtz-Sloan J, Bondy M, Cote ML, Funkhouser E, Peters E, Schwartz AG, Terry PD, Wallace K, Akushevich L, Wang F, Crankshaw S, Berchuck A, Schildkraut JM, Moorman PG. The Association Between Body Mass Index and Presenting Symptoms in African American Women with Ovarian Cancer. *Journal of women's health* 2016;**25**: 571-8.
8. Qin B, Moorman PG, Alberg AJ, Barnholtz-Sloan JS, Bondy M, Cote ML, Funkhouser E, Peters ES, Schwartz AG, Terry P, Schildkraut JM, Bandera EV. Dairy, calcium, vitamin D and ovarian cancer risk in African-American women. *British journal of cancer* 2016;**115**: 1122-30.
9. Peres LC, Bandera EV, Qin B, Guertin KA, Shivappa N, Hebert JR, Abbott SE, Alberg AJ, Barnholtz-Sloan J, Bondy M, Cote ML, Funkhouser E, Moorman PG, Peters ES, Schwartz AG, Terry PD, Camacho F, Wang F, Schildkraut JM. Dietary inflammatory index and risk of epithelial ovarian cancer in African American women. *International journal of cancer* 2017;**140**: 535-43.
10. Schildkraut JM, Murphy SK, Palmieri RT, Iversen E, Moorman PG, Huang Z, Halabi S, Calingaert B, Gusberg A, Marks JR, Berchuck A. Trinucleotide repeat polymorphisms in the androgen receptor gene and risk of ovarian cancer. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology* 2007;**16**: 473-80.

11. Grant DJ, Hoyo C, Akushevich L, Iversen ES, Whitaker R, Marks J, Berchuck A, Schildkraut JM. Vitamin D receptor (VDR) polymorphisms and risk of ovarian cancer in Caucasian and African American women. *Gynecologic oncology* 2013;**129**: 173-8.

12. Chornokur G, Lin HY, Tyrer JP, Lawrenson K, Dennis J, Amankwah EK, Qu X, Tsai YY, Jim HS, Chen Z, Chen AY, Permuth-Wey J, Aben KK, Anton-Culver H, Antonenkova N, Bruinsma F, Bandera EV, Bean YT, Beckmann MW, Bisogna M, Bjorge L, Bogdanova N, Brinton LA, Brooks-Wilson A, Bunker CH, Butzow R, Campbell IG, Carty K, Chang-Claude J, Cook LS, Cramer DW, Cunningham JM, Cybulski C, Dansonka-Mieszkowska A, du Bois A, Despierre E, Dicks E, Doherty JA, Dork T, Durst M, Easton DF, Eccles DM, Edwards RP, Ekici AB, Fasching PA, Fridley BL, Gao YT, Gentry-Maharaj A, Giles GG, Glasspool R, Goodman MT, Gronwald J, Harrington P, Harter P, Hein A, Heitz F, Hildebrandt MA, Hillemanns P, Hogdall CK, Hogdall E, Hosono S, Jakubowska A, Jensen A, Ji BT, Karlan BY, Kelemen LE, Kellar M, Kiemeny LA, Krakstad C, Kjaer SK, Kupryjanczyk J, Lambrechts D, Lambrechts S, Le ND, Lee AW, Lele S, Leminen A, Lester J, Levine DA, Liang D, Lim BK, Lissowska J, Lu K, Lubinski J, Lundvall L, Massuger LF, Matsuo K, McGuire V, McLaughlin JR, McNeish I, Menon U, Milne RL, Modugno F, Moysich KB, Ness RB, Nevanlinna H, Eilber U, Odunsi K, Olson SH, Orlow I, Orsulic S, Weber RP, Paul J, Pearce CL, Pejovic T, Pelttari LM, Pike MC, Poole EM, Risch HA, Rosen B, Rossing MA, Rothstein JH, Rudolph A, Runnebaum IB, Rzepecka IK, Salvesen HB, Schernhammer E, Schwaab I, Shu XO, Shvetsov YB, Siddiqui N, Sieh W, Song H, Southey MC, Spiewankiewicz B, Sucheston L, Teo SH, Terry KL, Thompson PJ, Thomsen L, Tangen IL, Tworoger SS, van Altena AM, Vierkant RA, Vergote I, Walsh CS, Wang-Gohrke S, Wentzensen N, Whittemore AS, Wicklund KG, Wilkens LR, Wu AH, Wu X, Woo YL, Yang H, Zheng W, Ziogas A, Hasmad HN, Berchuck A, Georgia C-T, group Am, Iversen ES, Schildkraut JM, Ramus SJ, Goode EL, Monteiro AN, Gayther SA,

Narod SA, Pharoah PD, Sellers TA, Phelan CM. Common Genetic Variation In Cellular Transport Genes and Epithelial Ovarian Cancer (EOC) Risk. *PLoS one* 2015;**10**: e0128106.

13. Kasai N, Sakaki T, Shinkyo R, Ikushiro S, Iyanagi T, Ohta M, Inouye K. Metabolism of 26,26,26,27,27,27-F6-1 alpha,23S,25-trihydroxyvitamin D3 by human UDP-glucuronosyltransferase 1A3. *Drug metabolism and disposition: the biological fate of chemicals* 2005;**33**: 102-7.

14. Kaeding J, Belanger J, Caron P, Verreault M, Belanger A, Barbier O. Calcitrol (1alpha,25-dihydroxyvitamin D3) inhibits androgen glucuronidation in prostate cancer cells. *Molecular cancer therapeutics* 2008;**7**: 380-90.

15. Shen Z, Zhang X, Tang J, Kasiappan R, Jinwal U, Li P, Hann S, Nicosia SV, Wu J, Zhang X, Bai W. The coupling of epidermal growth factor receptor down regulation by 1alpha,25-dihydroxyvitamin D3 to the hormone-induced cell cycle arrest at the G1-S checkpoint in ovarian cancer cells. *Molecular and cellular endocrinology* 2011;**338**: 58-67.

16. Nylen H, Bjorkhem-Bergman L, Ekstrom L, Roh HK, Bertilsson L, Eliasson E, Lindh JD, Diczfalusy U. Plasma levels of 25-hydroxyvitamin D3 and in vivo markers of cytochrome P450 3A activity in Swedes and Koreans: effects of a genetic polymorphism and oral contraceptives. *Basic & clinical pharmacology & toxicology* 2014;**115**: 366-71.

17. Huang M, Wang HM, Guo Y, Ping J, Chen M, Xu D, Wang H. Single nucleotide polymorphism of CYP3A4 intron 2 and its influence on CYP3A4 mRNA expression and liver enzymatic activity in human liver. *Journal of Huazhong University of Science and Technology Medical sciences = Hua zhong ke ji da xue xue bao Yi xue Ying De wen ban = Huazhong keji daxue xuebao Yixue Yingdewen ban* 2015;**35**: 502-7.

18. Carpenter TO. CYP24A1 loss of function: Clinical phenotype of monoallelic and biallelic mutations. *The Journal of steroid biochemistry and molecular biology* 2017.
19. Thacher TD, Levine MA. CYP2R1 mutations causing vitamin D-deficiency rickets. *The Journal of steroid biochemistry and molecular biology* 2016.
20. Rezaie Z, Taheri M, Kohan L, Sayad A. Down-regulation of CYP27B1 gene expression in Iranian patients with relapsing-remitting multiple sclerosis. *Human antibodies* 2016;**24**: 71-6.
21. Mateos-Munoz B, Garcia-Martin E, Torrejon MJ, Devesa-Medina MJ, Esguevillas G, Cardenas MC, Fernandez C, Carballo M, Agundez JA, Ladero JM. GC Gene Polymorphism and Unbound Serum Retinol-Binding Protein 4 Are Related to the Risk of Insulin Resistance in Patients With Chronic Hepatitis C: A Prospective Cross-Sectional Study. *Medicine* 2016;**95**: e3019.
22. Miller WL. Genetic disorders of Vitamin D biosynthesis and degradation. *The Journal of steroid biochemistry and molecular biology* 2017;**165**: 101-8.
23. Pike JW, Meyer MB, Benkusky NA, Lee SM, St John H, Carlson A, Onal M, Shamsuzzaman S. Genomic Determinants of Vitamin D-Regulated Gene Expression. *Vitamins and hormones* 2016;**100**: 21-44.
24. Cheng CY, Slominski AT, Tuckey RC. Hydroxylation of 20-hydroxyvitamin D3 by human CYP3A4. *The Journal of steroid biochemistry and molecular biology* 2016;**159**: 131-41.



25. Morales E, Sanchez-Solis M, Garcia-Marcos L. Vitamin D Metabolism Genes in Asthma and Atopy. *Mini reviews in medicinal chemistry* 2015;**15**: 913-26.
26. Mondul AM, Shui IM, Yu K, Weinstein SJ, Tsilidis KK, Joshi AD, Agudo A, Berg CD, Black A, Buring JE, Chasman DI, Gaudet MM, Haiman C, Hankinson SE, Henderson BE, Hoover RN, Hunter DJ, Khaw KT, Kuhn T, Kvaskoff M, Le Marchand L, Lindstrom S, McCullough ML, Overvad K, Peeters PH, Riboli E, Ridker PM, Stram DO, Sund M, Trichopoulos D, Tumino R, Weiderpass E, Willett W, Kraft P, Ziegler RG, Albanes D. Vitamin D-associated genetic variation and risk of breast cancer in the breast and prostate cancer cohort consortium (BPC3). *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology* 2015;**24**: 627-30.
27. Wang Z, Wong T, Hashizume T, Dickmann LZ, Scian M, Koszewski NJ, Goff JP, Horst RL, Chaudhry AS, Schuetz EG, Thummel KE. Human UGT1A4 and UGT1A3 conjugate 25-hydroxyvitamin D3: metabolite structure, kinetics, inducibility, and interindividual variability. *Endocrinology* 2014;**155**: 2052-63.
28. Shike M, Doane AS, Russo L, Cabal R, Reis-Filho JS, Gerald W, Cody H, Khanin R, Bromberg J, Norton L. The effects of soy supplementation on gene expression in breast cancer: a randomized placebo-controlled study. *Journal of the National Cancer Institute* 2014;**106**.
29. Maguire O, Pollock C, Martin P, Owen A, Smyth T, Doherty D, Campbell MJ, McClean S, Thompson P. Regulation of CYP3A4 and CYP3A5 expression and modulation of "intracrine" metabolism of androgens in prostate cells by liganded vitamin D receptor. *Molecular and cellular endocrinology* 2012;**364**: 54-64.

30. Barnholtz-Sloan JS, Raska P, Rebbeck TR, Millikan RC. Replication of GWAS "Hits" by Race for Breast and Prostate Cancers in European Americans and African Americans. *Frontiers in genetics* 2011;**2**: 37.
31. Khan AA, Dragt BS, Porte RJ, Groothuis GM. Regulation of VDR expression in rat and human intestine and liver--consequences for CYP3A expression. *Toxicology in vitro : an international journal published in association with BIBRA* 2010;**24**: 822-9.
32. Hashizume T, Xu Y, Mohutsky MA, Alberts J, Hadden C, Kalhorn TF, Isoherranen N, Shuhart MC, Thummel KE. Identification of human UDP-glucuronosyltransferases catalyzing hepatic 1alpha,25-dihydroxyvitamin D3 conjugation. *Biochemical pharmacology* 2008;**75**: 1240-50.
33. Matsubara T, Yoshinari K, Aoyama K, Sugawara M, Sekiya Y, Nagata K, Yamazoe Y. Role of vitamin D receptor in the lithocholic acid-mediated CYP3A induction in vitro and in vivo. *Drug metabolism and disposition: the biological fate of chemicals* 2008;**36**: 2058-63.
34. McCullough ML, Bostick RM, Mayo TL. Vitamin D gene pathway polymorphisms and risk of colorectal, breast, and prostate cancer. *Annual review of nutrition* 2009;**29**: 111-32.
35. Zeigler-Johnson CM, Walker AH, Mancke B, Spangler E, Jalloh M, McBride S, Deitz A, Malkowicz SB, Ofori-Adjei D, Gueye SM, Rebbeck TR. Ethnic differences in the frequency of prostate cancer susceptibility alleles at SRD5A2 and CYP3A4. *Human heredity* 2002;**54**: 13-21.

36. Shuch B, Mikhail M, Satagopan J, Lee P, Yee H, Chang C, Cordon-Cardo C, Taneja SS, Osman I. Racial disparity of epidermal growth factor receptor expression in prostate cancer. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2004;**22**: 4725-9.
37. Guillemette C, Millikan RC, Newman B, Housman DE. Genetic polymorphisms in uridine diphospho-glucuronosyltransferase 1A1 and association with breast cancer among African Americans. *Cancer research* 2000;**60**: 950-6.
38. Paris PL, Kupelian PA, Hall JM, Williams TL, Levin H, Klein EA, Casey G, Witte JS. Association between a CYP3A4 genetic variant and clinical presentation in African-American prostate cancer patients. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology* 1999;**8**: 901-5.
39. Grant DJ, Hoyo C, Oliver SD, Gerber L, Shuler K, Calloway E, Gaines AR, McPhail M, Livingston JN, Richardson RM, Schildkraut JM, Freedland SJ. Association of uridine diphosphate-glucuronosyltransferase 2B gene variants with serum glucuronide levels and prostate cancer risk. *Genetic testing and molecular biomarkers* 2013;**17**: 3-9.
40. Schildkraut JM, Iversen ES, Akushevich L, Whitaker R, Bentley RC, Berchuck A, Marks JR. Molecular signatures of epithelial ovarian cancer: analysis of associations with tumor characteristics and epidemiologic risk factors. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology* 2013;**22**: 1709-21.

41. Berchuck A, Schildkraut JM, Pearce CL, Chenevix-Trench G, Pharoah PD. Role of genetic polymorphisms in ovarian cancer susceptibility: development of an international ovarian cancer association consortium. *Advances in experimental medicine and biology* 2008;**622**: 53-67.
42. Li Y, Byun J, Cai G, Xiao X, Han Y, Cornelis O, Dinulos JE, Dennis J, Easton D, Gorlov I, Seldin MF, Amos CI. FastPop: a rapid principal component derived method to infer intercontinental ancestry using genetic data. *BMC bioinformatics* 2016;**17**: 122.
43. Lin P, Hartz SM, Zhang Z, Saccone SF, Wang J, Tischfield JA, Edenberg HJ, Kramer JR, A MG, Bierut LJ, Rice JP, Coga Collaborators Cogend Collaborators G. A new statistic to evaluate imputation reliability. *PloS one* 2010;**5**: e9697.
44. Wakefield J. A Bayesian measure of the probability of false discovery in genetic epidemiology studies. *American journal of human genetics* 2007;**81**: 208-27.
45. Mackenzie PI, Bock KW, Burchell B, Guillemette C, Ikushiro S, Iyanagi T, Miners JO, Owens IS, Nebert DW. Nomenclature update for the mammalian UDP glycosyltransferase (UGT) gene superfamily. *Pharmacogenetics and genomics* 2005;**15**: 677-85.
46. Steiman-Shimony A, Shtrikman O, Margalit H. Assessing the functional association of intronic miRNAs with their host genes. *Rna* 2018;**24**: 991-1004.

47. Martinez-Montiel N, Rosas-Murrieta NH, Anaya Ruiz M, Monjaraz-Guzman E, Martinez-Contreras R. Alternative Splicing as a Target for Cancer Treatment. *International journal of molecular sciences* 2018;**19**.
48. Ho MCW, Quintero-Cadena P, Sternberg PW. Genome-wide discovery of active regulatory elements and transcription factor footprints in *Caenorhabditis elegans* using DNase-seq. *Genome research* 2017;**27**: 2108-19.
49. Tourancheau A, Margaillan G, Rouleau M, Gilbert I, Villeneuve L, Levesque E, Droit A, Guillemette C. Unravelling the transcriptomic landscape of the major phase II UDP-glucuronosyltransferase drug metabolizing pathway using targeted RNA sequencing. *The pharmacogenomics journal* 2016;**16**: 60-70.
50. Perreault M, Gauthier-Landry L, Trottier J, Verreault M, Caron P, Finel M, Barbier O. The Human UDP-glucuronosyltransferase UGT2A1 and UGT2A2 enzymes are highly active in bile acid glucuronidation. *Drug metabolism and disposition: the biological fate of chemicals* 2013;**41**: 1616-20.
51. Sten T, Bichlmaier I, Kuuranne T, Leinonen A, Yli-Kauhaluoma J, Finel M. UDP-glucuronosyltransferases (UGTs) 2B7 and UGT2B17 display converse specificity in testosterone and epitestosterone glucuronidation, whereas UGT2A1 conjugates both androgens similarly. *Drug metabolism and disposition: the biological fate of chemicals* 2009;**37**: 417-23.
52. Itaaho K, Mackenzie PI, Ikushiro S, Miners JO, Finel M. The configuration of the 17-hydroxy group variably influences the glucuronidation of beta-estradiol and epiestradiol by human UDP-glucuronosyltransferases. *Drug metabolism and disposition: the biological fate of chemicals* 2008;**36**: 2307-15.

53. Bushey RT, Chen G, Blevins-Primeau AS, Krzeminski J, Amin S, Lazarus P. Characterization of UDP-glucuronosyltransferase 2A1 (UGT2A1) variants and their potential role in tobacco carcinogenesis. *Pharmacogenetics and genomics* 2011;**21**: 55-65.
54. Sneitz N, Court MH, Zhang X, Laajanen K, Yee KK, Dalton P, Ding X, Finel M. Human UDP-glucuronosyltransferase UGT2A2: cDNA construction, expression, and functional characterization in comparison with UGT2A1 and UGT2A3. *Pharmacogenetics and genomics* 2009;**19**: 923-34.
55. Bushey RT, Dluzen DF, Lazarus P. Importance of UDP-glucuronosyltransferases 2A2 and 2A3 in tobacco carcinogen metabolism. *Drug metabolism and disposition: the biological fate of chemicals* 2013;**41**: 170-9.
56. Heikkinen S, Vaisanen S, Pehkonen P, Seuter S, Benes V, Carlberg C. Nuclear hormone 1alpha,25-dihydroxyvitamin D3 elicits a genome-wide shift in the locations of VDR chromatin occupancy. *Nucleic acids research* 2011;**39**: 9181-93.
57. Bushey RT, Lazarus P. Identification and functional characterization of a novel UDP-glucuronosyltransferase 2A1 splice variant: potential importance in tobacco-related cancer susceptibility. *The Journal of pharmacology and experimental therapeutics* 2012;**343**: 712-24.
58. Faber MT, Kjaer SK, Dehlendorff C, Chang-Claude J, Andersen KK, Hogdall E, Webb PM, Jordan SJ, Australian Cancer S, Australian Ovarian Cancer Study G, Rossing MA, Doherty JA, Lurie G, Thompson PJ, Carney ME, Goodman MT, Ness RB, Modugno F, Edwards RP, Bunker CH, Goode EL, Fridley BL, Vierkant RA, Larson MC, Schildkraut J, Cramer DW, Terry KL, Vitonis AF, Bandera EV, Olson SH, King M, Chandran U, Kiemeny LA, Massuger LF, van Altena AM, Vermeulen SH, Brinton L, Wentzensen N, Lissowska J, Yang HP, Moysich KB, Odunsi K, Kasza K, Odunsi-Akanji O,

Song H, Pharaoh P, Shah M, Whittemore AS, McGuire V, Sieh W, Sutphen R, Menon U, Gayther SA, Ramus SJ, Gentry-Maharaj A, Pearce CL, Wu AH, Pike MC, Risch HA, Jensen A, Ovarian Cancer Association C. Cigarette smoking and risk of ovarian cancer: a pooled analysis of 21 case-control studies. *Cancer causes & control : CCC* 2013;**24**: 989-1004.

59. Kelemen LE, Abbott S, Qin B, Peres LC, Moorman PG, Wallace K, Bandera EV, Barnholtz-Sloan JS, Bondy M, Cartmell K, Cote ML, Funkhouser E, Paddock LE, Peters ES, Schwartz AG, Terry P, Alberg AJ, Schildkraut JM. Cigarette smoking and the association with serous ovarian cancer in African American women: African American Cancer Epidemiology Study (AACES). *Cancer causes & control : CCC* 2017.

60. Sheng Q, Liu J. The therapeutic potential of targeting the EGFR family in epithelial ovarian cancer. *British journal of cancer* 2011;**104**: 1241-5.

61. Gui T, Shen K. The epidermal growth factor receptor as a therapeutic target in epithelial ovarian cancer. *Cancer epidemiology* 2012;**36**: 490-6.

62. Reyes HD, Thiel KW, Carlson MJ, Meng X, Yang S, Stephan JM, Leslie KK. Comprehensive profiling of EGFR/HER receptors for personalized treatment of gynecologic cancers. *Molecular diagnosis & therapy* 2014;**18**: 137-51.

63. Showell R, Romano C, Valganon M, Lambros M, Trivedi P, Van Noorden S, Sriraksa R, El-Kaffash D, El-Etreby N, Natrajan R, Foroni L, Osborne R, El-Bahrawy M. The status of epidermal growth factor receptor in borderline ovarian tumours. *Oncotarget* 2016;**7**: 10568-77.

64. Ong JS, Cuellar-Partida G, Lu Y, Australian Ovarian Cancer S, Fasching PA, Hein A, Burghaus S, Beckmann MW, Lambrechts D, Van Nieuwenhuysen E, Vergote I, Vanderstichele A, Anne Doherty J, Anne Rossing M, Chang-Claude J, Eilber U, Rudolph A, Wang-Gohrke S, Goodman MT, Bogdanova N, Dork T, Durst M, Hillemanns P, Runnebaum IB, Antonenkova N, Butzow R, Leminen A, Nevanlinna H, Pelttari LM, Edwards RP, Kelley JL, Modugno F, Moysich KB, Ness RB, Cannioto R, Hogdall E, Hogdall CK, Jensen A, Giles GG, Bruinsma F, Kjaer SK, Hildebrandt MA, Liang D, Lu KH, Wu X, Bisogna M, Dao F, Levine DA, Cramer DW, Terry KL, Tworoger SS, Stampfer M, Missmer S, Bjorge L, Salvesen HB, Kopperud RK, Bischof K, Aben KK, Kiemeny LA, Massuger LF, Brooks-Wilson A, Olson SH, McGuire V, Rothstein JH, Sieh W, Whittemore AS, Cook LS, Le ND, Gilks CB, Gronwald J, Jakubowska A, Lubinski J, Kluz T, Song H, Tyrer JP, Wentzensen N, Brinton L, Trabert B, Lissowska J, McLaughlin JR, Narod SA, Phelan C, Anton-Culver H, Ziogas A, Eccles D, Campbell I, Gayther SA, Gentry-Maharaj A, Menon U, Ramus SJ, Wu AH, Dansonka-Mieszkowska A, Kupryjanczyk J, Timorek A, Szafron L, Cunningham JM, Fridley BL, Winham SJ, Bandera EV, Poole EM, Morgan TK, Risch HA, Goode EL, Schildkraut JM, Pearce CL, Berchuck A, Pharoah PD, Chenevix-Trench G, Gharahkhani P, Neale RE, Webb PM, MacGregor S. Association of vitamin D levels and risk of ovarian cancer: a Mendelian randomization study. *International journal of epidemiology* 2016;**45**: 1619-30.

65. Murphy SE, Park SS, Thompson EF, Wilkens LR, Patel Y, Stram DO, Le Marchand L. Nicotine N-glucuronidation relative to N-oxidation and C-oxidation and UGT2B10 genotype in five ethnic/racial groups. *Carcinogenesis* 2014;**35**: 2526-33.

66. Patel YM, Stram DO, Wilkens LR, Park SS, Henderson BE, Le Marchand L, Haiman CA, Murphy SE. The contribution of common genetic variation to nicotine and cotinine glucuronidation in multiple ethnic/racial populations. *Cancer*



*epidemiology, biomarkers & prevention* : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology 2015;**24**: 119-27.

Author Manuscript

APPENDIX

**Table 1. Top SNP P-values from gene regions associated with EOC in African American OncoArray analysis**

SNP ID (Effect / other allele) Nearest gene(s)	Effect Allele Frequency	N	OR	95% CI	P-value	Bayesian False Discovery Probability (BFDP)	Imputation quality
rs10017134 (C/T) UGT2A1/2 <sup>a*</sup>	0.73	1990	1.4	(1.2, 1.7)	1.2 x 10 <sup>-6</sup>	0.020	0.998
rs2288741 (T/G) UGT2A1/2*	0.73	1990	1.4	(1.2, 1.6)	1.9 x 10 <sup>-6</sup>	-----	-----
rs11939884 (T/G) UGT2A1/2 <sup>a*</sup>	0.14	1990	0.7	(0.5, 0.8)	1.7 x 10 <sup>-6</sup>	-----	-----

<sup>a</sup> Imputed

\*Bonferroni correction was applied to adjust for multiple SNPs comparisons. There were 433 SNPs in UGT2A1/2 gene. BFDP is reported based on a prior probability of association (pi0) equal to 0.6 \* 1/(Number of SNPs / 3).

**Table 2. Top SNP P-values from gene regions associated with HGSOC in African American OncoArray analysis**

SNP ID (Effect / other allele) Nearest gene(s)	Effect Allele Frequency	N	OR	95% CI	P-value	Bayesian False Discovery Probability (BFDP)	Imputation quality
rs114972508 (T/C) EGFR <sup>a*</sup>	0.04	1772	2.3	(1.2, 3.4)	1.6 x 10 <sup>-5</sup>	0.293	0.890
rs10017134 (C/T) UGT2A1/2 <sup>a+</sup>	0.72	1772	1.4	(1.2, 1.7)	2.3 x 10 <sup>-5</sup>	0.228	0.998
rs2288741 (T/G) UGT2A1/2 <sup>+</sup>	0.72	1772	1.4	(1.2, 1.7)	3.1 x 10 <sup>-5</sup>	-----	-----

<sup>a</sup> Imputed

This article is protected by copyright. All rights reserved

---

\*Bonferroni correction was applied to adjust for multiple SNPs comparisons. There were 824 SNPs in EGFR gene, and 413 SNPs in UGT2A1/2 gene. BFDP is reported based on a prior probability of association ( $\pi_0$ ) equal to  $0.5 * 1/(\text{Number of SNPs} / 3)$ .

Author Manuscript