

Mental health disorders, participation, and bullying in children with cerebral palsy

DANIEL G WHITNEY  | MARK D PETERSON | SETH A WARSCHAUSKY

Department of Physical Medicine and Rehabilitation, Michigan Medicine, University of Michigan, Ann Arbor, MI, USA.

Correspondence to Daniel G Whitney, Department of Physical Medicine and Rehabilitation, Michigan Medicine, University of Michigan, 325 E. Eisenhower Pkwy, Ann Arbor, MI 48108, USA. E-mail: dgwhit@umich.edu

This article is commented on by Paulus on page 862 of this issue.

PUBLICATION DATA

Accepted for publication 15th December 2018.

Published online 1st February 2019.

ABBREVIATION

NSCH National Survey of Children's Health

AIM To examine how social factors might mitigate the elevated risk of mental health disorders in children with cerebral palsy (CP).

METHOD This cross-sectional study included 6- to 17-year-olds with ($n=111$; 40.4% 6–11y, 59.6% 12–17y) and without ($n=29\ 909$; 50.2% 6–11y, 49.8% 12–17y) CP from the 2016 National Survey of Children's Health. Mental health disorders included depression, anxiety, behavior/conduct problems, and attention-deficit/hyperactivity disorder. Social factors included participation in activities, bully victimization, and difficulty with friendships.

RESULTS After adjusting for sociodemographic factors and the presence of chronic pain, children with CP had higher odds of anxiety (odds ratio [OR] 4.4; 95% confidence interval [CI] 1.9–8.5), behavior/conduct problems (OR 3.9; 95% CI 1.4–11.3), and multimorbidity (OR 2.8; 95% CI 1.1–7.0), but not depression (OR 1.4; 95% CI 0.6–3.8) or attention-deficit/hyperactivity disorder (OR 1.7; 95% CI 0.6–4.6), compared to controls. With adjustment for participation in activities, the odds of anxiety, behavior/conduct problems, and multimorbidity remained increased in children with CP. With adjustment for difficulty with friendships, the odds of anxiety, behavior/conduct problems, and multimorbidity were no longer increased in children with CP. With adjustment for bully victimization, the odds of behavior/conduct problems and multimorbidity were attenuated in children with CP; however, the odds of anxiety remained increased.

INTERPRETATION The elevated prevalence of certain mental health disorders in children with CP is partly associated with modifiable social factors.

Cerebral palsy (CP) refers to a group of motor disorders with a considerable range of motor dysfunction. It is the most common pediatric physical disability.¹ Individuals with CP are at a heightened risk for mental health disorders with prevalence rates up to 57%.^{2,3} This is concerning because mental health disorders are among the leading causes of the overall global burden of disease among children and adolescents.⁴ Further, psychopathology screening as part of a multidisciplinary follow-up in children and adolescents with CP has been recommended.³ Therefore, it is important to identify the risk factors associated with mental health disorders in children with CP.

There is evidence that physical problems and speech difficulties that are inherent to CP (e.g. motor dysfunction, dysarthria),^{5,6} in addition to medical comorbidities (e.g. sleep disorders),⁷ can affect mental health. Physical^{8–10} and communication⁵ impairments can alter the individual's interaction with the physical and social environment, thus affecting social development. Furthermore, social factors can have a profound impact on the individual's mental health status, such as quality of life and self-concept.¹⁰

Children with CP have less participation and enjoyment of social and recreational activities compared to typically developing children.^{11,12} Specifically, they spend less time with¹² and perceive lower social support from¹³ friends compared to the general adolescent population, providing a barrier to social development. Despite these social difficulties, children with CP place a high value and need for social and emotional networks,^{14–16} such as friendships.¹⁵ Friendships play an important role in the subjective well-being of children and have been shown to decrease the risk of developing mental health disorders.¹⁷ Evidence suggests that friendships may serve as a buffer to lessening the socially harmful effects of bullying,¹⁸ which is important because children with CP are at a heightened risk for being bullied.^{19–21} In the general population of adolescents, victims of bullying have a myriad of risk factors pertaining to poor mental health^{22,23} compared to non-victimized adolescents.

It is particularly important to examine mental health and associated risk factors in individuals with CP during the earlier phases of development. Therefore, the purpose of this study was to examine how social factors, including

participation in activities, difficulty with friendships, and bully victimization, in children with CP associate with several mental health disorders in this pediatric population. We hypothesized that each social factor would individually account for some of the elevated prevalence of mental health disorders in children with CP.

METHOD

The 2016 National Survey of Children's Health (NSCH) was conducted from June 2016 to February 2017 and was designed to generate state-level and national-level data on the health status of American children up to 17 years of age, their families, and their communities. Surveys were completed by mail and using online data collection instruments with a two-phase self-administered data collection design. Over 50 000 interviews of households that had children 0 to 17 years of age were completed across the USA. As previously described,²⁴ one eligible child per household was randomly selected for the detailed questionnaire and children with special healthcare needs were over-sampled to allow for more robust data estimates. The respondent was an adult (usually a parent) who had knowledge of the health status of the surveyed child. The overall weighted response rate was 40.7%. The completion rate for households that initiated the survey was 69.7%. The United States Census Bureau conducted a nonresponse bias analysis for the 2016 NSCH²⁵ and concluded that, although response was higher in geographical locations where income was higher, there is no strong or consistent evidence of non-response bias. Additional information about the NSCH methodology is available from the Data Resource Center for Child & Adolescent Health (<http://www.childhealthdata.org/>).

Participants

Details regarding participant selection have been previously described for this sample.²⁴ Briefly, the sample consisted of individuals with a current condition of CP and individuals who never had CP or any of the following conditions that were available in the 2016 NSCH: epilepsy, blood disorders, genetic disorders, cystic fibrosis, Down syndrome, developmental delay, intellectual disability, autism, or severe difficulty walking. Several variables used in the current investigation were not administered to children 5 years of age and younger. Therefore, the final sample was restricted to children aged 6 to 17 years old. Approval from an institutional review board was not necessary because the data used in this investigation are publicly available and deidentified.

Criteria commonly used to classify specific characteristics of CP were not available in the 2016 NSCH (e.g. the Gross Motor Function Classification System). On the other hand, the respondent's subjective report of the child's condition severity was available for all participants, and included either 'mild' or 'moderate or severe'.

What this paper adds

- Difficulty with friendships predicts an elevated prevalence of psychiatric conditions in children with cerebral palsy (CP).
- Bully victimization predicts an elevated prevalence of behavior/conduct problems in children with CP.
- Low participation does not predict mental health disorders in this population.

Outcome measures

Respondents answered the following prompt: 'Has a doctor or other healthcare provider ever told you that this child has [specified disorder]?' The child was considered to have a mental health disorder if the respondent reported a current condition of depression, anxiety problems, behavioral/conduct problems, or attention-deficit/hyperactivity disorder. The child was considered to not have a mental health disorder if the respondent reported 'no' to the condition or the child had the condition, but not at the time of the survey. Multimorbidity was defined as having two or more individual mental health disorders.

Predictor variables

A dichotomous variable (yes/no) for participation in activities was determined if the child participated in one or more extracurricular organized activities or lessons after school or on the weekends in the past 12 months. Activities included sports teams or lessons, clubs or organizations, and any other organized activities or lessons.

A dichotomous variable (yes/no) for difficulty with friendships was determined by the following prompt: 'Compared to other children his or her age, how much difficulty does this child have making or keeping friends?' The child was defined as having difficulty with friendships if the respondent answered 'a little difficulty/a lot of difficulty'. The child was defined as not having difficulty with friendships if the respondent answered 'no difficulty'.

A dichotomous variable (yes/no) for bully victimization was determined by how well the following phrase described the child: 'This child is bullied, picked on, or excluded by other children'. The child was defined as experiencing bully victimization if the respondent answered 'definitely true' or 'somewhat true'. The child was defined as not experiencing bully victimization if the respondent answered 'not true'.

Covariates

Chronic pain is a problem in individuals with CP and is associated with lower quality of life,¹⁰ behavioral and emotional problems,^{8,9} and other mental health disorders, such as anxiety and depression.⁸ Therefore, chronic pain was included as a covariate. A dichotomous variable for chronic pain (yes/no) was determined if the child had 'FREQUENT or CHRONIC difficulty' with 'repeated or chronic physical pain, including headaches or other back or body pain', as previously described. Other covariates included age, sex, ethnic group, and household poverty status as a ratio of the federal poverty line.

Statistical analysis

To account for the multistage and complex survey design of the 2016 NSCH, we used sampling strata and the primary sampling unit to obtain correct variance estimation, and weighted estimates to account for oversampling, survey non-response, and distribution of the target population. Statistical analysis was performed using SAS version 9.4 (SAS Institute, Cary, NC, USA) with statistical significance determined by p -values less than 0.05 and two-sided confidence intervals (CIs). Unweighted sample size and weighted estimates of all variables as percentages (95% CI) are presented. Group differences for all categorical variables were tested using χ^2 tests. Data regarding the severity of CP are not validated. Therefore, this variable is reported with descriptive characteristics to better characterize the CP sample, but has not been used to adjust the statistical analyses. We also examined the correlation between the three predictor variables for the total sample using the phi coefficient.

Weighted multivariable logistic regression analyses were performed with mental health disorders (yes/no) as the dependent variable and group (CP vs controls) as the primary exposure variable. Model 1 adjusted for age, sex, ethnic group, household poverty status, and chronic pain. Then, the predictor variables were added to the multivariable logistic model based on the significance of correlation among the three predictor variables for the total sample. If the three predictor variables were significantly correlated with one another, they were modeled individually in addition to the variables in model 1 to reduce collinearity. The interaction of group by predictor variables was examined for each outcome; if not significant, it was removed from the model. Goodness of fit for each model was examined using the C-statistic. Since multivariable logistic regression is event-driven, we followed the recommendation by Peduzzi et al.²⁶ and only performed multivariable logistic regression analysis when the number of events per variable was greater than 10 for each outcome to limit bias in estimating regression coefficients.

In the event that the odds of a mental health disorder were not significantly increased in children with CP after adjusting for the variables in model 1, subsequent models were not performed because the association between CP and that mental health disorder was accounted for. Interpretations are made based on the structure of the logistic regression models; that is, the predictors are the aforementioned social factors and the response variable are the individual mental health disorders. Individuals who had missing data for all variables in the logistic regression models, which represented less than 5% for CP and controls,²⁴ were included in the analyses.

RESULTS

The descriptive characteristics of participants are presented in Table I. Of the children with CP, just over half (56.6%) had moderate or severe disability. There was no statistical difference between groups for age, sex, or

Table I: Descriptive characteristics and social factors by group

	CP <i>n</i> =111 Percentage (95% CI)	Controls <i>n</i> =29 909 Percentage (95% CI)	<i>p</i>
Descriptive characteristics			
CP severity			
Mild	43.4 (25.3–61.6)		
Moderate or severe	56.6 (38.4–74.8)		
Age			0.31
6–11y	40.4 (21.9–58.9)	50.2 (48.9–51.6)	
12–17y	59.6 (41.1–78.1)	49.8 (48.4–51.1)	
Male	62.8 (46.4–79.2)	49.3 (48.0–50.6)	0.11
Ethnic group			<0.01
Non-Hispanic white	60.7 (41.7–79.7)	51.5 (50.1–52.8)	
Non-Hispanic black	27.1 (8.1–46.0)	12.5 (11.6–13.5)	
Hispanic	10.8 (0.0–22.0)	25.7 (24.2–27.2)	
Other	1.4 (0.2–2.7)	10.3 (9.6–11.0)	
Household poverty status			0.67
0–199%	50.6 (32.3–68.9)	43.7 (42.3–45.1)	
200–299%	9.7 (2.8–16.5)	15.0 (14.1–15.9)	
300–399%	13.2 (3.5–22.9)	11.0 (10.4–11.6)	
≥400%	26.6 (9.6–43.5)	30.3 (29.3–31.3)	
Chronic pain	39.0 (20.6–57.3)	7.6 (6.9–8.3)	<0.01
Social risk factors			
Participation in activities	49.6 (30.6–68.5)	80.5 (79.2–81.7)	<0.01
Difficulty with friendships	75.2 (60.5–89.9)	16.7 (15.8–17.7)	<0.01
Bully victimization	55.0 (36.7–73.2)	19.3 (18.2–20.3)	<0.01

Weighted data and analyses are presented. CP, cerebral palsy; CI, confidence interval.

household poverty status. There was a statistical difference between groups for the distribution of ethnic group ($p=0.005$), and children with CP had higher prevalence of chronic pain compared to controls ($p<0.001$). Children with CP had lower prevalence of participation in activities ($p<0.001$) and higher prevalence of difficulty with friendships ($p<0.001$) and bully victimization ($p<0.001$) compared to controls. In the total sample, participation in activities was significantly correlated with difficulty with friendships ($\phi=-0.14$, $p<0.001$) and bully victimization ($\phi=-0.10$, $p<0.001$); difficulty with friendships was significantly correlated with bully victimization ($\phi=0.44$, $p<0.001$).

Children with CP had higher prevalence of depression (7.8%, 2.7%; $p=0.017$), anxiety (30.2%, 6.2%; $p<0.001$), behavior/conduct problems (27.3%, 4.9%; $p<0.001$), attention-deficit/hyperactivity disorder (19.5%, 7.1%; $p=0.030$), and multimorbidity (22.3%, 5.2%; $p<0.001$) compared to controls.

The adjusted odds of mental health disorders are presented in Table II. After adjusting for the variables in model 1, the odds of depression (odds ratio [OR] 1.4; 95% CI 0.6–3.8; C-statistic 0.72) and attention-deficit/hyperactivity disorder (OR 1.7; 95% CI 0.6–4.6; C-statistic 0.64) were no longer significantly increased in children with CP; however, the odds of anxiety (OR 4.4; 95% CI 2.1–9.0; C-statistic 0.65), behavior/conduct problems (OR 3.9; 95% CI 1.4–11.3; C-statistic 0.66), and multimorbidity (OR 2.8;

Table II: Multivariable logistic regression models for mental health disorders in children with CP

	Model 1 ^a OR (95% CI)	Participation in activities and model 1 OR (95% CI)	Difficulty with friendships and model 1 OR (95% CI)	Bully victimization and model 1 OR (95% CI)
Depression				
CP (reference:controls)	1.4 (0.6–3.8)	Not performed	Not performed	Not performed
Anxiety				
CP (reference:controls)	4.4 (2.1–9.0)	4.0 (1.9–8.5)	2.1 (0.9–4.6)	3.4 (1.7–6.9)
Behavior/conduct problems				
CP (reference:controls)	3.9 (1.4–11.3)	3.7 (1.3–10.4)	1.7 (0.5–5.7)	2.5 (0.7–8.4)
ADHD				
CP (reference:controls)	1.7 (0.6–4.6)	Not performed	Not performed	Not performed
Multimorbidity				
CP (reference:controls)	2.8 (1.1–7.0)	2.6 (1.1–6.4)	1.1 (0.4–2.9)	1.9 (0.7–4.8)

^aModel 1: adjusted for age, sex, ethnic group, household poverty status, and chronic pain. Weighted data and analyses are presented. CP, cerebral palsy; OR, odds ratio; CI, confidence interval; ADHD, attention-deficit/hyperactivity disorder.

95% CI 1.1–7.0; C-statistic 0.61) were significantly increased in children with CP.

There was no significant group by predictor variable interaction (all $p > 0.05$). When participation in activities was added to model 1, the odds of anxiety (OR 4.0; 95% CI 1.9–8.5; C-statistic 0.66), behavior/conduct problems (OR 3.7; 95% CI 1.3–10.4; C-statistic 0.67), and multimorbidity (OR 2.6; 95% CI 1.1–6.4; C-statistic 0.63) remained significantly increased in children with CP. When difficulty with friendships was added to model 1, the odds of anxiety (OR 2.1; 95% CI 0.9–4.6; C-statistic 0.75), behavior/conduct problems (OR 1.7; 95% CI 0.5–5.7; C-statistic 0.80), and multimorbidity (OR 1.1; 95% CI 0.4–2.9; C-statistic 0.77) were no longer significantly increased (i.e. attenuated) in children with CP. When bully victimization was added to model 1, the odds of behavior/conduct problems (OR 2.5; 95% CI 0.7–8.4; C-statistic 0.75) and multimorbidity (OR 1.9; 95% CI 0.7–4.8; C-statistic 0.72) were attenuated, while the odds of anxiety (OR 3.4; 95% CI 1.7–6.9; C-statistic 0.72) remained significantly increased in children with CP.

DISCUSSION

The principal finding of this investigation is that social factors were associated with a higher prevalence of mental health disorders in children with CP after adjusting for sociodemographic variables and chronic pain. Specifically, difficulty with friendships accounted for the association between CP and anxiety, CP and behavior/conduct problems, and CP and multimorbidity. Bully victimization accounted for the association between CP and behavior/conduct problems and multimorbidity. Moreover, interactions of group by social factors were not significant for any of the mental health disorders in the present study, suggesting that children with CP demonstrate a similar pattern of association between these social factors and mental health disorders as their peers without CP.

Our finding of lower participation in activities in children with CP is consistent with previous studies.^{11,12} Many factors can lead to low participation for children with CP.

Physical factors (e.g. pain, low physical activity levels), developmental comorbidities (e.g. communication impairments), and environmental factors can independently or synergistically affect social participation. As potential evidence against our hypothesis, when participation in activities was added to model 1, the increased odds of selected mental health disorders persisted in children with CP (Table II). However, there is limited evidence to suggest that participation in and of itself may not affect peer relations in this population because participation is not synonymous with inclusion. For example, there is evidence that parents' efforts to facilitate social activities for children with CP are not associated with quantity and quality of friendships.²⁷

We found that 75% of respondents reporting on a child with CP reported that the child had difficulty with friendships compared to 17% of controls. Evidence suggests that children with CP value social and emotional networks,^{14–16} especially friendships.¹⁵ However, these children also report lower social support from friends compared to children without CP,¹³ which is consistent with a previous study of children born preterm having fewer friends.²⁸ In the current study, when difficulty with friendships was added to model 1, the increased odds of selected mental health disorders were attenuated in children with CP (Table II). These findings suggest an association between difficulty with friendships and mental health disorders in children with CP. This is consistent with a longitudinal study that found friendships mitigated the risk of developing mental health disorders in adolescents without CP.¹⁷ Therefore, it is important to identify the factors contributing to the gap between the desire and need for friendships^{14–16} and the perceived fulfillment of friendships¹³ in children with CP.

The importance of friendships on lessening the burden of mental health disorders includes an additional provision on the protective effects on being bullied.¹⁸ Children and adolescents who have higher perceived support of friends have a lower propensity to bully or be bullied.²⁹ In the current study, children with CP had a higher rate of bully

victimization compared to controls, with a prevalence in over half (55%) of the CP group. Children with CP are at an additional risk for bully victimization because of the physical vulnerabilities and social stigma associated with disability.^{19–21} In the general adolescent population, bully victimization has detrimental effects on mental health, suicidal ideation, self-esteem, and overall quality of life.^{22,23} Further, the association between bully victimization and mental health problems has been shown to be bidirectional.³⁰ In the current study, when bully victimization was added to model 1, the increased odds of anxiety persisted, but the odds of behavior/conduct problems and multimorbidity were attenuated in children with CP (Table II). These data suggest that bully victimization accounts for the CP-behavior/conduct problems and CP-multimorbidity associations. The findings support the need to improve bullying policies and develop programs that promote resilience to bully victimization in this population to improve mental health profiles.

Study limitations

First, this is a cross-sectional and observational study; thus, we could not determine causality or directionality in association between exposures and outcomes, as well as unmeasured confounding variables. Other comorbidities (e.g. intellectual disability, communication impairments, epilepsy) are prevalent in the CP population and are associated with risk factors for mental health disorders. Second, the NSCH survey method uses parent/guardian proxy reporting instead of self-reporting. Further, it was not possible to verify conditions through diagnostic evaluation or review of medical records. Therefore, conditions are subject to bias and misreporting. Third, commonly used criteria to classify CP characteristics, such as the Gross Motor Function Classification System, are not available in the NSCH survey, as previously noted. Pinquart¹⁹ conducted a systematic review of bullying in children and found that children with more visible chronic physical conditions were more likely to have reported bully victimization compared to children with less visible chronic physical conditions. Future studies incorporating the degree and/or visibility of motor impairment (e.g. the Gross Motor Function Classification System) as a factor for the development of mental health disorders may

provide further insight. Fourth, there were large differences in sample size between groups. However, the prevalence rate of CP in the 2016 NSCH is similar to previous reports from the USA, and the comparison group is a nationally representative group of children without developmental disabilities. Therefore, statistical analyses incorporating ‘group’ as the primary exposure variable in this study is representative of the CP population in terms of prevalence and provides clinical insight about the deviation of outcomes measures (e.g. anxiety) from what is typical in a non-disabled population of children.

CONCLUSION

The concerning prevalence of mental health disorders in children with CP²⁴ can be accounted for by modifiable social factors. Social development is a complex, dynamic, and interactive construct that is affected by many factors associated with CP. Lower participation was not associated with mental health disorders in children with CP in the current study; however, participation is not synonymous with full inclusion. Friendships are important to master developmental tasks and for psychosocial functioning in adulthood. Further, friendships protect against the mentally harmful effects of bully victimization, which is also associated with adverse mental health. These data may suggest the need for social and emotional support of friendships, promoting inclusivity among peers, and mitigating bully victimization or increasing resilience to bullying in children with CP.

ACKNOWLEDGEMENTS

DGW is supported by the University of Michigan Advanced Rehabilitation Research Training Program in Community Living and Participation from the National Institute on Disability, Independent Living, and Rehabilitation Research (90AR5020-0200). SAW is funded by a National Institutes of Health grant (5 UL1 TR002240-05), the Cerebral Palsy Alliance, and the Mildred E. Swanson Foundation. Funders had no role in the design or conduct of the study; the collection, management, analysis, or interpretation of the data; the preparation, review, or approval of the manuscript; or the decision to submit the manuscript for publication. The authors declare no conflicts of interest.

REFERENCES

- Oskoui M, Coutinho F, Dykeman J, Jetté N, Pringsheim T. An update on the prevalence of cerebral palsy: a systematic review and meta-analysis. *Dev Med Child Neurol* 2013; **55**: 509–19.
- Downs J, Blackmore AM, Epstein A, et al. The prevalence of mental health disorders and symptoms in children and adolescents with cerebral palsy: a systematic review and meta-analysis. *Dev Med Child Neurol* 2018; **60**: 30–8.
- Rackauskaite G, Bilenberg N, Bech BH, Uldall P, Østergaard JR. Screening for psychopathology in a national cohort of 8- to 15-year-old children with cerebral palsy. *Res Dev Disabil* 2016; **49–50**: 171–80.
- Kyu HH, Pinho C, Wagner JA, et al. Global and national burden of diseases and injuries among children and adolescents between 1990 and 2013: findings from the Global Burden of Disease 2013 study. *JAMA Pediatr* 2016; **170**: 267–87.
- Hidecker MJC, Slaughter J, Abeysekara P, et al. Early predictors and correlates of communication function in children with cerebral palsy. *J Child Neurol* 2018; **33**: 275–85.
- Christensen D, Van Naarden Braun K, Doernberg NS, et al. Prevalence of cerebral palsy, co-occurring autism spectrum disorders, and motor functioning – Autism and Developmental Disabilities Monitoring Network, USA, 2008. *Dev Med Child Neurol* 2014; **56**: 59–65.
- Zuculo GM, Knap CC, Pinato L. Correlation between sleep and quality of life in cerebral palsy. *Codas* 2014; **26**: 447–56.

8. Yamaguchi R, Nicholson Perry K, Hines M. Pain, pain anxiety and emotional and behavioural problems in children with cerebral palsy. *Disabil Rehabil* 2014; **36**: 125–30.
9. Novak I, Hines M, Goldsmith S, Barclay R. Clinical prognostic messages from a systematic review on cerebral palsy. *Pediatrics* 2012; **130**: e1285–312.
10. Russo RN, Miller MD, Haan E, Cameron ID, Crotty M. Pain characteristics and their association with quality of life and self-concept in children with hemiplegic cerebral palsy identified from a population register. *Clin J Pain* 2008; **24**: 335–42.
11. Chiarello LA, Palisano RJ, McCoy SW, et al. Child Engagement in Daily Life: a measure of participation for young children with cerebral palsy. *Disabil Rehabil* 2014; **36**: 1804–16.
12. Michelsen SI, Flachs EM, Damsgaard MT, et al. European study of frequency of participation of adolescents with and without cerebral palsy. *Eur J Paediatr Neurol* 2014; **18**: 282–94.
13. Colver A, Rapp M, Eisemann N, et al. Self-reported quality of life of adolescents with cerebral palsy: a cross-sectional and longitudinal analysis. *Lancet* 2015; **385**: 705–16.
14. Majnemer A, Shikako-Thomas K, Chokron N, et al. Leisure activity preferences for 6- to 12-year-old children with cerebral palsy. *Dev Med Child Neurol* 2010; **52**: 167–73.
15. Vinson J, Shank L, Thomas PD, Warschausky S. Self-generated domains of quality of life in children with and without cerebral palsy. *J Dev Phys Disabil* 2010; **22**: 497–508.
16. Shikako-Thomas K, Shevell M, Lach L, et al. Picture me playing – a portrait of participation and enjoyment of leisure activities in adolescents with cerebral palsy. *Res Dev Disabil* 2013; **34**: 1001–10.
17. van Harmelen AL, Gibson JL, St Clair MC, et al. Friendships and family support reduce subsequent depressive symptoms in at-risk adolescents. *PLoS ONE* 2016; **11**: e0153715.
18. Kochel KP, Ladd GW, Bagwell CL, Yabko BA. Bully/victim profiles' differential risk for worsening peer acceptance: the role of friendship. *J Appl Dev Psychol* 2015; **41**: 38–45.
19. Pinquart M. Systematic review: bullying involvement of children with and without chronic physical illness and/or physical/sensory disability – a meta-analytic comparison with healthy/nondisabled peers. *J Pediatr Psychol* 2017; **42**: 245–59.
20. Nadeau L, Tessier R. Social adjustment of children with cerebral palsy in mainstream classes: peer perception. *Dev Med Child Neurol* 2006; **48**: 331–6.
21. Dawkins JL. Bullying, physical disability and the paediatric patient. *Dev Med Child Neurol* 1996; **38**: 603–12.
22. Bowes L, Wolke D, Joinson C, Lereya ST, Lewis G. Sibling bullying and risk of depression, anxiety, and self-harm: a prospective cohort study. *Pediatrics* 2014; **134**: e1032–9.
23. Patrick DL, Bell JF, Huang JY, Lazarakis NC, Edwards TC. Bullying and quality of life in youths perceived as gay, lesbian, or bisexual in Washington State, 2010. *Am J Public Health* 2013; **103**: 1255–61.
24. Whitney DG, Warschausky SA, Peterson MD. Mental health disorders and physical risk factors in children with cerebral palsy: a cross-sectional study. *Dev Med Child Neurol* 2019. <https://doi.org/10.1111/dmcn.14083>. [Epub ahead of print]
25. 2016 National Survey of Children's Health. Nonresponse Bias Analysis [Internet]. Washington, DC: United States Department of Commerce. Available at: <https://census.gov/content/dam/Census/programs-surveys/nscch/tech-documentation/nonresponse-bias-analysis/NSCH%202016%20Nonresponse%20Bias%20Analysis.pdf> (accessed 3 January 2019).
26. Peduzzi P, Concato J, Kemper E, Holford TR, Feinstein AR. A simulation study of the number of events per variable in logistic regression analysis. *J Clin Epidemiol* 1996; **49**: 1373–9.
27. Thomas PD, Warschausky S, Golin R, Meiners K. Direct parenting methods to facilitate the social functioning of children with cerebral palsy. *J Dev Phys Disabil* 2008; **20**: 167–74.
28. Heuser KM, Jaekel J, Wolke D. Origins and predictors of friendships in 6- to 8-year-old children born at neonatal risk. *J Pediatr* 2018; **193**: 93–101 e5.
29. Kendrick K, Jutengren G, Stattin H. The protective role of supportive friends against bullying perpetration and victimization. *J Adolesc* 2012; **35**: 1069–80.
30. Reijntjes A, Kamphuis JH, Prinzie P, Telch MJ. Peer victimization and internalizing problems in children: a meta-analysis of longitudinal studies. *Child Abuse Negl* 2010; **34**: 244–52.



Editor's Choice

My Editor's Choice for the August 2019 issue is this study of factors associated with mental health problems in children with cerebral palsy. This is a particularly important but overlooked issue, with a great potential for both prevention and specific management. The study also highlights bullying, as does Kathy Zebracki in her editorial on social media (p. 856), and resilience factors such as friendships – an 'F-word' previously identified by Peter Rosenbaum and Jan Willem Gorter as an essential social factor for the development of personhood. These new results call for urgent action to promote mental health, and reduce peer exclusion and bullying victimization.

RESUMEN**ENFERMEDADES DE SALUD MENTAL, PARTICIPACIÓN Y BULLYING EN NIÑOS CON PARÁLISIS CEREBRAL**

OBJETIVOS Examinar cómo los factores sociales podrían mitigar el riesgo elevado de trastornos de salud mental en niños con parálisis cerebral (PC).

METODO Este estudio transversal incluyó niños de 6 a 17 años de edad con PC ($n=11$; 40,4% 6-11 años, 59,6% 12-17 años) y sin PC ($n=29.909$; 50,2% 6-11 años, 49,8% 12-17 años) de la Encuesta Nacional de Salud Infantil 2016. Los trastornos de la salud mental incluyeron depresión, ansiedad, problemas de conducta/comportamiento, y déficit de atención/hiperactividad. Los factores sociales incluyeron participación en actividades, victimización de Bullying y dificultades con amistades.

RESULTADOS Luego de ajustar con los factores sociodemográficos y la presencia de dolor crónico, los niños con PC tuvieron mayores probabilidades de presentar ansiedad (odds ratio [OR] 4,4; 95% intervalo de confianza [IC] 1,9-8,5), problemas de conducta/comportamiento (OR 3,9; 95%IC 1,4-11,3), y multimorbilidad (OR 2,8; 95% CI 1,1-7,0), pero no mayor depresión (OR 1,4; 95% CI 0,6-3,8) o déficit de atención/hiperactividad (OR 1,7; 95%CI 0,6-4,6), comparado con los controles. Con el ajuste para participación en actividades, la probabilidad de ansiedad, problemas de conducta/comportamiento, y multimorbilidad permanecieron más alta en niños con PC. Ajustando para dificultades con amistades, la probabilidad de ansiedad, problemas de conducta/comportamiento, y multimorbilidad no fueron más altas en niños con PC. Ajustando para la victimización de Bullying, la probabilidad de problemas de comportamiento/conducta y multimorbilidad se atenuaron en niños con PC; sin embargo la probabilidad de presentar ansiedad permaneció aumentada.

INTERPRETACION La elevada prevalencia de ciertos trastornos de salud mental en niños con PC se asocia en parte con factores sociales modificables.

RESUMO**TRANSTORNOS DE SAÚDE MENTAL, PARTICIPAÇÃO E BULLYING EM CRIANÇAS COM PARALISIA CEREBRAL**

OBJETIVO Examinar como fatores sociais podem mitigar o elevado risco de transtornos de saúde mental em crianças com paralisia cerebral (PC).

MÉTODO Este estudo transversal incluiu pessoas entre 6 e 17 anos de idade com ($n=111$; 40,4% 6-11a, 59,6% 12-17a) e sem ($n=29.909$; 50,2% 6-11a; 49,8% 12-17a) PC do Levantamento Nacional de Saúde da Criança de 2016. Transtornos de saúde mental incluíram depressão, ansiedade, problemas de comportamento/conduita, e transtorno de déficit de atenção e hiperatividade. Fatores sociais incluíram participação em atividades, ser vítima de bullying e dificuldade com amizades.

RESULTADOS Após ajustar para fatores sócio-demográficos e presença de dor crônica, crianças com PC tiveram maior risco de ansiedade (odds ratio [OR] 4,4; intervalo de confiança [IC] a 95% 1,9-8,5), problemas de comportamento/conduita (OR 3,9; IC 95% 1,4-11,3), e multimorbidade (OR 2,8; IC 95% 1,1-7,0), mas não depressão (OR 1,4; IC 95% 0,6-3,8) ou transtorno do déficit de atenção e hiperatividade (OR 1,7; IC 95% 0,6-4,6), em comparação com controles. Ajustando para participação em atividades, os riscos de ansiedade, problemas de comportamento/conduita, e multimorbidade permaneceram aumentados em crianças com PC. Ajustando para a dificuldade com amizades, o risco de ansiedade, problemas de comportamento/conduita, e multimorbidade não foram mais aumentados em crianças com PC. Com ajuste para a presença de bullying, os riscos de problemas de comportamento/conduita e multimorbidade foram atenuados em crianças com PC; no entanto, os riscos de ansiedade permaneceram aumentados.

INTERPRETAÇÃO A elevada prevalência de certos transtornos de saúde mental em crianças com PC é parcialmente associado a fatores sociais modificáveis.