

Article type : Full Length

2019 EULAR/ACR classification criteria for systemic lupus erythematosus

CRITERIA+ Add Section head+ Add Section head

Martin Aringer,¹ Karen Costenbader,² David Daikh,³ Ralph Brinks,⁴ Marta Mosca,⁵ Rosalind Ramsey-Goldman,⁶ Josef S Smolen,⁷ David Wofsy,⁸ Dimitrios T Boumpas,^{9,10} Diane L Kamen,¹¹ David Jayne,¹² Ricard Cervera,¹³ Nathalie Costedoat-Chalumeau,¹⁴ Betty Diamond,¹⁵ Dafna D Gladman,¹⁶ Bevra Hahn,¹⁷ Falk Hiepe,¹⁸ Søren Jacobsen,¹⁹ Dinesh Khanna,²⁰ Kirsten Lerstrøm,²¹ Elena Massarotti,^{22,23} Joseph McCune,²⁰ Guillermo Ruiz-Irastorza,²⁴ Jorge Sanchez-Guerrero,^{25,26} Matthias Schneider,²⁷ Murray Urowitz,²⁸ George Bertsias,²⁹ Bimba F Hoyer,^{18,30} Nicolai Leuchten,¹ Chiara Tani,³¹ Sara K Tedeschi,^{23,32} Zahi Touma,³³ Gabriela Schmajuk,³ Branimir Anic,³⁴ Florence Assan,³⁵ Tak Mao Chan,³⁶ Ann Elaine Clarke,³⁷ Mary K Crow,³⁸ László Czirják,³⁹ Andrea Doria,⁴⁰ Winfried Graninger,⁴¹ Bernadett Halda-Kiss,³⁹ Sarfaraz Hasni,⁴² Peter M Izmirlly,⁴³ Michelle Jung,³⁷ Gábor Kumánovics,³⁹ Xavier Mariette,^{44,45} Ivan Padjen,³⁴ José M Pego-Reigosa,⁴⁶ Juanita Romero-Diaz,⁴⁷ Íñigo Rúa-Figueroa,⁴⁸ Raphaèle Seror,³⁵ Georg H Stummvoll,⁴⁹ Yoshiya Tanaka,⁵⁰ Maria G Tektonidou,⁵¹ Carlos Vasconcelos,⁵² Edward M Vital,^{53,54} Daniel J Wallace,⁵⁵ Sule Yavuz,⁵⁶ Pier Luigi Meroni,⁵⁷ Marvin J Fritzler,⁵⁸ Ray Naden,⁵⁹ Thomas Dörner,¹⁸ Sindhu R Johnson^{60,61}

¹Medicine III, University Medical Center and Faculty of Medicine Carl Gustav Carus, TU Dresden, Dresden, Germany; ²Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA; ³University of California at San Francisco and VA Medical Center, San Francisco, California, USA; ⁴Policlinic and Hiller Research Unit for Rheumatology, Medical Faculty, Heinrich-Heine-University Duesseldorf, Duesseldorf, Germany; ⁵Azienda Ospedaliero Universitaria Pisana, University of Pisa, Pisa, Italy; ⁶Northwestern University Feinberg School of Medicine, Chicago, Illinois, USA; ⁷Medical University of Vienna, Vienna, Austria; ⁸Russell/Engleman Rheumatology Research Center, University of California at San Francisco, San Francisco, California, USA; ⁹Medical School, National and Kapodestrian University of Athens, and Biomedical Research Foundation of the Athens Academy, **This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/ART.40930](https://doi.org/10.1002/ART.40930)**

This article is protected by copyright. All rights reserved

Athens, Greece; ¹⁰Medical School, University of Cyprus, Nicosia, Cyprus; ¹¹Medical University of South Carolina, Charleston, South Carolina, USA; ¹²Department of Medicine, University of Cambridge, Cambridge, UK; ¹³Department of Autoimmune Diseases, Hospital Clínic, University of Barcelona, Barcelona, Catalonia, Spain; ¹⁴Cochin Hospital, Internal Medicine Department, Centre de référence maladies auto-immunes et systémiques rares d'île de France, Paris, France; ¹⁵The Feinstein Institute for Medical Research, Manhasset, New York, USA; ¹⁶Division of Rheumatology, Department of Medicine, Toronto Western Hospital, University of Toronto, Toronto, Ontario, Canada; ¹⁷University of California at Los Angeles, Los Angeles, California, USA; ¹⁸Department of Rheumatology and Clinical Immunology, Charité – Universitätsmedizin Berlin, Corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Berlin, Germany; ¹⁹Copenhagen Lupus and Vasculitis Clinic, Rigshospitalet, Copenhagen University Hospital, Copenhagen, Denmark; ²⁰Division of Rheumatology, University of Michigan, Ann Arbor, Michigan, USA; ²¹Lupus Europe, Co-Opted Trustee for Research, Essex, UK; ²²Medicine, Brigham and Women's Hospital, Boston, Massachusetts, USA; ²³Harvard Medical School, Boston, Massachusetts, USA; ²⁴Biocruces Bizkaia Health Research Institute, Hospital Universitario Cruces, UPV/EHU, Bizkaia, The Basque Country, Spain; ²⁵Division of Rheumatology, Department of Medicine, Mount Sinai Hospital/University Health Network, University of Toronto, Toronto, Ontario, Canada; ²⁶Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City, Mexico; ²⁷Policlinic and Hiller Research Unit for Rheumatology, Medical Faculty, Heinrich-Heine-University, Duesseldorf, Germany; ²⁸Center for Prognosis Studies in the Rheumatic Diseases, Toronto Western Hospital, University of Toronto, Lupus Clinic, Toronto, Ontario, Canada; ²⁹Rheumatology, Clinical Immunology and Allergy, University of Crete Medical School, Heraklion, Crete, Greece; ³⁰University of Schleswig-Holstein at Kiel, Kiel, Germany; ³¹Rheumatology Unit, Azienda Ospedaliero Universitaria Pisana, University of Pisa, Pisa, Italy; ³²Brigham and Women's Hospital, Boston, Massachusetts, USA; ³³Division of Rheumatology, Department of Medicine, Toronto Western Hospital, Institute of Health Policy, Management and Evaluation, University of Toronto, Toronto, Ontario, Canada; ³⁴Division of Clinical Immunology and Rheumatology, University of Zagreb School of Medicine and University Hospital Centre Zagreb, Zagreb, Croatia; ³⁵Université Paris Sud, Hôpitaux Universitaires Paris-Sud, AP-HP, INSERM UMR 1184, Le Kremlin-Bicêtre, France; ³⁶Department of Medicine, University of Hong Kong, Hong Kong, China; ³⁷Division of Rheumatology, Cumming School of Medicine, University of Calgary, Calgary, Alberta, Canada; ³⁸Hospital for Special Surgery, New York, New York, USA; ³⁹Department of Rheumatology and Immunology, University of Pécs, Pécs, Hungary; ⁴⁰Rheumatology Unit, Department of Medicine (DIMED), University of Padova, Padova, Italy; ⁴¹Internal Medicine, Medical University of Graz, Graz, Austria; ⁴²Lupus Clinical Research Program, Office of the Clinical Director, National Institute of Arthritis and Musculoskeletal and Skin Diseases, National Institutes of Health, Bethesda, Maryland, USA; ⁴³Rheumatology, New York University School of Medicine, New York, New York, USA; ⁴⁴Rheumatology, Assistance Publique-Hôpitaux de Paris (AP-HP), Hôpitaux universitaires Paris-

Sud – Hôpital Bicêtre, Le Kremlin Bicêtre, France; ⁴⁵3Université Paris-Sud, Center for Immunology of Viral Infections and Auto-immune Diseases (IMVA), Institut pour la Santé et la Recherche Médicale (INSERM) UMR 1184, Université Paris-Saclay, Le Kremlin Bicêtre, France; ⁴⁶University Hospital of Vigo, IRIDIS Group, Instituto de Investigación Sanitaria Galicia Sur (IISGS), Vigo, Spain; ⁴⁷Department of Immunology and Rheumatology, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City, Mexico; ⁴⁸Rheumatology, Doctor Negrín University Hospital, Las Palmas de Gran Canaria, Las Palmas, Spain; ⁴⁹Rheumatology, Medical University of Vienna, Vienna, Austria; ⁵⁰First Department of Internal Medicine, University of Occupational and Environmental Health, Japan, Kitakyushu, Japan; ⁵¹Medical School, National and Kapodistrian University of Athens, Athens, Greece; ⁵²Centro Hospitalar do Porto, ICBAS, University of Porto, Porto, Portugal; ⁵³Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds, Leeds, UK; ⁵⁴NIHR Leeds Biomedical Research Centre, Leeds Teaching Hospitals NHS Trust, Leeds, UK; ⁵⁵Division of Rheumatology, Cedars-Sinai Medical Center, Los Angeles, California, USA; ⁵⁶Istanbul Bilim University, Istanbul, Turkey; ⁵⁷Clinical Immunology and Rheumatology Unit, IRCCS Istituto Auxologico Italiano, Milan, Italy; ⁵⁸Faculty of Medicine, Cumming School of Medicine, University of Calgary, Calgary, Alberta, Canada; ⁵⁹Maternal-Fetal Medicine, McMaster University Faculty of Health Sciences, Hamilton, Ontario, Canada; ⁶⁰Medicine, Toronto Western Hospital, University Health Network, Mount Sinai Hospital, University of Toronto, Toronto Scleroderma Research Program, Toronto, Ontario, Canada; ⁶¹Institute of Health Policy, Management and Evaluation, University of Toronto, Toronto, Ontario, Canada

Correspondence: Martin Aringer

Medicine III, University Medical Center and Faculty of Medicine Carl Gustav Carus, TU Dresden, Dresden D-01307, Germany

Email martin.aringer@uniklinikum-dresden.de

251120182704201901052019

Keywords: systemic lupus erythematosus , lupus , classification criteria , consensus methods , multi-criteria decision analysis , validation

Related Article:

Cite as: Aringer M, Costenbader K, Daikh D, et al. *Ann Rheum Dis* 2019;:1–9. doi:

[annrheumdis-2018-214819](https://doi.org/10.1136/annrheumdis-2018-214819)

© Author(s) (or their employer(s)) 2019. No commercial re-use. See rights and permissions. Published by BMJ.

2019

Objective: To develop new classification criteria for systemic lupus erythematosus (SLE) jointly supported by the European League Against Rheumatism (EULAR) and the American College of Rheumatology (ACR).

Methods: This international initiative had four phases. (1) Evaluation of antinuclear antibody (ANA) as an entry criterion through systematic review and meta-regression of the literature and criteria generation through an international Delphi exercise, an early patient cohort and a patient survey. (2) Criteria reduction by Delphi and nominal group technique exercises. (3) Criteria definition and weighting based on criterion performance and on results of a multi-criteria decision analysis. (4) Refinement of weights and threshold scores in a new derivation cohort of 1001 subjects and validation compared with previous criteria in a new validation cohort of 1270 subjects.

Results: The 2019 EULAR/ACR classification criteria for SLE include positive ANA at least once as obligatory entry criterion; followed by additive weighted criteria grouped in seven clinical (constitutional, haematological, neuropsychiatric, mucocutaneous, serosal, musculoskeletal, renal) and three immunological (antiphospholipid antibodies, complement proteins, SLE-specific antibodies) domains, and weighted from 2 to 10. Patients accumulating ≥ 10 points are classified. In the validation cohort, the new criteria had a sensitivity of 96.1% and specificity of 93.4%, compared with 82.8% sensitivity and 93.4% specificity of the ACR 1997 and 96.7% sensitivity and 83.7% specificity of the Systemic Lupus International Collaborating Clinics 2012 criteria.

Conclusion: These new classification criteria were developed using rigorous methodology with multidisciplinary and international input, and have excellent sensitivity and specificity. Use of ANA entry criterion, hierarchically clustered and weighted criteria reflect current thinking about SLE and provide an improved foundation for SLE research.

Introduction

Systemic lupus erythematosus (SLE) is a complex autoimmune disease with variable clinical features.¹ SLE manifestations are associated with multiple autoantibodies, ensuing immune complex formation and deposition, and other immune processes.^{2 3} This complex clinical presentation and pathogenesis makes SLE a difficult disease to grasp and define. Classification criteria are essential for the identification of relatively homogeneous groups of patients for inclusion in research studies and trials.⁴ The 1982 revised American College of Rheumatology (ACR) SLE classification criteria⁶ and their

1997 revision⁷ have been used worldwide. Since then, our understanding of the disease has advanced. Additional specific skin manifestations were described, some clinical symptoms were better understood, and immunological tests, such as diminished levels of serum complement components C3 and C4 or testing for anti- β 2 glycoprotein I antibodies, entered routine clinical practice. Better understanding of organ system involvement, such as mucocutaneous abnormalities, led to questions about whether some of the independently counted criteria were in fact manifestations of the same phenomenon.⁸

The 2012 Systemic Lupus International Collaborating Clinics (SLICC) classification criteria addressed many of these issues.⁹ Mucocutaneous and neuropsychiatric manifestations were added, as were hypocomplementemia and new antiphospholipid antibody tests; and criteria definitions were refined. The SLICC criteria emphasised that SLE is primarily an autoantibody disease, requiring at least one immunological criterion to be present, and categorised histology-proven nephritis compatible with SLE as sufficient for classification, if antinuclear antibodies (ANAs) or antibodies to double-stranded DNA (dsDNA) were present. While achieving their goal of increasing sensitivity, the SLICC criteria have lower specificity than the 1997 ACR criteria.^{9 10}

Existing SLE classification criteria perform better in patients with longstanding disease than in new-onset SLE,¹¹ and there is an increasing recognition and demand that subjects with early SLE should be included in clinical studies and trials. We therefore attempted to enrich our sample populations for early SLE in several phases of the project.

In parallel with improved understanding of SLE, the field of classification criteria development has also seen advances.^{4 12–14} In order to minimise investigator bias, it is now recommended that the cohorts in which the criteria are tested are from independent centres.⁴ Other methodological recommendations include a balanced use of both expert-based and data-driven methods, and inclusion of the patient perspective.^{13 14} The approach chosen for these 2019 European League Against Rheumatism (EULAR)/ACR SLE classification criteria was specifically designed to maintain this balance and to uphold rigorous methodology.

Methods

Methodological overview

Using a methodological approach based on measurement science the criteria were developed in four phases¹⁰: (1) criteria generation, (2) criteria reduction, (3) criteria definition and weighting and (4) refinement and validation (figure 1). The whole initiative was overseen by a 12-member steering committee (MA, KHC, DD, MM, RR-G, JSS, DW, DB, DK, DJ, TD and SRJ) nominated by EULAR and the ACR in equal numbers, based on SLE and/or methodological experience and previous involvement in international projects.

Figure 1 Development and validation of SLE classification criteria. ANA, antinuclear antibody; SLE, systemic lupus erythematosus.

The current project, jointly supported by the EULAR and the ACR, was originally based on two key concepts. One, we hypothesised that the presence of ANA would be better employed as an entry criterion than as a classification criterion.¹⁰ Such an approach was thought to reflect underlying SLE pathogenesis, and take into account ANA test characteristics of high sensitivity and limited specificity. Two, we expected individual criteria would not be of equal utility (weight) for the classification of SLE,¹⁵ for example, mucosal ulcers versus biopsy-proven lupus nephritis. Accordingly, the validity of using positive ANA as an entry criterion was explicitly addressed in phase I of the current activity.¹⁶ Likewise, methodological strategies to develop weighted criteria were used.

Phase I. Criteria generation

The purpose of phase I was to test ANA as a potential entry criterion and identify candidate criteria that should be considered for SLE classification using both data-based and expert-based methods, including the patient perspective. *Phase Ia* comprised a systematic literature review of Medline, Embase and the Cochrane databases with meta-regression to evaluate the operating characteristics of ANA testing for consideration as an entry criterion.¹⁶ *Phase Ib* consisted of a Delphi exercise of international SLE experts from the America, Europe and Asia.¹⁷ These experts included rheumatologists, dermatologists, nephrologists, paediatricians and non-clinical SLE researchers, providing a broad perspective. The Delphi participants were asked to nominate a broad set of items potentially useful in the classification of SLE.¹⁷ In round 2 and 3, participants rated the items from 1 (not at all appropriate) to 9 (completely appropriate) for classification of SLE. Criteria were retained if they reached a median rating of ≥ 6.5 ; that is, at least 50% of the ratings in the high range (7, 8 or 9). Participants were also asked about the importance of ANA and histopathology for classification of SLE. *Phase Ic* established an international cohort of patients with early SLE or conditions mimicking SLE to identify criteria that may discriminate subjects with early (less than 12 months) disease.¹⁸ *Phase Id* comprised a cross-sectional survey of SLE patients, administered via the quarterly journal of the German SLE patient organisation, which asked about symptoms within 1 year before and after the patient's diagnosis of SLE.¹⁹ While at a risk of recall bias and not necessarily representative of other regions worldwide, this survey was done to explicitly take a patient standpoint into account.

For phase II and III, additional renowned European and North American SLE experts were nominated by the steering committee and invited to participate.

Phase II. Criteria reduction

Phase IIa. The objective of this phase was to select a set of criteria from phase I that maximised the likelihood of accurate classification of SLE, particularly of early disease. An independent panel of seven of the international SLE experts (RC, NC-C, DDG, BHH, FH, EM and JS-G) ranked the candidate criteria from phase I. A consensus meeting of 19 international SLE experts (n=7 nominal group technique (NGT) experts+steering committee+DK (moderator)) using NGT was conducted to reduce the list of criteria.²⁰ Data for each candidate criterion were reviewed and discussed until consensus was achieved. The NGT experts voted on items to be retained. *Phase IIb.* NGT participants pointed out that some criteria could be correlated. With the idea of potentially clustering criteria into domains, associations between candidate criteria were evaluated separately in two cohorts, the phase Ic early SLE and the Euro-lupus cohorts.²¹

Phase III. Criteria definition and weighting

Phase IIIa. The operating characteristics of the retained candidate criteria were evaluated by literature review. Candidate criteria were hierarchically organised into clinical and immunological domains, and definitions for the candidate criteria were iteratively refined. SLE patient advocates participated in the review of data and the steering committee discussions.²²

Phase IIIb. 164 case vignettes reflecting broad SLE clinical presentation were sampled from SLE centres across several countries. A panel of six of the international experts not involved in earlier phases of the project (BD, SJ, WJMcC, GR-I, MS and MBU) and 11 members of the steering committee assessed and ranked a representative sample of the cases. Subsequently, at a face-to-face meeting, this panel of 17 international SLE experts iteratively compared pairs of criteria, using multicriteria decision analysis facilitated by 1000minds software.²³ The panel unanimously agreed to further reduce the list of criteria. Based on the results, provisional criteria weights were assigned and a provisional threshold score for classification was determined as the lowest score at which the expert panel had achieved consensus on classifying a case vignette as SLE.²⁴

Phase IV. Refinement and validation

International SLE experts not involved in phase II or phase III panels were asked to contribute cases diagnosed as SLE and controls with conditions mimicking SLE sampled from patients evaluated at their centres. Each centre was asked to contribute up to 100 cases and an equal number of controls, preferentially sampling those with early disease, and regardless of their specific clinical or immunological manifestations. Pseudonymised data on the criteria were collected using a standardised data collection form. Ethics committee approval and informed consent were obtained as per local

requirements. The status ('SLE' or not) of each case underwent independent adjudication by three of four SLE experts (GB, BFH, NL and CT) from different centres. Queries were sent back to the submitting investigator for clarification. Of this cohort, 501 SLE and 500 control subjects were randomly selected to comprise the derivation cohort, while the remaining 696 SLE and 574 control subjects formed the validation cohort.

Refinement. The performance of the draft criteria set was iteratively tested in the derivation cohort. A data-driven threshold for classification was determined by receiver operating characteristics (ROC) analysis and compared with the provisional expert-based consensus threshold. The data of SLE subjects below the threshold (misclassified) were reviewed for groups of patients with unequivocal SLE who still missed classification, and criteria weights adjusted slightly, while preserving the weighting hierarchy (details below in Results, Phase IV section). Sensitivity and specificity was tested against the ACR 1997 and the SLICC 2012 criteria. In addition, ANA as an entry criterion was tested against not having an entry criterion. Finally, the criteria weights were simplified to whole numbers. Refinements to the criteria set were presented to the steering committee and phase III expert panel, and unanimously endorsed.

Validation. The sensitivity and specificity of the final criteria were tested in the validation cohort and compared with previous SLE criteria sets.

Statistical analysis. Descriptive statistics were used to summarise the data. CIs were calculated using the bias-corrected and accelerated bootstrap method (BCa method) with B=2000 bootstrap samples. The BCa method resamples the input data B times (with replacement) and calculates the required statistics (sensitivity, specificity, AUC). Based on the B bootstraps samples, the bias-correction is applied and the associated 95% CIs for the statistics are estimated. The BCa method has proven to yield very accurate coverage of estimated CIs.²⁵ The number B of bootstrap resamples is recommended to be at least B=1000. We have chosen B=2000 and additionally checked if B=5000 bootstraps changed the estimated confidence bounds, which was not the case. Statistical analyses were performed using R, V.3.4.0 (The R Foundation of Statistical Computing).

Results

Phase I: Criteria generation

Phase Ia. ANA as an entry criterion. A systematic review of MEDLINE, EMBASE and the Cochrane database identified 13 080 patients from 64 studies reporting ANA by immunofluorescence on HEp-2 cells. Meta-regression of the operating characteristics of ANA found a sensitivity of 97.8% (95% CI 96.8% to 98.5%) for ANA of $\geq 1:80$ supporting use of ANA as an entry criterion.¹⁶ Since some SLE centres do not have access to HEp-2 ANA, and in view of ongoing work on the standardisation of

serology and potential future advances in the field, the steering committee and additional autoantibody consultants (MJF and PLM) recommended the provision ‘or an equivalent positive ANA test. Testing by immunofluorescence on HEp-2 cells or a solid phase ANA screening immunoassay with at least equivalent performance is highly recommended’.

Phase Ib. Delphi exercise. One hundred and forty-seven international SLE experts nominated 145 candidate criteria.¹⁷ By rating the appropriateness for SLE classification, the participants in the second and third Delphi round reduced the list to 40 candidate criteria (online supplementary table 1).

Phase Ic. International early SLE cohort. The cohort comprised 616 subjects who had been referred for possible SLE with a disease duration of less than 1 year (n=389 early SLE and n=227 mimicking diseases) from North America, Europe, Asia and South America.¹⁸ In addition to supporting many of the 40 candidate criteria derived from the Delphi exercise, the comparison between early SLE and non-SLE patients showed that fever occurred more frequently (34.5% vs 13.7%, p<0.001) in SLE, while SLE patients less commonly suffered from arthralgias (20.3% vs 42.7%, p=0.001) and fatigue (28.3% vs 37%, p=0.02).

Phase Id. Patient survey. 339 SLE patients (>99% Caucasian, 93% female) responded to the survey.¹⁹ More than half of these patients reported mucocutaneous findings in the first year of their disease (online supplementary table 1), but also fatigue (89%), joint pain (87%) and fever (54%).¹⁹ Given that these items were highlighted both in the early SLE cohort and the patient survey, fever, fatigue and arthralgias were forwarded to the next phase in addition to the 40 Delphi items. Accordingly, phases Ia–Id resulted in a total of 43 candidate criteria for consideration (online supplementary table 1).

Phase II. Criteria reduction

Phase IIa. The expert panel NGT exercise reduced the candidate criteria from 43 to 21.²⁶ The panel distinguished potential ‘entry criteria’, which would be required for classification, from potential ‘additive criteria’. They endorsed ‘positive ANA (\geq 1:80 by HEp-2 immunofluorescence)’ as an entry criterion. The 20 remaining additive criteria included: lupus nephritis by renal biopsy, autoantibodies, cytopenias, fever, arthritis, serositis, mucocutaneous and neuropsychiatric manifestations (online supplementary table 1).

Phase IIb. Associations between the candidate criteria were evaluated in 389 subjects in the early SLE cohort and the 1000 SLE subjects of the Euro-lupus cohort. Modest statistically significant correlations were limited to the mucocutaneous (r=0.22–0.30), neurological (r=0.22) and immunological (r=0.33) domains in the early SLE cohort, and this modest correlation was replicated in the Euro-lupus cohort.²¹ Given these associations, criteria were clustered within domains, so that only one criterion within each domain would be counted.

Phase III. Criteria definition and weighting

Phase IIIa. Based on the literature, definitions of the 20 candidate additive criteria were refined, using a data-driven evaluation of operating characteristics,²² retaining only feasible items with a prevalence of at least 1% according to literature. Literature-review led to the consensus decision to evaluate five different candidate criteria within the neuropsychiatric domain (delirium, psychosis, seizure, mononeuropathy, cranial neuropathy) and potential separation of acute pericarditis from pleural or pericardial effusions; and between diminished C3 or C4 versus diminished C3 *and* C4 (online supplementary table 1). The resulting 23 candidate criteria (online supplementary table 1) were organised into seven clinical and three immunological domains, with hierarchical clustering.²² Only the highest-ranking item in each domain was to be counted. Instead of devising exclusion definitions for each criterion, the decision was made to attribute any item to SLE only if no more likely explanation was present. For leucopenia and joint involvement, it was decided to formally test alternative definitions in the derivation cohort. Given the importance of testing for antibodies, particularly for anti-dsDNA, for which tests of relatively low specificity are in use, great care was taken to precisely define testing (table 1).

Table 1 Definitions of SLE classification criteria

Criteria	Definition
Antinuclear antibodies (ANA)	ANA at a titre of $\geq 1:80$ on HEp-2 cells or an equivalent positive test at least once. Testing by immunofluorescence on HEp-2 cells or a solid phase ANA screening immunoassay with at least equivalent performance is highly recommended
Fever	Temperature $>38.3^{\circ}\text{C}$
Leucopenia	White blood cell count $<4000/\text{mm}^3$
Thrombocytopenia	Platelet count $<100\ 000/\text{mm}^3$
Autoimmune haemolysis	Evidence of haemolysis, such as reticulocytosis, low haptoglobin, elevated indirect bilirubin, elevated LDH
Delirium	Characterised by (1) change in consciousness or level of arousal with reduced ability to focus, (2) symptom development over hours to <2 days, (3) symptom fluctuation throughout the day, (4) either (4a) acute/subacute change in cognition (eg, memory deficit or disorientation), or (4b) change in behaviour, mood, or affect (eg, restlessness, reversal of sleep/wake cycle)
Psychosis	Characterised by (1) delusions and/or hallucinations without insight and

	(2) absence of delirium
Seizure	Primary generalised seizure or partial/focal seizure
Non-scarring alopecia	Non-scarring alopecia observed by a clinician*
Oral ulcers	Oral ulcers observed by a clinician*
Subacute cutaneous or discoid lupus	<p>Subacute cutaneous lupus erythematosus observed by a clinician*: Annular or papulosquamous (psoriasiform) cutaneous eruption, usually photodistributed</p> <p>Discoid lupus erythematosus observed by a clinician*: Erythematous-violaceous cutaneous lesions with secondary changes of atrophic scarring, dyspigmentation, often follicular hyperkeratosis/plugging (scalp), leading to scarring alopecia on the scalp</p> <p>If skin biopsy is performed, typical changes must be present. Subacute cutaneous lupus: interface vacuolar dermatitis consisting of a perivascular lymphohistiocytic infiltrate, often with dermal mucin noted. Discoid lupus: interface vacuolar dermatitis consisting of a perivascular and/or periappendageal lymphohistiocytic infiltrate. In the scalp, follicular keratin plugs may be seen. In longstanding lesions, mucin deposition and basement membrane thickening may be noted</p>
Acute cutaneous lupus	<p>Malar rash or generalised maculopapular rash observed by a clinician*</p> <p>If skin biopsy is performed, typical changes must be present (acute cutaneous lupus: interface vacuolar dermatitis consisting of a perivascular lymphohistiocytic infiltrate, often with dermal mucin noted. Perivascular neutrophilic infiltrate may be present early in the course</p>
Pleural or pericardial effusion	Imaging evidence (such as ultrasound, X-ray, CT scan, MRI) of pleural or pericardial effusion, or both
Acute pericarditis	≥2 of (1) pericardial chest pain (typically sharp, worse with inspiration, improved by leaning forward), (2) pericardial rub, (3) EKG with new widespread ST-elevation or PR depression, (4) new or worsened pericardial effusion on imaging (such as ultrasound, X-ray, CT scan, MRI)
Joint involvement	EITHER (1) synovitis involving two or more joints characterised by swelling or effusion OR (2) tenderness in two or more joints and at least 30 min of morning stiffness
Proteinuria >0.5 g/24 hours	Proteinuria >0.5 g/24 hours by 24 hours urine or equivalent spot urine protein-to-creatinine ratio
Class II or V lupus	Class II: mesangial proliferative lupus nephritis: purely mesangial

nephritis on renal biopsy according to ISN/RPS 2003 classification	hypercellularity of any degree or mesangial matrix expansion by light microscopy, with mesangial immune deposit. A few isolated subepithelial or subendothelial deposits may be visible by immunofluorescence or electron microscopy, but not by light microscopy Class V: membranous lupus nephritis: global or segmental subepithelial immune deposits or their morphological sequelae by light microscopy and by immunofluorescence or electron microscopy, with or without mesangial alterations
Class III or IV lupus nephritis on renal biopsy according to ISN/RPS 2003	Class III: focal lupus nephritis: active or inactive focal, segmental or global endocapillary or extracapillary glomerulonephritis involving <50% of all glomeruli, typically with focal subendothelial immune deposits, with or without mesangial alterations Class IV: diffuse lupus nephritis: active or inactive diffuse, segmental or global endocapillary or extracapillary glomerulonephritis involving ≥50% of all glomeruli, typically with diffuse subendothelial immune deposits, with or without mesangial alterations. This class includes cases with diffuse wire loop deposits but with little or no glomerular proliferation
Positive antiphospholipid antibodies	Anticardiolipin antibodies (IgA, IgG, or IgM) at medium or high titer (>40 APL, GPL or MPL, or >the 99th percentile) or positive anti-β2GP1 antibodies (IgA, IgG, or IgM) or positive lupus anticoagulant
Low C3 OR low C4	C3 OR C4 below the lower limit of normal
Low C3 AND low C4	Both C3 AND C4 below their lower limits of normal
Anti-dsDNA antibodies OR anti-Smith (Sm) antibodies.	Anti-dsDNA antibodies in an immunoassay with demonstrated ≥90% specificity for SLE against relevant disease controls OR anti-Sm antibodies

Notes: ISN/RPS International Society of Nephrology/Renal Pathology Society.

Notes: *This may include physical examination or review of a photograph.

Abbreviations: SLE, systemic lupus erythematosus; dsDNA, double-stranded DNA.

Phase IIIb. The 1.5 day in-person consensus meeting using multicriteria decision analysis involved 74 decisions between pairs of criteria. Criteria weights were calculated by the 1000minds software based on these decisions (table 2). International Society of Nephrology/Renal Pathology Society class III or IV nephritis consistently attained higher weight than class II or V nephritis, so lupus nephritis by histology was separated into two different criteria. Class VI lupus nephritis as an end stage manifestation was

unanimously eliminated. Likewise, the experts unanimously voted to not retain mononeuropathy and cranial neuropathy, which had been included into the set of potential neuropsychiatric items in phase IIIa but turned out to add little to SLE classification. The use of weighted criteria led to a sum score that is a measure of the relative probability of a subject having SLE, with higher scores indicating higher likelihood. Experts reached full consensus on a classification of SLE at a provisional threshold score of >83 of a theoretical maximum of 305.24

Table 2 Relative weights of the additive classification criteria items

Domain	Item	Original	Modification	Revised	Simplified
Constitutional	Fever	13		13	2
Haematological	Leucopenia	12	+7	19	3
	Thrombocytopenia	26		26	4
	Autoimmune haemolysis	28		28	4
Neuropsychiatric	Delirium	12		12	2
	Psychosis	20		20	3
	Seizure	34		34	5
Mucocutaneous	Alopecia	13		13	2
	Oral ulcers	14		14	2
	SCLE/DLE	29		29	4
	ACLE	38		38	6
Serosal	Effusion	34		34	5
	Acute pericarditis	38		38	6
Musculoskeletal	Joint involvement	34	+4	38	6
Renal	Proteinuria	27		27	4
	Class II/V	55		55	8
	Class III/IV	74		74	10
APL antibodies	Antiphospholipid	13		13	2

Complements	C3 or C4 low	19		19	3
	C3 and C4 low	27		27	4
SLE-specific antibodies	Anti-Sm	40		40	6
	Anti-dsDNA	38		38	6

Notes: Weights derived from the phase III consensus meeting with multicriteria decisions analysis (original), added points for leucopenia and joint involvement (modification), the resulting weights (revised) and the final simplified weights (simplified).

Abbreviations: SLE, systemic lupus erythematosus; anti-Sm, anti-Smith; dsDNA, double-stranded DNA.

Phase IV. Refinement and validation

Twenty-one centres from the USA, Canada, Mexico, Austria, Croatia, France, Germany, Greece, Hungary, Italy, Portugal, Spain, the UK, Turkey, Hong Kong and Japan submitted a total of 2339 cases from their cohorts. 1197 SLE and 1074 non-SLE diagnoses (table 3) were verified by three adjudicators blinded to the proposed classification criteria system. Due to lack of consensus during adjudication, 68 subjects (2.9%) were excluded from the analysis.

Table 3 Demographic characteristics of the derivation and validation cohorts

	Derivation cohort		Validation cohort	
	SLE	Non-SLE	SLE	Non-SLE
n	501	500	696	574
Female/male	447/54	421/79	608/88	490/84
Age (mean±SD) years	45±14	54±16	45±14	56±16
Disease duration (mean±SD) years	11±8	9±8	11±8	9±8
Ethnicity				
Black	29	10	56	12
East Asian	36	29	53	34
Hispanic	59	48	73	51
South/South East Asian	16	6	21	11
White	355	404	480	461
Other	6	3	13	5

SLE	501		696	
Non-SLE		500		574
Adult onset still's disease		2		11
Autoimmune thyroiditis		6		5
Behcet's disease		7		9
Cancer		2		3
Inflammatory myositis		37		27
Fibromyalgia		6		3
Membranous nephritis		11		14
Osteoarthritis		2		NA
Mixed connective tissue disease		9		15
Primary antiphospholipid antibody syndrome		45		48
Psoriatic arthritis		12		9
Rheumatoid arthritis		94		110
Sarcoidosis		2		2
Sjögren's syndrome		112		124

Spondyloarthritis		5		5
Systemic sclerosis		99		120
Tuberculosis		0		2
Undifferentiated connective tissue disease		16		20
Vasculitis		9		13
Viral infection		5		5
Other		19		29

Notes: Inflammatory myositis includes dermatomyositis, polymyositis and juvenile dermatomyositis.

Abbreviations: SLE, systemic lupus erythematosus.

Derivation cohort. Of the 2271 triple-adjudicated cases, 501 SLE and 500 non-SLE cases were randomly assigned to the derivation cohort. The provisional weighting system derived from phase III was tested in the derivation cohort. ROC analysis suggested a data-driven threshold of ≥ 70 (of a maximum of 305), with a sensitivity of 95.4% and a specificity of 95.2%, which was superior to the consensus-derived provisional threshold of > 83 that had high specificity (98.8%), but lower sensitivity (81.6%). Review of subjects below the threshold of 70 identified a subgroup of SLE subjects with joint involvement and/or leucopenia. Thus, weights for leucopenia and joint involvement were each adjusted (table 2) to reduce misclassification. When alternative definitions for leucopenia and joint involvement were tested, leucopenia defined as a white blood cell count (WBC) $< 4000/\text{mm}^3$ at least once⁹ also had a slightly higher sensitivity +specificity (1.944 vs 1.942) than leucopenia defined as WBC $< 4000/\text{mm}^3$ on two or more occasions.^{6 26} Joint involvement defined as EITHER ‘synovitis involving two or more joints, characterised by swelling or effusion’, OR ‘tenderness in two or more joints and at least 30 min of morning stiffness’⁹ had a higher combined sensitivity and specificity than arthritis defined simply as synovitis of two or more joints (1.944 vs 1.900). When retested, the revised criteria had increased sensitivity, and maintained sensitivity +specificity. Evaluating ANA as an entry criterion, the criteria

with the ANA entry criterion had better performance than without (sensitivity +specificity 1.944 vs 1.930). Next, the weights were simplified by division to whole numbers to achieve a threshold of 10 (table 2). In the derivation cohort, the sensitivity and specificity of the final criteria set (figure 2) were reaching the performance benchmarks set for this project (table 4).

Figure 2 Classification criteria for systemic lupus erythematosus.

Table 4 Operating characteristics of the new classification criteria compared with the ACR 1997 and SLICC 2012 classification criteria in the derivation and the validation cohorts

	ACR 1997 criteria	SLICC 2012 criteria	EULAR/ACR 2019 criteria
Derivation			
Sensitivity (95% CI)	0.85 [0.81 to 0.88]	0.97 [0.95 to 0.98]	0.98 [0.97 to 0.99]
Specificity (95% CI)	0.95 [0.93 to 0.97]	0.90 [0.87 to 0.92]	0.96 [0.95 to 0.98]
Combined (95% CI)	1.80 [1.76 to 1.83]	1.87 [1.84 to 1.90]	1.94 [1.92 to 1.96]
Validation			
Sensitivity (95% CI)	0.83 [0.80 to 0.85]	0.97 [0.95 to 0.98]	0.96 [0.95 to 0.98]
Specificity (95% CI)	0.93 [0.91 to 0.95]	0.84 [0.80 to 0.87]	0.93 [0.91 to 0.95]
Combined (95% CI)	1.76 [1.73 to 1.80]	1.80 [1.77 to 1.84]	1.90 [1.87 to 1.92]

Abbreviations: ACR, American College of Rheumatology; EULAR, European League Against Rheumatism; SLICC, Systemic Lupus International Collaborating Clinics.

Validation. The validation cohort, that is, the full cohort minus the derivation cohort, comprised 1270 triple adjudicated subjects (n=696 SLE, n=574 controls). The criteria, with positive ANA as an entry criterion, weighted criteria in seven clinical domains (constitutional, hematological, neuropsychiatric, mucocutaneous, serosal, musculoskeletal, renal) and three immunological domains (antiphospholipid antibodies, low complements, anti-Smith (anti-Sm) and anti-dsDNA as SLE-specific antibodies) and a classification threshold score of ≥ 10 (out of a theoretical maximum of 51) (figure 2), had a sensitivity of 96.1% and a specificity of 93.4% (table 4). It demonstrated improved performance compared with the ACR 1997 and SLICC 2012 criteria.

Discussion

New SLE classification criteria were developed with support by both the ACR and EULAR. Through a four-phase, iterative process, we have defined an additive, weighted multicriteria system that produces a measure of the relative probability that an individual can be classified as SLE. The system defines a threshold above which experts would classify cases as SLE for the purpose of research studies. We have carefully defined the criteria to improve reliability and precision; and have grouped the criteria into ten hierarchical domains. We have validated the criteria against a large number of cases, including many patients with manifestations that resemble SLE but who do not have SLE. This approach, as well as the resulting criteria system, represents a paradigm shift for the classification of SLE.

We have defined positive ANA at any time as required entry criterion. There were three possible ways to deal with ANA testing. The previous criteria sets have treated ANA the same as the much more specific antibodies against Sm and dsDNA, which we considered suboptimal given important differences in sensitivity and specificity. We could have excluded ANA completely in classifying lupus, but we still consider ANA a useful test and concept. We therefore decided to test ANA as an entry criterion, which reflects the use of ANA as a highly sensitive screening test.

Criteria using ANA as entry criterion had better performance. During the phase I Delphi exercise, 58% of SLE experts did not feel comfortable and an additional 19% were uncertain about classifying a patient with SLE in the absence of ever having a positive ANA.¹⁷ The systematic literature review and meta-regression of data on 13 080 subjects demonstrated ANA $\geq 1:80$ have a sensitivity of 98% with a lower limit of the 95% CI at 97%.¹⁶ In the phase I early SLE cohort, 99.5% of the 389 SLE patients were ANA positive.¹⁸ The frequencies of ANA positive SLE patients in the derivation and validation cohorts (99.6% and 99.3%, respectively) were in the same range. Since both in the early SLE cohort and in the derivation and validation cohorts, patients were included in many centres worldwide independent of ANA positivity, the latter data provide additional support for ANA as an entry criterion.

Using ANA as entry criterion means the new criteria cannot classify SLE among patients who are persistently ANA negative. While possibly also distinguished by lower cytokine levels²⁷ and lower efficacy of immunomodulatory treatment,²⁸ such a subgroup of patients exists. Although small, it may vary in size in different populations.¹⁶ This patient subset needs to be put high on the scientific agenda for further investigation. Additional characterisation of this phenomenon may lead to an alternative entry criterion for this small group of patients. For the moment, we still think it is acceptable to exclude ANA negative patients from clinical trials.

Molecular classification criteria were also considered during the development of these criteria.²⁹ Many novel biomarkers were nominated, such as increased circulating B lymphocyte stimulator (BLyS), IFN γ induced protein 10 kD (IP-10), monocyte chemoattractant protein-1 (MCP-1), TNF- α , type I interferon

signature, or increased Th17 and plasma cell populations. They were all voted out in the expert Delphi exercise, largely because of limited availability in the clinical setting and/or insufficient evidence.⁵ However, inclusion of novel biomarkers, beyond autoantibodies, may ultimately further improve the specificity of SLE classification, increase alignment of classification with underlying disease pathogenesis and improve the performance and information content of clinical trials. Thus, testing of biomarkers against these criteria is an important area for future research.

A new clinical criterion, unexplained fever, turned out to be common and remarkably characteristic for SLE. However, since infections are a major cause of death in SLE, it is of utmost importance to stress that fever, like all other criteria manifestations, should only be counted if no better explanation exists, and that infections have to be suspected first in any patient with (potential) SLE, particularly when CRP is elevated.³⁰ The concept that all criteria are only to be counted if SLE is thought to be the most likely cause of the manifestation (ie, no other more likely cause exists) is central to these new EULAR/ACR criteria, and is explicitly stated as an overarching principle. Some criteria, such as delirium, psychosis and acute pericarditis, were in part redefined based on existing scientific definitions.²² Where alternative definitions were used, the performance of the alternative definitions was comparatively evaluated in the derivation cohort.

The differential weighting of criteria better represents their relative contribution to an individual's classification of SLE. For SLE, renal biopsy with Class III or IV lupus nephritis carries the most weight and in the presence of a positive ANA is enough to classify a patient as SLE. This further develops a concept of the SLICC criteria⁹ and reflects the current thinking of SLE experts; in the Delphi exercise, 85% would classify SLE on renal pathology alone.¹⁷ Renal biopsy with class II or V lupus nephritis still carries a large weight (eight points) but is not by itself sufficient for the classification of SLE.

The numerical goal of this project was to keep the specificity similar to the specificity of the ACR 1997 criteria, but increase the sensitivity to the high sensitivity level of the SLICC criteria, if possible. The validation cohort data suggest that this goal has been achieved. From our data, it appears that the SLICC criteria increase in sensitivity was to a significant degree founded in accepting renal histology and adding subacute cutaneous lupus and low complement levels. These three advances are mirrored in the current criteria. Many of the other additional symptoms of the SLICC criteria were of very low frequency. Specificity was increased by weighting of criteria, by the NGT expert panel decision to not allow lymphopenia to go forward, and, importantly, by the decision that no criterion be counted if better explained by another condition.

The new criteria provide a simple, directed and highly accurate method for classifying SLE. An electronic 'app' is in preparation, which will assist in the use of these criteria. However, it is important to stress that classification criteria are not designed for diagnosis or treatment decisions.⁵ They should never be used to exclude patients who do not fully meet these criteria from receiving appropriate

therapies. This is also pertinent to patients with ANA-negative SLE discussed above. Diagnosis of SLE remains the purview of an appropriately trained physician evaluating an individual patient.⁵

The new SLE classification system also provides new research opportunities. With much interest in early or latent SLE,^{31 32} the additive point system and the relative probability of classification it produces, allows for systematic study of individuals who fall below the classification threshold. This will facilitate studies of disease evolution and early intervention. Furthermore, the use of an additive scoring system will allow for studying the idea of ‘ominosity’, that is, the potential implications of having very high scores on disease severity and subsequent prognosis. This work would need to reconsider the relative contribution of individual criteria (weights) and consider additional criteria that potentially contribute to ominosity.

It is anticipated that other groups will test these criteria, which will constitute important external validation. This will be particularly important for paediatric SLE and those with organ dominant, for example, skin dominant disease, since it is a limitation of this criteria project that the patient cohorts do not represent these subgroups. Similar limitations also pertain to several ethnic groups and to men with SLE, each only included in lower numbers. It is important to independently test the EULAR/ACR criteria in these subgroups. Investigators doing so are reminded about the critical importance of the correct attribution of each criterion. Criteria can only be counted when not better explained by another condition. The attribution process requires diligence and clinical experience.

In summary, our multiphase methodological approach and ensuing classification system using ANA as an entry criterion and weighted, hierarchically clustered criteria, constitute a paradigm shift in the classification of SLE. These criteria have excellent performance characteristics and face validity, as the structure and weighting were designed to reflect current thinking about SLE. The inclusion of fever assists with the classification of early SLE. The separation of renal biopsy findings reflects their differential impact on the probability of SLE classification. These criteria have strong operating characteristics, with excellent sensitivity and specificity. This classification system was built using rigorous methodology that was both data-driven and expert-based. With the inclusion of over 200 SLE experts from multiple countries and medical disciplines, methodologists, patient advocates and over 4000 subjects, this work is the largest international, collaborative SLE classification effort to date.

Footnotes

supplement

Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/annrheumdis-2018-214819>).

Prepub

Original reference

EAHP Statement

Citation

Presented at

Twitter

Handling editor

David S Pisetsky

Acknowledgements

This body of work was jointly supported by the European League Against Rheumatism and the American College of Rheumatology. One part of the derivation and validation cohort was supported by the Intramural Research Program of the National Institute of Arthritis and Musculoskeletal and Skin Diseases of the National Institutes of Health. The authors wish to acknowledge the diligent work of Banita Aggarwal and Keshini Devakandan in data entry, data cleaning, queries to submitting investigators, data cutting and maintenance of the derivation and validation cohorts; and of Corine Sinnette, Massachusetts, in the preparatory work for the multicriteria decision analysis exercise.

Contributors

MA, SRJ, TD, KC, DD, RB, MM, RR-G, JSS, DW, DB, DLK, DJ and RN have been involved in the planning and execution of the project and in writing the manuscript. RC, NC-C, BD, DDG, BH, FH, SJ, DK, KL, EM, W. JM, GR-I, JS-G, MS, MU, GB, BFH, NL, CT, SKT, ZT, GS, BA, FA, TMC, AEC,

This article is protected by copyright. All rights reserved

MKC, LC, AD, WG, BH-K, SH, PMI, MJ,GK, XM, IP, JMP-R, JR-D, IR-F, RS, GHS, YT, MGT, CV, EMV, DJW, SY, PLM and MJF have each significantly contributed to the body of work of the project and involved in correcting and finalising the manuscript.

Author note

Disclaimer

Competing interests

None declared.

Patient and public involvement statement

Patient consent for publication

Not required.

Ethics approval

Local ethics committees where applicable for chart based review of data.

Provenance and peer review

Not commissioned; externally peer reviewed.

Data availability statement

Press Release

Collaborators

Correction notice

Permission

Editor's note

Republishing

Guest Chief Editor

Deceased

References

1. Bertias GK, Pamfil C, Fanouriakis A, et al. Diagnostic criteria for systemic lupus erythematosus: has the time come? *Nat Rev Rheumatol* 2013;9:687–94.
2. Rahman A, Isenberg DA. Systemic lupus erythematosus. *N Engl J Med* 2008;358:929–39.
3. Tsokos GC. Systemic lupus erythematosus. *N Engl J Med* 2011;365:2110–21.
4. Johnson SR, Goek O-N, Singh-Grewal D, et al. Classification criteria in rheumatic diseases: a review of methodologic properties. *Arthritis Rheum* 2007;57:1119–33.
5. Aggarwal R, Ringold S, Khanna D, et al. Distinctions between diagnostic and classification criteria? *Arthritis Care Res* 2015;67:891–7.
6. Tan EM, Cohen AS, Fries JF, et al. The 1982 revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum* 1982;25:1271–7.
7. Hochberg MC. Updating the American College of rheumatology revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum* 1997;40:1705–9.
8. Albrecht J, Berlin JA, Braverman IM, et al. Dermatology position paper on the revision of the 1982 ACR criteria for systemic lupus erythematosus. *Lupus* 2004;13:839–49.
9. Petri M, Orbai A-M, Alarcón GS, et al. Derivation and validation of the systemic lupus international collaborating clinics classification criteria for systemic lupus erythematosus. *Arthritis Rheum* 2012;64:2677–86.
10. Aringer M, Dörner T, Leuchten N, et al. Toward new criteria for systemic lupus erythematosus—a standpoint. *Lupus* 2016;25:805–11.

11. Inês L, Silva C, Galindo M, et al. Classification of systemic lupus erythematosus: systemic lupus international collaborating clinics versus American College of rheumatology criteria. A comparative study of 2,055 patients from a real-life, International systemic lupus erythematosus cohort. *Arthritis Care Res* 2015;67:1180–5.
12. Felson DT, Anderson JJ. Methodological and statistical approaches to criteria development in rheumatic diseases. *Baillière's Clinical Rheumatology* 1995;9:253–66.
13. Singh JA, Solomon DH, Dougados M, et al. Development of classification and response criteria for rheumatic diseases. *Arthritis Rheum* 2006;55:348–52.
14. Dougados M, Gossec L. Classification criteria for rheumatic diseases: why and how? *Arthritis Rheum* 2007;57:1112–5.
15. Costenbader KH, Karlson EW, Mandl LA. Defining lupus cases for clinical studies: the Boston weighted criteria for the classification of systemic lupus erythematosus. *J Rheumatol* 2002;29:2545–50.
16. Leuchten N, Hoyer A, Brinks R, et al. Performance of antinuclear antibodies for classifying systemic lupus erythematosus: a systematic literature review and meta-regression of diagnostic data. *Arthritis Care Res* 2018;70:428–38.
17. Schmajuk G, Hoyer BF, Aringer M, et al. Multicenter Delphi exercise to identify important key items for classifying systemic lupus erythematosus. *Arthritis Care Res* 2018;70:1488–94.
18. Mosca M, Costenbader KH, Johnson SR, et al. Brief report: how do patients with newly diagnosed systemic lupus erythematosus present? A multicenter cohort of early systemic lupus erythematosus to inform the development of new classification criteria. *Arthritis Rheumatol* 2019;71:91–8.
19. Leuchten N, Milke B, Winkler-Rohlfing B, et al. Early symptoms of systemic lupus erythematosus (SLE) recalled by 339 SLE patients. *Lupus* 2018;27:1431–6.
20. Nair R, Aggarwal R, Khanna D. Methods of formal consensus in classification/diagnostic criteria and guideline development. *Semin Arthritis Rheum* 2011;41:95–105.
21. Touma Z, Cervera R, Brinks R, et al. Associations among classification criteria items within systemic lupus erythematosus. *Arthritis Rheumatol* 2017;69.
22. Tedeschi SK, Johnson SR, Boumpas D, et al. Developing and refining new candidate criteria for systemic lupus erythematosus classification: an international collaboration. *Arthritis Care Res* 2018;70:571–81.

23. Johnson SR, Naden RP, Fransen J, et al. Multicriteria decision analysis methods with 1000Minds for developing systemic sclerosis classification criteria. *J Clin Epidemiol* 2014;67:706–14.
24. Tedeschi SK, Johnson SR, Boumpas DT, et al. Multicriteria decision analysis process to develop new classification criteria for systemic lupus erythematosus. *Ann Rheum Dis* 2019;78:634–40.
25. Efron B, Tibshirani RJ. *An introduction to the bootstrap*. Boca Raton: Chapman & Hall / CRC, 1994.
26. Johnson SR, Khanna D, Daikh D, et al. Use of consensus methodology to determine candidate items for systemic lupus erythematosus classification criteria. *J Rheumatol* 2018.
27. Torell F, Eketjäll S, Idborg H, et al. Cytokine profiles in autoantibody defined subgroups of systemic lupus erythematosus. *J Proteome Res* 2019;18:1208–17.
28. Wallace DJ, Stohl W, Furie RA, et al. A phase II, randomized, double-blind, placebo-controlled, dose-ranging study of belimumab in patients with active systemic lupus erythematosus. *Arthritis Rheum* 2009;61:1168–78.
29. Johnson SR, Hinchcliff M, Asano Y. Controversies: molecular vs. clinical systemic sclerosis classification. *J Scleroderma Relat Disord* 2016;1:277–85.
30. Littlejohn E, Marder W, Lewis E, et al. The ratio of erythrocyte sedimentation rate to C-reactive protein is useful in distinguishing infection from flare in systemic lupus erythematosus patients presenting with fever. *Lupus* 2018;27:1123–9.
31. Ganczarczyk L, Urowitz MB, Gladman DD, et al. "Latent lupus".. *J Rheumatol* 1989;16:475–8.
32. Costenbader KH, Schur PH. We need better classification and terminology for "people at high risk of or in the process of developing lupus". *Arthritis Care Res* 2015;67:593–6.



online supplementary file 1



online supplementary file 2