



COMMENTARY

Is a soft tissue graft harvested from the maxillary tuberosity the approach of choice in an isolated site?

Lorenzo Tavelli¹ | Shayan Barootchi¹ | Henry Greenwell² | Hom-Lay Wang¹

¹Department of Periodontics & Oral Medicine, University of Michigan School of Dentistry, Ann Arbor, MI

²Department of Periodontics, University of Louisville, School of Dental Medicine, Louisville, KY

Correspondence

Hom-Lay Wang, DDS, MS, PhD, Department of Periodontics and Oral Medicine, University of Michigan School of Dentistry, 1011 North University Avenue, Ann Arbor, Michigan 48109-1078, USA.

Email: homlay@umich.edu

Abstract

Soft tissue augmentation procedures are becoming more popular these days. Different soft tissue graft harvesting approaches have been proposed. Nonetheless, the location of the donor site (whether anterior-, lateral-, superficial-, deep-palate or the maxillary tuberosity) can affect the graft shape and its composition. Soft tissue grafts from the maxillary tuberosity are rich in connective tissue fibers, with minimal presence of fatty or glandular components. Clinical, histological, and molecular evidence shows that a soft tissue graft obtained from the maxillary tuberosity has unique properties. In addition, harvesting from this area presents minimal risk for intra- or postoperative complications, leading to reduced patient morbidity. The aim of this commentary is to discuss the advantages and disadvantages of harvesting a soft tissue graft from the tuberosity and to compare it with the traditional palatal graft, while highlighting functional, esthetic, and patient-related outcomes.

KEYWORDS

connective tissue graft, dental implants, gingival recession, maxillary tuberosity, soft tissue augmentation, surgical flaps, tooth root

1 | INTRODUCTION

The connective tissue graft (CTG) is considered the material of choice in treating gingival or mucosal recessions around teeth and implants because it provides an increased marginal soft tissue thickness, protects postsurgical healing by the first intention, and maintains the innate tissue texture as of adjacent teeth or implants.¹⁻⁴ Initially, soft tissue graft was introduced for re-establishing an adequate width of keratinized tissue and mucosa (KT/KM),⁵ and since then, the free gingival graft (FGG) has also been advocated for increasing the vestibular depth,⁶ root coverage,⁷ and for augmenting the KT and KM before crown placement.⁸ Nowadays, a FGG is mainly used to recreate/augment KT width,^{9,10} which is most important for dental implants. Indeed, in the 2017 World Workshop, the beneficial role of KM around implants for patient comfort and plaque control was confirmed.¹¹

The introduction of the CTG^{12,13} and the progressive changeover from the FGG to CTG was identified by Zuhr and colleagues as the catalyst for the transition from traditional mucogingival surgery (focused to a greater extent on increas-

ing the tissue thickness and KT width) to periodontal plastic surgery (more esthetic- and patient-centered).¹ CTG acts as a biologic scaffold that improves the stabilization of the flap to the root surface, promoting a greater soft tissue thickness and KT width at the same time.¹⁴ Indeed, it has been shown that CTG is able to induce the keratinization of the overlying epithelium,¹⁵ especially if mainly composed by lamina propria and collagen fibers. On the contrary, a CTG from the deep palate seems not to have the same potential of inducing keratinization as superficial CTG,¹⁶ which may be because of the large amount of adipose and glandular tissue that may act as barriers to the plasmatic diffusion and vascularization during the first phase of healing.¹⁷

With this in mind, it remains controversial in determining the best location for obtaining a graft and the ideal harvesting technique that minimizes a patient's morbidity. Indeed, patient opinions and preferences have slowly influenced decision making in the daily practice and clinical-oriented considerations.¹⁸ Although graft substitutes seem to provide less stable long-term outcomes when compared to the autologous soft tissue graft,¹⁹⁻²¹ efforts have been made for



developing techniques for harvesting a soft tissue graft that minimizes patient discomfort, as well as one that allows healing by the first intention.²²⁻²⁴ Despite the fact that first intention healing is one of the main goals of these approaches, overthinning of the palatal flap is often encountered, leading to wound sloughing and increased patient morbidity.^{22,23,25} Zucchelli et al. showed that a CTG can also be obtained by de-epithelialization of a FGG with similar postoperative morbidity compared to the traditional harvesting approach.²⁵

Nevertheless, the choice of harvesting technique is usually dictated by the anatomy of the site (such as, the palatine artery, shape of the palatal vault, palatal thickness), the required graft thickness, and the clinician's preference.^{1,26}

It has been suggested that the location of the donor site (whether anterior, lateral, superficial, deep palate, or the maxillary tuberosity) can affect the graft shape and its composition.¹ In particular, concerns have been raised regarding the presence and amount of adipose tissue in the CTG that may act as a barrier to plasmatic circulation and impairs the revascularization during the early healing phase.¹⁷ The importance of these consequences has led some clinicians to prefer harvesting from the superficial palate or from the maxillary tuberosity because of the high amount of lamina propria and minimal submucosal tissue (adipose and glandular tissue).^{9,27}

Soft tissue grafts from the tuberosity are increasingly gaining popularity because of their ease in harvesting, the presence of low fatty or glandular tissue, and the richness in connective tissue fibers.^{1,28,29} Additionally, harvesting from the tuberosity presents minimal risk of intraoperative or postoperative complications, resulting in reduced patient morbidity.^{1,30-32} The aim of this commentary is, therefore, to discuss the advantages and disadvantages of harvesting soft tissue grafts from the tuberosity and compare it to the traditional palatal grafts, while highlighting functional-, esthetic-, and patient-related outcomes.

2 | HARVESTING APPROACHES FROM MAXILLARY TUBEROSITY

Although a FGG from the maxillary tuberosity (tFGG) can only be harvested by performing a gingivectomy, a CTG from the tuberosity (tCTG) can be obtained with a distal wedge procedure^{1,33} or by removing a gingival cuff and performing the de-epithelialization extraorally.^{28,31}

The distal wedge technique allows for approximating the mesial and distal flap margins post-harvesting, achieving healing of the donor site by the first intention. However, there is no doubt that performing a gingivectomy is a faster and simpler procedure.³¹ Several authors have described the harvesting of a gingival cuff from the tuberosity, followed by extraoral de-

epithelialization and trimming to perfectly adapt to the recipient site.^{28,31} In addition, tCTG can be split and “opened like a book” to increase the graft width for the treatment of multiple defect areas.³²

However, there are situations in which the amount of tCTG that can be harvested from maxillary tuberosity is limited, such as in the presence of a third molar or following a past periodontal surgery, therefore, harvesting from the lateral palate may be a better choice in these conditions.

It may be concluded that the healing of the tuberosity donor site, whether by first or second intention, is not crucial to patients' postoperative morbidity because the maxillary tuberosity is less exposed to friction during eating and does not come in direct contact with the tongue compared to the palate.³⁰ Moreover, it should be mentioned that greater consumption of analgesics was found to be related to a lower residual soft tissue thickness from the donor site.²⁵ This may also explain the reduced postoperative pain after harvesting from the maxillary tuberosity, which is the thickest area of the oral masticatory mucosa.³⁴

3 | SOFT TISSUE GRAFT FROM THE TUBEROSITY AROUND NATURAL TEETH

The simplicity of the procedure, the limited risk of complications, and the minimal graft shrinkage during healing^{1,31} are main advantages that have contributed to the popularity of tCTG among clinicians. In addition, a tCTG can also be obtained during a distal wedge procedure from the posterior maxillary region for the purpose of treating gingival recessions in other areas as well.³² When compared to a palatal CTG in the bilaminar technique, the tCTG has shown similar mean root coverage outcomes and a greater gain in tissue thickness.³⁰ This tendency was also observed for the FGG.³⁰ Moreover, given the increasing attention to patient perception and their subjective reported outcomes in clinical practice and clinical trials,^{18,35} it is important to highlight that the tuberosity donor site may heal faster than the palatal donor site and leads to much less morbidity (based on visual analog scales and painkiller consumption).³⁰ This may be because of a greater thickness remaining over the bone after the harvesting procedure on the tuberosity sites compared to the palatal donor sites, which is associated with better tolerance of postoperative pain.²⁵

Although a greater tissue thickness provided by a tCTG can be considered beneficial in soft tissue augmentation,³⁶ a tendency for a fibrotic response that can lead to unesthetic outcomes (requiring an additional plastic surgery) has also often been observed following the tCTG.^{31,36} Table 1 summarizes the outcomes of the studies that compared a soft tissue graft from palate and from the maxillary tuberosity.

TABLE 1 Outcomes of the studies comparing soft tissue graft from palate and from maxillary tuberosity

Treatment groups	mRC (mean ± SD) (%)	KT gain (mean ± SD) (mm)	Thickness (mean ± SD) (mm)	Final PES (mean ± SD)	Pain (mean ± SD) during the first 2 weeks (VAS)	Pain (mean ± SD) at 4 and 8 weeks (VAS)	References
tCTG vs SCTG	NA	NA	6.8 ± 1.1 vs 4.9 ± 0.6 (at 1 year)	NA	NA	NA	Dellavia et al. ³⁶
tCTG vs SCTG	NA	0.83 ± 0.61 vs 0.22 ± 0.48*	Comparable from 1 to 5 mm apical to the healing abutment. Significantly higher in favor of tCTG at 6–7 mm apical to the healing abutment	9.15 ± 2.34 vs 10.07 ± 2.19	NA	NA	Rojo et al. ³³
tCTG vs DGG	67 ± 12 vs 62 ± 13	NA	1.29 ± 0.5 vs 2.3 ± 0.6 (at 8 weeks) [†]	NA	2.6 ± 2.2 vs 5.9 ± 2.7 [‡]	0 ± 0 vs 0 ± 0	Amin et al. ³⁰
tFGG vs pFGG	NA	NA	2.7 ± 0.7 vs 2.1 ± 0.7 (at 8 weeks) [†]	NA			Amin et al. ³⁰

Note. tCTG: connective tissue graft from tuberosity; SCTG: subepithelial connective tissue graft (from palate); DGG: de-epithelialized gingival graft (from the palate); tFGG: free gingival graft from tuberosity; pFGG: free gingival graft from palate; mRC: mean root coverage; PES: pink esthetic score; VAS: visual analog scale.

* $P < 0.01$; [†] $P < 0.05$; [‡] $P < 0.001$.

4 | TUBEROSITY SOFT TISSUE GRAFT FOR DENTAL IMPLANTS

The treatment of soft tissue dehiscences around implants poses more challenges and often results in lower outcomes than root coverage in natural teeth.^{37,38} It has been suggested that the graft quality and composition may play a crucial role.^{4,27,28} Indeed, outcomes of 96.3% and 89.6% in mean dehiscence coverage were obtained with a CTG from the superficial palate²⁷ and tCTG,²⁸ respectively (both mainly composed of lamina propria), as opposed to clinical trials that used a subepithelial connective tissue graft (SCTG) from the deep palate (rich in fatty and glandular tissue) that reported a tendency of graft shrinkage over time, as well as less percentage of mean dehiscence coverage.^{37,38} Additionally, the quality of the graft has also been shown to play a role in the long-term stability of the results.^{4,39}

In addition, when soft tissue augmentation is performed for correcting peri-implant volume deficiencies, tCTG was shown to provide a greater KT width gain than the SCTG (and a greater tissue thickness apically)³³ (Table 1). Once again, this result may be explained by the composition of the graft with tCTG being more stable and collagen-rich than the SCTG,³³ that better enables inducing keratinization of the overlying epithelium.^{15,40} In particular, having an adequate amount of KM has been shown to be crucial for maintaining implant health.^{9,41}

To increase tissue thickness, using tCTG can also be successfully applied during an immediate implant placement in the esthetic area, where thickening marginal soft tissue

can prevent recession of the midfacial mucosa.^{42,43} tCTG has demonstrated a mean gain of 0.1 mm in the midbuccal mucosal tissue versus a loss of 0.5 mm in the no-graft site during immediate implant placement.⁴³ Nevertheless, no differences in the pink esthetic score and patient satisfaction were found.⁴³

5 | HISTOLOGIC AND MOLECULAR ANALYSIS OF TUBEROSITY SOFT TISSUE GRAFT

The reason for the different clinical performance between the SCTG and tCTG has also been investigated in histologic studies and at the molecular level.^{29,36} Compared to the SCTG, the tCTG was found to have a denser but less vascularized lamina propria (72.79% for the tCTG vs. 51.08% for the SCTG) whereas the SCTG showed to be richer in submucosal tissue including glandular and adipose components (25.75% vs. 4.89%).^{29,36} Given the evidence suggesting that high amounts of submucosa tissue in the graft may result in more graft shrinkage, less volume gain, and no or minimal effect on epithelium keratinization induction,^{1,33,40} these findings further support the rationale to consider tCTG as a viable (or of higher quality) alternative to palatal CTG, especially when comparing to grafts harvested from the deep palate.¹⁶

On a molecular level, a tendency for decrease in mRNA levels of collagen type I and III (COL-I and COL-III) was



observed in tCTG compared to SCTG and the overall long lysyl hydroxylase 2 (LH2b) mRNA levels was upregulated.³⁶ In addition, a four-fold increase in LH2b/COL-1 ratio has been reported in tCTG, suggesting that tuberosity-derived collagen is less subject to degradation by metalloproteinases.³⁶ This may be the mechanism responsible for collagen accumulation in sites augmented with tCTG. In addition, tCTG was found to exhibit a higher expression of LLH2 antibodies, which are overexpressed in fibrotic tissues. Last, the tendency for higher cytokeratin formation at the epithelia in tCTGs has been speculated that it is related to the potential of inducing keratinization of the peri-implant mucosa.²⁹

Therefore, it can be summarized that tCTG and SCTG illustrate not only different percentages of lamina propria and submucosa, but also gene expression that may explain the variability in clinical outcomes (KT width gain, volume gain, and hyperplastic healing). In particular, tCTG resembles a more a fibrotic tissue with a tendency for a hyperplastic response and, therefore, as suggested by Dellavia et al.,³⁶ its thickness should be limited to <3 mm.

6 | CONCLUSIONS

Within its limitation, the present article highlights the advantages and disadvantages of a soft tissue graft from the maxillary tuberosity, describing its application around natural teeth and dental implants. Clinical, histological, and molecular analyses have shown a different behavior and composition compared with the palatal soft tissue graft, especially from the deep palate. Therefore, clinicians should bear in mind that the maxillary tuberosity is a valid donor site for harvesting a soft tissue graft and it may provide advantages compared with the palate, particularly, when increased KT width, gain in soft tissue thickness, and reduced patient morbidity are primary aims. Nevertheless, this procedure may result in an unesthetic hyperplastic response and in cases with a thick phenotype or when esthetic concerns are of primary concern, a CTG harvested from the palate may provide superior outcomes.

ACKNOWLEDGMENT

This paper was partially supported by the University of Michigan Periodontal Graduate Student Research Fund.

CONFLICT OF INTEREST AND SOURCE OF FUNDING

The authors do not have any financial interests, either directly or indirectly, in the products or information mentioned in the paper.

REFERENCES

- Zuhr O, Baumer D, Hurzeler M. The addition of soft tissue replacement grafts in plastic periodontal and implant surgery: critical elements in design and execution. *J Clin Periodontol*. 2014;41(Suppl 15):S123-S142.
- Cairo F, Rotundo R, Miller PD, Pini Prato GP. Root coverage esthetic score: a system to evaluate the esthetic outcome of the treatment of gingival recession through evaluation of clinical cases. *J Periodontol*. 2009;80:705-710.
- Chambrone L, Tatakis DN. Periodontal soft tissue root coverage procedures: a systematic review from the AAP Regeneration Workshop. *J Periodontol*. 2015;86:S8-S51.
- Zucchelli G, Felice P, Mazzotti C, et al. 5-year outcomes after coverage of soft tissue dehiscence around single implants: a prospective cohort study. *Eur J Oral Implantol*. 2018;11:215-224.
- Nabers JM. Free gingival grafts. *Periodontics*. 1966;4:243-245.
- Fagan F. Clinical comparison of the free soft tissue autograft and partial thickness apically positioned flap—preoperative gingival or mucosal margins. *J Periodontol*. 1975;46:586-595.
- Mlinek A, Smukler H, Buchner A. The use of free gingival grafts for the coverage of denuded roots. *J Periodontol*. 1973;44:248-254.
- Haggerty PC. The use of a free gingival graft to create a healthy environment for full crown preparation. Case history. *Periodontics*. 1966;4:329-331.
- Rocuzzo M, Grasso G, Dalmaso P. Keratinized mucosa around implants in partially edentulous posterior mandible: 10-year results of a prospective comparative study. *Clin Oral Implants Res*. 2016;27:491-496.
- Agudio G, Chambrone L, Pini Prato G. Biologic remodeling of periodontal dimensions of areas treated with gingival augmentation procedure: a 25-year follow-up observation. *J Periodontol*. 2017;88:634-642.
- Berglundh T, Armitage G, Araujo MG, et al. Peri-implant diseases and conditions: consensus report of workgroup 4 of the 2017 world workshop on the classification of periodontal and peri-implant diseases and conditions. *J Periodontol*. 2018;89(Suppl 1):S313-S318.
- Langer B, Calagna LJ. The subepithelial connective tissue graft. A new approach to the enhancement of anterior cosmetics. *Int J Periodontics Restorative Dent*. 1982;2:22-33.
- Langer B, Langer L. Subepithelial connective tissue graft technique for root coverage. *J Periodontol*. 1985;56:715-720.
- Zucchelli G, Mounssif I. Periodontal plastic surgery. *Periodontol 2000*. 2015;68:333-368.
- Karring T, Lang NP, Loe H. The role of gingival connective tissue in determining epithelial differentiation. *J Periodontol Res*. 1975;10:1-11.
- Sculean A, Gruber R, Bosshardt DD. Soft tissue wound healing around teeth and dental implants. *J Clin Periodontol*. 2014;41(Suppl 15):S6-S22.
- Sullivan HC, Atkins JH. Free autogenous gingival grafts. I. Principles of successful grafting. *Periodontics*. 1968;6:121-129.
- McGuire MK, Scheyer ET, Gwaltney C. Commentary: incorporating patient-reported outcomes in periodontal clinical trials. *J Periodontol*. 2014;85:1313-1319.

19. Cairo F, Nieri M, Pagliaro U. Efficacy of periodontal plastic surgery procedures in the treatment of localized facial gingival recessions. A systematic review. *J Clin Periodontol.* 2014;41(Suppl 15):S44-S62.
20. Thoma DS, Buranawat B, Hammerle CH, Held U, Jung RE. Efficacy of soft tissue augmentation around dental implants and in partially edentulous areas: a systematic review. *J Clin Periodontol.* 2014;41(Suppl 15):S77-S91.
21. Tonetti MS, Cortellini P, Pellegrini G, et al. Xenogenic collagen matrix or autologous connective tissue graft as adjunct to coronally advanced flaps for coverage of multiple adjacent gingival recession: randomized trial assessing non-inferiority in root coverage and superiority in oral health-related quality of life. *J Clin Periodontol.* 2018;45:78-88.
22. Edel A. Clinical evaluation of free connective tissue grafts used to increase the width of keratinised gingiva. *J Clin Periodontol.* 1974;1:185-196.
23. Harris RJ. The connective tissue with partial thickness double pedicle graft: the results of 100 consecutively-treated defects. *J Periodontol.* 1994;65:448-461.
24. Hurzeler MB, Weng D. A single-incision technique to harvest subepithelial connective tissue grafts from the palate. *Int J Periodontics Restorative Dent.* 1999;19:279-287.
25. Zucchelli G, Mele M, Stefanini M, et al. Patient morbidity and root coverage outcome after subepithelial connective tissue and de-epithelialized grafts: a comparative randomized-controlled clinical trial. *J Clin Periodontol.* 2010;37:728-738.
26. Tavelli L, Barootchi S, Ravidà A, Oh T-J, Wang H-L. What is the safety zone for palatal soft tissue graft harvesting based upon the locations of the greater palatine artery and foramen? A systematic review. *J Oral Maxillofac Surg.* 2018. <https://doi.org/10.1016/j.joms.2018.10.002>.
27. Zucchelli G, Mazzotti C, Mounssif I, Mele M, Stefanini M, Montebugnoli L. A novel surgical-prosthetic approach for soft tissue dehiscence coverage around single implant. *Clin Oral Implants Res.* 2013;24:957-962.
28. Rocuzzo M, Gaudio L, Bunino M, Dalmaso P. Surgical treatment of buccal soft tissue recessions around single implants: 1-year results from a prospective pilot study. *Clin Oral Implants Res.* 2014;25:641-646.
29. Sanz-Martin I, Rojo E, Maldonado E, Stroppa G, Nart J, Sanz M. Structural and histological differences between connective tissue grafts harvested from the lateral palatal mucosa or from the tuberosity area. *Clin Oral Investig.* 2018. <https://doi.org/10.1007/s00784-018-2516-9>.
30. Amin PN, Bissada NF, Ricchetti PA, Silva APB, Demko CA. Tuberosity versus palatal donor sites for soft tissue grafting: a split-mouth clinical study. *Quintessence Int.* 2018;49:589-598.
31. Jung UW, Um YJ, Choi SH. Histologic observation of soft tissue acquired from maxillary tuberosity area for root coverage. *J Periodontol.* 2008;79:934-940.
32. Hirsch A, Attal U, Chai E, Goultschin J, Boyan BD, Schwartz Z. Root coverage and pocket reduction as combined surgical procedures. *J Periodontol.* 2001;72:1572-1579.
33. Rojo E, Stroppa G, Sanz-Martin I, Gonzalez-Martin O, Alemany AS, Nart J. Soft tissue volume gain around dental implants using autogenous subepithelial connective tissue grafts harvested from the lateral palate or tuberosity area. A randomized controlled clinical study. *J Clin Periodontol.* 2018;45:495-503.
34. Studer SP, Allen EP, Rees TC, Kouba A. The thickness of masticatory mucosa in the human hard palate and tuberosity as potential donor sites for ridge augmentation procedures. *J Periodontol.* 1997;68:145-151.
35. Cairo F, Barbato L, Tonelli P, Batalocco G, Pagavino G, Nieri M. Xenogenic collagen matrix versus connective tissue graft for buccal soft tissue augmentation at implant site. A randomized, controlled clinical trial. *J Clin Periodontol.* 2017;44:769-776.
36. Dellavia C, Ricci G, Pettinari L, Allievi C, Grizzi F, Gagliano N. Human palatal and tuberosity mucosa as donor sites for ridge augmentation. *Int J Periodontics Restorative Dent.* 2014;34:179-186.
37. Burkhardt R, Joss A, Lang NP. Soft tissue dehiscence coverage around endosseous implants: a prospective cohort study. *Clin Oral Implants Res.* 2008;19:451-457.
38. Anderson LE, Inglehart MR, El-Kholy K, Eber R, Wang HL. Implant associated soft tissue defects in the anterior maxilla: a randomized control trial comparing subepithelial connective tissue graft and acellular dermal matrix allograft. *Implant Dent.* 2014;23:416-425.
39. Rocuzzo M, Dalmaso P, Pittoni D, Rocuzzo A. Treatment of buccal soft tissue dehiscence around single implant: 5-year results from a prospective study. *Clin Oral Investig.* 2018. <https://doi.org/10.1007/s00784-018-2634-4>.
40. Ouhayoun JP, Sawaf MH, Gofflaux JC, Etienne D, Forest N. Re-epithelialization of a palatal connective tissue graft transplanted in a non-keratinized alveolar mucosa: a histological and biochemical study in humans. *J Periodontol Res.* 1988;23:127-133.
41. Zigdon H, Machtei EE. The dimensions of keratinized mucosa around implants affect clinical and immunological parameters. *Clin Oral Implants Res.* 2008;19:387-392.
42. Cosyn J, Eghbali A, Hermans A, Vervaeke S, De Bruyn H, Cleymaet R. A 5-year prospective study on single immediate implants in the aesthetic zone. *J Clin Periodontol.* 2016;43:702-709.
43. Zuiderveld EG, Meijer HJA, den Hartog L, Vissink A, Raghoobar GM. Effect of connective tissue grafting on peri-implant tissue in single immediate implant sites: a RCT. *J Clin Periodontol.* 2018;45:253-264.

How to cite this article: Tavelli L, Barootchi S, Greenwell H, Wang HL. Is a soft tissue graft harvested from the maxillary tuberosity the approach of choice in an isolated site? *J Periodontol.* 2019;90:821-825. <https://doi.org/10.1002/JPER.18-0615>