

**REVIEW**

# What's retinoic acid got to do with it? Retinoic acid regulation of the neural crest in craniofacial and ocular development

Antionette L. Williams | Brenda L. Bohnsack 

Department of Ophthalmology and Visual Sciences, Kellogg Eye Center, University of Michigan, Ann Arbor, Michigan

**Correspondence**

Department of Ophthalmology and Visual Sciences, Kellogg Eye Center, University of Michigan, 1000 Wall Street, Ann Arbor, MI 48105.

Email: [brendabo@med.umich.edu](mailto:brendabo@med.umich.edu)

**Summary**

Retinoic acid (RA), the active derivative of vitamin A (retinol), is an essential morphogen signaling molecule and major regulator of embryonic development. The dysregulation of RA levels during embryogenesis has been associated with numerous congenital anomalies, including craniofacial, auditory, and ocular defects. These anomalies result from disruptions in the cranial neural crest, a vertebrate-specific transient population of stem cells that contribute to the formation of diverse cell lineages and embryonic structures during development. In this review, we summarize our current knowledge of the RA-mediated regulation of cranial neural crest induction at the edge of the neural tube and the migration of these cells into the craniofacial region. Further, we discuss the role of RA in the regulation of cranial neural crest cells found within the frontonasal process, periocular mesenchyme, and pharyngeal arches, which eventually form the bones and connective tissues of the head and neck and contribute to structures in the anterior segment of the eye. We then review our understanding of the mechanisms underlying congenital craniofacial and ocular diseases caused by either the genetic or toxic disruption of RA signaling. Finally, we discuss the role of RA in maintaining neural crest-derived structures in postembryonic tissues and the implications of these studies in creating new treatments for degenerative craniofacial and ocular diseases.

**KEYWORDS**

auditory development, cranial neural crest, craniofacial development, ocular development, retinoic acid signaling, vitamin A


## 1 | INTRODUCTION

Retinoic acid (RA) is derived from the liposoluble essential dietary nutrient vitamin A (retinol; Cunningham & Duester, 2015; de la Cruz, Sun, Vangvanichyakorn, & Desposito, 1984; Deltour, Ang, & Duester, 1996; Rosa, Wilk, & Felsey, 1986). Many functions have been assigned to this vitamin, and studies in avian, rodent, frog, and fish models have established that the tight control of RA levels is essential for normal embryonic development, as both vitamin A deficiency and toxic exposure to retinoids lead to a complex spectrum of abnormalities (Cunningham & Duester, 2015; Das et al., 2014; de la Cruz et al.,

1984; Deltour et al., 1996; Rosa et al., 1986; Vieux-Rochas et al., 2010). At the tissue level RA gradients are mediated through the differential and dynamic expression patterns of specific synthesizing (i.e., Raldh2, Raldh3, and Raldh4) and metabolizing enzymes (Cyp26a1, Cyp26b1, and Cyp26c1) (Hernandez, Putzke, Myers, Margaretha, & Moens, 2007; McCaffery, Wagner, O'Neil, Petkovich, & Drager, 1999; Molotkov, Molotkova, & Duester, 2006; Suzuki et al., 2000). RA primarily acts as a ligand for switching the transcription factors of the nuclear retinoid acid receptor (RAR $\alpha/\beta/\gamma$ ) and retinoic X receptor (RXR $\alpha/\beta/\gamma$ ) superfamily from potential repressors to transcriptional activators (Hale et al., 2006; Linville, Radtke, Waxman,

Yelon, & Schilling, 2009; Nicholas Matt et al., 2005; Nicholas Matt, Ghyselinck, Pellerin, & Dupe, 2008; N. Matt, Ghyselinck, Wendling, Chambon, & Mark, 2003). Vertebrate models, from zebrafish to mouse, and experimental tools, such as reporter transgenes, inhibitors of RA synthesis, and selective antagonists for RARs, have been successfully used to decipher retinoid functions during development at the cellular and molecular levels.

Numerous examples from animal models and human diseases highlight the importance of RA in regulating neural crest contributions to craniofacial and ocular structures. The neural crest is a set of transient embryonic stem cells unique to vertebrates that originate at the border of the neural plate spanning the embryo from the diencephalon to the lumbosacral spinal cord. Neural crest cells undergo extensive and coordinated movements away from folds of the neural



**FIGURE 1** Neural crest derivatives in the craniofacial region. (a) The neural crest is a transient population of embryonic stem cells that delaminate from the edge of the neural tube spanning from the diencephalon (Di) to the lumbosacral spinal cord (SpC). Neural crest cells that originate from the diencephalon and anterior mesencephalon (AM) migrate dorsal and ventral to the eye to populate the POM and frontonasal process (FNP). These cells migrate toward regions of high RA levels within the telencephalon (Te), POM, and frontonasal process. Neural crest cells from the posterior mesencephalon (PM) migrate into the first pharyngeal arch. Neural crest cells from the rhombencephalon, which are patterned by a RA gradient, migrate into the first through fourth pharyngeal arches in an anterior (AR) to posterior (PR) pattern. (b) The cranial neural crest is important in the development of the craniofacial skeleton. Neural crest cells in the frontonasal process give rise to the frontal bone (Fr), nasal bone (Na), and philtrum (Ph) in the midline of the face. The anterior portion of the first pharyngeal arch forms the maxillary (Mx) and zygomatic (Zy) bones while the posterior aspect gives rise to the mandible (Mn). The third and fourth pharyngeal arches both contribute to the hyoid (Ny) bone in the neck. (c) Neural crest cells, which are derived from the anterior mesenchyme and diencephalon, populate the periocular mesenchyme and migrate into the anterior segment of the eye. Ocular structures derived from the neural crest include the corneal stroma (CS), corneal endothelium (CEn), trabecular meshwork (TM), sclera (Scl), iris stroma (IS), uveal melanocytes (me), ciliary body muscle (CBM), and the tendons of the extraocular muscles (Tn). The corneal epithelium (CEp) and lens are derived from surface ectoderm while the ciliary body epithelium (CBE), retina (Ret), and retinal pigmented epithelium (RPE) arise from neuroepithelium. Schlemm's canal (SC) and extraocular muscles (EOM) originate from mesoderm. (d) Neural crest cells are also important in middle ear development. Vibrations are transmitted from the external ear, which consists of the pinna, auditory canal, and the tympanic membrane (Tym). The tympanic membrane is attached to the malleus in the middle ear. Both the malleus and the incus originate from the first pharyngeal arch. The vibration is then transmitted to the stapes, which arises from the second pharyngeal arch. The eustachian tube (Eus) connects the middle ear to the nasopharynx

ectoderm to different regions of the embryo to give rise to a broad range of tissues, including myofibroblasts, melanocytes, endocrine cells, neurons, glial cells, cartilage and bone (Barembaum & Bronner-Fraser, 2005; Gammill & Bronner-Fraser, 2003; Knecht & Bronner-Fraser, 2002; Minoux & Rijli, 2010; Morales, Barbas, & Nieto, 2005). Cranial neural crest cells that originate from the diencephalon and anterior mesencephalon form the frontal bone, connective tissue of the orbit, and tendons of the extraocular muscles (Figure 1a–c). Further, a subset of these neural crest cells contribute to the sclera, iris, cornea, ciliary body, and aqueous humor outflow pathways of the eye (Bohnsack & Kahana, 2013; Chawla, Schley, Williams, & Bohnsack, 2016; Gong, 2014; Trainor, 2005; Trainor & Tam, 1995). Cranial neural crest cells that delaminate from the posterior mesencephalon and rhombencephalon give rise to the cartilage and bones of the midface, middle ear, jaw, and neck (Figure 1a,b,d). Cells from the posterior mesencephalon enter into the first pharyngeal arch and eventually form the maxillary and zygomatic bones of the midface. Neural crest cells from the rhombencephalon, which are organized into seven separate rhombomeres, migrate into the corresponding first through fourth pharyngeal arches and ultimately develop into the mandible, teeth, middle ear, and hyoid (Barembaum & Bronner-Fraser, 2005; Gong, 2014; Hong & Saint-Jeannet, 2005; Minoux & Rijli, 2010; Steventon, Carona-Fontaine, & Mayor, 2005; Trainor, 2005; Williams & Bohnsack, 2015). Multiple complex pathways and factors, including RA, regulate neural crest induction, migration, survival and differentiation (Barembaum & Bronner-Fraser, 2005; Gammill & Bronner-Fraser, 2003; Gong, 2014; Hong & Saint-Jeannet, 2005; Minoux & Rijli, 2010; Morales et al., 2005; Steventon et al., 2005; Paul A. Trainor, 2005). Here, we review the main functions of retinoid signaling in regulating the neural crest in the context of craniofacial and ocular development.

## 2 | RA AND CRANIAL NEURAL CREST INDUCTION

Neural crest cell specification occurs between the neural plate border and the non-neural ectoderm during and immediately after gastrulation. As the neural folds come together at the dorsal midline, the neural crest cells that now lie between the overlying epidermis and the neural tube, undergo an epithelial to mesenchymal transition and delaminate from the neuroepithelium (Barembaum & Bronner-Fraser, 2005; Grenier, Teillet, Grifone, Kelly, & Duprez, 2009; Knecht & Bronner-Fraser, 2002). Studies in rodents, birds, frogs, and zebrafish have shown that the neural plate border is primed by BMP signaling gradients, but a second signal, including RA and certain members of the FGF or Wnt family proteins, are required for the induction of *Gbx2*, one of the earliest neural crest markers (Barembaum & Bronner-Fraser, 2005; Knecht & Bronner-Fraser, 2002; LaBonne & Bronner-Fraser, 1998; Li, Lao, & Joyner, 2005; Mayor, Morgan, & Sargent, 1995; Streit & Stern, 1999; von Bubnoff, Schmidt, & Kimelman, 1996; Wawersik, Evola, & Whitman, 2005).

Specifically, RA is important in the anteroposterior patterning of the neural tube and the demarcation of the anterior and posterior neural plates (Papalopulu & Kitner, 1996; Villanueva, Glavic, Ruiz, & Mayor, 2002; Wu, Yang, & Klein, 2005). In frogs, RA suppresses the expression of *ag1* at the anterior border and induces the expression of *snai2*, an important regulator of the epithelial-mesenchymal transition in premigratory neural crest cells at the posterior border (Villanueva et al., 2002). To define the telencephalon, which does not give rise to neural crest cells, endogenous RA levels are tightly regulated by the localized expression of *Cyp26a1* and *Cyp26c1* (Alexandre et al., 1996; Dubey, Rose, Jones, & Saint-Jeannet, 2018; Yu et al., 2016). At the posterior neural plate, *Cyp26a1* expression is inhibited in a RA-dependent fashion via FGF and Wnt signaling resulting in increased RA levels (Kudoh, Wilson, & Dawid, 2002). Exogenous treatment with RA in zebrafish, frogs, and rats during and immediately after gastrulation disrupts the anteroposterior patterning and alters cranial neural crest induction from the diencephalon, mesencephalon, and rhombencephalon (Alexandre et al., 1996; Chawla et al., 2016; Gitton et al., 2010; Lee et al., 1995). In addition, in double *Cyp26a1*  $-/-$  and *Cyp26c1*  $-/-$  knockout mice, the posteriorization of the anterior neural tube results in the absence of neural crest cells from the diencephalon and mesencephalon (Uehara et al., 2007). In these animal studies, increased RA leads to indistinct populations of cranial neural crest cells that are unable to migrate properly from the edge of the neural tube. Therefore, the tight control of RA levels within the developing brain is critical for neural crest induction and the initiation of cell migration into the craniofacial region.

## 3 | RA REGULATION OF CRANIAL NEURAL CREST CELLS IN THE PERIOCCULAR MESENCHYME AND FRONTO NASAL PROCESS

Following induction, cranial neural crest cells undergo epithelial-mesenchymal transition and migrate along defined pathways from the neural tube into the craniofacial region and pharyngeal arches (Trainor, 2005). In zebrafish, time-lapse imaging demonstrated that cranial neural crest cells from the diencephalon follow a migratory stream dorsal to the developing eye while cells originating from the anterior mesencephalon track ventral to the eye; these cells populate the periocular mesenchyme (POM) and then converge in the frontonasal process, which is ventral to the prosencephalon (Figure 1a; Bohnsack & Kahana, 2013; Bohnsack, Kasprick, Kish, Goldman, & Kahana, 2012; Chawla et al., 2016).

During neural crest cell migration, RA activity is highest within the prosencephalon and developing eye (Chawla et al., 2016; Niederreither, Vermot, Fraulob, Chambon, & Dolle, 2002). In zebrafish, the cranial neural crest cells via  $RAR\alpha$  and  $RAR\gamma$  migrate toward the frontonasal process and developing eye, areas of high RA activity, but circumvent areas of low RA activity. Exogenous RA or treatment with  $RAR\alpha$  or  $RAR\gamma$  agonists at the initial stage of cell migration decreases survival and inhibits the migration of the remaining neural crest into the POM

and frontonasal process. This inhibition results in underdeveloped corneas and irides and abnormal midline facial structures. This effect of exogenous RA is in part due to the RAR $\alpha$ -mediated downregulation of the homeobox transcription factor, *pitx2* in neural crest cells migrating from the anterior mesencephalon. In zebrafish, knockdown of *pitx2* inhibits the ventral but not the dorsal stream of cranial neural crest cells (Chawla et al., 2016). As a result, there are fewer neural crest cells in the POM and frontonasal process in *pitx2* knockdown fish. This finding gives insight into the midface hypoplasia and anterior segment dysgenesis observed in humans with Axenfeld-Rieger syndrome (OMIM:180500) due to *PITX2* mutations (Dressler et al., 2010; Hjalt & Semina, 2005; Idrees et al., 2006; Tumer & Bach-Holm, 2009).

The frontonasal process gives rise to midline facial structures, including part of the maxillary bone (philtrum region), the frontal bone and the avian beak. Through RAR $\alpha$  and RAR $\gamma$ , RA regulates the frontonasal skeleton. In mice, the combined knockout of RAR $\alpha$  and RAR $\gamma$  causes midfacial clefts (absence of and incomplete midline fusion of portions of the frontal, maxillary, and sphenoid bones) due to the mislocalization of apoptosis in midline neural crest cells (Dupe & Pellerin, 2009). Further, RA is required for osteoblast differentiation from neural crest cells in the presumptive frontal bone. However, exogenous RA alters the balance between osteoblast and osteoclast activity in bone remodeling resulting in a thickened calvarium (Ferguson, Devarajan, & Atit, 2018). In chick, RA is concentrated in the frontonasal process and induces FGF and SHH signaling in the surrounding ectoderm to initiate beak formation (Schneider & Helms, 2003; Schneider, Hu, Rubenstein, Maden, & Helms, 2001). Again, the tight control of RA levels is critical, as increased RA in the frontonasal process leads to the doubling of midline structures, including the beak (Lee, Fu, Hui, & Richman, 2001).

Few studies have investigated the molecular regulation of neural crest migration and the differentiation of these cells into structures in the anterior segment of the eye. In zebrafish, neural crest cells migrate in two pathways: (a) between the optic cup and surface ectoderm and (b) through the ocular fissure. These neural crest cells coalesce around the lens and give rise to the corneal stroma, the iris stroma, and the aqueous outflow channels (Eason, Williams, Chawla, Apsey, & Bohnsack, 2017). In humans there are three separate migratory waves of neural crest cells that sequentially contribute first to the cornea, then to the iris, and finally to the aqueous outflow tract. However, in the chick, there are two waves of migrating cells, and in zebrafish and mice, there is one continuous influx of neural crest cells into the developing eye. Despite these species differences, the general sequence of anterior segment formation remains consistent (Eason et al., 2017; Gage, Rhoades, Prucka, & Hjalt, 2005; Hay, 1979; Hay & Revel, 1969; Pei & Rhodin, 1970).

In humans, birds, mice, and zebrafish, the genetic or pharmacologic disruption of RA signaling causes corneal, iris and aqueous outflow tract defects which can lead to blindness (Aldahmesh, Khan, Hijazi, & Alkuraya, 2013; Bohnsack et al., 2012; Casey et al., 2011; Fares-Taie et al., 2013; Golzio et al., 2007; Pasutto et al., 2007; Roos et al., 2014; Srour et al., 2013). The local source of RA is the eye itself, as RA-synthesizing enzymes are expressed in the dorsal and ventral


retina and are separated by cranial and caudal *Cyp26c1* expression (Bohnsack et al., 2012; Cvekl & Wang, 2009; Duester, 2009; Molotkov et al., 2006; Williams & Bohnsack, 2015). This specific spatial gene expression pattern creates dorsal and ventral RA gradients centered in and around the eye that through RAR $\beta$  and RAR $\gamma$ , directly target the POM but not the optic cup (fig. 1A0 (Matt et al., 2003, 2005, 2008; Williams & Bohnsack, 2015). In mice and zebrafish, RA increases *Pitx2* expression within the POM, unlike in the early ventral neural crest. This increased *Pitx2* expression in turn regulates the migration and differentiation of ocular neural crest cells through several downstream pathways, including *dlx2* and the canonical Wnt signaling system (Bohnsack et al., 2012; Gage, Qian, Wu, & Rosenberg, 2008; Kumar & Duester, 2010; Zacharias & Gage, 2010).

Taken together, the cranial neural crest that originates from the diencephalon and anterior mesencephalon requires RA at numerous steps during the migration and differentiation of these cells into structures in the midline of the face and anterior segment of the eye. Alteration in RA levels, either increased or decreased, results in congenital malformations such as cleft lip, cleft palate, and anterior segment dysgenesis of the eye.

#### 4 | RA AND PHARYNGEAL ARCH DEVELOPMENT

The pharyngeal arches are a set of six outpouchings that develop on both sides of the pharynx and give rise to numerous head and neck structures including the maxilla, mandible, teeth, middle ear bones, striated facial muscles, and mucous membranes lining the pharynx (Graham, 2003; Graham & Smith, 2001; Mark, Ghyselinck, & Chambon, 2004). Each arch comprises a mesodermal core surrounded by endoderm on the pharynx side and epithelium on the outside. Then, neural crest cells from the posterior mesencephalon and consecutive rhombomeres of the rhombencephalon migrate into the specific arches and interact with the three surrounding germ layers (Figure 2) (Graham, 2003; Graham & Smith, 2001; Mark et al., 2004).

The neural crest cells within the first pharyngeal arch give rise to the maxillary and mandibular bones of the face. In mice and zebrafish, maxillary bone formation is dependent on the epithelial-expressed FGF8-mediated induction of *Dlx1* and *Dlx2* expression in the neural crest (Chai & Maxson, 2006; Minoux & Rijli, 2010; Park et al., 2004; Shigetani et al., 2002). In contrast, the mandible is regulated by the endothelin-1 (Et1)-induced expression of *Dlx5* and *Dlx6* (Charite et al., 2001; Ozeki, Kurihara, Tonami, Watatani, & Kurihara, 2004; Ruest, Xiang, Lim, Levi, & Clouthier, 2004). In zebrafish, exogenous treatment with RA after neural crest induction and delamination suppresses mandible formation and, to a lesser extent, maxilla development. Further, in mice, a pulse of RA during somitogenesis altered the mandible into a maxilla-like structure. In these studies, RA decreased both *Fgf8* and *Et1* expression within their restricted domains in the pharyngeal arch. However, RA had a greater effect on suppressing the expression of *Dlx5* and *Dlx6* compared to that of *Dlx1* and *Dlx2*, which may account for the alteration in neural crest cell fate (Abe, Maeda, &



**FIGURE 2** RA signaling and pharyngeal arch formation. (a) Expression of RA synthesis (*Raldh2*) and degradation (*Cyp26*) enzymes within the rhombencephalon creates a gradient in which RA levels are lowest anteriorly and highest posteriorly. The maxillary versus mandibular fate of neural crest cells within the first pharyngeal arch is dictated by *Dlx1/2* and *Dlx5/6*, respectively. The RA gradient along with spatial expression of RARs regulates the Hox genes within the second through seventh rhombomeres (R2–R7). The specific combination of Hox gene expression patterns the premigratory neural crest cells with the rhombencephalon and confers the identity of the second through fourth pharyngeal arches that is necessary for subsequent differentiation into their respective skeletal, cartilaginous, and nerve elements

Wakisaka, 2008; Ellies, Langille, Martin, Akimenko, & Ekker, 1997; Vieux-Rochas et al., 2007)

The patterning of the neural crest cells within the second through the fourth pharyngeal arches is determined by the expression of Hox (homeobox) transcription factors from premigratory cells in the rhombomeres (Figure 2; Hunt, Gulisano, et al., 1991; Hunt, Whiting, et al., 1991; Lumsden & Krumlauf, 1996; Trainor & Krumlauf, 2000). For example, across mice, frogs, and fish, *Hoxa2* is expressed up to the border of the first and second rhombomeres and is critical in conferring the identity of the second pharyngeal arch. The knockout or downregulation of *Hoxa2* results in the transformation of the second arch into a first arch, leading to the loss of the hyoid bone and the redundancy of the first arch-derived maxillary and mandibular structures (Baltzinger, Ori, Pasqualetti, Nardi, & Rijli, 2005; Gendon-Maguire, Mallo, Zhang, & Gridley, 1993; Hunter & Prince, 2002; Rijli et al., 1993). Similarly, *Hoxa3* and *Hoxa4* (and their paralogs) are important in conferring the identity of the third and fourth pharyngeal arches (Chisaka & Capecchi, 1991; Condie & Capecchi, 1994; Manley & Capecchi, 1995, 1997). However, in the more posterior arches, it is the specific combination of Hox genes that ultimately directs the development of the neural crest from each pharyngeal arch

into their respective skeletal structures (Minoux, Antonarakis, Kmita, Duboule, & Rijli, 2009).

RA is a key regulator of Hox genes within the rhombomeres. The expression of *Cyp26a1*, *Cyp26b1*, and *Cyp26c1* spatially and temporally patterns the rhombencephalon to create a RA gradient with the lowest levels expressed anteriorly and the highest levels expressed posteriorly (Figures 2; Dubey et al., 2018; Mark et al., 2004). In frogs, the differential sensitivity of the various Hox-2 family genes to RA establishes the anterior to posterior identity of the neural crest cells and directs their migration from the rhombomeres into the corresponding pharyngeal arches (Dekker et al., 1992). Further, in zebrafish, chick, and mice, exogenous retinoic acid treatment during gastrulation and early somitogenesis alters the Hox code, transforming rhombomeres 2 and 3 into a more posterior identity. This transformation results in the enlargement of the neural crest-derived hyoid cartilage in zebrafish and the conversion of the trigeminal nerve into a facial nerve in mice (Alexandre et al., 1996; Mark et al., 1995; Marshall et al., 1992; Plant, MacDonald, Grad, Ritchie, & Richman, 2000).

Following neural crest migration from the rhombencephalon into the pharyngeal arches, RA primarily targets the endoderm and ectoderm and, through interactions with the adjacent neural crest, has subsequent effects on these cells (Mark et al., 2004; Matt et al., 2003; Muller et al., 1996; Veitch, Begbie, Schilling, Smith, & Graham, 1999; Wending, Dennefeld, Chambon, & Mark, 2000). The RA gradient established within the rhombencephalon is maintained within the pharyngeal arches, with the lowest concentrations of RA found more anteriorly. Thus, the second pharyngeal arch is only mildly sensitive to RA deficiency, as this structure is absent in *Raldh2* knockout mice, but not in *raldh2* null zebrafish or in *RARα/RARγ* null mice (Begemann, Schilling, Rauch, Geisler, & Ingham, 2001; Niederreither, Subbarayan, Dolle, & Chambon, 1999; Wendling, Ghyselinck, Chambon, & Mark, 2001). On the other hand, exposure to isotretinoin in humans or exogenous RA in mice during gastrulation results in the fusion and hypoplasia of the first and second arches, which is mediated by *RARβ/RXR* heterodimers (Lammer et al., 1985; Matt et al., 2003). The third through the sixth pharyngeal arches are highly sensitive to RA deficiency in humans, rodents, birds, frogs, and zebrafish. Vitamin A deficiency in chick or the genetic knockout of *RARα/RARβ/RARγ* or *Raldh2* in mice and zebrafish causes hypoplasia of the four posterior pharyngeal arches (Begemann et al., 2001; Dupe, Ghyselinck, Wendling, Chambon, & Mark, 1999; Maden, Gale, Kostetskii, & Zile, 1996; Mendelsohn et al., 1994; Niederreither et al., 1999; Niederreither et al., 2003; Vermot, Niederreither, Garnier, Chambon, & Dolle, 2003; Wendling et al., 2001; White, Highland, Kaiser, & Clagett-Dame, 2000). In humans, the effects of vitamin A deficiency on the posterior pharyngeal arches most commonly manifest as cardiac anomalies, such as aortic arch, aorticopulmonary septal, and ventricular septal defects (Mark et al., 2004).

RA deficiency in humans and animals is also associated with congenital hearing loss due to malformations of the bones of the middle ear (Emmett & West, 2014; Romand, 2003). In rodents, birds, frogs, and fish, abnormal rhombencephalon patterning due to alterations in

RA signaling also affects the malleus and incus, which originate from the mandibular portion of the first arch, and the stapes, which arises from the second arch (Kil et al., 2005; Maden et al., 1996; Niederreither et al., 1999; Pittlik, Domingues, Meyer, & Begemann, 2008; White et al., 2000). However, in mice RA-induced transformation of the second arch identity did not cause duplication of the malleus and incus and loss of the stapes. Instead, exogenous RA treatment during early somitogenesis resulted in absence of the malleus and incus and malformation of the stapes. These studies showed that independent of Hox gene-induction, RA directly regulates migration of neural crest cells destined for the middle ear (Mallo, 1997).

Taken together, RA is critical for pharyngeal arch development and subsequent formation of the head and neck. RA is required at multiple steps within the rhombomeres of the brain and within the individual pharyngeal arches. Disruption of RA in humans and animal models can result in craniofacial malformations and hearing loss.

## 5 | CONGENITAL CRANIOFACIAL AND OCULAR DISEASES RESULTING FROM THE DISRUPTION OF RA SIGNALING

Similar to animal studies, both decreases and increases in RA signaling are associated with congenital craniofacial, ear, and ocular anomalies in humans (Gitton et al., 2010; Lampert et al., 2003; Sandell et al., 2007). (Akhtar et al., 2013; Jans et al., 2015; West, 2003; Zile, 1998). The embryo is dependent on maternal sources of vitamin A, which are in the form of retinol, retinyl esters, and  $\beta$ -carotene (Spiegler, Youn-Kyung, Wassef, Shete, & Quadro, 2012). Although uncommon in developed countries, fetal vitamin A deficiency is typically associated with cleft lip, cleft palate, and microphthalmia (small disorganized eye). As detailed throughout this review, animal models including rodents, birds, frogs, and zebrafish have yielded important insight into the pathogenesis of these findings. For example, RAR knockout mice demonstrate that decreased RA signaling impairs midline fusion of facial bones arising from the frontonasal process and the first pharyngeal arch resulting in cleft lip and palate (Begemann et al., 2001; Bohnsack et al., 2012; Bohnsack, Lai, Dolle, & Hischi, 2004; Lohnes et al., 1994; Maden et al., 1996; Matt et al., 2003; Mendelsohn et al., 1994; Niederreither et al., 2002; Niederreither et al., 2003; See & Clagett-Dame, 2009; White et al., 2000)

Decreased RA signaling is also hypothesized to contribute to the craniofacial and ocular defects caused by prenatal alcohol exposure. The most severe form, fetal alcohol syndrome, is characterized by a thin vermilion (thin upper lip), shortened palpebral fissures (decreased eyelid opening), smooth philtrum (no groove between the nose and upper lid), low nasal bridge, micrognathia (small jaw), microphthalmia, and in severe cases, cleft lip and cleft palate (Foroud et al., 2012; Hoyme et al., 2005; Kligenberg et al., 2010; Popova et al., 2016; Riley, Infante, & Warren, 2011). The variability of phenotype in this congenital disease is due to differences in the timing and amount of alcohol exposure during embryogenesis. Bird, rodent, frog, and zebrafish models of fetal alcohol syndrome utilize different exposure

times and at various stages to understand the mechanisms underlying tissue damage (Joya, Garcia-Algar, Vall, & Pujades, 2014; Kiecker, 2016; Petrelli, Bendelac, Hicks, & Fainsod, 2019; Riley et al., 2011; Sulik, Johnson, & Webb, 1981; Sulik, Lauder, & Dehart, 1984; Zhang et al., 2014). The original hypothesis proposed that ethanol is a competitive inhibitor of the aldehyde dehydrogenase enzymes that are also important in the conversion of retinol to RA (Deltour et al., 1996; Duester, 1991). Consistent with this idea, exogenous RA partially rescues the effects of alcohol in zebrafish embryos (Marrs et al., 2010; Muralidharan, Sarmah, & Marrs, 2015). In addition, similar to the disruption of RA signaling, ethanol exposure in chick decreases neural crest cell migration from the rhombencephalon into the pharyngeal arches, resulting in mandibular and maxillary defects (Kiecker, 2016). However, differences in molecular pathways and phenotypes suggest that ethanol may increase or decrease RA in a tissue-specific manner (Reijntjies, Blentic, Gale, & Maden, 2005). In addition, ethanol has been shown to have a number of RA-independent effects, including interfering with SHH, Wnt, and FGF signaling pathways and increasing oxidative stress which may account for the differences in phenotypes between vitamin A deficiency and fetal alcohol syndrome (Ahlgren, Thakur, & Bronner-Fraser, 2002; Aoto, Shikata, Higashiyama, Shiota, & Motoyama, 2008; Ashwell & Zhang, 1996; Chen & Sulik, 1996; Chrisman et al., 2004; Chu, Tong, & de la Monte, 2007; Garic, Flentke, Amberger, Hernandez, & Smith, 2011; Heaton, Paiva, Mayer, & Miller, 2002; Henderson, Devi, Perez, & Schenker, 1995; Koch, Galeotti, Bartoli, & Boveris, 1991; Ornoy & Ergaz, 2010; Videla, Fraga, Koch, & Boveris, 1983; Yelin, Kot, Yelin, & Fainsod, 2007).

Excess maternal vitamin A intake or, more commonly, exposure to RA analogs, such as isotretinoin, cause craniofacial anomalies, including microtia (small ears), hypertelorism (wide spaced eyes), cleft palate, cleft lip, micrognathia (small jaw), and midface hypoplasia (Maden, 2001; Rosa et al., 1986; See & Clagett-Dame, 2009; Zile, 1998). Animal models of RA embryopathy, as described throughout this review, have pinpointed specific time frames in rodents, birds, frogs, and zebrafish, during which different cranial neural crest populations are sensitive to increased RA exposure. For example, increased RA during gastrulation and early somitogenesis decreases neural crest cell migration from the diencephalon and mesencephalon. Consequently, the decrease in neural crest cells in the frontonasal process and first pharyngeal arch accounts for the midface hypoplasia and micrognathia. (Alexandre et al., 1996; Chawla et al., 2016; Gitton et al., 2010; Lee et al., 1995; Uehara et al., 2007). Taken together, these studies demonstrate that short pulses of RA throughout gastrulation and somitogenesis have deleterious effects on the pharyngeal arches and frontonasal process, resulting in irreversible jaw, middle ear, and midface abnormalities that recapitulate the human findings (Abe et al., 2008; Alexandre et al., 1996; Chawla et al., 2016; Dekker et al., 1992; Ellies et al., 1997; Gitton et al., 2010; Hart, McCue, Ragland, Winn, & Unger, 1990; Lee et al., 2001; Lee et al., 1995; Mallo, 1997; Mark et al., 1995; Marshall et al., 1992; Plant et al., 2000; Vieux-Rochas et al., 2007).

In addition to the pharmacologic manipulation of RA levels, genetic alterations in RA signaling are important causes of congenital craniofacial and ocular defects (Adams & Lammer, 1995; de la Cruz et al., 1984; Lammer et al., 1985; Maden, 2001; Rosa et al., 1986). Mutations in genes that are important for regulating RA synthesis (*ALDH1A3*, OMIM:600463; *CHD7*, OMIM:608892), degradation (*CYP26B1*, OMIM:605207), transport (*STRA6*; OMIM:610745), and signaling (*RAR $\beta$* ; OMIM:180220) have all been identified in congenital diseases (Aldahmesh et al., 2013; Casey et al., 2011; Chassaing et al., 2013; Fares-Taie et al., 2013; Golzio et al., 2007; Laue et al., 2011; Micucci et al., 2014; Pagon, Graham Jr., Zonana, & Young, 1981; Pasutto et al., 2007; Roos et al., 2014; Srour et al., 2013; Yahyavi et al., 2013; Yao et al., 2018). Further, known downstream targets of RA within neural crest cells, including *RAI1* (OMIM:607642), *FGFR1* (OMIM:136350), *COL2A1* (OMIM:120140), *TBX1* (OMIM:602054), and *PITX2* (OMIM:601542), have also been associated with craniofacial and ocular anomalies (Bellus et al., 1996; Bohnsack et al., 2012; Chen et al., 2016; Dressler et al., 2010; Gage et al., 2008; Goodship, Cross, LiLing, & Wren, 1998; Greenberg et al., 1996; Harris et al., 2004; Hjalt & Semina, 2005; Janesick, Shiotsugu, Taketani, & Blumberg, 2012; Kumar & Duester, 2010; Lee et al., 2012; Okubo et al., 2011; Patil & Bartley, 1984; Smith et al., 1986; Stickler, Hughes, & Houchin, 2001; Zacharias & Gage, 2010). This list will no doubt continue to expand as additional genetic testing associates other genes important in the RA signaling pathway with rare congenital craniofacial and ocular diseases.

## 6 | RA IN MAINTAINING NEURAL CREST-DERIVED STRUCTURES IN THE ADULT

Studying the molecular mechanisms that underlie development is not only important for understanding the basis of congenital anomalies but also gives insight into the pathogenesis of the degenerative diseases that affect these tissues. Notably, RA is paramount in postnatal tissues. Vitamin A deficiency, reflecting poor diet, malabsorption, or liver disease, is characterized by impaired immunity and hematopoiesis and causes skin rashes and ocular problems, for example, nyctalopia (night blindness), xerophthalmia (decreased tear production), and keratomalacia (corneal thinning and opacification) (Cañete, Cano, Muñoz-Chápuli, & Carmona, 2017; Lai, Ng, & Srinivasan, 2014; Parafita-Fernández et al., 2015). In bone, RA maintains mineral density, as both increased and decreased levels are associated with increased risk of fractures in elderly individuals (Maggio et al., 2006; Melhus et al., 1998; Opatowsky, Bilezikian, & Nif-u, 2004; Promislow, Goodman-Gruen, Slymen, & Barrett-Connor, 2002). RA regulates the balance between osteoblast and osteoclast differentiation and activity; however, the specific effect is dependent on the bone type and location (Lind et al., 2017).

In a recent study, we showed that in neural crest-derived craniofacial and ocular tissues, there is a continued need for RA in maintaining the function and integrity of these structures (Chawla, Swain, Williams, & Bohnsack, 2018). In adult zebrafish, the neural crest-

derived craniofacial bones, including the jaw and frontal bone, are highly sensitive to exogenous RA and the inhibition of endogenous RA synthesis. Within 5 days of altering RA levels, morphogenic changes due to bone remodeling and apoptosis in the jaw and frontal bone were observed. As expected, RA deficiency caused blindness due to the deprivation of substrate for phototransduction in the adult zebrafish retina. However, both increased and decreased RA had additional effects on neural crest-derived anterior segment structures, including the cornea and iridocorneal angle, which resulted in decreased aqueous outflow from the eyes (Chawla et al., 2018). This study represents the first evidence for the tight control of RA levels to maintain the normal structure and function of the anterior segment of the adult eye. Additional studies are required to determine whether RA plays a role in degenerative diseases affecting the anterior segment of the eye, such as primary open angle glaucoma or keratoconus. Moreover, it would also be interesting to determine whether these effects are also observed in mammals. Nevertheless, RA plays a continued role in maintaining neural crest-derived structures in post-embryonic tissues. Further, the deleterious effects of alterations in RA levels have implications in the therapeutic use of retinoids in humans and the environmental toxic exposures to animals.

## 7 | FUTURE DIRECTIONS AND CHALLENGES

RA is critical for craniofacial and ocular development. As a nutrient-based signaling factor, RA levels can be increased or decreased which can unfortunately result in severe birth defects. Vitamin A deficiency continues to be a serious global health issue, especially in regions with limited access to fresh fruits and vegetables and supplemental vitamins. Further, the therapeutic use of retinoids and excess vitamin A is also a public health concern. Numerous studies have worked to characterize both the essential and detrimental roles of RA during embryogenesis. Within craniofacial and ocular development, RA regulates almost all steps of neural crest induction, migration, proliferation, survival, and differentiation. Thus, pharmacologic or genetic alterations in RA signaling cause congenital craniofacial, ear, or eye defects that result in disfiguration, deafness, and blindness. Improving our understanding of the mechanisms that underlie the RA-mediated regulation of neural crest development will lead to increased knowledge of the pathogenesis of these birth defects and provide breakthroughs in treatments for affected individuals.

In addition, studying the signaling pathways important for the migration and differentiation of embryonic neural crest cells may lead to new treatments for degenerative diseases that affect craniofacial and ocular neural crest-derived structures. The signals that are important during embryogenesis, such as RA, have recently been shown to play roles in adult tissues. Modifying or enhancing these pathways may yield new therapeutic targets for adult diseases. In addition, one of the greatest therapeutic challenges within the anterior segment of the eye is the lack of a known neural crest-derived stem cell population that can regenerate cells for cornea, iris, or aqueous outflow

tracts. Determining embryonic markers and signaling pathways within embryonic ocular neural crest cells will lead to the identification of an adult neural crest stem cell population that could potentially restore vision.

## ORCID

Brenda L. Bohnsack  <https://orcid.org/0000-0001-8961-6343>

## REFERENCES

- Abe, M., Maeda, T., & Wakisaka, S. (2008). Retinoic acid affects craniofacial patterning by changing *Fgf8* expression in the pharyngeal ectoderm. *Development, Growth & Differentiation*, *50*, 717–729.
- Adams, J., & Lammer, L. E. (1995). Human isotretinoin exposure: The teratogenesis of a syndrome of congenitive defects. *Neurotoxicology and Teratology*, *17*, 386.
- Ahlgren, S. C., Thakur, V., & Bronner-Fraser, M. (2002). Sonic hedgehog rescues cranial neural crest from cell death induced by ethanol exposure. *Proceedings of the National Academy of Sciences of the United States of America*, *99*, 10476–10481.
- Akhtar, S., Ahmed, A., Randhawa, M. A., Atukorala, S., Arlappa, N., Ismail, T., & Ali, Z. (2013). Prevalence of vitamin a deficiency in South Asia: Causes, outcomes, and possible remedies. *Journal of Health, Population, and Nutrition*, *31*(4), 413–423.
- Aldahmesh, M. A., Khan, A. O., Hijazi, H., & Alkuraya, F. S. (2013). Mutations in *ALDH1A3* cause microphthalmia. *Clinical Genetics*, *84*, 128–131.
- Alexandre, D., Clarke, J. D., Oxtoby, E., Yan, Y. L., Jowett, T., & Holder, N. (1996). Ectopic expression of *Hoxa-1* in the zebrafish alters the fate of the mandibular arch neural crest and phenocopies a retinoic acid-induced phenotype. *Development*, *122*(3), 735–746.
- Aoto, K., Shikata, Y., Higashiyama, D., Shiota, K., & Motoyama, J. (2008). Fetal ethanol exposure activates protein kinase a and impairs *Shh* expression in prechordal mesendoderm cells in the pathogenesis of holoprosencephaly. *Birth Defects Research. Part A, Clinical and Molecular Teratology*, *82*, 224–231.
- Ashwell, K. W., & Zhang, L. L. (1996). Forebrain hypoplasia following acute prenatal ethanol exposure: Quantitative analysis of effects on specific forebrain nuclei. *Pathology*, *23*, 161–166.
- Baltzinger, M., Ori, M., Pasqualetti, M., Nardi, I., & Rijli, F. M. (2005). *Hoxa2* knockdown in *Xenopus* results in hyoid to mandibular homeosis. *Developmental Dynamics*, *234*, 858–867.
- Barembaum, M., & Bronner-Fraser, M. (2005). Early steps in neural crest specification. *Seminars in Cell and Developmental Biology*, *16*, 642–646.
- Begemann, G., Schilling, T. F., Rauch, G.-J., Geisler, R., & Ingham, P. W. (2001). The zebrafish neckless mutation reveals a requirement for *raldh2* in mesodermal signals that pattern the hindbrain. *Development*, *128*, 3081–3094.
- Bellus, G. A., Gaudenz, K., Zackai, E. H., Clarke, L. A., Szabo, J., Francomano, C. A., & Muenke, M. (1996). Identical mutations in three different fibroblast growth factor receptor genes in autosomal dominant craniosynostosis syndromes. *Nature Genetics*, *14*, 174–176.
- Bohnsack, B. L., & Kahana, A. (2013). Thyroid hormone and retinoic acid interact to regulate zebrafish craniofacial neural crest development. *Developmental Biology*, *373*, 300–309.
- Bohnsack, B. L., Kasprick, D., Kish, P. E., Goldman, D., & Kahana, A. (2012). A zebrafish model of Axenfeld-Rieger syndrome reveals that *pitx2* regulation by retinoic acid is essential for ocular and craniofacial development. *Investigative Ophthalmology & Visual Science*, *53*(1), 7–22.
- Bohnsack, B. L., Lai, L., Dolle, P., & Hischi, K. K. (2004). Signaling hierarchy downstream of retinoic acid that independently regulates vascular remodeling and endothelial cell proliferation. *Genes & Development*, *18*, 1345–1358.
- Cañete, A., Cano, E., Muñoz-Chápuli, R., & Carmona, R. (2017). Role of vitamin A/retinoic acid in regulation of embryonic and adult hematopoiesis. *Nutrients*, *9*, e159.
- Casey, J., Kawaguchi, R., Morrissey, M., Sun, H., McGettigan, P., Nielsen, J. E., ... Ennis, S. (2011). First implication of *STRA6* mutations in isolated anophthalmia, microphthalmia, and coloboma: A new dimension to the *STRA6* phenotype. *Human Mutation*, *32*, 1417–1426.
- Chai, Y., & Maxson, R. E. J. (2006). Recent advances in craniofacial morphogenesis. *Developmental Dynamics*, *235*, 2353–2375.
- Charite, J., McFadden, D. G., Merlo, G., Levi, G., Clouthier, D. E., Yanagisawa, M., ... Olson, E. N. (2001). Role of *Dlx6* in regulation of an endothelin-1-dependent *dHAND* branchial arch enhancer. *Genes & Development*, *15*, 3039–3049.
- Chassaing, N., Rague, N., Kariminejad, A., Buffet, A., Ghaderi-Sohi, S., Martinovic, J., & Calvas, P. (2013). Mutation analysis of the *STRA6* gene in isolated and non-isolated anophthalmia/microphthalmia. *Clinical Genetics*, *83*, 244–250.
- Chawla, B., Schley, E., Williams, A. L., & Bohnsack, B. L. (2016). Retinoic acid and *pitx2* regulate early neural crest survival and migration in craniofacial and ocular development. *Birth Defects Research. Part B, Developmental and Reproductive Toxicology*, *107*, 126–135. <https://doi.org/10.1002/bdrb.21177>
- Chawla, B., Swain, W., Williams, A. L., & Bohnsack, B. L. (2018). Retinoic acid maintains function of neural crest-derived ocular and craniofacial structures in adult zebrafish. *Investigative Ophthalmology & Visual Science*, *59*, 1924–1935.
- Chen, L., Tao, Y., Song, F., Yuan, X., Wang, J., & Saffen, D. (2016). Evidence for genetic regulation of mRNA expression of the dosage-sensitive gene retinoic acid induced-1 (*RAI1*) in human brain. *Scientific Reports*, *6*, 19010.
- Chen, S. Y., & Sulik, K. K. (1996). Free radical and ethanol-induced cytotoxicity in neural crest cells. *Alcoholism, Clinical and Experimental Research*, *20*(6), 1071–1076.
- Chisaka, O., & Capecchi, M. R. (1991). Regionally restricted developmental defects resulting from targeted disruption of the mouse homeobox gene *hox-1.5*. *Nature*, *350*, 473–479.
- Chrisman, K., Kenney, R., Comin, J., Thal, T., Suchocki, L., Yueh, Y. G., & Gardner, D. P. (2004). Gestational ethanol exposure disrupts the expression of *FGF8* and sonic hedgehog during limb patterning. *Birth Defects Research. Part A, Clinical and Molecular Teratology*, *70*, 163–171.
- Chu, J., Tong, M., & de la Monte, S. M. (2007). Chronic ethanol exposure causes mitochondrial dysfunction and oxidative stress in immature central nervous system neurons. *Acta Neuropathologica*, *113*, 659–673.
- Condie, B. G., & Capecchi, M. R. (1994). Mice with targeted disruptions in the paralogous genes *hoxa-3* and *hoxd-3* reveal synergistic interactions. *Nature*, *370*, 304–307.
- Cunningham, T. J., & Duester, G. (2015). Mechanisms of retinoic acid signaling and its roles in organ and limb development. *Nature Reviews. Molecular Cell Biology*, *16*(2), 110–123.
- Cvekl, A., & Wang, W.-L. (2009). Retinoic acid signaling in mammalian eye development. *Experimental Eye Research*, *89*, 280–291.
- Das, B. C., Thapa, P., Karki, R., Das, S., Mehapatra, S., Lui, T.-C., ... Evans, T. (2014). Retinoic acid signaling pathways in development and diseases. *Bioorganic & Medicinal Chemistry*, *22*(2), 673–683.
- de la Cruz, E., Sun, S., Vangvanichyakorn, K., & Desposito, F. (1984). Multiple congenital malformations associated with maternal isotretinoin therapy. *Pediatrics*, *74*(3), 428–430.
- Dekker, E.-J., Pannese, M., Houtzager, E., Timmermans, A., Boncinelli, E., & Durston, A. (1992). *Xenopus Hox-2* genes are expressed sequentially after the onset of gastrulation and are differentially inducible by retinoic acid. *Development*, *121*, 339–350.



- Deltour, L., Ang, H. L., & Duester, G. (1996). Ethanol inhibition of retinoic acid synthesis as a potential mechanism for fetal alcohol syndrome. *The FASEB Journal*, 10(9), 1050–1057.
- Dressler, S., Meyer-Marcotty, P., Weisschuh, N., Jablonski-Momeni, A., Pieper, K., Gramer, G., & Gramer, E. (2010). Dental and craniofacial anomalies associated with Axenfeld-Rieger syndrome with PITX2 mutation. *Case Reports in Medicine*, 2010, 621984.
- Dubey, A., Rose, R. E., Jones, D. R., & Saint-Jeannet, J.-P. (2018). Generating retinoic acid gradients by local degradation during craniofacial development: One cell's cue is another cell's poison. *Genesis*, 56(2), e23091.
- Duester, G. (1991). A hypothetical mechanism for fetal alcohol syndrome involving ethanol inhibition of retinoic acid synthesis at the alcohol dehydrogenase step. *Alcoholism, Clinical and Experimental Research*, 15, 568–572.
- Duester, G. (2009). Keeping an eye on retinoic acid signaling during eye development. *Chemico-Biological Interactions*, 178, 178–181.
- Dupe, V., Ghyselinck, N. B., Wendling, O., Chambon, P., & Mark, M. (1999). Key roles of retinoic acid receptors alpha and beta in the patterning of the caudal hindbrain, pharyngeal arches and otocyst in the mouse. *Development*, 126, 5051–5059.
- Dupe, V., & Pellerin, I. (2009). Retinoic acid receptors exhibit cell-autonomous functions in cranial neural crest cells. *Developmental Dynamics*, 238, 2701–2711.
- Eason, J., Williams, A. L., Chawla, B., Apsey, C., & Bohnsack, B. L. (2017). Differences in neural crest sensitivity to ethanol account for the infrequency of anterior segment defects in the eye compared with craniofacial anomalies in a zebrafish model of fetal alcohol syndrome. *Birth Defects Research*, 109, 1212–1227.
- Ellies, D. L., Langille, R. M., Martin, C. C., Akimenko, M. A., & Ekker, M. (1997). Specific craniofacial cartilage dysmorphogenesis coincides with a loss of *dlx* gene expression in retinoic acid-treated zebrafish embryos. *Mechanisms of Development*, 61, 23–36.
- Emmett, S. D., & West, K. P. J. (2014). Gestational vitamin A deficiency: A novel cause of sensorineural hearing loss in the developing world? *Medical Hypotheses*, 82, 6–10.
- Fares-Taie, L. S., G., Chassing, N., Clayton-Smith, J., Hanein, S., Silva, E., ... Rozet, J.-M. (2013). *ALDH1A3* mutations cause recessive anophthalmia and microphthalmia. *American Journal of Human Genetics*, 92, 265–270.
- Ferguson, J. W., Devarajan, M., & Atit, R. P. (2018). Stage-specific roles of *Ezh2* and retinoic acid signaling ensure calvarial bone lineage commitment. *Developmental Biology*, 443, 173–187.
- Foroud, T., Wetherill, L., Vinci-Booher, S., Moore, E. S., Ward, R. E., Hoyme, H. E., ... Jacobson, S. W. (2012). Relation over time between facial measurements and cognitive outcomes in fetal alcohol-exposed children. *Alcoholism, Clinical and Experimental Research*, 36(9), 1634–1646.
- Gage, P. J., Qian, M., Wu, D., & Rosenberg, K. I. (2008). The canonical Wnt signaling antagonist *DKK2* is an essential effector of *PITX2* function during normal eye development. *Developmental Biology*, 317, 310–324.
- Gage, P. J., Rhoades, W., Prucka, S. K., & Hjalt, T. (2005). Fate maps of neural crest and mesoderm in the mammalian eye. *Investigative Ophthalmology & Visual Science*, 46(11), 4200–4208.
- Gammill, L. S., & Bronner-Fraser, M. (2003). Neural crest specification: Migrating into genomics. *Nature Reviews. Neuroscience*, 4, 795–805.
- Garic, A., Flentke, G. R., Amberger, E., Hernandez, M., & Smith, S. M. (2011). *CaMKII* activation is a novel effector of alcohol's neurotoxicity in neural crest stem/progenitor cells. *Journal of Neurochemistry*, 118, 646–657.
- Gendon-Maguire, M., Mallo, M., Zhang, M., & Gridley, T. (1993). *Hoxa-2* mutant mice exhibit homeotic transformation of skeletal elements derived from cranial neural crest. *Cell*, 75, 1317–1331.
- Gitton, Y., Heude, E., Vieux-Rochas, M., Benouaiche, L., Fontaine, A., Sato, T., ... Levi, G. (2010). Evolving maps in craniofacial development. *Seminars in Cell & Developmental Biology*, 21(3), 301–308.
- Golzio, C., Martinovic-Bouriel, J., Thomas, S., Mougou-Zrelli, S., Grattagliano-Bessieres, B., Bonniere, M., ... Etchevers, H. C. (2007). Matthew-wood syndrome is caused by truncating mutations in the retinol-binding protein receptor gene *STRA6*. *American Journal of Human Genetics*, 80, 1179–1187.
- Gong, S.-G. (2014). Cranial neural crest: Migratory cell behavior and regulatory networks. *Experimental Cell Research*, 325, 90–95.
- Goodship, J., Cross, I., LiLing, J., & Wren, C. (1998). A population study of chromosome 22q11 deletions in infancy. *Archives of Disease in Childhood*, 79, 348–351.
- Graham, A. (2003). Development of the pharyngeal arches. *American Journal of Medical Genetics. Part A*, 119A(3), 251–256.
- Graham, A., & Smith, A. (2001). Patterning of the pharyngeal arches. *BioEssays*, 23(1), 54–61.
- Greenberg, F., Lewis, R. A., Potocki, L., Glaze, D., Parke, J., Killian, J., ... Lupski, J. R. (1996). Multi-disciplinary clinical study of Smith-Magenis syndrome (deletion 17p11.2). *American Journal of Medical Genetics*, 62, 247–254.
- Grenier, J., Teillet, M. A., Grifone, R., Kelly, R. G., & Duprez, D. (2009). Relationship between neural crest cells and cranial mesoderm during head muscle development. *PLoS One*, 4(2), e4381.
- Hale, L. A., Tallafuss, A., Yan, Y. L., Dudley, L., Elsen, J. S., & Postlethwait, J. H. (2006). Characterization of the retinoic acid receptor genes *raraa*, *rarab* and *rarg* during zebrafish development. *Gene Expression Patterns*, 6(5), 546–555.
- Harris, L. R., Kamarainen, O.-P., Sevakiivi, M., Miller, G. C., Clarke, J. W., Potter, J. L., & Bridgewater, L. C. (2004). A novel retinoic acid-response element requires an enhancer element mediator for transcriptional activation. *The Biochemical Journal*, 383(Pt1), 37–43.
- Hart, R. C., McCue, P. A., Ragland, W. L., Winn, K. J., & Unger, E. R. (1990). Avian model for 13-cis-retinoic acid embryopathy: Demonstration of neural crest related defects. *Teratology*, 41, 463–472.
- Hay, E. (1979). Development of the vertebrate cornea. *International Review of Cytology*, 63, 263–322.
- Hay, E. D., & Revel, J. P. (1969). Fine structure of the developing avian cornea. *Monographs in Developmental Biology*, 1, 1–144.
- Heaton, M. B., Paiva, M., Mayer, J., & Miller, R. (2002). Ethanol-mediated generation of reactive oxygen species in developing rat cerebellum. *Neuroscience Letters*, 334, 83–86.
- Henderson, G. I., Devi, G. B., Perez, A., & Schenker, S. (1995). In utero ethanol exposure elicits oxidative stress in the rat fetus. *Alcoholism, Clinical and Experimental Research*, 19, 714–720.
- Hernandez, R. E., Putzke, A. P., Myers, J. P., Margaretha, L., & Moens, C. B. (2007). *Cyp26* enzymes generate the retinoic acid response pattern necessary for hindbrain development. *Development*, 134(1), 177–187.
- Hjalt, T. A., & Semina, E. V. (2005). Current molecular understanding of Axenfeld-Rieger. *Expert Reviews in Molecular Medicine*, 7(25), 1–17.
- Hong, C.-S., & Saint-Jeannet, J.-P. (2005). Sox proteins and neural crest development. *Seminars in Cell and Developmental Biology*, 16, 694–703.
- Hoyme, H. E., May, P. A., Kalberg, W. O., Kodituwakku, P., Gossage, J. P., Trujillo, P. M., ... Robinson, L. K. (2005). A practical clinical approach to diagnosis of fetal alcohol syndrome disorders: Clarification of the 1996 institute of medicine criteria. *Pediatrics*, 115(1), 39–47.
- Hunt, P., Gulisano, M., Cook, M., Sham, M. H., Faiella, A., Wilkinson, D., ... Krumlauf, R. (1991). A distinct Hox code for the branchial region of the vertebrate head. *Nature*, 353, 861–864.
- Hunt, P., Whiting, J., Nonchev, S., Sham, M. H., Marshall, H., Graham, A., ... Krumlauf, R. (1991). The branchial Hox code and its implications for gene regulation, patterning of the nervous system and head evolution. *Development*, 2, 63–77.

- Hunter, M. P., & Prince, V. E. (2002). Zebrafish hox paralogue group 2 genes function redundantly as selector genes to pattern the second pharyngeal arch. *Developmental Biology*, 247, 367–389.
- Idrees, F., Bloch-Zupan, A., Free, S. L., Vaideanu, D., Thompson, P. J., Ashley, P., ... Sowden, J. C. (2006). A novel homeobox mutation in the PITX2 gene in a family of Axenfeld-Rieger syndrome associated with brain, ocular, and dental phenotypes. *American Journal of Medical Genetics. Part B, Neuropsychiatric Genetics*, 141B(2), 184–191.
- Janesick, A., Shiotsugu, J., Taketani, M., & Blumberg, G. (2012). RIPPLY3 is a retinoic acid-inducible repressor required for setting the borders of the pre-placodal ectoderm. *Development*, 139, 1213–1224.
- Jans, G., Matthys, C., Bogaerts, A., Lannoo, M., Verhaeghe, J., Van der Schueren, B., & Devlieger, R. (2015). Maternal micronutrient deficiencies and related adverse neonatal outcomes after bariatric surgery: A systematic review. *Advances in Nutrition*, 6, 420–429.
- Joya, X., Garcia-Algar, O., Vall, O., & Pujades, C. (2014). Transient exposure to ethanol during zebrafish embryogenesis results in defects in neuronal differentiation: An alternative model system to study FASD. *PLoS One*, 9(11), e112851.
- Kiecker, C. (2016). The chick embryo as a model for the effects of prenatal exposure to alcohol on craniofacial development. *Developmental Biology*, 415, 314–325.
- Kil, S.-H., Streit, A., Brown, S. T., Agrawal, N., Collazo, A., Zile, M. H., & Groves, A. K. (2005). Distinct roles for hindbrain and paraxial mesoderm in the induction and patterning of the inner ear revealed by a study of vitamin-A-deficient quail. *Developmental Biology*, 285, 252–271.
- Klingenberg, C. P., Wetherill, L., Rogers, J., Moore, E., Ward, R., Autti-Rämö, I., ... Consortium, C. (2010). Prenatal alcohol exposure alters the patterns of facial asymmetry. *Alcohol*, 44(7–8), 649–657.
- Knecht, A. K., & Bronner-Fraser, M. (2002). Induction of the neural crest: A multigene process. *Nature Reviews*, 3, 453–461.
- Koch, O. R., Galeotti, T., Bartoli, G. M., & Boveris, A. (1991). Alcohol-induced oxidative stress in rat liver. *Xenobiotica*, 21, 1077–1084.
- Kudoh, T., Wilson, S. W., & Dawid, I. B. (2002). Distinct roles for Fgf, Wnt and retinoic acid in posteriorizing the neural ectoderm. *Development*, 129, 4335–4346.
- Kumar, S., & Duester, G. (2010). Retinoic acid signaling in perioptic mesenchyme represses Wnt signaling via induction of Pitx2 and Dkk2. *Developmental Biology*, 340, 67–74.
- LaBonne, C., & Bronner-Fraser, M. (1998). Neural crest induction in *Xenopus*: Evidence for a two-signal model. *Development*, 125, 2403–2414.
- Lai, K. L., Ng, J. Y., & Srinivasan, S. (2014). Xerophthalmia and keratomalacia secondary to diet-induced vitamin a deficiency in Scottish adults. *Canadian Journal of Ophthalmology*, 49(1), 109–112.
- Lammer, E. J., Chen, D. T., Hoar, R. M., Agnish, N. D., Benke, P. J., Braun, J. T., ... Sun, S. C. (1985). Retinoic acid embryopathy. *The New England Journal of Medicine*, 313, 837–841.
- Lampert, J. M., Holzschuh, J., Hessel, S., Driever, W., Vogt, K., & von Lintig, J. (2003). Provitamin a conversion to retinal via the beta,beta-carotene-15,15'-oxygenase (bcox) is essential for pattern formation and differentiation during zebrafish embryogenesis. *Development*, 130(10), 2173–2186.
- Laue, K., Pogoda, H. M., Daniel, P. B., van Haeringen, A., Alanay, Y., von Ameln, S., ... Robertson, S. P. (2011). Craniosynostosis and multiple skeletal anomalies in humans and zebrafish result from a defect in the localized degradation of retinoic acid. *American Journal of Human Genetics*, 89(5), 595–606.
- Lee, S. H., Fu, K. K., Hui, J. N., & Richman, J. M. (2001). Noggin and retinoic acid transform the identity of avian facial prominences. *Nature*, 414, 909–912.
- Lee, Y. M., Osumi-Yamashita, N., Ninomiya, Y., Moon, C. K., Eriksson, U., & Eto, K. (1995). Retinoic acid stage-dependently alters the migration and identity of hindbrain neural crest cells. *Development*, 212(3), 825–837.
- Lee, Y.-W., Terranova, C., Birkaya, B., Narla, S., Kehoe, D., Parikh, A., ... Stachowiak, M. K. (2012). A novel nuclear FGF Receptor-1 partnership with retinoid and Nur receptors during developmental gene programming of embryonic stem cells. *Journal of Cellular Biochemistry*, 113, 2920–2936.
- Li, J. Y. H., Lao, Z., & Joyner, A. L. (2005). New regulatory interactions and cellular responses in the isthmus organizer region revealed by altering "Gbx2" expression. *Development*, 132, 1971–1981.
- Lind, T., Öhman, C., Calounova, G., Rasmusson, A., Andersson, G., Pejler, G., & Melhus, H. (2017). Excessive dietary intake of vitamin a reduces skull bone thickness in mice. *PLoS One*, 12(4), e0176217.
- Linville, A., Radtke, K., Waxman, J. S., Yelon, D., & Schilling, T. F. (2009). Combinatorial roles for zebrafish retinoic acid receptors in the hindbrain, limbs and pharyngeal arches. *Developmental Biology*, 325, 60–70.
- Lohnes, D., Mark, M., Mendelsohn, C., Dolle, P., Dierich, A., Gorry, P., ... Chambon, P. (1994). Function of the retinoic acid receptors (RARs) during development (I). Craniofacial and skeletal abnormalities in RAR double mutants. *Development*, 120(10), 2723–2748.
- Lumsden, A., & Krumlauf, R. (1996). Patterning the vertebrate neuraxis. *Science*, 274, 1109–1115.
- Maden, M. (2001). Vitamin a and the developing embryo. *Postgraduate Medical Journal*, 77(910), 489–491.
- Maden, M., Gale, E., Kostetskii, I., & Zile, M. (1996). Vitamin A-deficient quail embryos have half a hindbrain and other neural defects. *Current Biology*, 6, 417–426.
- Maggio, D., Polidori, M. C., Barabani, M., Tufi, A., Ruggiero, C., Cecchetti, R., ... Cherubini, A. (2006). Low levels of carotenoids and retinol in involutional osteoporosis. *Bone*, 38, 244–248.
- Mallo, M. (1997). Retinoic acid disturbs mouse middle ear development in a stage-dependent fashion. *Developmental Biology*, 184, 175–186.
- Manley, N. R., & Capecchi, M. R. (1995). The role of Hoxa-3 in mouse thymus and thyroid development. *Development*, 121, 1989–2003.
- Manley, N. R., & Capecchi, M. R. (1997). Hox group 3 paralogous genes act synergistically in the formation of somitic and neural crest-derived structures. *Developmental Biology*, 192, 274–288.
- Mark, M., Ghyselinck, N. B., & Chambon, P. (2004). Retinoic acid signaling in the development of branchial arches. *Current Opinion in Genetics & Development*, 14(5), 591–598.
- Mark, M., Lohnes, D., Mendelsohn, C., Dupe, V., Vonesch, J.-L., Kastner, P., ... Chambon, P. (1995). Roles of retinoic acid receptors and of Hox genes in the patterning of the teeth and of the jaw skeleton. *The International Journal of Developmental Biology*, 39, 111–125.
- Marrs, J. A., Clendenon, S. G., Ratcliffe, D. R., Fielding, S. M., Liu, Q., & Bosron, W. F. (2010). Zebrafish fetal alcohol syndrome model: Effects of ethanol are rescued by retinoic acid supplement. *Alcohol*, 44, 707–715.
- Marshall, H., Nonchev, S., Sham, M. H., Muchamore, I., Lumsden, A., & Krumlauf, R. (1992). Retinoic acid alters hindbrain Hox code and induces transformation of rhombomeres 2/3 into a 4/5 identity. *Nature*, 360, 737–739.
- Matt, N., Dupe, V., Garnier, J.-M., Dennefeld, C., Chambon, P., Mark, M., & Ghyselinck, N. B. (2005). Retinoic acid-dependent eye morphogenesis is orchestrated by neural crest cells. *Development*, 132, 4789–4800.
- Matt, N., Ghyselinck, N. B., Pellerin, I., & Dupe, V. (2008). Impairing retinoic acid signalling in the neural crest cells is sufficient to alter entire eye morphogenesis. *Developmental Biology*, 320, 140–148.
- Matt, N., Ghyselinck, N. B., Wendling, O., Chambon, P., & Mark, M. (2003). Retinoic acid-induced developmental defects are mediated by RARbeta/RXR heterodimers in the pharyngeal endoderm. *Development*, 130(10), 2083–2093.
- Mayor, R., Morgan, R., & Sargent, M. G. (1995). Induction of the prospective neural crest of *Xenopus*. *Development*, 130, 6441–6452.

- McCaffery, P., Wagner, E., O'Neil, J., Petkovich, M., & Drager, U. C. (1999). Dorsal and ventral retinal territories defined by retinoic acid synthesis, break-down and nuclear receptor expression. *Mechanisms of Development*, 82(1-2), 119-130.
- Melhus, H., Michaëlsson, K., Kindmark, A., Bertgröm, R., Holmberg, L., Mallmin, H., ... Ljunghall, S. (1998). Excessive dietary intake of vitamin a is associated with reduced bone mineral density and increased risk for hip fracture. *Annals of Internal Medicine*, 129(10), 770-778.
- Mendelsohn, C., Lohnes, D., Decimo, D., Lufkin, T., LeMeur, M., Chambon, P., & Mark, M. (1994). Function of the retinoic acid receptors (RARs) during development (II). Multiple abnormalities at various stages of organogenesis in RAR double mutants. *Development*, 120, 2749-2771.
- Micucci, J. A., Layman, W. S., Hurd, E. A., Sperry, E. D., Frank, S. F., Durham, M. A., ... Martin, D. M. (2014). CHD7 and retinoic acid signaling cooperate to regulate neural stem cell and inner ear development in mouse models of CHARGE syndrome. *Human Molecular Genetics*, 23, 434-448.
- Minoux, M., Antonarakis, G. S., Kmita, M., Duboule, E., & Rijli, F. M. (2009). Rostral and caudal pharyngeal arches share a common neural crest ground pattern. *Development*, 136, 637-645.
- Minoux, M., & Rijli, F. M. (2010). Molecular mechanisms of cranial neural crest cell migration and patterning in craniofacial development. *Development*, 137, 2605-2621.
- Molotkov, A., Molotkova, N., & Duester, G. (2006). Retinoic acid guides eye morphogenetic movements via paracrine signaling but is unnecessary for retinal dorsoventral patterning. *Development*, 133(10), 1901-1910.
- Morales, A. V., Barbas, J. A., & Nieto, M. A. (2005). How to become neural crest: From segregation to delamination. *Seminars in Cell & Developmental Biology*, 16, 655-662.
- Muller, T. S., Ebersperger, C., Neubuser, A., Koseki, H., Balling, R., Christ, B., & Wilting, J. (1996). Expression of avian Pax1 and Pax9 is intrinsically regulated in the pharyngeal endoderm, but depends on environmental influences in the paraxial mesoderm. *Developmental Biology*, 178, 403-417.
- Muralidharan, P., Sarmah, S., & Marrs, J. A. (2015). Zebrafish retinal defects induced by ethanol exposure are rescued by retinoic acid and folic acid supplement. *Alcohol*, 49, 149-163.
- Niederreither, K., Subbarayan, V., Dolle, P., & Chambon, P. (1999). Embryonic retinoic acid synthesis is essential for early mouse post-implantation development. *Nature Genetics*, 21, 444-448.
- Niederreither, K., Vermot, J., Fraulob, V., Chambon, P., & Dolle, P. (2002). Retinaldehyde dehydrogenase 2 (RALDH2)-independent patterns of retinoic acid synthesis in the mouse embryo. *Proceedings of the National Academy of Sciences of the United States of America*, 99(25), 16111-16116.
- Niederreither, K., Vermot, J., Le Roux, I., Schuhbaur, B., Chambon, P., & Dolle, P. (2003). The regional pattern of retinoic acid synthesis by RALDH2 is essential for the development of posterior pharyngeal arches and the enteric nervous system. *Development*, 130, 2525-2534.
- Okubo, T., Kawamura, A., Takahashi, J., Yagi, H., Morishima, M., Matsuo, R., & Takada, S. (2011). Ripply3, a Tbx1 repressor, is required for development of the pharyngeal apparatus and its derivatives in mice. *Development*, 138, 339-348.
- Opatowsky, A. R., Bilezikian, J. P., & Nif-u, S. (2004). Serum vitamin a concentration and the risk of hip fracture among women 50 to 74 years old in the United States: A prospective analysis of the NHANES I follow-up study. *American Journal of Medical Genetics*, 117(3), 169-174.
- Ornoy, A., & Ergaz, A. (2010). Alcohol abuse in pregnant women: Effects on the fetus and newborn, mode of action and maternal treatment. *International Journal of Environmental Research and Public Health*, 7, 364-379.
- Ozeki, H., Kurihara, H., Tonami, K., Watatani, S., & Kurihara, H. (2004). Endothelin-1 regulates the dorsoventral branchial arch patterning in mice. *Mechanisms of Development*, 121, 387-395.
- Pagon, R. A., Graham, J. M., Jr., Zonana, J., & Young, S. L. (1981). Coloboma, congenital heart disease and choanal atresia with multiple anomalies: CHARGE association. *The Journal of Pediatrics*, 99, 223-227.
- Papalopulu, N., & Kitner, C. (1996). A posteriorising factor, retinoic acid, reveals that anteroposterior patterning controls the timing of neuronal differentiation in *Xenopus* neuroectoderm. *Development*, 122, 3409-3418.
- Parafita-Fernández, A., Escalona-Fermin, M. M., Sampil, M., Moraña, N., Viso, E., & Fernández-Vila, P. C. (2015). Acquired night blindness due to bad eating patterns. *European Journal of Clinical Nutrition*, 69, 752-754.
- Park, B. K., Sperber, S. M., Choudhury, A., Ghanem, N., Hatch, G. T., Sharpe, P. T., ... Ekker, M. (2004). Intergenic enhancers with distinct activities regulate *dlx* gene expression in the mesenchyme of the branchial arches. *Developmental Biology*, 268, 532-545.
- Pasutto, F., Sticht, H., Hammersen, G., Gillessen-Kaesbach, G., FitzPatrick, D. R., Nurnberg, G., ... Rauch, A. (2007). Mutations in *STRA6* cause a broad spectrum of malformations including anophthalmia, congenital heart defects, diaphragmatic hernia, alveolar capillary dysplasia, lung hypoplasia, and mental retardation. *American Journal of Human Genetics*, 80, 550-560.
- Patil, S. R., & Bartley, J. A. (1984). Interstitial deletion of the short arm of chromosome 17. *Human Genetics*, 67, 237-238.
- Pei, Y. F., & Rhodin, J. A. (1970). The prenatal development of the mouse eye. *The Anatomical Record*, 168, 105-125.
- Petrelli, B., Bendelac, L., Hicks, G. G., & Fainsod, A. (2019). Insights into retinoic acid deficiency and the induction of craniofacial malformations and microcephaly in fetal alcohol spectrum disorder. *Genesis*, 57, e23278.
- Pittlik, S., Domingues, S., Meyer, A., & Begemann, G. (2008). Expression of zebrafish *aldh1a3* (*raldh3*) and absence of *aldh1a1* in teleosts. *Gene Expression Patterns*, 8, 141-147.
- Plant, M. R., MacDonald, M. E., Grad, L. I., Ritchie, S. J., & Richman, J. M. (2000). Locally released retinoic acid repatterns the first branchial arch cartilages in vivo. *Developmental Biology*, 222, 12-26.
- Popova, S., Lange, S., Shield, K., Mihic, A., Chudley, A. e., Mukherjee, R. A. S., ... Rehm, J. (2016). Comorbidity of fetal alcohol spectrum disorder: A systemic review and meta-analysis. *Lancet*, 387, 978-987.
- Promislow, J. H., Goodman-Gruen, D., Slymen, D. J., & Barrett-Connor, E. (2002). Retinol intake and bone mineral density in the elderly: The rancho Bernardo study. *Journal of Bone and Mineral Research*, 17(8), 1349-1358.
- Reijntjes, S., Blentic, A., Gale, E., & Maden, M. (2005). The control of morphogen signaling: Regulation of the synthesis and catabolism of retinoic acid in the developing embryo. *Developmental Biology*, 285, 224-237.
- Rijli, F. M., Mark, M., Lakkaraju, S., Dierich, A., Dolle, P., & Chambon, P. (1993). A homeotic transformation is generated in the rostral branchial region of the head by disruption of *Hoxa-2*, which acts as a selector gene. *Cell*, 75, 1333-1349.
- Riley, E. P., Infante, M. A., & Warren, K. R. (2011). Fetal alcohol spectrum disorders: An overview. *Neuropsychology Review*, 21, 73-80.
- Romand, R. (2003). The roles of retinoic acid during inner ear development. *Current Topics in Developmental Biology*, 57, 261-291.
- Roos, L., Fang, M., Dali, C., Jensen, H., Christoffersen, N., Wu, B., ... Tumer, Z. (2014). A homozygous mutation in a consanguineous family consolidates the role of *ALDH1A3* in autosomal recessive microphthalmia. *Clinical Genetics*, 86, 276-281.
- Rosa, F. W., Wilk, A. L., & Felsey, F. O. (1986). Teratogen update: vitamin A congeners. *Teratology*, 33(3), 355-364.

- Ruest, L. B., Xiang, X., Lim, K. C., Levi, G., & Clouthier, D. E. (2004). Endothelin-a receptor-dependent and -independent signaling pathways in establishing mandibular identity. *Development*, *131*, 4413–4423.
- Sandell, L. L., Sanderson, B. W., Moiseyev, G., Johnson, T., Mushegian, A., Young, K., ... Trainor, P. A. (2007). RDH10 is essential for synthesis of embryonic retinoic acid and is required for limb, craniofacial, and organ development. *Genes & Development*, *21*(9), 1113–1124.
- Schneider, R. A., & Helms, J. A. (2003). The cellular and molecular origins of beak morphology. *Science*, *299*, 565–568.
- Schneider, R. A., Hu, D., Rubenstein, J. L., Maden, M., & Helms, J. A. (2001). Local retinoid signaling coordinates forebrain and facial morphogenesis by maintaining FGF8 and SHH. *Development*, *128*, 1605–1613.
- See, A. W., & Clagett-Dame, M. (2009). The temporal requirement for vitamin a in the developing eye: Mechanism of action in optic fissure closure and new roles for the vitamin in regulating cell proliferation and adhesion in the embryonic retina. *Developmental Biology*, *325*, 94–105.
- Shigetani, Y., Sugahara, F., Kawakami, Y., Murakami, Y., Hirano, S., & Kuratani, S. (2002). Heterotopic shift of epithelial-mesenchymal interactions in vertebrate jaw evolution. *Science*, *296*, 1316–1319.
- Smith, A. C. M., McGavran, L., Robinson, J., Waldstein, G., Macfarlane, J., Zonana, J., ... Magenis, E. (1986). Interstitial deletion of (17)(p11.2p11.2) in nine patients. *American Journal of Medical Genetics*, *24*, 393–414.
- Spiegler, E., Youn-Kyung, K., Wassef, L., Shete, V., & Quadro, L. (2012). Maternal-fetal transfer and metabolism of vitamin a and its precursor b-carotene in the developing tissues. *Biochimica et Biophysica Acta*, *1821*, 88–98.
- Srour, M., Chitayat, D., Caron, V., Chassing, N., Bitoun, P., Patry, L., ... Michaud, J. L. (2013). Recessive and dominant mutations in retinoic acid receptor beta in cases with microphthalmia and diaphragmatic hernia. *American Journal of Human Genetics*, *93*, 765–772.
- Steventon, B., Carona-Fontaine, C., & Mayor, R. (2005). Genetic network during neural crest induction: From cell specification to cell survival. *Seminars in Cell and Developmental Biology*, *16*, 647–654.
- Stickler, G. B., Hughes, W., & Houchin, P. (2001). Clinical features of hereditary progressive arthroophthalmopathy (Stickler syndrome): A survey. *Genetics in Medicine*, *3*, 192–196.
- Streit, A., & Stern, C. D. (1999). Establishment and maintenance of the border of the neural plate in the chick: Involvement of FGF and BMP activity. *Mechanisms of Development*, *82*, 51–66.
- Sulik, K. K., Johnson, M. C., & Webb, M. A. (1981). Fetal alcohol syndrome: Embryogenesis in a mouse model. *Science*, *214*, 936–938.
- Sulik, K. K., Lauder, J. M., & Dehart, D. B. (1984). Brain malformations in prenatal mice following acute maternal ethanol administration. *International Journal of Developmental Neuroscience*, *2*, 203–214.
- Suzuki, R., Shintani, T., Sakuta, H., Kato, A., Ohkawara, T., Osumi, N., & Noda, M. (2000). Identification of RALDH-3 a novel retinaldehyde dehydrogenase, expressed in the ventral region of the retina. *Mechanisms of Development*, *98*(1–2), 37–50.
- Trainor, P. A. (2005). Specification of neural crest cell formation and migration in mouse embryos. *Seminars in Cell and Developmental Biology*, *16*, 683–693.
- Trainor, P. A., & Krumlauf, R. (2000). Patterning the cranial neural crest: Hindbrain segmentation and Hox gene plasticity. *Nature Reviews. Neuroscience*, *1*, 116–124.
- Trainor, P. A., & Tam, P. P. L. (1995). Cranial paraxial mesoderm and neural crest of the mouse embryo-codistribution in the craniofacial mesenchyme but distinct segregation in the branchial arches. *Development*, *121*, 2569–2582.
- Tumer, Z., & Bach-Holm, D. (2009). Axenfeld-Rieger syndrome and spectrum of Pitx2 and Foxc1 mutations. *European Journal of Human Genetics*, *17*, 1527–1539.
- Uehara, M., Yashiro, K., Mamiya, S., Nishino, J., Chambon, P., Dolle, P., & Sakai, Y. (2007). CYP26A1 and CYP26C1 cooperatively regulate anterior-posterior patterning of the developing brain and the production of migratory cranial neural crest cells in the mouse. *Developmental Biology*, *302*, 399–411.
- Veitch, E., Begbie, J., Schilling, T. F., Smith, M. M., & Graham, A. (1999). Pharyngeal arch patterning in the absence of neural crest. *Current Biology*, *9*, 1481–1484.
- Vermot, J., Niederreither, K., Garnier, J. M., Chambon, P., & Dolle, P. (2003). Decreased embryonic retinoic acid synthesis results in a DiGeorge syndrome phenotype in newborn mice. *Proceedings of the National Academy of Sciences of the United States of America*, *100*, 1763–1768.
- Videla, L. A., Fraga, C. G., Koch, O. R., & Boveris, A. (1983). Chemiluminescence of the in situ rat liver after acute ethanol intoxication. Effect of (+)-cyanidanol-3. *Biochemical Pharmacology*, *32*, 2822–2825.
- Vieux-Rochas, M., Bouhali, K., Baudry, S., Fontaine, A., Coen, L., & Levi, G. (2010). Irreversible effects of retinoic acid pulse on *Xenopus* jaw morphogenesis: New insight into cranial neural crest specification. *Birth Defects Research Part B*, *89*, 493–503.
- Vieux-Rochas, M., Coen, L., Sato, T., Kurihara, H., Gitton, Y., Barbieri, O., ... Levi, G. (2007). Molecular dynamics of retinoic acid-induced craniofacial malformations: Implications for the origin of gnathostome jaws. *PLoS One*, *2*, e510.
- Villanueva, S., Glavic, A., Ruiz, P., & Mayor, R. (2002). Posteriorization by FGF, Wnt, and retinoic acid is required for neural crest induction. *Developmental Biology*, *241*, 289–301.
- von Bubnoff, A., Schmidt, J. E., & Kimelman, D. (1996). The *Xenopus laevis* homeogene *Xbbx-2* is an early marker of anteroposterior patterning in the ectoderm. *Mechanisms of Development*, *54*, 149–160.
- Wawersik, S., Evola, C., & Whitman, M. (2005). Conditional BMP inhibition in *Xenopus* reveals stage-specific roles for BMPs in neural and neural crest induction. *Developmental Biology*, *277*, 425–442.
- Wending, O., Dennefeld, C., Chambon, P., & Mark, M. (2000). Retinoid signaling is essential for patterning the endoderm of the third and fourth pharyngeal arches. *Development*, *127*, 1553–1562.
- Wending, O., Ghyselinck, N. B., Chambon, P., & Mark, M. (2001). Roles of retinoic acid receptors in early embryonic morphogenesis and hind-brain patterning. *Development*, *128*, 2031–2038.
- West, K. P. J. (2003). Vitamin a deficiency disorders in children and women. *Food and Nutrition Bulletin*, *24*(4 Suppl), S78–S90.
- White, J. C., Highland, M., Kaiser, M., & Clagett-Dame, M. (2000). Vitamin a deficiency results in the dose-dependent acquisition of anterior character and shortening of the caudal hindbrain of the rat embryo. *Developmental Biology*, *220*, 263–284.
- Williams, A. L., & Bohnsack, B. L. (2015). Neural crest derivatives in ocular development: Discerning the eye of the storm. *Birth Defects Research. Part C, Embryo Today*, *105*(2), 87–95.
- Wu, J., Yang, J., & Klein, P. S. (2005). Neural crest induction by the canonical Wnt pathway can be dissociated from anterior-posterior neural patterning in *Xenopus*. *Developmental Biology*, *279*, 220–232.
- Yahyavi, M., Abouzeid, H., Gawdat, G., de Preux, A.-S., Xiao, T., Bardakjian, T., ... Slavotinek, A. M. (2013). ALDH1A3 loss of function causes bilateral anophthalmia/microphthalmia and hypoplasia of the optic nerve and optic chiasm. *Human Molecular Genetics*, *22*, 3250–3258.
- Yao, H., Hill, S. F., Skidmore, J. M., Sperry, E. D., Swiderski, D. L., Sanchez, G. J., ... Martin, D. M. (2018). CHD7 represses the retinoic acid synthesis enzyme ALDH1A3 during inner ear development. *JCI Insight*, *3*, e97440.
- Yelin, R., Kot, H., Yelin, D., & Fainsod, A. (2007). Early molecular effects of ethanol during vertebrate embryogenesis. *Differentiation*, *75*, 393–403.
- Yu, S.-B., Umair, Z., Kumar, S., Lee, U., Lee, S.-H., Kim, J.-I., ... Kim, J. (2016). xCyp26c induced by inhibition of BMP signaling is involved in

- anterior-posterior neural patterning of *Xenopus laevis*. *Molecules and Cells*, 39, 352–357.
- Zacharias, A. L., & Gage, P. J. (2010). Canonical Wnt/Beta-catenin signaling is required for maintenance but not activation of Pitx2 expression in neural crest during eye development. *Developmental Dynamics*, 239 (12), 3215–3225.
- Zhang, C., Frazier, J. M., Chen, H., Liu, Y., Lee, J.-A., & Cole, G. J. (2014). Molecular and morphological changes in zebrafish following transient ethanol exposure during defined developmental stages. *Neurotoxicology and Teratology*, 44, 70–80.
- Zile, M. H. (1998). Vitamin a and embryonic development: An overview. *The Journal of Nutrition*, 128, 455S–458S.

**How to cite this article:** Williams AL, Bohnsack BL. What's retinoic acid got to do with it? Retinoic acid regulation of the neural crest in craniofacial and ocular development. *genesis*. 2019;57:e23308. <https://doi.org/10.1002/dvg.23308>