

# 2018 International Atmospheric Rivers Conference: Multi-disciplinary studies and high-impact applications of atmospheric rivers

**Short title:** IARC2018 Meeting Report

**Alexandre M. Ramos<sup>(1\*)</sup>, Anna Wilson<sup>(2)</sup>, Michael J. DeFlorio<sup>(2)</sup>, Michael D. Warner<sup>(3)</sup>, Elizabeth Barnes<sup>(4)</sup>, Rene Garreaud<sup>(5)</sup>, Irina V. Gorodetskaya<sup>(6)</sup>, David A. Lavers<sup>(7)</sup>, Benjamin Moore<sup>(8)</sup>, Ashley Payne<sup>(9)</sup>, Chris Smallcomb<sup>(10)</sup>, Harald Sodemann<sup>(11)</sup>, Michael Wehner<sup>(12)</sup>, F. Martin Ralph<sup>(2)</sup>**

(1) Instituto Dom Luiz (IDL), Faculdade de Ciências, Universidade de Lisboa, Portugal

(2) Center for Western Weather and Water Extremes, Scripps Institution of Oceanography, University of California San Diego, USA

(3) US Army Corps of Engineers, Seattle District, USA

(4) Colorado State University, USA

(5) Universidad de Chile, Chile

(6) Centre for Environmental and Marine Studies (CESAM), Department of Physics, University of Aveiro, Portugal

(7) ECMWF, UK

(8) CIRES and NOAA ESRL/PSD, USA

(9) University of Michigan, USA

(10) NWS Reno, USA

(11) University of Bergen, Norway

(12) Lawrence Berkeley National Lab, USA

\* Corresponding author:

Alexandre M. Ramos (amramos@fc.ul.pt)  
Instituto Dom Luiz,  
Faculdade de Ciências, Universidade de Lisboa,  
Campo Grande, Edif. C8, Piso 3, Sala 8.3.1,  
1749-016 Lisboa  
Portugal

Submitted to: **Atmospheric Science Letters**

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1002/asl.935](https://doi.org/10.1002/asl.935)

**Keywords:** Atmospheric Rivers, International Atmospheric Rivers Conference, Meeting Report

**Abstract**

Atmospheric rivers (ARs) play a vital role in shaping the hydroclimate of many regions globally, and can substantially impact water resource management, emergency response planning, and other socioeconomic entities. The second International Atmospheric Rivers Conference took place at the Scripps Institution of Oceanography, University of California San Diego, during 25-28 June 2018, in La Jolla, California, USA. It was sponsored by the Center for Western Weather and Water Extremes (CW3E). A total of 120 people attended the Conference with 94 abstracts submitted and 30 participating students.

In addition to the conference, the Student Forecasting Workshop was organized in the same week. During this workshop, students were exposed to AR forecasting tools, and learned examples of how these tools could be used to make decisions for various applications.

The main goals of this conference were to bring together experts from across the fields of hydrology, atmospheric, oceanic, and polar sciences, as well as water management, civil engineering, and ecology to advance the state of AR science and to explore the future directions for the field. The conference was organized into traditional oral and poster presentations, along with panel discussions and Breakout Groups. This format allowed enhanced interaction between participants, driving progress within the scientific community and the enhanced communication of societal needs by various stakeholders. Several emerging topics of research were highlighted, including subseasonal-to-seasonal (S2S) prediction of ARs and an overview of the AR Reconnaissance campaign. In addition to providing a forum to disseminate and debate new results from scientific talks and posters, the conference was equally effective and useful in linking scientists to

users and decision makers that require improved knowledge on ARs to manage resources and prepare for hazards.

The third International Atmospheric Rivers Conference will be held in Chile in 2020, and hosted by the University of Chile, Santiago.

## **1. Background**

Regions around the globe face challenges in water management due to droughts and/or floods (IPCC, 2018). Since the seminal work of Newell et al. (1992) and Zhu and Newell (1998), research on atmospheric rivers (ARs) has emerged as an interdisciplinary convergence of hydrologists and atmospheric scientists on the transport mechanisms and impacts of precipitation extremes and other significant impacts caused by AR landfall (e.g. Ralph et al 2004; Lavers and Villarini, 2013; Gorodetskaya et al 2014; Paltan et al., 2017; Ramos et al., 2018; Nash et al 2018; Neff 2018, Chen et al., 2018). ARs are a focal point of research and operations within the Center for Western Weather and Water Extremes (CW3E; cw3e.ucsd.edu), as they project strongly onto interannual variations in precipitation over the western U.S. (Dettinger et al., 2011), but their hydrometeorological impacts are substantial in other regions of the globe (Ralph et al., 2017a; Espinoza et al., 2018), including over western Europe (Lavers et al., 2011; Lavers and Villarini, 2013; Ramos et al., 2015; Ramos et al., 2016), western South America (DeFlorio et al., 2018; Viale et al., 2018), polar regions (Gorodetskaya et al., 2014; Nash et al., 2018), and other regions. Emerging research topics in AR science include an intercomparison of AR detection methods (Shields et al., 2018), subseasonal-to-seasonal (S2S) AR prediction (Baggett et al. 2017; DeFlorio et al., 2018; Mundhenk et al., 2018; Nardi et al., 2018; DeFlorio et al., 2019a; DeFlorio et al., 2019b) and the creation of an AR scale to characterize societal impacts (Ralph et al., 2019), among many others.

A large community of scientists coalesced to create a formal definition for ARs by participating in debates at scientific conferences, Town Halls at the American Meteorological Society and American Geophysical Union Annual Meetings, as well as a panel discussion at the 1<sup>st</sup> International Atmospheric Rivers Conference (IARC2016, Ralph et al. 2017a). After years of these discussions and deliberations, the definition of ARs was finally submitted and made available to the Glossary of Meteorology of the American Meteorological Society (**Figure 1**, Ralph et al., 2018a). The definition states that an atmospheric river is: “a long, narrow, and transient corridor of strong horizontal water vapour transport that is typically associated with a low-level jet stream ahead of the cold front of an extratropical cyclone. The water vapour in atmospheric rivers is supplied by tropical and/or extratropical moisture sources. ARs frequently lead to heavy precipitation where they are forced upward—for example, by mountains or by ascent in the warm conveyor belt. Horizontal water vapour transport in the mid-latitudes occurs primarily in ARs and is focused in the lower troposphere. ARs are the largest "rivers" of fresh water on Earth, transporting on average more than double the flow of the Amazon River”. IARC2016 catalysed the preparation of this definition via an energetic panel discussion, which was requested by the American Meteorological Society. In addition, attendees of the IARC2016 strongly supported the idea of holding another IARC at the Scripps Institution of Oceanography in the summer of 2018, which ultimately occurred in June.

## **2. The 2018 International Atmospheric Rivers Conference**

The 2018 International Atmospheric Rivers Conference (IARC2018, <http://cw3e.ucsd.edu/IARC2018/>) took place at the Seaside Forum of the Scripps Institution of Oceanography, 25-28 June, in La Jolla, California, USA. It was hosted and sponsored by CW3E. A total of 120 people attended the IARC2018, in which 94 abstracts were submitted, corresponding to an increase in both the number of attendees and of abstracts when compared with the IARC2016 (105 and 78 respectively, Ralph et al., 2017a). A total of 30 students participated in the IARC2018, 15 of whom received a student scholarship waiving their registration fee.

The main goal of this conference was to bring together experts across the fields of atmospheric, hydrologic, oceanic, and polar sciences, as well as water management, civil engineering, and ecology, to advance the state of the AR science and to explore new directions, improved means of disseminating AR forecast information, and upgrades to existing monitoring techniques.

In pursuit of this goal, the conference was organized with oral presentations (61), poster presentations (33), panel discussions (2), and Breakout Groups (2). The topic of the two panel discussions were “Advances in AR Research for Water Management”, and “AR Definition and New Directions”. Moreover, the two Breakout Groups allowed a thorough discussion on the topics of: “AR Reconnaissance and Data Assimilation” and “Subseasonal to Seasonal (S2S) Challenges and Ways Forward”. The full agenda can be downloaded at the conference webpage: <http://cw3e.ucsd.edu/IARC2018/>.

## 2.1. Sessions summary

The oral presentations, after two introductory talks (one on recent advances on ARs and the other focusing on AR impacts on the Atlantic Ocean), were organized in 12 sessions. These sessions stressed the importance of bringing together climate scientists, engineers, social scientists, impact modellers, and decision-makers to paint a full portrait of complex events. Due to the exceptional U.S. West Coast *Winter of 2016-2017*, a dedicated session was organized, where the large-scale dynamics of extreme precipitation was analysed in terms of atmospheric and oceanic forcing, which led to persistent ARs. The *Applications and Communications* session weighed in on strategic engagement as scientists-communicators in water management, media, and the general population. In addition, the *ARs and Hydrologic Impacts* session’s focus was on the role of ARs in not only extreme precipitation and consequently floods, but also the lack of precipitation, with emphasis on the social-economic impacts of these types of events, e.g. the influence on reservoir storage.

The *Airborne Observations of ARs* session discussed the role of dropsondes in AR predictability, and the advantages of airborne Global Navigation Satellite System -

Radio Occultation observations data assimilation on numerical forecast weather models. Considering the ARs short-range impacts on extreme precipitation and floods, a session was devoted to the *Weather Forecasting of ARs*, where the skill of different numerical weather forecast models, mainly WRF and the ECMWF forecast systems, was quantified for case studies in California and the Iberian Peninsula. In terms of long-term forecast, the *Subseasonal to Seasonal (S2S) Forecasting of ARs* session showed the importance of seasonal forecasting in water management decisions. Several presentations were dedicated to hindcast analysis and experimental real-time forecasting efforts on the prediction skill of ARs at a global and regional scale using different forecasting models.

The ongoing Atmospheric River Tracking Method Intercomparison Project's (ARTMIP) main goal is to understand and quantify uncertainties in AR science based on the choice of the detection/tracking methodology. The climatological characteristics of ARs, such as AR frequency, duration, intensity, and seasonality, are all strongly dependent on the method used to identify ARs (Shields et al., 2018; Ralph et al. 2018b). Taking this into account, a session dedicated to AR Tracking was included in the IARC2018, where the latest results of the ARTMIP project were presented. The remaining presentations focused on novel AR detection schemes made by object-based algorithms and machine learning techniques.

ARs are a global phenomenon, and their frequency is highest in the midlatitude storm track region during a given hemisphere's winter (Guan and Waliser, 2015, Ramos et al., 2016, DeFlorio et al. 2018, 2019a, Viale et al., 2018). In the Regional Perspectives on ARs session, the presentations discussed the importance of ARs in polar regions, as well as extreme precipitation events on the west coast of South America and South Africa, and non-coastal regions of the USA such as Texas or the southern Appalachian Mountains. The *AR Dynamics* session showed the importance of large-scale synoptic dynamics (e.g. Rossby wave breaking or extra-tropical cyclones) to the genesis and evolution of ARs.

The *ARs and Climate Variability: Past, Present, and Future* session focused on paleoclimate trends from model output (Paleoclimate Modelling Intercomparison

Project Phase III, PMIP3), and on geological observations. Present variability was also analysed with a focus on the western USA. Finally, several presentations addressed ARs and future climate scenarios from CMIP5 and high-resolution future climate model simulations, not only at a global scale but also at the regional level.

In the AR Microphysics, Aerosols, and Chemistry and Emerging Directions sessions, several presentations focused on the role of local vs long-range transport of aerosols in precipitation formation within an AR. Furthermore, several presentations showed the utility of stable isotope analysis from precipitation or water vapor in determining the major moisture sources of ARs (local evaporation vs long-range moisture transport). In addition to these topics, another presentation focused on floods following wild fires and showed the relationship between ARs and debris flows in Southern California. This type of event is a clear example of a “compound event” (Zscheischler et al., 2018), where the processes that cause debris flows result from the interaction of two separate extreme events that are spatially and temporally dependent.

## **2.2. Breakout groups and Panel Discussions**

In the “Atmospheric River Reconnaissance and Data Assimilation” group, an update was provided on plans for upcoming seasons. The major conclusions from the breakout group were the following: 1) there is an excellent system in place for flight planning and data generation following the 2016 and 2018 missions; 2) an interagency team of experts has formed to guide work on modelling and data assimilation efforts in ARs with reconnaissance observations; and 3) appropriate time and resources are essential to follow through on the first two items. In the second case, an increase in the storm sample size is necessary. For the last case, additional research is needed in order to choose metrics and data assimilation methods, tested with a variety of models (Reynolds et al., 2019).

The “Subseasonal to Seasonal (S2S) Challenges and Ways Forward” breakout group started the discussion by defining an S2S forecast as a prediction of a variable in the climate system at lead times ranging from 2 weeks to 2 months (Vitart et al. 2017). The major conclusions from this breakout group were: 1) there is high demand and high

potential value to stakeholders and the applications community for skilful S2S forecasts of weather and climate variables (e.g. atmospheric rivers and their associated precipitation); 2) S2S forecasts typically involve an exchange of spatiotemporal forecast precision for potential increased forecast skill at longer lead times; and 3) significant progress has been made in the research community in developing S2S forecast systems and identifying key physical processes (e.g. climate mode teleconnections and atmospheric ridging) that may provide conditional increases in longer-lead S2S forecast skill.

The first panel discussion entitled “AR Research for Water Managers” was moderated by Mike Anderson and panellists included Ben Hatchett, Jeanine Jones, Nina Oakley, and Jonathan Rutz. They discussed ongoing projects in California such as Forecast Informed Reservoir Operations (FIRO), and various projects in water limited areas of the desert southwest, where ARs can have a large impact on water resources, but present challenges for management due to inconsistency in the number of annual ARs.

The second panel discussion was focused on the recently established official AR Definition for the American Meteorological Society (AMS), as well as New Directions for AR science. The panel was moderated by Duane Waliser and panellists included Lance Bosart, Mike Dettinger, Rene Garreaud, F. Martin Ralph, Alexandre Ramos, and Natalia Tilinina. A wide array of new directions for AR science were discussed, including: 1) other types of extreme events in AR dominated areas, like floods following wild fires illustrating the relationship between ARs and debris flows, 2) the use of high temporal and spatial resolution new reanalysis datasets like ERA-5 from ECMWF which can enable the study of physical processes between ARs and the ocean surface and 3) studies using isotope analysis during ARs in order to analyse their sources and the transport of water vapour, which can validate studies that use Lagrangian models to investigate water vapour transport (e.g. Ramos et al., 2016). The discussion then shifted to a more dynamically based one where it was noted that IVT as a detection method has several limitations and that new detection methods should also



include the AR in relationship to large-scale features such as cyclonic activity or linkage with the associated frontal zone (e.g. Viale et al., 2018).

### **2.3 Atmospheric River Forecasting Workshop**

A Student Forecasting Workshop sponsored by the CW3E was organized directly following the IARC2018 on 29-30 June 2018. The workshop brought a smaller group of students (12 out of 30 students) that participated in the IARC2018 together with AR scientists and forecasters to gain hands-on experience with predictions focused on practical and scientific applications. The expected outcomes for participants included improved understanding of modern AR prediction tools and methods, and how AR forecasting supports selected examples of decision-making.

Students participated in two separate interactive lectures and hands-on sessions. The first was on AR predictions and was led by the National Weather Service (Reno and San Diego offices), the San Diego Swift Water Rescue Team, and CW3E. In this session, students learned about forecasting AR impacts as well as how to appropriately communicate various risks to local stakeholders. Students developed their own plans to help decision makers mitigate dangers associated with the forecasted AR event. In the second section, students used CW3E's AR forecasting tools to plan an AR Reconnaissance mission. This section of the workshop gave students an operational perspective on the many challenges and opportunities associated with organizing a large-scale field campaign.

In addition, the Student Forecasting Workshop included a visit to the National Weather Service in San Diego, with a tour given by the Warning Coordination Meteorologist, and a Radiosonde launch from the Scripps pier, where the students participated in preparing and releasing the balloon.

### **3. Outlook**

Overarching conference outcomes include informing the planning of future conferences and colloquia that will best serve the community in terms of ensuring

participation from various disciplines that are impacted by the development of AR science, tools, and applications; cementing collaborative relationships in this new and fast growing community; and linking scientists to users and decision makers that can incorporate improved knowledge on ARs to manage resources and prepare for hazards.

### **3.1 Planning future conferences and colloquium**

The large increase in scientific publications that discuss ARs (Ralph et al., 2017a) and the success of the IARC2018 led to a decision to continue holding the IARC conference on biennial basis, with plans for IARC2020 to be hosted at the Universidad de Chile in Santiago, Chile.

Students who participated in the Student Forecasting Workshop were invited by the organizing committee (Chris Smallcomb, Alexandre M. Ramos, Meredith Fish, Anna Wilson, and Irina Gorodetskaya) to provide feedback that could be applied to the upcoming AR Colloquium Summer School at CW3E. The overarching goal of the Atmospheric Rivers Colloquium Summer School 2019 is to provide the next generation of atmospheric, hydrology and other climate-related scientists with an in-depth look at the cutting-edge techniques in understanding, monitoring, and predicting ARs and their associated high-impact weather. Input from the students was essential to development of the Colloquium framework. Outcomes for participants will include improved understanding of (1) the fundamental dynamics and physics associated with ARs, including their role in the water cycle and impacts on different regions across the globe; (2) the techniques to detect, observe, model, and forecast ARs at all relevant time scales, including in future climate scenarios; and (3) applications of AR science to water management, engineering, and hazard resilience. Fifteen expert instructors, including four from institutions outside of the U.S., will give lectures and lead hands-on exercises. Thirty students are registered from institutions in eleven countries.

### **3.2. Cementing collaborations**

The AR Monograph entitled “Atmospheric Rivers”, which has been developed and written over the past several years, is to be published by Springer International and will be released in 2019. The book is co-edited by: F. Martin Ralph (Chief Editor), Michael D. Dettinger, Jonathan J. Rutz, and Duane Waliser, with the contribution of

international experts on AR research as chapter authors. This Monograph also forms the basic framework of topics covered at the AR Colloquium.

There are many examples of collaborative efforts begun within this community. Multiple side meetings were held with participants in these joint efforts. These include side meetings on Atmospheric River Reconnaissance efforts (in addition to the breakout group), the Atmospheric River Tracking Method Intercomparison Project (ARTMIP), and others. Since IARC2016 these groups have released publications together (Shields et al., 2018; Ralph et al., 2018b; Reynolds et al., 2019; Lavers et al., 2018) and more are in process. These are just a small sample of the collaborative activities fostered by the continued engagement with and dedication to group efforts by members of the AR community.

### **3.3. Linking scientists to users and decision-makers**

Improved understanding, modelling, and prediction of atmospheric rivers is critical to support emergency and water management decisions in the many locations around the globe where they are associated with a majority of the precipitation accumulated on a yearly scale in just a few events (Blamey et al., 2018, Viale et al., 2018; Dettinger et al., 2011). In particular, on the U.S. West Coast significant investment has recently been made aimed at improving prediction of ARs for water resource benefits at time scales ranging from shorter (0-3 day) to subseasonal to seasonal, where the latter would allow for much more robust decision making around resource allocation. At the shorter time scales, if forecasts can be shown to have enough skill, there has been interest in exploring the potential to use them to inform reservoir operations (FIRO, 2017). Given this interest from the applications community, as in IARC2016, several sessions and panels were dedicated to using scientific advances as bases for decision making, developing decision support tools, and supporting sound data driven policy. In 2018 these included, among others, the AR Research for Water Managers panel and Applications and Communications session. The AR Colloquium Summer School has several planned sessions on these topics, given by meteorologists and engineers from the US Army Corps of Engineers and CA Department of Water Resources, who were specifically invited in order to communicate with students about

their needs and the importance of advances in research topics such as S2S scale forecasting. Having this sector integrated into each IARC benefits both the scientists and the stakeholders, as this fortifies lines of communication, builds trust and relationships, and allows for the coproduction of ideas (Vano et al., 2017).

In short, the international community focusing on the topic of ARs continues to grow more vibrant and connected, and we look forward to being involved in its continued development and maturation.

### **Acknowledgments**

The IARC2018 Organizing Committee acknowledges the University of California, San Diego's Scripps Institution of Oceanography's Center for Western Weather and Water Extremes for support, as well as the proactive group of graduate students and postdoctoral scholars, who provided invaluable logistical support throughout the conference.

This publication and work were supported by FCT project UID/GEO/50019/2019 – Instituto Dom Luiz and project IMDROFLOOD funded by Fundação para a Ciência e a Tecnologia, Portugal (FCT, WaterJPI/0004/2014). A. M. Ramos was supported by the Scientific Employment Stimulus 2017 from FCT (CEECIND/00027/2017).

We would like to thank the reviewers for their thoughtful comments and efforts towards improving the manuscript.

## References

Baggett, C.F., Barnes, E., Maloney, E., Mundhenk, B. (2017) Advancing atmospheric river forecasts into subseasonal-to-seasonal time scales. *Geophysical Research Letters*, 44, 7528–7536. doi:10.1002/2017GL074434.

Blamey, R.C., Ramos, A.M., Trigo, R.M., Tome, R., Reason, C.J.C. (2018) The influence of atmospheric rivers over the South Atlantic on winter rainfall in South Africa. *Journal of Hydrometeorology*, 19, 127– 142.

Chen, X., Leung, L. R., Gao, Y., Liu, Y., Wigmosta, M., & Richmond, M. ( 2018). Predictability of extreme precipitation in western U.S. watersheds based on atmospheric river occurrence, intensity, and duration. *Geophysical Research Letters*, 45, 11,693–11,701. <https://doi.org/10.1029/2018GL079831>

DeFlorio, M.J., Waliser, D., Ralph, F.M., Guan, B., Goodman, A., Gibson, P.B., Asharaf, S., Delle Monache, L., Zhang, Z., Subramanian, A.C., Vitart, F., Lin, H., and Kumar, A. (2019b) Experimental Subseasonal-to-Seasonal (S2S) Forecasting of Atmospheric Rivers in a Multi-Model Framework over the Western United States. *Journal of Geophysical Research – Atmospheres*, in prep.

DeFlorio, M.J., Waliser, D., Guan, B., Ralph, F.M., Vitart, F. (2019a) Global evaluation of atmospheric river subseasonal prediction skill. *Climate Dynamics*. doi:10.1007/s00382-018-4309-x.

DeFlorio, M.J., Waliser, D. E., Guan, B., Lavers, D.A., Ralph, F.M. and Vitart F. (2018), Global assessment of atmospheric river prediction skill. *Journal of Hydrometeorology*, 19, 409-426, doi:10.1175/JHM-D-17-0135.1.

Dettinger, M., Ralph, F. M., and Lavers, D. (2015) Setting the stage for a global science of atmospheric rivers. *Eos Transactions American Geophysical Union*, 96. doi:10.1029/2015EO038675

Dettinger, M. D., Ralph, F. M., Das, T., Neiman, P. J., & Cayan, D. R. (2011). Atmospheric rivers, floods and the water resources of California. *Water*, 3(2), 445-478.

Espinoza, V., Waliser, D. E., Guan, B., Lavers, D. A., & Ralph, F. M. (2018). Global analysis of climate change projection effects on atmospheric rivers. *Geophysical Research Letters*, 45(9), 4299-4308.

FIRO, Forecast Informed Reservoir Operations, & Scripps Institution of Oceanography. (2017). Preliminary Viability Assessment of Lake Mendocino. Retrieved from <https://escholarship.org/uc/item/66m803p2>

Gorodetskaya, I.V., Tsukernik, M., Claes, K., Ralph, M.F., Neff, W.D., and Van Lipzig, N.P.M. (2014) The role of atmospheric rivers in anomalous snow accumulation in East Antarctica. *Geophysical Research Letters*, 41(17), 6199–6206. doi:10.1002/2014GL060881

IPCC (2018) Global warming of 1.5°C. An IPCC special report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty [V. Masson-Delmotte, P. Zhai, H. O. Pörtner, D. Roberts, J. Skea, P.R. Shukla, A. Pirani, Y. Chen, S. Connors, M. Gomis, E. Lonnoy, J. B. R. Matthews, W. Moufouma-Okia, C. Péan, R. Pidcock, N. Reay, M. Tignor, T. Waterfield, X. Zhou (eds.)]. In Press.

Lamjiri, M.A., Dettinger, M.D., Ralph, F.M., and Guan, B. (2017) Hourly Storm Characteristics along the U.S. West Coast: Role of Atmospheric Rivers in Extreme Precipitation. *Geophysical Research Letters*, 44. doi:10.1002/2017GL074193

Lavers, D. A., Allan, R. P., Wood, E. F., Villarini, G., Brayshaw, D. J., & Wade, A. J. (2011). Winter floods in Britain are connected to atmospheric rivers. *Geophysical Research Letters*, 38(23).

Lavers, D. A., and Villarini, G. (2013) The nexus between atmospheric rivers and extreme precipitation across Europe, *Geophysical Research Letters*, 40, 3259–3264. doi:10.1002/grl.50636

Lavers, D.A., M.J. Rodwell, D.S. Richardson, F.M. Ralph, J.D. Doyle, C.A. Reynolds, V. Tallapragada, and F. Pappenberger, 2018: The Gauging and Modeling of Rivers in the Sky, *Geophysical Research Letters*, 45, 7828–7834. doi:10.1029/2018GL079019.

Mundhenk, B. D., Barnes, E. A., Maloney, E. D., & Baggett, C. F. (2018). Skillful empirical subseasonal prediction of landfalling atmospheric river activity using the Madden–Julian oscillation and quasi-biennial oscillation. *npj Climate and Atmospheric Science*, 1(1), 7. <https://doi.org/10.1038/s41612-017-0008-2>.

Nardi, K. M., Barnes, E. A., & Ralph, F. M. (2018). Assessment of numerical weather prediction model reforecasts of the occurrence, intensity, and location of atmospheric rivers along the West Coast of North America. *Monthly Weather Review*, 146(10), 3343–3362. <https://doi.org/10.1175/MWR-D-18-0060.1>.

Nash, D., Waliser, D., Guan, B., Ye, H., and Ralph, F. M. (2018) The Role of Atmospheric Rivers in Extratropical and Polar Hydroclimate. *Journal of Geophysical Research: Atmospheres*, 123(13), 6804–6821. doi:10.1029/2017JD028130.

Neff, W. (2018). Atmospheric rivers melt Greenland. *Nature Climate Change*, 8(10), 857–858. <http://doi.org/10.1038/s41558-018-0297-4>

Newell, R. E., Newell, N. E., Zhu, Y., and Scott, C. (1992) Tropospheric rivers? A pilot study. *Geophysical Research Letters*, 19, 2401–2404. doi:10.1029/92GL02916

Paltan, H., Waliser, D., Lim, W.H., Guan, B., Yamazaki, D., Pant, R., and Dadson, S. (2017) Global floods and water availability driven by atmospheric rivers. *Geophysical Research Letters*, 44. doi:10.1002/2017GL07488

Ralph, F.M., Neiman, P.J., and Wick, G.A. (2004) Satellite and CALJET Aircraft Observations of Atmospheric Rivers over the Eastern North Pacific Ocean during the Winter of 1997/98. *Monthly Weather Review*, 132(7), 1721–1745.

Ralph, F.M., Dettinger, M., Lavers, D., Gorodetskaya, I.V., Martin, A., Viale, M., White, A.B., Oakley, N, Rutz, J., Spackman, J.R., Wernli, H. and Cordeira J. (2017a)

Atmospheric Rivers Emerge as a Global Science and Applications Focus. *Bulletin of the American Meteorological Society*, 98, 1969–1973. doi:10.1175/BAMS-D-16-0262.1

Ralph, F.M., S.F. Iacobellis, P.J. Neiman, J.M. Cordeira, J.R. Spackman, D.E. Waliser, G.A. Wick, A.B. White, and C. Fairall (2017b) Dropsonde Observations of Total Integrated Water Vapor Transport within North Pacific Atmospheric Rivers. *Journal of Hydrometeorology*, 18, 2577–2596.

Ralph, F.M., Dettinger, M.D., Cairns, M.M., Galarneau, T.J., and Eylander, J. (2018a) Defining “Atmospheric River”: How the Glossary of Meteorology Helped Resolve a Debate. *Bulletin of the American Meteorological Society*, 99, 837–839. doi:10.1175/BAMS-D-17-0157.1

Ralph, F. M., A. M. Wilson, T. Shulgina, B. Kawzenuk, S. Sellars, J. J. Rutz, M. Asgari-Lamjiri, E. A. Barnes, A. Gershunov, B. Guan, K. Nardi, T. Osborne, and G. A. Wick (2018b) Comparison of Atmospheric River Detection Tools: How Many Atmospheric Rivers Hit Northern California’s Russian River Watershed? *Climate Dynamics*. 10.1007/s00382-018-4427-5.

Ralph, F. M., Rutz, J. J., Cordeira, J. M., Dettinger, M., Anderson, M., Reynolds, D., Schick, L. J., and Smallcomb, C. (2019) A scale to characterize the strength and impacts of atmospheric rivers. *Bulletin of the American Meteorological Society*, 100, 269–289. doi: 10.1175/BAMS-D-18-0023.1

Ramos, A. M., Trigo, R. M., Liberato, M. L., & Tomé, R. (2015). Daily precipitation extreme events in the Iberian Peninsula and its association with atmospheric rivers. *Journal of Hydrometeorology*, 16(2), 579–597.

Ramos, A.M., Nieto, R., Tomé, R., Gimeno, L., Trigo, R.M., Liberato, M.L.R., and Lavers, D.A. (2016) Atmospheric rivers moisture sources from a Lagrangian perspective. *Earth System Dynamics*, 7, 371–384. doi:10.5194/esd-7-371-2016

Ramos, A.M., Blamey, R.C., Algarra, I., Nieto, R., Gimeno, L., Tomé, R., Reason, C.J., and Trigo, R.M. (2018) From Amazonia to southern Africa: atmospheric moisture transport through low-level jets and atmospheric rivers. *Annals of the New York Academy of Sciences*, 1436: 217–230 doi:10.1111/nyas.13960

Reynolds, C. A., J. D. Doyle, F. M. Ralph and R. Demirdjian, 2019: Adjoint sensitivity of North Pacific atmospheric river forecasts. *Mon. Wea. Rev.* (in press Feb 2019).

Shields, C. et al (2018) Atmospheric river tracking method intercomparison project (ARTMIP): project goals and experimental design. *Geoscientific Model Development*, 11:2455–2474.



Vano, J.A., D. Behar, P.W. Mote, D.B. Ferguson, and Panday, R. (2017) Partnerships drive science to action across the AGU community. EOS, doi:10.1029/2017EO088041

Viale, M., Valenzuela, R., Garreaud, R. D., & Ralph, F. M. (2018). Impacts of atmospheric rivers on precipitation in southern South America. *Journal of Hydrometeorology*, 19(10), 1671-1687. <https://doi.org/10.1175/JHM-D-18-0006.1>.


Vitart, F. et al (2017) The subseasonal to seasonal (S2S) prediction project database. *Bulletin of the American Meteorological Society*, 98(1),163–176. doi:10.1175/BAMS-D-16-0017.1

Zhu, Y. and Newell, R.E. (1998) A Proposed Algorithm for Moisture Fluxes from Atmospheric Rivers. *Monthly Weather Review*, 126, 725–735, doi:10.1175/1520-0493(1998)126<0725:APAFMF>2.0.CO;2

Zscheischler, J., Westra, S., van den Hurk, B., Seneviratne, S., Ward, P., Pitman, A., AghaKouchak, A., Bresch, D., Leonard, M., Wahl, T., and Zhang, X. (2018) Future climate risk from compound events. *Nature Climate Change* 8 (6):469-477.

## Figure Captions

**Figure 1.** Schematic summary of the structure and strength of an atmospheric river based on dropsonde measurements deployed from research aircraft across many atmospheric rivers and on corresponding reanalyses that provide the plan-view context. Magnitudes of variables represent an average midlatitude atmospheric river. Average width is based on atmospheric river boundaries defined by vertically integrated water vapour transport (IVT; from surface to 300 hPa) lateral boundary threshold of  $250 \text{ kg m}^{-1} \text{ s}^{-1}$ . Depth corresponds to the altitude below which 75% of IVT occurs. The total water vapour transport (a.k.a. flux) corresponds to the transport along an atmospheric river, bounded laterally by the positions of  $\text{IVT} = 250 \text{ kg m}^{-1} \text{ s}^{-1}$  and vertically by the surface and 300 hPa. (a) Plan view including parent low pressure system and associated cold, warm, and warm-occluded surface fronts. IVT is shown by colour fill (magnitude;  $\text{kg m}^{-1} \text{ s}^{-1}$ ) and direction in the core (white arrow). Vertically integrated water vapour (IWV; cm) is contoured. A representative length scale is shown. The position of the cross section shown in (b) is denoted by the dashed line A–A'. (b) Vertical cross-section perspective, including the core of the water vapour transport in the atmospheric river (orange contours and colour fill) and the pre-cold-frontal low-level jet (LLJ), in the context of the jet-front system and tropopause. Water vapour mixing ratio (green dotted lines;  $\text{g kg}^{-1}$ ) and cross-section-normal isotachs (blue contours;  $\text{m s}^{-1}$ ) are shown. (source: Ralph et al., 2017b, 2018a, [http://glossary.ametsoc.org/wiki/Atmospheric\\_river](http://glossary.ametsoc.org/wiki/Atmospheric_river))



ASL2\_935\_Figure 1.tif