

# Child stunting is associated with child, maternal, and environmental factors in Vietnam

Ty Beal,<sup>1,2</sup> Le Danh Tuyen,<sup>3</sup> Huong Trinh Thi,<sup>4,5</sup> Dharani Burra,<sup>4</sup> Tuyen Huynh,<sup>4</sup> Thanh Thi Duong,<sup>4</sup> Truong Tuyet Mai,<sup>3</sup> Nguyen Duy Son,<sup>3</sup> Kien Tri Nguyen,<sup>4</sup> Stef de Haan,<sup>4</sup> Andrew D Jones<sup>6</sup>

<sup>1</sup>Knowledge Leadership, Global Alliance for Improved Nutrition (GAIN), Washington DC, USA;

<sup>2</sup>Department of Environmental Science and Policy, University of California, Davis, CA, USA;

<sup>3</sup>National Institute of Nutrition (NIN), Hanoi, Vietnam;

<sup>4</sup>International Center for Tropical Agriculture (CIAT)-Asia Office, Hanoi, Vietnam;

<sup>5</sup>Department of Mathematics and Statistics, Thuongmai University, Hanoi, Vietnam; and

<sup>6</sup>Department of Nutritional Sciences, School of Public Health, University of Michigan, Ann Arbor, MI, USA

Address correspondence to Ty Beal: 729 15th Street NW, 8th Floor | Washington, DC 20005, USA. Phone: 602-481-5211. Email: [tbeal@gainhealth.org](mailto:tbeal@gainhealth.org)

Abstract: 250 words

Main text: 4088 words

References: 38

Tables: 1

Figures: 3

Acknowledgements: Authors Le Danh Tuyen, Truong Tuyet Mai and Nguyen Duy Son acknowledge Dr. Tran Thanh Do, former head of the Nutrition Surveillance and Policy Department, NIN for his devoted work and contribution to developing and completing the NSS database.

Source of funding: This study was supported with funding from the Graham Sustainability Institute of the University of Michigan and the CGIAR Research Program on Agriculture for Nutrition and Health (CRP-A4NH). We also acknowledge the Vietnamese Ministry of Health for having funded the primary data collection by NIN and staff from the Centers of Preventive Medicine in each of 63 provinces in Vietnam.

Conflict of interest statement: None declared

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1111/mcn.12826](https://doi.org/10.1111/mcn.12826)

Contributor statement: TB conceptualized the study, drafted the initial manuscript, and had primary responsibility for the final content of the manuscript; LDT, TTM, and NDS designed and facilitated data collection; TB, HTT, and DB analyzed the data; and all authors read and approved the final manuscript.

## Abstract

Child stunting in Vietnam has reduced substantially since the turn of the century but has remained relatively high for several years. We analyzed data on children 6–59 months ( $n = 85932$ ) from the Vietnam Nutritional Surveillance System, a nationally representative cross-sectional survey. Multivariable Poisson regression models were used to estimate relative risk of stunting, stratified by child age and ecological region. Covariates at the child, maternal, household, and environmental levels were included based on available data and the WHO conceptual framework on child stunting. Among children 6–23 months, the strongest associations with child stunting were child age in years (RR: 2.49; 95% CI: 2.26, 2.73), maternal height  $< 145$  cm compared to  $\geq 150$  cm (RR: 2.04; 95% CI: 1.85, 2.26), living in the Northeast compared to the Southeast (RR: 2.01; 95% CI: 1.69, 2.39), no maternal education compared to a graduate education (RR: 1.77; 95% CI: 1.44, 2.16), and birthweight  $< 2500$  g (RR: 1.75; 95% CI: 1.55, 1.98). For children 24–59 months, the strongest associations with child stunting were no maternal education compared to a graduate education (RR: 2.07; 95% CI: 1.79, 2.40), living in the Northeast compared to the Southeast (RR: 1.94; 95% CI: 1.74, 2.16), and maternal height  $< 145$  cm compared to  $\geq 150$  cm (RR: 1.81; 95% CI: 1.69, 1.94). Targeted approaches that address the strongest stunting determinants among vulnerable populations are needed and discussed. Multifaceted approaches outside the health sector are also needed to reduce inequalities in socioeconomic status.

Key words: child stunting, linear growth faltering, determinants, etiology, causes, Vietnam

## Introduction

Child stunting has immediate and long-term consequences including increased morbidity and mortality as well as detrimental impacts on child development and adult health, and it contributes to the intergenerational cycle of malnutrition, hindering economic development (Black et al., 2013; Stewart, Iannotti, Dewey, Michaelsen, & Onyango, 2013). Reducing child stunting is a high priority on the global nutrition and development agenda, a key indicator in the Global Nutrition Targets for 2025 (WHO, 2012) and the Sustainable Development Goal of Zero Hunger (United Nations, Department of Economic and Social Affairs, 2016). Stunting determinants are complex and extensive and vary substantially by local context. To optimally reduce stunting through appropriate policies and programs, there must be an understanding of which stunting determinants are most important nationally and for targetable subpopulations.

The prevalence of child stunting in Vietnam has reduced from 43% in 2000 to 25% in 2015) (Ministry of Health, National Institute of Nutrition, 2012a; Vietnam National Institute of Nutrition, 2018). Vietnam is one of the few countries in Southeast Asia to have reduced malnutrition at nearly the rates set in the Millennium Development Goals (Ministry of Health, National Institute of Nutrition, 2012b). Rapid economic development has helped to improve the nutritional status of the population along with specific nutritional efforts by the Vietnamese government (Ministry of Health, National Institute of Nutrition, 2012b; Trinh Thi, Simioni, & Thomas-Agnan, 2018). For example, the national nutrition strategy outlines goals to reduce hunger and improve dietary quality and recommends the development of nutrition interventions

that target vulnerable populations, including women and children, the poor, and ethnic minorities (Ministry of Health, National Institute of Nutrition, 2012b). Political will to improve nutrition and reduce stunting is strong in Vietnam but more evidence is needed on effective interventions, especially since stunting reduction appears to be stagnating (Vietnam National Institute of Nutrition, 2018).

There is little nationally representative evidence of specific stunting determinants in Vietnam and how they vary by subpopulation. Stunting prevalence varies substantially by region (Viet Nam National Institute of Nutrition, UNICEF, Alive & Thrive, 2014), and regional disparities in socioeconomic conditions and access to education and other services may be driving these differences. A recent analysis of children from select Vietnamese provinces found the following determinants accounted for 90% of the change in height-for-age Z-scores (HAZs) between 2010 and 2014: socioeconomic status (26%), prenatal visits (25%), hygiene (19%), child birth weight (10%), maternal education (7%), and household food insecurity (3%) (P. H. Nguyen et al., 2017). To our knowledge, no multivariable analyses of stunting determinants at the national level have been conducted in Vietnam. We aimed to improve understanding of the determinants of child stunting in Vietnam nationally and among vulnerable subpopulations by assessing associations of hypothesized child-, maternal-, household- and environment-level determinants of stunting in children 6–59 months using nationally representative data from the 2015 Vietnam Nutrition Surveillance System. We used a modified version of the WHO conceptual framework on child stunting (Beal, Tumilowicz, Sutrisna, Izwardy, & Neufeld,

2018), and hypothesized that relationships with stunting would exist at all levels among young Vietnamese children, but child- and maternal-level characteristics would have the strongest relationships, especially among children 6-23 m.

## Key messages

- Under-five child stunting in Vietnam is strongly associated with maternal, child, and environmental factors, particularly short maternal height, low maternal education, and living in a northern region.
- Among children 6-23 months, child age and birthweight are also strongly associated with child stunting.
- There are striking differences in stunting risk factors between regions, which suggests policies and programs aimed at reducing stunting would have more impact if they were tailored geographically.
- Multifaceted approaches outside of the health sector are needed to reduce inequalities in socioeconomic status, which are inextricably linked with malnutrition in Vietnam.

## Methods

### Sampling design and participants

The 2015 Vietnam Nutrition Surveillance System surveyed 99450 children 0-59 months, following a three-stage cluster sampling design to obtain a nationally representative sample. In

the first stage, 30 clusters (communes/wards) were selected from each of 63 provinces using probability proportion to size. Ha Noi and Ho Chi Minh were first stratified by urban and rural before selecting 30 clusters in each locality for both cities. In the second stage, three villages were randomly selected from each cluster using a lottery system or table of random numbers. In the third stage, two children 0–5 months, five children 6–23 months and 10 children 24–59 months were selected from each village using simple random sampling (Viet Nam National Institute of Nutrition, UNICEF, Alive & Thrive, 2014). Sample weights were calculated for each cluster and province using population data from the 2009 Census (General Statistics Office (Vietnam) & United Nations Population Fund (UNFPA), 2009). No sample weights were calculated at the village level because administrative codes to facilitate the match between the census population and survey sample were unavailable. **Figure 1** shows the participant flow chart. Our analysis included 85,932 children 6–59 m. Execution of the National Nutrition Surveillance System in Vietnam has been approved as an integral part of the 2001-2010 National Nutrition Strategy, which was approved by the Prime Minister of Vietnam on 22 February 2001 (Document no. 21/2011/QĐ TTG). The National Institute of Nutrition's Scientific and Ethical Committee and the Ministry of Health's Scientific and Ethical Committee approved the current tools and methodology in 2009.

## Variables

### Outcome

Child stunting ( $HAZ < -2$ ) was the outcome variable in this analysis. Length of children 6-23 months was measured using an infantometer and height of children 24-59 months with a stadiometer to the nearest 0.1 cm. HAZs were calculated using the 2006 WHO child growth standards (WHO, 2006).

### Covariates

An expanded version of the WHO conceptual framework on child stunting (Beal et al., 2018) was used to identify potential covariates from the available data. A total of 28 variables were categorized at the child, maternal, household, and environmental levels. Child characteristics included age, sex, ethnicity, birthweight, if the child had a fever or diarrhea, respectively, in past two weeks, and if the child was given deworming medicine or vitamin A supplements, respectively, in the past six months. Child feeding variables were included for children 6–23 months including currently breastfed; consumed a minimum acceptable diet (MAD) in the past 24 hours; consumed meat, fish, eggs, or milk in the past 24 hours; and consumed vitamin A-rich fruits and vegetables in the past 24 hours. MAD was calculated based on the 2010 WHO guidelines (World Health Organization, 2010), but counting breast milk as a dairy food in the 7-point score for minimum dietary diversity to eliminate the advantage of non-breastfed children (UNICEF, FANTA, USAID, & WHO, 2017). Maternal characteristics included age, height, BMI, education, occupation (farmer or not), washed hands before preparing foods and feeding



child, took iron-folic acids tablets in the past six months, and met with health staff in the past three months. Household-level variables included number of children in the home ( $> 2$ ) and whether iodized salt was currently used. Environmental variables included locality (urban/rural), poor commune, mountainous commune, and ecological region.

### Statistical analysis

All analyses accounted for the study design and sample weights so that results are representative of the national population of children 6–59 months in Vietnam. We excluded biologically implausible values ( $HAZ > +6$  or  $< -6$ ) according to WHO guidelines (Organization, 2009). Children 6–23 months were analyzed separately from children 24–59 months (except for in regional analyses which were limited by sample size) to allow for inclusion of child feeding variables for the younger age group and for a different maternal age categorical cutoff for the two child age groups. All continuous independent variables except for child age were coded into commonly reported categorical levels. Four variables had relatively high levels of missingness but all less than 10%: consumption of vitamin A-rich fruits and vegetables, maternal education, poor commune, and mountainous commune. To include observations with missing values for these categorical variables, we created a new factor level for answers that were not applicable (NAs) but did not report results for this level.

We conducted relative risk (RR) regression with a quasi-Poisson log link function and model-robust standard error estimates (Lumley, Kronmal, & Ma, 2006) to calculate RRs with corresponding 95% confidence intervals (CIs) and P-values, using the ‘survey’ package (Thomas

Lumley & Maintainer Thomas Lumley, 2018) in R (version 3.5.1). We first conducted bivariate analyses for all available variables identified in the conceptual framework (Beal et al., 2018) and then built full multivariable models using these variables. We selected the variables for the final models using a combination of approaches, most of which considered sample weights and complex survey design. Approaches that considered sample weights and complex survey design included bivariate analyses, likelihood ratio tests from the ‘survey’ package (Thomas Lumley & Maintainer Thomas Lumley, 2018), and stepwise (forwards/backwards) model selection from the ‘mass’ package (Ripley et al., 2013) in R (v 3.5.1) using both Akaike information criterion (AIC) and Bayesian information criterion (BIC) (Zhang, 2016). We also used the feature selection approach with the random forest classifier, which did not consider sample weights and complex survey designs (Kursa & Rudnicki, 2010; Liaw & Wiener, 2002). For the random forest-based feature selection, we obtained an optimum value of “mtry” by tuning the full model with 1000 trees, with the improvement parameter set at 1e-05. We partitioned the dataset into 60% training and 40% for model validation using the ‘caret’ package (Max Kuhn et al., 2018) in R (v 3.5.1) and deployed the random forest classifier on the training dataset to calculate a “variable importance” metric for each variable in the full model. Variables obtained from these multiple approaches were assessed and compared with prior knowledge of stunting determinants globally (Beal et al., 2018; Stewart et al., 2013), based on which we selected the final set of variables. For children 6–23 months we included the interaction between breastfeeding status and child age to

account for the influence of reverse causality of breastfeeding and stunting (Kramer, Moodie, Dahhou, & Platt, 2011; Marquis, Habicht, Lanata, Black, & Rasmussen, 1997).

Because the final models differed between the two child age groups, we also built an alternative model with the same variables for each age group. This allowed us to compare effect sizes of included covariates between age groups as well as conduct stratified regional analyses. This model dropped variables that were not available for both age groups (child feeding variables) and retained variables that were included in the final model of either age group. Using this alternative model, we conducted stratified analyses of children 6-59 months for each of the eight regions in Vietnam to understand how stunting determinants varied regionally.

## Results

Among 30771 children 6–23 months, mean HAZ was  $-0.73$  (95% CI:  $-0.77, -0.70$ ) and prevalence of stunting 17.5% (95% CI: 16.8, 18.2; **Table 1**). About half (53%) of children 6–23 months were male, 18% ethnic minorities, 63% from rural areas, and 44% currently breastfed. Among 55161 children 24–59 months, mean HAZ was  $-1.05$  (95% CI:  $-1.08, -1.03$ ) and prevalence of stunting 21.4% (95% CI: 20.7, 22.1). Background characteristics were similar between children 6–23 months and 24–59 months, with the exception of the child experiencing fever in the past two weeks (15% versus 10%, respectively), not being given deworming medication within the past six months (97% versus 56%, respectively), and not being given vitamin A supplements in the past six months (24% versus 30%, respectively).

## Risk factors of stunting among children 6–23 months

In the multivariable analysis, variables with the strongest association with stunting in children 6–23 months included child age, birthweight, maternal height, maternal education, and region (**Figure 2**). An increase in child age of 1 y was associated with a more than two-fold increase in stunting risk (RR: 2.49; 95% CI: 2.26, 2.73). Having a birthweight below 2500 g was associated with a 75% increase in stunting risk (RR: 1.75; 95% CI: 1.55, 1.98). Children of mothers with a height < 145 cm had twice the stunting risk (RR: 2.04; 95% CI: 1.85, 2.26) and height 145–149 cm a 62% increase in risk (RR: 1.62; 95% CI: 1.50, 1.74) compared to children of mothers with a height  $\geq$  150 cm. The risk of stunting increased with decreasing maternal education. Children of mothers with no education (RR: 1.77; 95% CI: 1.44, 2.16) or only primary education (RR: 1.77; 95% CI: 1.48, 2.11) had a 77% increase in stunting risk compared to children of mothers with a graduate education. Compared to children from the Southeast, children from all other regions had significantly increased stunting risk. Risk of stunting in the Northeast was double that in the Southeast (RR: 2.01; 95% CI: 1.69, 2.39), 83% higher in the Northwest (RR: 1.83; 95% CI: 1.49, 2.26), and ~75% higher in the Central North (RR: 1.75; 95% CI: 1.46, 2.10) and Central Highlands (RR: 1.73; 95% CI: 1.43, 2.08). Other variables with moderately strong associations with child stunting included not currently breastfeeding (RR: 1.44; 95% CI: 1.14, 1.82), male sex (RR: 1.35; 95% CI: 1.27, 1.43), ethnic minority (RR: 1.31; 95% CI: 1.19, 1.44), and maternal BMI < 18.5 (RR: 1.20; 95% CI: 1.12, 1.30; Figure 2). The interaction of child age (in years) and

not currently breastfeeding was associated with a reduced risk of stunting (RR: 0.65; 95% CI: 0.55, 0.75).

### Risk factors of stunting among children 24–59 months

Variables with the strongest association with stunting in children 24–59 months in the multivariable analysis included maternal height, maternal education, and region (**Figure 3**).

Children of mothers with a height < 145 cm had an 81% increase in stunting risk (RR: 1.81; 95% CI: 1.69, 1.94) and height 145–149 cm a 49% increase in risk (RR: 1.49; 95% CI 1.41, 1.57) compared to children of mothers with a height  $\geq$  150 cm. Compared to children of mothers with a graduate education, mothers with all other levels of schooling, including a college or university education, had a significantly higher risk of stunting, and risk increased with decreasing education. Risk of stunting for children of mothers with no schooling was double that of children from mothers with a graduate education (RR: 2.07; 95% CI: 1.79, 2.40). Children from the Northeast had twice the stunting risk (RR: 1.94; 95% CI: 1.74, 2.16), Central North a 78% increase (RR: 1.78, 95% CI: 1.56, 2.03), Northwest a 72% increase (RR: 1.72; 95% CI: 1.49, 1.98), and Central Highlands a 68% increase (RR: 1.68; 95% CI: 1.47, 1.91) in risk compared to children from the Southeast. Variables with moderately strong associations with child stunting included ethnic minority (RR: 1.30; 95% CI: 1.21, 1.39), birthweight < 2500 g (RR: 1.43; 95% CI: 1.33, 1.53), maternal age < 20 y (RR: 1.20; 95% CI: 1.15, 1.26), maternal BMI < 18.5 (RR: 1.17; 95% CI: 1.11, 1.25), and > 2 children in the household (RR: 1.18; 95% CI: 1.12, 1.24; Figure 3).

### Comparing risk factors of stunting between younger and older children

The alternative models, which used the same variables for both age groups, allowed us to compare stunting risks between children 6–23 months and 24–59 months (**Supplemental Figures 1 and 2**). We do not report exact risk values because the models included a suboptimal set of covariates, but they were suitable for comparing risks between age groups. Child age, male sex, and birthweight < 2500 g were much more strongly associated with stunting among children 6–23 months than 24–59 months—child age was not meaningfully associated with stunting in the older age group. Sex differences in stunting risk disappeared by age ~3 y (RR in children 36–59 m: 1.02; 95% CI: 0.97, 1.08). Short maternal height was a larger risk factor and maternal education a smaller risk factor for children 6–23 months than 24–59 m. Risk of stunting regionally maintained a similar pattern between children 6–23 months and 24–59 months, but risk was lower relative to the Southeast among the older age group for all regions except the Central North, and substantially lower for the Mekong River Delta and Central South, whose RRs became insignificant in children 24–59 m.

### Comparing risk factors of stunting across regions

Stratified analyses by region of the combined age group of children 6–59 months using the alternative models revealed substantial heterogeneity in stunting risk factors between regions (**Supplemental Figures 3-10**). Ethnic minorities had double the stunting risk (RR: 1.99; 95% CI: 1.72, 2.31) in the Central Highlands but no increased risk in the Mekong River Delta (RR: 1.00; 95% CI: 0.86, 1.17). Birthweight < 2500 g was associated with a two-fold increase in stunting

risk in the Southeast (RR: 2.06; 95% CI: 1.61, 2.63) and only a 17% increase in the Northeast (RR: 1.17; 95% CI: 1.04, 1.31). Stunting risk among children of mothers with only a primary school education compared to mothers with a graduate education increased by three-fold in the Southeast (RR: 3.01; 95% CI: 1.77, 5.12) and only ~55% in the Northwest (RR: 1.55; 95% CI: 0.77, 3.11) and Central North (RR: 1.54; 95% CI: 1.18, 2.01). Interestingly, living in a mountainous commune was associated with a 25% decreased risk of stunting in the Central Highlands (RR: 0.75; 95% CI: 0.67, 0.85), a 63% increased risk in the Southeast (RR: 1.63; 95% CI: 1.38, 1.93) and a two-fold increased risk in the Northwest (RR: 1.91; 95% CI: 1.04, 3.49).

## Discussion

Our multivariable analysis of child stunting and its determinants adjusted for a range of covariates at the child, maternal, household, and environmental levels. While correlations in cross-sectional data do not prove causal relationships, we included variables that have a strong theoretical basis as determinants of stunting (Stewart et al., 2013). The variables with the strongest relationships with stunting among children 6–59 months were maternal height and education, and region—child stunting risk was roughly double among children of stunted mothers (height < 145 cm) compared to those not stunted (height  $\geq$  150 cm), mothers with no education or primary school only compared to those with a graduate education, and living in the Northeast compared to the Southeast. Maternal height and education have been shown to be strongly associated with child growth in Vietnam (P. H. Nguyen et al., 2017; Young et al., 2018) and other countries in South and Southeast Asia (Beal et al., 2018; Ikeda, Irie, & Shibuya, 2013;

P. H. Nguyen et al., 2017) as well as globally (Black et al., 2013; Danaei et al., 2016). About 15% of Vietnamese mothers had no education or primary school only, highlighting the need to improve access to education for women, which has potential to reduce stunting and provides other benefits (Black et al., 2013). The strong relationship between maternal short stature and child stunting in Vietnam indicates the importance of nutrition interventions before and during pregnancy, particularly because maternal short stature is clearly linked to fetal growth restriction and newborn stunting (Black et al., 2013; Victora et al., 2015). Research is needed to determine which nutrition interventions among mothers with short stature can most effectively improve birth outcomes and prevent children from following the same path as their mothers.

Unsurprisingly, the child-level variables age, sex, and birthweight were more strongly associated with stunting among younger than older children. Our findings correspond with recent global evidence that sex differences in child stunting diminish with age and disappear by about age 40 months (Alderman & Headey, 2018). Among children 6-23 months, discontinuing breastfeeding in older child age was associated with reduced risk of stunting but this was likely due to children's poor growth leading to increased breastfeeding, not the other way around (Kramer et al., 2011; Marquis et al., 1997).

Risk of stunting was highest in the Northeast, Northwest, Central North, and Central Highlands. This geographic pattern clearly represents variations in socioeconomic status, proportion of ethnic minority groups, and access to food, health services, and clean water (Dearden et al., 2017). The small size of landholdings, consequent carrying capacity of farming



systems to provide sufficient quality food, level of inclusion in profitable agricultural value chains, and opportunities for off-farm employment pose a challenge for the socioeconomic development in these regions (Minot & Baulch, 2005; Phat, 2012; Pimhidzai, 2018). Stunting reduction efforts should target regions at the highest risk. Regional variations in stunting risk factors were striking and suggest that tailoring policies and programs by region will more effectively address the most relevant stunting determinants. For example, ethnic minorities in the Central Highlands, children with low birthweight and of poorly educated mothers in the Southeast, and mountainous communes in the Northwest could be prioritized. Risk factors even varied substantially between regions with similar stunting risk. For example, in the Northwest, living in a mountainous commune was the strongest risk factor, while in the Central North, maternal height and education were the strongest risk factors, and living in a mountainous commune was not a risk factor.

We found little evidence that dietary factors such as consumption of animal-source foods, vitamin A-rich fruits and vegetables, and a MAD, measured by recent dietary intake or feeding practices, influenced stunting risk. It is unlikely that dietary factors are unimportant for child growth in Vietnam, however, given the dietary inadequacies of child diets (21). Methodological limitations may explain why dietary factors appeared unimportant. Recent intake in the past 24 hours does not necessarily reflect usual consumption in earlier childhood, and nutritional insults must have occurred over a period of time to influence stunting. Additionally, quantities of foods consumed were not measured, so the dose of the foods may not have been sufficient to

demonstrate a change in stunting. Living in a poor commune was associated with only a small increase in stunting risk. Socioeconomic status is clearly a strong driver of child stunting (P. H. Nguyen et al., 2017), but our analysis only included poverty at the commune level, and household wealth can vary substantially within communes of various socioeconomic status. Furthermore, while stunting and other indicators of malnutrition have improved, socioeconomic and ethnic inequalities have increased, highlighting the need for policies and comprehensive measures beyond the health sector that aim to reduce socioeconomic disparities (Benjamin, Brandt, & McCaig, 2017; Kien et al., 2016).

Our study has many strengths. Variables were selected from a rigorous conceptual framework and thus have a strong theoretical basis for being stunting determinants in Vietnam. Model selection was informed by this framework along with a combination of sophisticated statistical model selection approaches. The large, recent nationally representative sample of Vietnamese children 6-59 months allows for the generalization of findings across Vietnam and inference that is applicable to the current context. Analyzing children 6-23 months separately from 24-59 months allowed for richer models for each age group and the ability to compare determinants between younger and older children. Stratified analyses by region provides insight into how to best tailor interventions geographically to be most effective.

There are also important limitations. Although variables were selected from a strong conceptual framework, it is impossible to know with certainty the direction of the relationships, and there were likely important confounding factors omitted from the analysis. Many of the

determinants listed in the conceptual framework were unavailable in the dataset. Poverty was only included at the commune level and would have likely had a much stronger relationship with child stunting if it were included at the household level (P. H. Nguyen et al., 2017). Sample weights were unavailable at the village level, so any differences in population sizes between villages were not reflected in weighting. Lastly, child stunting is an imperfect proxy indicator for child growth failure (Perumal, Bassani, & Roth, 2018), and an unknown proportion of Vietnamese children with a  $HAZ < -2$  are growing normally and  $HAZ \geq -2$  experiencing growth faltering. This suggests our findings underestimated associations between child growth failure and its determinants.

In conclusion, our cross-sectional analysis of children 6-59 months found strong relationships between variables at the maternal and environmental levels with child stunting, especially short maternal height, low maternal education, and living in a northern region. Among children 6-23 months, child-level variables also had strong relationships with stunting, particularly age and birthweight. Regional analyses revealed strong differences in stunting risk factors between regions, which suggests policies and programs aimed at reducing stunting would have more impact if they were tailored geographically. Important progress has been made to reduce stunting since 2000, which has largely been due to economic development, but more specialized approaches may be required to reach the most vulnerable populations, where prevalence remains much higher than the national average. At the same time, multifaceted

approaches outside of the health sector are needed to reduce inequalities in socioeconomic status, which are inextricably linked with malnutrition.

## References

- Alderman, H., & Headey, D. (2018). The timing of growth faltering has important implications for observational analyses of the underlying determinants of nutrition outcomes. *PLOS ONE*, *13*(4), e0195904. <https://doi.org/10.1371/journal.pone.0195904>
- Beal, T., Tumilowicz, A., Sutrisna, A., Izwardy, D., & Neufeld, L. M. (2018). A review of child stunting determinants in Indonesia. *Maternal & Child Nutrition*, *14*(4), e12617. <https://doi.org/10.1111/mcn.12617>
- Benjamin, D., Brandt, L., & McCaig, B. (2017). Growth with equity: income inequality in Vietnam, 2002–14. *The Journal of Economic Inequality*, *15*(1), 25–46. <https://doi.org/10.1007/s10888-016-9341-7>
- Black, R. E., Victora, C. G., Walker, S. P., Bhutta, Z. A., Christian, P., de Onis, M., ... Uauy, R. (2013). Maternal and child undernutrition and overweight in low-income and middle-income countries. *The Lancet*, *382*(9890), 427–451. [https://doi.org/10.1016/S0140-6736\(13\)60937-X](https://doi.org/10.1016/S0140-6736(13)60937-X)
- Danaei, G., Andrews, K. G., Sudfeld, C. R., Fink, G., McCoy, D. C., Peet, E., ... Fawzi, W. W. (2016). Risk Factors for Childhood Stunting in 137 Developing Countries: A Comparative Risk Assessment Analysis at Global, Regional, and Country Levels. *PLOS Medicine*, *13*(11), e1002164. <https://doi.org/10.1371/journal.pmed.1002164>
- Dearden, K. A., Schott, W., Crookston, B. T., Humphries, D. L., Penny, M. E., Behrman, J. R., ... The Young Lives Determinants and Consequences of Child Growth Project Team.

- (2017). Children with access to improved sanitation but not improved water are at lower risk of stunting compared to children without access: a cohort study in Ethiopia, India, Peru, and Vietnam. *BMC Public Health*, *17*(1), 110. <https://doi.org/10.1186/s12889-017-4033-1>
- General Statistics Office (Vietnam), & United Nations Population Fund (UNFPA). (2009). *Vietnam Population and Housing Census 2009*.
- Ikeda, N., Irie, Y., & Shibuya, K. (2013). Determinants of reduced child stunting in Cambodia: analysis of pooled data from three Demographic and Health Surveys. *Bulletin of the World Health Organization*, *91*(5), 341–349. <https://doi.org/10.2471/BLT.12.113381>
- Kien, V. D., Lee, H.-Y., Nam, Y.-S., Oh, J., Giang, K. B., & Minh, H. V. (2016). Trends in socioeconomic inequalities in child malnutrition in Vietnam: findings from the Multiple Indicator Cluster Surveys, 2000–2011. *Global Health Action*, *9*(s1), 29263. <https://doi.org/10.3402/gha.v9.29263>
- Kramer, M. S., Moodie, E. E. M., Dahhou, M., & Platt, R. W. (2011). Breastfeeding and Infant Size: Evidence of Reverse Causality. *American Journal of Epidemiology*, *173*(9), 978–983. <https://doi.org/10.1093/aje/kwq495>
- Kursa, M. B., & Rudnicki, W. R. (2010). Feature selection with the Boruta package. *J Stat Softw*, *36*(11), 1–13.
- Liaw, A., & Wiener, M. (2002). Classification and regression by randomForest. *R News*, *2*(3), 18–22.

Lumley, T., Kronmal, R., & Ma, S. (2006). Relative Risk Regression in Medical Research: Models, Contrasts, Estimators, and Algorithms, 25.

Marquis, G. S., Habicht, J. P., Lanata, C. F., Black, R. E., & Rasmussen, K. M. (1997). Association of breastfeeding and stunting in Peruvian toddlers: an example of reverse causality. *International Journal of Epidemiology*, 26(2), 349–356.  
<https://doi.org/10.1093/ije/26.2.349>

Max Kuhn. Contributions from Jed Wing, Steve Weston, Andre Williams, Chris Keefer, Allan Engelhardt, Tony Cooper, Zachary Mayer, Brenton Kenkel, the R Core Team, Michael Benesty, Reynald Lescarbeau, Andrew Ziem, Luca Scrucca, Yuan Tang, Can Candan, and Tyler Hunt. (2018). Package “caret.”

Ministry of Health, National Institute of Nutrition. (2012a). *General Nutrition Survey 2009 – 2010*. Hanoi, Vietnam: Medical Publishing House.

Ministry of Health, National Institute of Nutrition. (2012b). *National Nutrition Strategy for 2011–2020, with a Vision Towards 2030*. Hanoi, Vietnam: Medical Publishing House.

Minot, N., & Baulch, B. (2005). Spatial patterns of poverty in Vietnam and their implications for policy. *Food Policy*, 30(5), 461–475. <https://doi.org/10.1016/j.foodpol.2005.09.002>

Nguyen, B. K. L., Thi, H. L., Do, V. A. N., Thuy, N. T., Huu, C. N., Do, T. T., ... Khouw, I. (2013). Double burden of undernutrition and overnutrition in Vietnam in 2011: results of the SEANUTS study in 0.5–11-year-old children. *British Journal of Nutrition*, 110(S3), S45–S56. <https://doi.org/10.1017/S0007114513002080>

- Nguyen, P. H., Headey, D., Frongillo, E. A., Tran, L. M., Rawat, R., Ruel, M. T., & Menon, P. (2017). Changes in Underlying Determinants Explain Rapid Increases in Child Linear Growth in Alive & Thrive Study Areas between 2010 and 2014 in Bangladesh and Vietnam. *The Journal of Nutrition*, jn243949. <https://doi.org/10.3945/jn.116.243949>
- Organization, W. H. (2009). WHO AnthroPlus for personal computers Manual: Software for assessing growth of the world's children and adolescents. *Geneva: WHO*.
- Perumal, N., Bassani, D. G., & Roth, D. E. (2018). Use and Misuse of Stunting as a Measure of Child Health. *The Journal of Nutrition*, 148(3), 311–315. <https://doi.org/10.1093/jn/nxx064>
- Phat, N. T. (2012). Geographical barriers impacts on the efficiency of poverty alleviation policy in Vietnam. *Journal of Research in International Business Management*, 2(3), 073–076.
- Pimhidzai, O. (2018). *Climbing the ladder* ~~poverty~~ *poverty reduction and shared prosperity in Vietnam* (No. 124916) (pp. 1–49). The World Bank. Retrieved from <http://documents.worldbank.org/curated/en/206981522843253122/Climbing-the-ladder-poverty-reduction-and-shared-prosperity-in-Vietnam>
- Ripley, B., Venables, B., Bates, D. M., Hornik, K., Gebhardt, A., Firth, D., & Maintainer Brian Ripley. (2013). Package 'mass.' *CRAN Repos. Httpcran R-Proj. OrgwebpackagesMASSMASS Pdf*.



Stewart, C. P., Iannotti, L., Dewey, K. G., Michaelsen, K. F., & Onyango, A. W. (2013).

Contextualising complementary feeding in a broader framework for stunting prevention.

*Maternal & Child Nutrition*, 9, 27–45. <https://doi.org/10.1111/mcn.12088>

Thomas Lumley, & Maintainer Thomas Lumley. (2018). *Package 'survey.'*

Trinh Thi, H., Simioni, M., & Thomas-Agnan, C. (2018). Assessing the nonlinearity of the calorie-income relationship: An estimation strategy – With new insights on nutritional transition in Vietnam. *World Development*, 110, 192–204.

<https://doi.org/10.1016/j.worlddev.2018.05.030>

UNICEF, FANTA, USAID, & WHO. (2017). *Meeting Report: Reconsidering, Refining, and Extending the World Health Organization Infant and Young Child Feeding Indicators (June 20-22, 2017)*. New York, NY. Retrieved from

<https://reliefweb.int/report/world/meeting-report-reconsidering-refining-and-extending-world-health-organization-infant>

United Nations, Department of Economic and Social Affairs. (2016). *Goal 2: End hunger, achieve food security and improved nutrition and promote sustainable agriculture* (Sustainable Development Knowledge Platform). Retrieved from [sustainabledevelopment.un.org/sdg2](https://sustainabledevelopment.un.org/sdg2)

Victora, C. G., Villar, J., Barros, F. C., Ismail, L. C., Chumlea, C., Papageorghiou, A. T., ...

Kennedy, S. H. (2015). Anthropometric Characterization of Impaired Fetal Growth: Risk

Factors for and Prognosis of Newborns With Stunting or Wasting. *JAMA Pediatrics*, 169(7), e151431–e151431. <https://doi.org/10.1001/jamapediatrics.2015.1431>

Viet Nam National Institute of Nutrition, UNICEF, Alive & Thrive. (2014). *Nutrition Surveillance Profiles 2013*. Ha Noi, Viet Nam.

Vietnam National Institute of Nutrition. (2018). Statistical data on child nutritional status over the years. Retrieved October 16, 2018, from <http://viendinhduong.vn/vi/suy-dinh-duong-tre-em/so-lieu-thong-ke-ve-tinh-trang-dinh-duong-tre-em-qua-cac-nam-106.html>

WHO. (2006). The WHO Child Growth Standards. Retrieved March 31, 2017, from <http://www.who.int/childgrowth/standards/en/>

WHO. (2012). *Maternal, infant and young child nutrition*. (WHO, Ed.). Geneva, Switzerland: The Sixty-fifth World Health Assembly WHA65.6.

World Health Organization. (2010). Indicators for assessing infant and young child feeding practices: part 2: measurement.

Young, M. F., Nguyen, P. H., Casanova, I. G., Addo, O. Y., Tran, L. M., Nguyen, S., ...

Ramakrishnan, U. (2018). Role of maternal preconception nutrition on offspring growth and risk of stunting across the first 1000 days in Vietnam: A prospective cohort study. *PLOS ONE*, 13(8), e0203201. <https://doi.org/10.1371/journal.pone.0203201>

Zhang, Z. (2016). Variable selection with stepwise and best subset approaches. *Annals of Translational Medicine*, 4(7), 136–136. <https://doi.org/10.21037/atm.2016.03.35>

## Tables

**TABLE 1**

Background characteristics of Vietnamese children by age

Variable	Children 6-23 months		Children 24-59 months	
	N	Mean/Prevalence (95% CI)	N	Mean/Prevalence (95% CI)
<b>Child characteristics</b>				
Height-for-age Z-score	30771	-0.73 (-0.77, -0.70)	55161	-1.05 (-1.08, -1.03)
Child stunting (HAZ < -2)	5823	17.5% (16.8, 18.2)	12432	21.4% (20.7, 22.1)
Child age (months)				
6-11	9136	29.4% (28.6, 30.0)	—	—
12-23	21635	70.6% (69.8, 71.0)	—	—
24-35	—	—	24023	44.3% (43.6, 45.0)
36-59	—	—	31138	55.7% (55.0, 56.4)
Child sex, male	16307	53.0% (52.2, 54.0)	29216	52.6% (52.0, 53.2)
Ethnic minority	6912	18.3% (17.0, 20.0)	12975	19.2% (18.0, 20.5)

Birthweight < 2,500 g	1421	4.5% (4.2, 5.0)	2798	4.8% (4.6, 5.1)
Child fever in past two weeks	4502	15.3% (14.6, 16.0)	5348	10.0% (9.6, 10.5)
Child not given deworming medication in past six months	29758	96.7% (96.4, 97.0)	30215	55.9% (54.9, 56.8)
Child not given vitamin A supplements in past six months	7118	23.5% (22.4, 25.0)	16468	30.0% (29.0, 31.0)
Child diarrhea in past two weeks	1226	4.3% (3.9, 5.0)	1285	2.3% (2.1, 2.5)
Currently breastfed	13275	43.6% (42.7, 44.0)	—	—
Child did not consume a minimum acceptable diet in past 24 hours	17017	53.1% (51.9, 54.0)	—	—
Child did not consume meat, fish, eggs, or milk products in past 24 hours	3296	10.2% (9.3, 11.0)	—	—
Child did not consume vitamin A-rich fruits and vegetables in past 24 hours	7047	24.4% (23.5, 25.0)	—	—
<b>Maternal characteristics</b>				
Maternal age < 18 years	4814	14.3% (13.7, 15.0)	—	—

Maternal age < 20 years	—	—	9316	15.8% (15.2, 16.4)
<hr/>				
Maternal height (cm)				
<hr/>				
< 145	1347	4.1% (3.8, 4.0)	2118	3.6% (3.3, 3.9)
<hr/>				
145–149.9	4454	13.8% (13.2, 14.0)	7246	12.5% (12.0, 13.0)
<hr/>				
≥ 150	24970	82.1% (81.3, 83.0)	45797	83.9% (83.2, 84.5)
<hr/>				
Maternal Body Mass Index (BMI)				
<hr/>				
< 18.5	4541	14.4% (13.9, 15.0)	6519	11.6% (11.2, 12.1)
<hr/>				
Maternal education (years)				
<hr/>				
None	1149	3.3% (2.9, 4.0)	2525	4.1% (3.7, 4.5)
<hr/>				
Primary (grades 1-5)	3278	9.8% (9.2, 10.0)	6836	11.4% (10.9, 12.0)
<hr/>				
Secondary (grades 6-9)	11435	35.7% (34.6, 37.0)	21326	37.4% (36.4, 38.4)
<hr/>				
High school (grades 10-12)	9079	30.5% (29.5, 31.0)	15533	29.3% (28.5, 30.1)
<hr/>				
College or university	3360	12.2% (11.5, 13.0)	3372	10.3% (9.7, 11.0)
<hr/>				
Graduate school	2266	8.6% (7.9, 9.0)	5083	7.5% (6.9, 8.1)
<hr/>				
Mother's occupation, farmer	11398	31.9% (30.3, 34.0)	22200	35.2% (33.7, 36.7)
<hr/>				
Mother did not wash hands both	17065	54.7% (53.2, 56.0)	32161	56.8% (55.4, 58.2)

before preparing foods and  
feeding child

---

Mother did not take iron or iron-

folate tablets in 3 months before or after birth	5846	18.1% (17.2, 19.0)	—	—
---	------	--------------------	---	---

---

Caregiver did not meet with  
health staff in past 3 months

	5425	18.9% (17.7, 20.0)	12144	22.9% (21.9, 24.0)
--	------	--------------------	-------	--------------------

---

**Household characteristics**

More than 2 children in the  
household

	4102	13.5% (12.9, 14.0)	8373	15.5% (14.9, 16.0)
--	------	--------------------	------	--------------------

Household does not use iodized  
salt

	3516	11.7% (10.9, 13.0)	6113	11.9% (11.0, 12.7)
--	------	--------------------	------	--------------------

---

**Environmental characteristics**

Rural

	21978	63.2% (60.3, 66.0)	40477	66.1% (63.5, 68.6)
--	-------	--------------------	-------	--------------------

Poor commune

	5371	14.7% (13.1, 16.0)	9982	15.2% (13.7, 16.8)
--	------	--------------------	------	--------------------

Mountainous commune

	9497	26.9% (25.0, 29.0)	17364	28.1% (26.3, 29.9)
--	------	--------------------	-------	--------------------


---

Region

Red River Delta


	4152	15.6% (14.6, 17.0)	7820	16.9% (15.8, 18.1)
--	------	--------------------	------	--------------------

Northeast	6066	16.0% (15.0, 17.0)	11009	16.5% (15.6, 17.5)
Northwest	1717	4.6% (4.1, 5.0)	3410	5.1% (4.6, 5.6)
Central North	2800	13.0% (11.8, 14.0)	5023	12.8% (11.7, 13.9)
Central South	2853	8.6% (8.0, 9.0)	5195	9.0% (8.4, 9.7)
Central Highlands	1933	6.4% (5.8, 7.0)	3483	6.4% (5.8, 7.0)
Southeast	5461	16.0% (14.6, 17.0)	8167	16.0% (14.6, 17.4)
Mekong River Delta	5789	16.2% (15.3, 17.0)	11054	17.3% (16.4, 18.2)




**FIGURE 1** Participant flow chart. HAZ, height-for-age Z-score; MAD, minimum acceptable diet.





**FIGURE 2** Adjusted relative risks of stunting determinants at the child, maternal, household, and environmental levels among 30771 Vietnamese children 6–23 m. Error bars show 95% confidence intervals. The multivariable model adjusted for all variables displayed. BMI, body

mass index; CI, confidence interval; MAD, minimum acceptable diet; RR, relative risk; veg, vegetables; vit, vitamin.



**FIGURE 3** Adjusted relative risks of stunting determinants at the child, maternal, household, and environmental levels among 55161 Vietnamese children 24–59 m. Error bars show 95% confidence intervals. The multivariable model adjusted for all variables displayed. BMI, body mass index; CI, confidence interval; RR, relative risk; vit, vitamin.

