


Chaperone OsmY facilitates the biogenesis of a major family of autotransporters

Zhen Yan,¹ Sunyia Hussain,^{2,†} Xu Wang,²
Harris D. Bernstein² and James C. A. Bardwell ^{1*}

¹Howard Hughes Medical Institute and Department of Molecular, Cellular & Development Biology, University of Michigan, Ann Arbor, MI 48109, USA.

²Genetics and Biochemistry Branch, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD 20892, USA.

Summary

OsmY is a widely conserved but poorly understood 20 kDa periplasmic protein. Using a folding biosensor, we previously obtained evidence that OsmY has molecular chaperone activity. To discover natural OsmY substrates, we screened for proteins that are destabilized and thus present at lower steady-state levels in an *osmY*-null strain. The abundance of an outer membrane protein called antigen 43 was substantially decreased and its β -barrel domain was undetectable in the outer membrane of an *osmY*-null strain. Antigen 43 is a member of the diffuse adherence family of autotransporters. Like strains that are defective in antigen 43 production, *osmY*-null mutants failed to undergo cellular autoaggregation. In vitro, OsmY assisted in the refolding of the antigen 43 β -barrel domain and protected it from added protease. Finally, an *osmY*-null strain that expressed two members of the diffuse adherence family of autotransporters that are distantly related to antigen 43, EhaA and TibA, contained reduced levels of the proteins and failed to undergo cellular autoaggregation. Taken together, our results indicate that OsmY is involved in the biogenesis of a major subset of autotransporters, a group of proteins that play key roles in bacterial pathogenesis.

Introduction

The cell envelope of Gram-negative bacteria is composed of an outer membrane and an inner membrane that enclose a compartment called the periplasm (Ruiz *et al.*, 2006). The periplasm is an aqueous but crowded compartment that occupies ~20% of total cell volume; it contains a thin layer of peptidoglycan and ~300 different proteins (Van Wielink and Duine, 1990). Periplasmic proteins are capable of performing diverse functions, including envelope biogenesis, signal transduction, the absorption and transportation of nutrients, the efflux of toxic substances and the determination of cell shape and virulence (Miller and Salama, 2018).

In vivo, the folding of proteins is commonly assisted by chaperones (Kim *et al.*, 2013). This is true in the periplasm as well as in the cytoplasm (Stull *et al.*, 2018). Periplasmic chaperones can assist protein folding independent of ATP, a remarkable feature that differentiates them from most cytoplasmic chaperones (Goemans *et al.*, 2014). Periplasmic chaperones have been shown to be involved in two important processes: (1) the biogenesis of outer membrane β -barrel proteins (OMPs) and (2) the protection of the periplasmic proteome from unfolding and/or aggregation under stress conditions. Nascent polypeptides for OMPs are secreted into the periplasm through an inner membrane channel complex called SecYEG (Wickner *et al.*, 1991); periplasmic chaperones then bind to them, maintain them in a partially unfolded form, and escort them to the Bam complex, which inserts β -barrel proteins into the outer membrane (Konovalova *et al.*, 2017; Noinaj *et al.*, 2017). The outer membrane serves as a semi-permeable barrier between the bacteria and the external environment. The proteins embedded in this layer help control which small molecules and proteins are allowed into the periplasmic space and which are excluded. These OMPs are a diverse group of proteins that commonly fold into β -rich structures often consisting of a barrel like structure, with 8–36 β -strands integrated into the outer membrane. Some possess periplasmic or extracellular domains. In some proteins, the core is open, forming a pore, while in others the core is filled. A number of periplasmic chaperones are thought to maintain the solubility and assist in the folding of OMPs as they transit

Accepted 22 July, 2019. *For correspondence. E-mail jbardwell@umich.edu; Tel. +1 734-764-8028; Fax +1 734-615-4226. Present addresses: [†]TetraGenetics, Inc., 91 Mystic St., Arlington, MA 02474, USA.

the periplasm. These include SurA, Skp and DegP. These proteins appear to have at least somewhat overlapping and redundant roles (Lazar and Kolter, 1996; Missiakas *et al.*, 1996; Spiess *et al.*, 1999). The absence of SurA, Skp or DegP results in decreased levels of certain OMPs and a minor outer membrane biogenesis defect (Vertommen *et al.*, 2009; Denoncin *et al.*, 2012), whereas deletions of both SurA and Skp or SurA and DegP leads to inviability (Rizzitello *et al.*, 2001; Denoncin *et al.*, 2012). Since the outer membrane is permeable, the periplasm is vulnerable to changes in the external environment; periplasmic proteins must therefore be able to cope with harsh conditions. Several periplasmic chaperones have been shown to function specifically under stressful conditions; these include HdeA and HdeB, which act in response to exposure to acidic conditions (Hong *et al.*, 2005; Kern *et al.*, 2007), and Spy, which is induced by protein unfolding agents such as butanol and tannins (Quan *et al.*, 2011).

We recently found the 20 kDa periplasmic protein OsmY to be a molecular chaperone through a genetic selection that forces cells to optimize unstable protein folding *in vivo* (Lennon *et al.*, 2015). OsmY was first discovered due to its strong induction by osmotic stress conditions (Yim and Villarejo, 1992). The OsmY sequence is composed of two repeated conserved regions, each of which contain a bacterial OsmY nodulation (BON) domain. The BON domain is a conserved domain that is typically ~60 residues long and arranged in an $\alpha\beta\beta\alpha\beta$ fold (Yeats and Bateman, 2003). *In vitro*, we found OsmY could inhibit the aggregation of a number of proteins commonly used for assaying chaperone activity, including lactate dehydrogenase, luciferase and α -lactalbumin (Lennon *et al.*, 2015). Consistent with our findings, several studies previously reported that OsmY, when fused to the N or C terminus of difficult-to-express or poorly folded recombinant proteins, allows large quantities of properly folded proteins to be exported into the extracellular media (Qian *et al.*, 2008; Bokinsky *et al.*, 2011; Kotsch *et al.*, 2011; Zheng *et al.*, 2012; Gupta *et al.*, 2013; Cheng *et al.*, 2014). These findings suggest that OsmY can function as a chaperone *in cis* (i.e. when present as part of the same molecule), in keeping with our finding that OsmY can function as a chaperone *in trans*. The genetic selection through which OsmY was discovered to be a chaperone involves the stabilization of a tripartite fusion protein consisting of an unstable mutant of maltose binding protein that is inserted into β -lactamase; other *in vivo* clients for OsmY's chaperone activity remain unknown.

One particularly interesting class of OMPs are the so called autotransporters, a large family of virulence-linked OMPs that are present in numerous Gram-negative bacteria (Henderson *et al.*, 2004; Dautin and Bernstein, 2007; Leyton *et al.*, 2012). In addition to a C-terminal β -barrel domain, autotransporters contain an N-terminal

extracellular (α) domain. The two domains are connected by a linker that traverses the pore of the β -barrel domain. Although the α -domains diverge greatly in sequence, size and function, they usually fold into a repetitive β -helical structure that promotes virulence (Henderson and Nataro, 2001; Celik *et al.*, 2012). The C-terminal domains also show minimal sequence conservation, but form nearly superimposable 12-stranded β -barrels (Oomen *et al.*, 2004; Barnard *et al.*, 2007; van den Berg, 2010; Zhai *et al.*, 2011). Autotransporters in *Escherichia coli* mainly fall into three groups based on homology: (1) serine protease autotransporters of the *Enterobacteriaceae* (SPATEs), (2) adhesins involved in diffuse adherence (AIDA-I) type autotransporters and (3) trimeric autotransporter adhesins (TAAs) (Wells *et al.*, 2010; Vo *et al.*, 2017).

Autotransporters were originally thought to be self-contained secretion systems, i.e., these proteins were thought to encode all the information and machinery necessary to transport their α -domain across the outer membrane (Pohlner *et al.*, 1987). Although this is no longer thought to be the case, it is still unclear what ensemble of host factors are involved in autotransporter biogenesis. Based primarily on studies on EspP, a member of the SPATE family, there is now evidence that after being translocated by the Sec complex into the periplasm, both domains interact with molecular chaperones, including SurA, Skp, DegP and FkpA (Ieva and Bernstein, 2009; Ruiz-Perez *et al.*, 2009; Ieva *et al.*, 2011). The exact function of the chaperones is unclear because individual chaperones can be deleted without affecting viability, possibly due to redundancy (Rizzitello *et al.*, 2001; Sklar *et al.*, 2007). Nevertheless, they are thought to maintain autotransporters in an assembly-competent conformation by preventing their misfolding and aggregation in the periplasm (Ieva *et al.*, 2011; Bernstein, 2015; Bernstein, 2019). DegP may also be required for the inner membrane translocation of the autotransporter pertactin (Braselmann *et al.*, 2016).

Available evidence indicates that the β -barrel domain of EspP begins to fold in the periplasm (Ieva *et al.*, 2008; Hussain and Bernstein, 2018). The protein is subsequently targeted to the Bam complex, a five protein heterooligomer (BamABCDE) that promotes both the insertion of the β -barrel domain into the outer membrane and the translocation of the α -domain across the membrane (Ieva and Bernstein, 2009). The mechanism of α -domain secretion is unclear, but it has been proposed to occur through a hybrid channel consisting of the β -barrels of both autotransporter and BamA in an open conformation (Pavlova *et al.*, 2013; Fan *et al.*, 2016).

Other factors, however, may play important roles in the assembly of autotransporters that do not belong to the SPATE family. For example, the Translocation and Assembly Module TamAB has been reported to contribute to the maturation of the autotransporter Ag43

(Selkrig *et al.*, 2012). Here, we report findings indicating that OsmY plays an important role in the biogenesis of a number of AIDA-I group autotransporters, including Ag43, EhaA and TibA.

Results

OsmY homologs are widespread within Proteobacteria

A PSI-BLAST search of the nonredundant database using *E. coli* MG1655 OsmY as the query sequence retrieved thousands of sequences with greater than 30% identity to OsmY. Most of these are present in Proteobacteria. *E. coli* OsmY contains two bacterial BON domains that share 43% identity to each other. Many organisms, however, are predicted to produce related proteins containing either just one or more than two BON domains, but the BON domain is also found in proteins with more complex protein architectures, as described in the Pfam database entry for this protein family (<http://pfam.xfam.org/family/BON>).

*Ag43 is poorly expressed in an *osmY*-null strain*

We previously isolated OsmY as a protein with chaperone activity (Lennon *et al.*, 2015). To better define the *in vivo* function of OsmY and to screen for its *in vivo* substrates, we compared the steady-state levels of proteins expressed in wild-type (WT) and *osmY*-null mutants through quantitative proteomics. We reasoned that proteins that require OsmY for their proper folding may be destabilized and thus decreased in abundance in $\Delta osmY$ strains. The only cell envelope protein that we found to be significantly reduced in the $\Delta osmY$ strain was the autotransporter Ag43, encoded by a gene called '*flu*' originally designated for the 'fluffing' phenotype it produces (Diderichsen, 1980). Presumably following the completion of assembly, Ag43 is cleaved by an unknown mechanism into a ~60 kDa N-terminal fragment that contains most of the surface exposed α -domain (hereafter referred to as the ' α domain') and a ~53 kDa C-terminal fragment that likely contains not only the outer membrane integrated β -barrel domain but also a portion of the extracellular domain (hereafter referred to as the ' β -barrel domain') (Owen *et al.*, 1996). Both domains were found to decrease five- to six-fold in MS spectral counts in a $\Delta osmY$ strain relative to the levels found in a WT strain (Fig. 1A). Western blotting using antisera raised against Ag43's α - and β -domains verified these proteomics results. Because antibody against the α -domain is not sensitive enough to detect trace amounts of α -domain in the $\Delta osmY$ strain, we quantified the β -domain bands. We performed three independent experiments and found that β -domain levels

were decreased by 8.5 ± 1.7 -fold in the $\Delta osmY$ strain compared to the WT strain. Like many other OMPs, the β -barrel domain remained folded in the absence of heat and migrated much more rapidly than its predicted molecular weight on SDS-PAGE (at 37 kDa), as reported previously (Owen *et al.*, 1996) (Fig. 1B). In addition, the $\Delta osmY$ strain showed a phenotype associated with a loss of Ag43 function in that it failed to mediate cellular autoaggregation (Fig. 1C). This phenotype is at least partially complemented by expression of *osmY* from a plasmid (Fig. S1). Co-expression of *osmY* and its downstream gene *ytjA* did not increase Ag43-mediated aggregation, ruling out the possibility that the poor expression of Ag43 is due to a polar effect in the $\Delta osmY$ strain (Fig. S1). We next used quantitative RT-PCR to determine the steady-state levels of Ag43 mRNA. Although these experiments did show a two- to three-fold decrease in steady-state levels of Ag43 mRNA in the *osmY*-null mutant, this decrease was insufficient to entirely explain the decrease in protein levels and was much smaller than the decrease observed in cells that have a reduced concentration of SecA (Mee-Ngan and Bernstein, 2013) (Fig. 1D). In addition, the level of two regulators of *flu* gene transcription for Ag43, OxyR and Dam (van der Woude and Henderson, 2008), did not significantly change in the *osmY*-null mutant as determined by quantitative proteomics (Table S3). We therefore conclude that Ag43 is destabilized in the *osmY* deletion mutant, which is consistent with the idea that OsmY is a chaperone that assists in either the folding or stabilization of Ag43.

Overproduction of Ag43 in a $\Delta osmY$ strain results in its partial proteolysis

To gain insight into the role OsmY plays in Ag43 maturation and export, we cloned the Ag43 gene (*flu*) into a pTrc vector to generate pTrcflu and, using this vector, overexpressed Ag43 in Δflu and $\Delta flu \Delta osmY$ strains. The Δflu strain can be complemented by Ag43 expression of pTrcflu which restores the ability of Ag43 to mediate cellular autoaggregation (Fig. 2B), and a band that reacted to Ag43 antibody and migrated at the position of the Ag43 α -domain was prominently observed using western blotting (Fig. 2A). Overexpression of pTrcflu in $\Delta flu \Delta osmY$ did not cause cellular aggregation (Fig. 2B), suggesting that these strains are phenotypically Ag43 minus, but a band migrating at ~42 kDa was observed in western blots using antiserum against Ag43 α -domain N-terminal peptide (Fig. 2A). This band is thus likely to be a proteolytic fragment that contains only an N-terminal portion of the Ag43 α -domain. Both the α -domain and its fragment were found in the periplasmic/outer membrane fraction (Quan *et al.*, 2013) (Fig. 2A).




Fig. 1. Comparison of Ag43 steady-state levels in WT and *osmY*-null strains.

A. WT and $\Delta osmY$ cells were assayed for quantitative proteomics using LC-MS/MS; shown are normalized Ag43 α - and β -domain spectral counts.

B. WT and $\Delta osmY$ cell pellets were resuspended in SDS-reducing sample buffer. After boiling at 95°C for 5 min or not, equal volumes were analyzed by SDS-PAGE and western blot using antiserum raised against the indicated proteins.

C. Autoaggregation of WT and $\Delta osmY$ cells was assayed by taking samples 1 cm below the liquid surface and measuring optical density at 600 nm.

D. Graph shows the quantitative RT-PCR analysis of Ag43 α - and β -domain mRNA levels in $\Delta osmY$ relative to WT. [Colour figure can be viewed at wileyonlinelibrary.com]

We then used a protease treatment approach to characterize the Ag43 α -domain and its fragment. The α -domain was relatively resistance to proteolysis in cell lysates, remaining undigested by trypsin at concentrations from 20 to 200 $\mu\text{g ml}^{-1}$ and proteinase K at 10 $\mu\text{g ml}^{-1}$ and was only partially digested at higher proteinase K concentrations (Fig. 2C and D), indicating it is well folded, consistent with previous reports (Babu *et al.*, 2018). However, the α -domain fragment that is present in the $\Delta osmY$ strain was much more protease sensitive at all concentrations of trypsin and proteinase K used, indicating it is not properly folded (Fig. 2C and D). The small amount of α -domain fragment remaining after proteolysis is similar to the amount of the protease sensitive control protein SurA that remains at all protease concentrations. The apparently protease resistant subpopulations of these two proteins are very likely due to a residual population of unlysed cells.

Whether the Ag43 α -domain and its fragment were surface exposed or not were determined by dot blot assays,

which determine whether or not antibodies to a protein can react with unlysed cells (Cho *et al.*, 2014; Konovalova *et al.*, 2014). A strong signal for the α -domain of Ag43 was detected in the cells of Δflu containing pTrcflu, whereas virtually no signal for the α -domain fragment that was present in the $\Delta flu\Delta osmY$ cells containing pTrcflu was detected unless the cells were sonicated (Fig. 2E).

Cellular fractionation using ultracentrifugation revealed that most of the Ag43 α -domain is found in the membrane fraction, whereas most of the fragment is found in the soluble fraction (Fig. 2F). Cells overexpressing Ag43 autoaggregated, presumably via the surface exposure of the Ag43 α -domain, but cells overexpressing Ag43 from pTrcflu in $\Delta flu\Delta osmY$ strains did not autoaggregate, even though they expressed a large amount of the Ag43 fragment (Fig. 2A and B). This indicates that the Ag43 fragment present in $\Delta osmY$ strains is not functional. No Ag43 β -domain cross-reacting material was observable when we attempted Ag43 overexpression from pTrcflu in $\Delta flu\Delta osmY$ strains (Fig. 2A), suggesting that the

β -domain undergoes complete proteolysis. This proteolysis could occur either before or after insertion into the outer membrane, thereby leaving an α -domain fragment to accumulate in the periplasm in an unfolded, misfolded or partially folded state. One possible scenario is that the β -barrel portion of Ag43 misfolds and cannot integrate

into the outer membrane. Ag43 is therefore retained in the periplasm and the N terminus adopts a non-native but relatively stable conformation. The less well-folded C terminus is then clipped off and degraded, leaving the better-folded N terminus behind in the periplasm. It seems unlikely that the N terminus is completely unfolded – if it




Fig. 2. Overproduction of Ag43 in the *osmY*-null mutant results in the production of a ~42 kDa α -domain proteolytic fragment. Strains deleted for Ag43 (Δflu) and those deleted for both Ag43 and OsmY ($\Delta flu\Delta osmY$) harboring either pTrcflu to overexpress Ag43 or an empty vector pTrc as a control, were cultured as described in Experimental Procedures.

A. Whole cell lysates and periplasmic preparations extracted with 1 mg ml⁻¹ polymyxin were boiled at 95°C for 5 min and analyzed by SDS-PAGE and western blotting using antiserum raised against the indicated proteins.

B. Autoaggregation assays by taking samples 1 cm below the liquid surface and measuring optical density at 600 nm. ●, Ag43 overexpressing pTrcflu in Δflu . ○, Ag43 overexpressing pTrcflu in $\Delta flu\Delta osmY$. ▲, empty vector (pTrc) in Δflu . △, empty vector (pTrc) in $\Delta flu\Delta osmY$.

C. Proteinase K (PK) and trypsin-treated experiments were performed as described in Experimental Procedures. Samples were analyzed by SDS-PAGE and western blot using antiserum raised against the indicated proteins.

D. Ag43 α -domain and fragment bands were quantified using Image J, and intensities were normalized to MBP band intensities, which were used as a protease resistant loading control. The amount of Ag43 α -domain, fragment and SurA bands at no added protease were set to one.

E. Dot blot assays were performed as described in Experimental Procedures. Membranes were blocked and probed with the indicated antibodies. A high signal that is unaffected by sonication indicates surface exposure, a low signal in unsonicated cells that is enhanced by sonication indicates the protein is contained within the cell, low but visible signals in the absence of sonication can be due to some cell lysis. OppA and DegP were used as periplasmically localized control proteins.

F. Cell fractionation were performed as described in Experimental Procedures. Each fraction was then analyzed by SDS-PAGE and western blot using antiserum raised against the indicated proteins. [Colour figure can be viewed at wileyonlinelibrary.com]

were, it would probably be degraded by periplasmic proteases. Independent of the exact model, our results suggest that OsmY is important for the folding and/or insertion competence of the Ag43 β -domain.

OsmY specifically stabilizes the Ag43 β -domain in vitro

To study the refolding of the Ag43 β -barrel domain in vitro, we overexpressed it in the *E. coli* cytosol and purified it from inclusion bodies. We then dissolved the β -domain in 8 M urea and attempted to refold it in the presence or absence of OsmY. To do this, we diluted the urea-dissolved Ag43 β -domain into a buffer containing 0.5% of the detergent N,N-dimethyldodecylamine N-oxide (LDAO), which has been used to refold β -domains of other autotransporters (Zhai *et al.*, 2011; Yuan *et al.*, 2018), and monitored refolding over time using gel mobility on SDS-PAGE. This method exploits the observation that heating alters the mobility of OMPs in a conformationally dependent manner. Two forms of Ag43 migrated more rapidly than the urea-denatured form; these were detected as a prominent species by western blotting after only 1 min of refolding. The upper band of these two fast-migrating species decreased over time, possibly due to chasing into the lower band or proteolysis. Either possibility implies that the upper band of the doublet is less well folded than the more rapidly migrating species. However, since boiling the refolded sample caused both of these fast-migrating forms to disappear, we deduce that both bands are at least partially folded (Fig. 3A). Quantification of these two refolded bands suggests that the addition of OsmY initially slows the folding of the β -barrel domain but results in a higher folded yield (Fig. 3B). The abundance of both the unfolded and folded forms decreased with time in the absence of OsmY, suggesting that proteolytic degradation may be occurring due to protease contamination of our partially pure Ag43 preparation. We reasoned that since there are probably similar amounts of protease in the samples incubated in the presence or absence of OsmY,

and since degradation was much more prominent in its absence, OsmY may be protecting the Ag43 β -barrel domain from degradation. To test this hypothesis, we added a fixed amount of proteinase K into refolding mixtures that were supplemented either with OsmY or not. The results revealed that OsmY efficiently protects both the unfolded and folded Ag43 β -barrel domain from proteolysis. We then tested several well-known periplasmic chaperones to see if they could also stabilize Ag43, using lysozyme as an additional protein control. Among the chaperones shown to play roles in OMP biogenesis, only SurA and Skp were able to protect the Ag43 β -barrel domain from proteolysis as well as OsmY does (Fig. 4). Proteolysis by trypsin gave similar results (Fig. S2). Since the Ag43 β -domain belongs to the β -barrel OMP family, we next examined the effect of added OsmY on the proteinase K digestion of several other unfolded OMPs in detergent micelles. OsmY was able to prevent degradation for all those tested (Fig. S3). This effect may be specific to β -barrels, as OsmY was unable to mitigate the degradation of the unfolded Ag43 α -domain (Fig. S4). We also found that OsmY barely affects α -domain refolding from a urea-denatured form, although it is able to inhibit the time-dependent aggregation of the well-folded α -domain (Fig. 5), as might be expected for a protein possessing broad anti-aggregation activity against a number of commonly used, but admittedly heterogeneous substrates (Lennon *et al.*, 2015). In summary, our in vitro results suggest that OsmY plays a critical role in Ag43 β -barrel domain refolding and stabilization in detergent micelles, which would impact α -domain maturation in vivo.

*Display to cell surface of AIDA-I type autotransporters is impaired in the *osmY*-null mutant*

To determine whether any AIDA-I family autotransporters besides Ag43 are also dependent on OsmY for their activity, we expressed EhaA from enterohemorrhagic *E. coli* and TibA from enterotoxigenic *E. coli* in Δflu and




Fig. 3. Refolding of the Ag43 β -domain into LDAO micelles.

A. Purified Ag43 β -domain held in 8 M urea was diluted 10 times into 0.5% LDAO buffer to a final concentration of 5 μ M at 37°C; buffer either contained 25 μ M OsmY or 25 μ M BSA. Equal volumes of sample were removed from the reaction mixture at the indicated time points, followed by heating at 95°C for 5 min or not, then analyzed by SDS-PAGE and western blot using antiserum raised against the β -domain. Images are representative of three independent experiments.

B. The folded Ag43 β -domain bands of three independent experiments were quantified using Image J and plotted against time. In each experiment, the intensities of the folded β -domain were normalized to urea-denatured unfolded β -domain band intensities, which were first normalized to Ag43 β -domain concentrations used in each experiment. \blacktriangle , refolding of β -domain without OsmY. \bullet , refolding of β -domain with OsmY. [Colour figure can be viewed at wileyonlinelibrary.com]

$\Delta flhA\Delta osmY$ strains. We found similar defects for these autotransporters in an *osmY* deletion strain as we did for Ag43, both in terms of expression, as detected by western blot, and the cell aggregation phenotype, as detected by assaying static culture optical density (Fig. 6). Because three AIDA-I autotransporters showed defects in an *osmY*-null mutant, but EspP, which does not belong to the AIDA-I group, did not show biogenesis defects in the $\Delta osmY$ strain (Fig. S5), we tentatively conclude that OsmY affects the biogenesis of a range of AIDA-I type autotransporters.

Discussion

In this report, we show that OsmY plays a key role in the assembly of Ag43 and other members of the AIDA-I family of bacterial autotransporters. Whereas Ag43 is normally cleaved into two stable fragments after the β -barrel domain is inserted into the outer membrane and the α -domain is secreted, disruption of the *osmY* gene leads to the accumulation of an N-terminal α -domain fragment in the

periplasm and the almost complete disappearance of the β -barrel domain. This phenotype is striking given that: (1) proteolytic fragments of autotransporters have not been reported to accumulate following the depletion of BamA (Jain and Goldberg, 2007), and (2) the loss of a single periplasmic chaperone does not always strongly affect autotransporter assembly (Ieva and Bernstein, 2009; Ruiz-Perez *et al.*, 2009). The results suggest a scenario in which OsmY is specifically required to maintain the Ag43 β -barrel domain in an insertion-competent state. In the absence of OsmY, Ag43 remains in the periplasm where the C terminus of the protein is eventually digested by proteases. Presumably because the protein resides in the periplasm for an abnormally long time, an N-terminal segment that may correspond to one arm of the 'L' structure of the α -domain (Heras *et al.*, 2014) has an opportunity to fold into a protease-resistant conformation. Consistent with this interpretation, we found that OsmY promotes the refolding of the purified Ag43 β -barrel domain in vitro. Furthermore, OsmY protects the β -barrel domain, but not the α -domain, from digestion by exogenous proteases. OsmY was also required for the stable expression of two




Fig. 4. Proteinase K (PK) digestion of the Ag43 β -domain in LDAO micelles. 5 μ M samples of purified Ag43 β -domain were refolded in 0.5% LDAO preincubated with 25 μ M BSA, lysozyme, Spy, SurA, Skp, DegP S210A or OsmY, respectively, for 20 min at 37°C. PK was added to the reaction mixture, and equal volumes of sample were removed into 10 mM PMSF at the indicated time points. Samples were analyzed by SDS-PAGE and western blot using antiserum raised against the β -domain. In all experiments, the molar ratio of Ag43 β -domain to PK was 2000:1. [Colour figure can be viewed at wileyonlinelibrary.com]

other members of the AIDA-I family of autotransporters, EhaA and TibA. Our results are noteworthy because they may shed some light on the cellular factors that could promote autotransporter assembly.

Given the multiplicity of periplasmic chaperones and their apparent functional redundancy, it should be interesting to determine why Ag43 specifically requires OsmY for assembly. Unlike the assembly of Ag43, the assembly of the SPATE protein EspP was not impaired in a Δ osmY strain (Fig. S5). The two proteins are also distinct in that the efficient assembly of only Ag43 appears to require TamA/TamB (Selkrig *et al.*, 2012; Kang'ethe and Bernstein, 2013). In addition to having

an unusual L-shaped α -domain, Ag43 has a β -barrel domain that is very distantly related to the EspP β -barrel domain (< 20% identity). It is conceivable that the Ag43 β -barrel domain has structural elements that distinguish it from the C-terminal domain of other autotransporters. Perhaps unique features of the α -domain and/or the β -barrel domain require the recruitment of additional assembly factors. As previously proposed, TamB might modulate α -domain folding in the periplasm (Bamert *et al.*, 2017; Babu *et al.*, 2018) and might maintain the Ag43 α -domain in a secretion-competent conformation. Indeed, it is possible that the stable α -domain fragment that we observed in the absence of OsmY results from




Fig. 5. OsmY prevents Ag43 α -domain aggregation following acid induced unfolding, but has no effect on its refolding in vitro. **A.** 50 μ M samples of purified Ag43 α -domain in 50 mM Tris pH 8.0, 150 mM NaCl buffer were diluted 20 times into the same buffer at pH 4.0 that had been preincubated with different concentrations of OsmY. Acid induced protein aggregation over time was monitored by measuring light scattering at 360 nm using a fluorescence spectrophotometer at 25°C. **B.** 50 μ M samples of purified Ag43 α -domain were denatured with 8 M urea, and then diluted 50 times into 50 mM Tris pH 8.0, 150 mM NaCl in the presence or absence of 10 μ M OsmY. Refolding was monitored by measuring tryptophan fluorescence excitation at 295 nm and emission at 350 nm using a Cary Eclipse fluorimeter at 25°C. Refolding of 1 μ M samples of purified Ag43 α -domain (native form) was also monitored at the same time using the same refolding assay in the presence or absence of 10 μ M OsmY. Solid black, Ag43 α -domain refolding. Solid red, Ag43 α -domain refolding with OsmY. Dotted black, native Ag43 α -domain. Dotted red, native Ag43 α -domain with OsmY. [Colour figure can be viewed at wileyonlinelibrary.com]

its interaction with TamB. Consistent with current models (Albenne and Ieva, 2017), BamA and TamA might also function consecutively or cooperatively to catalyze the efficient insertion of the Ag43 β -barrel domain. Interestingly, an interaction between OsmY and the Tam complex has been reported (Babu *et al.*, 2018). This observation suggests that OsmY may target proteins to TamA/TamB. In any case, it seems reasonable to speculate that OsmY and TamA/TamB function in a pathway that is parallel to the canonical pathway (i.e. the chaperones SurA, Skp and DegP plus the Bam complex)

and that is required for the biogenesis of a subset of autotransporters.

Our finding that OsmY protects a variety of *E. coli* OMPs from degradation in vitro suggests that it has a broad affinity for β -barrels. Although the structure of OsmY is unknown, it is conceivable that the hydrophobic regions of the BON domains interact with exposed hydrophobic surfaces of partially folded β -barrel proteins and prevent them from aggregating. The interaction of OsmY and the hydrophobic amino acid phenylalanine has been reported previously (Piazza *et al.*, 2018). OsmY may be analogous to Skp in forming a cage that provides a protective environment for OMPs (Walton *et al.*, 2009; Burmann *et al.*, 2013; Schiffrin *et al.*, 2016). However, the finding that the level of most OMPs is similar in WT and Δ osmY strains strongly suggests that other chaperones can effectively substitute for OsmY under physiological conditions, perhaps analogous to how Skp and SurA can substitute for each other depending on the protein and growth conditions (Stull *et al.*, 2018).

The high conservation of OsmY in Proteobacteria strongly suggests that it may facilitate the biogenesis of other AIDA-I family autotransporters, which are also widespread in Proteobacteria. Whether OsmY is responsible for biogenesis of autotransporters besides those belonging to the AIDA-I family requires further study, autotransporters are intimately involved in virulence (Henderson *et al.*, 2004). Our finding that OsmY is an indispensable factor in Ag43 maturation is interesting in light of several recent reports that OsmY is also involved in bacterial virulence. One study showed that an *osmY*-null mutant of *Yersinia ruckeri* failed to be infectious in fish (Mendez *et al.*, 2018), and a second study suggested that *osmY* is linked to virulence factors that promote biofilm formation and flagellar motility in *Cronobacter sakazakii* (Ye *et al.*, 2015). OsmY has also been proposed to be indirectly associated with virulence in *Salmonella typhimurium* and *E. coli* (Bader *et al.*, 2003; Dong and Schellhorn, 2009). Taken together with our results, these studies suggest that OsmY plays a common role in the assembly of a subset of specialized OMPs that differ considerably in structure from the generic porins that dominate the outer membrane of laboratory strains of *E. coli*.

Experimental procedures

Bacterial strains and plasmids

All the strains and plasmids used in this study are listed in Table S1. Deletion of the *osmY* gene was performed as previously described (Datsenko and Wanner, 2000). The inserted antibiotic cassette generated using this procedure was excised from the chromosome using pCP20. The




Fig. 6. Deletion of the *osmY* gene inhibits EhaA- and TibA-mediated cellular autoaggregation. Δflu and $\Delta flu \Delta osmY$ strains harboring pBADEhaA, pBADTibA with a C-terminal MycHis tag, and empty vector (EV), respectively, were cultured in LB media with $100 \mu\text{g ml}^{-1}$ ampicillin until the OD_{600} reached 0.6. L-arabinose (0.002%) was added or not and cells were induced for 3 h.

A. Whole cell lysates were analyzed by SDS-PAGE followed by western blotting using antiserum raised against DnaK and the C-Myc tag.

B. Autoaggregation of different cells was assayed by taking samples 1 cm below the liquid surface for optical density readings at 600 nm. ○, pBADEhaA in Δflu . ●, pBADEhaA in Δflu with arabinose. ○, pBADEhaA in $\Delta flu \Delta osmY$. ●, pBADEhaA in $\Delta flu \Delta osmY$ with arabinose. △, pBADTibA in Δflu . ▲, pBADTibA in Δflu with arabinose. △, pBADTibA in $\Delta flu \Delta osmY$. ▲, pBADTibA in $\Delta flu \Delta osmY$ with arabinose. □, EV in Δflu . ■, EV in Δflu with arabinose. [Colour figure can be viewed at wileyonlinelibrary.com]

chromosomal insertion/deletion of the *flu* gene encoding Ag43 was transferred from the strain BW25133 (Keio collection) to MC4100 and $\Delta osmY$ strains by P1 transduction (Baba *et al.*, 2006). Vectors expressing Ag43 and OsmY were made by amplifying their respective genes with PCR from MC4100 and directly cloning into pTrc- or pBAD-based vectors. The genes for EhaA and TibA were amplified from *E. coli* O157:H7 and H10407 genomic DNA, respectively. All the vectors were constructed using In-Fusion HD Cloning kits (TaKaRa); all the oligo primers used are listed in Table S2.

Cell growth

Liquid cultures were grown in Luria–Bertani (LB) media at 37°C shaking at 200 r.p.m. Overnight cultures were diluted 1:100 into fresh LB media. If necessary, $100 \mu\text{g ml}^{-1}$ ampicillin was added to maintain the pTrc- and pBAD-based vectors. Expression from the lac promoter on pTrc-based vectors was done by adding IPTG to 0.5 mM final concentration followed by 3 h of induction prior to harvesting. Expression from the arabinose promoter in pBAD-based

vectors was similarly done but by using 0.002% arabinose final concentration.

Quantification of protein levels

WT and $\Delta osmY$ strains were grown until the OD_{600} reached 1.0. 1.0 ml of the cells was centrifuged at $3000\times g$ for 10 min, washed by resuspending in PBS buffer (Na_2HPO_4 10 mM, KH_2PO_4 1.8 mM, KCl 2.7 mM, NaCl 137 mM, pH 7.4) of a volume equal to the culture volume and then recentrifuged. The cell pellets were provided to MS Bioworks (Ann Arbor) who performed Mass Spec analysis as follows. The cell pellets were lysed by resuspension in 500 μl of 2% SDS, 150 mM NaCl and 50 mM Tris pH 8.0 containing one tablet of Roche Complete Protease Inhibitor Cocktail, followed by sonication for three cycles of 20 s on ice (Fisherbrand Model 505). The amount of protein present in the lysate was quantified by Qubit fluorometry (Invitrogen). Lysates containing 10 μg of protein were processed by SDS-PAGE using a 10% Bis-Tris NuPAGE Novex mini gel (Thermo) and the supplied MES buffer system. The region of the gel containing stained proteins was excised

and then processed by in-gel digestion with trypsin using a ProGest robot and the following protocol: (1) The gel slice was washed twice with 50 μ l of 25 mM ammonium bicarbonate followed by a wash with 50 μ l of acetonitrile; (2) proteins in the gel slice were reduced using 40 μ l of 10 mM dithiothreitol at 60°C followed by alkylation using 40 μ l of 50 mM iodoacetamide at room temperature; (3) proteins were digested by addition of 200 ng of sequencing grade trypsin (Promega) at 37°C for 4 h; and (4) digestion was stopped by the addition of 30 μ l of trifluoroacetic acid. Each gel digest was then analyzed by nano LC-MS/MS with a Waters NanoAcquity HPLC system interfaced to a ThermoFisher Q Exactive. Peptides were loaded on a trapping column and eluted over a 75 μ m analytical column. Both columns were packed with Luna C18 resin (Phenomenex). Peptides were eluted at 350 nl min⁻¹ with a 2 h binary reverse phase gradient. Buffer A was 0.1% formic acid; buffer B was 0.1% formic acid in acetonitrile. The gradient was at 0 min 98% A, at 1 min 95% A, at 95 min 75% A, at 110 min 65% A, at 112 min 10% A, at 113 min 98% A and at 120 min maintained at 98% A. The mass spectrometer was operated in data-dependent mode, with the Orbitrap operating at 70,000 full width at half maximum (FWHM) and 17,500 FWHM for MS and MS/MS respectively. The 15 most abundant ions were selected for MS/MS.

SDS-PAGE and western blots

The indicated volumes of cells were centrifuged at 3000 \times *g* for 10 min and washed with amounts of PBS buffer equal to that of the culture volume. Cells were resuspended in SDS-reducing sample buffer and, if indicated, boiled at 95°C for 5 min. SDS-PAGE gels were done using NuPAGE 4–12% Bis-Tris gel (Invitrogen) or 4–20% Mini-PROTEAN TGX stain free gels (Bio-Rad) as specified in the figure legends. After electrophoresis, gels were transferred to a turbo polyvinylidene difluoride (PVDF) membrane (Bio-Rad) using a Trans-Blot Turbo apparatus (Bio-Rad). The blotted PVDF membranes were then blocked using 5% nonfat dried milk in TBST (20 mM Tris, 150 mM NaCl, 0.1% Tween 20) for 1 h at room temperature and probed with the following primary antibodies at these dilutions in 5% nonfat dried milk for 1 h at room temperature: rabbit-derived OsmY polyclonal antibody (Pacific Immunology), 1:5000; rabbit-derived Ag43 α -domain polyclonal antibody (a gift from Begona Heras, La Trobe University), 1:3000; rabbit-derived Ag43 β -domain polyclonal antibody (Pacific Immunology), 1:5000; mouse-derived maltose-binding protein (MBP) monoclonal antibody (Biolabs), 1:15000; and mouse-derived DnaK monoclonal antibody (Enzo), 1:15000. The membranes were then washed 3 times (10 min each) with shaking by TBST and probed with fluorescence dye-conjugated goat anti-rabbit and goat anti-mouse secondary antibodies (1:15000) (LI-COR Biosciences). Imaging was performed using LI-COR Odyssey CLX.

Dot blot assay

Bacterial strains were grown until the OD₆₀₀ reached ~1.0. 1 ml of these cells was centrifuged at 3000 \times *g* for 10 min and resuspended in one volume of PBS buffer (137 mM NaCl, 2.7 mM KCl, 10 mM Na₂HPO₄, 1.8 mM KH₂PO₄). Half

of these cells were then lysed by three cycles of 20 s sonication on ice (Fisherbrand Model 505), with the other half serving as the intact cell sample. 2 μ l of intact cells or lysed cells was spotted on a nitrocellulose membrane and air-dried. Membranes were blocked with 2% nonfat dried milk in PBS for 30 min at room temperature and probed with the following primary antibodies in 5% nonfat dried milk for 1 h at room temperature: rabbit-derived OsmY polyclonal antibody, 1:5000; rabbit-derived Ag43 α -domain polyclonal antibody, 1:3000; rabbit-derived OppA polyclonal antibody, 1:5000 (Pacific Immunology); rabbit-derived DegP polyclonal antibody, 1:5000 (a gift from Michael Erhmann). The procedures for probing with secondary antibodies and imaging were done using the same protocol as described for the western blot.

Membrane fractionation

Hundred milliliters of Δ *flu* and Δ *flu* Δ *osmY* cells harboring pTrcflu were centrifuged at 3000 \times *g* for 10 min, washed with 10 mM HEPES buffer pH 7.5 and 2 mM MgCl₂ and resuspended in 20 ml of the same buffer supplemented with 1 mg of DNase and 1 mg of RNase. Cells were lysed by passing through a French press at 12,000 psi. Cell debris was removed by centrifugation at 4200 \times *g* at 4°C for 5 min. Sixteen milliliters of the supernatant were then loaded on top of a two-step sucrose gradient (2.3 ml 2.02 M sucrose and 6.6 ml 0.77 M sucrose). The samples were centrifuged at 130,000 \times *g* at 4°C for 3 h in a Type 70.1 Ti rotor (Beckman Coulter). After centrifugation, 500 μ l samples were removed from the top to the bottom of the gradient and analyzed by SDS-PAGE and western blotting.

Quantitative RT-PCR

WT and Δ *osmY* strains were grown to the late log phase (OD₆₀₀ of 1.0) then harvested by centrifugation at 3000 \times *g* for 10 min. Total RNA was then isolated using a NucleoSpin RNA kit (Macherey-Nagel, Düren, Germany). DNA contamination was eliminated by the use of DNase treatment and removal reagents in a DNA removal kit (Ambion by Life Technologies, AM1906). cDNAs were then synthesized with a primeScript First-strand cDNA synthesis kit (Takara) using the supplied mixture of random primers. Quantitative PCRs were performed using the Eppendorf Realplex® PCR detection system in a triplicate reaction. The reaction mixture contained Radiant™ Green qPCR Mix Lo-ROX, 400 nM primers and 100 ng cDNA. PCR was performed using the following program: 95°C for 2 min followed by 40 cycles of 95°C for 5 s and 60°C for 1 min. The threshold cycle (*C*_T) was determined using the manufacturer's software. Primers 11–12, 13–14 and 15–16 were used to amplify Ag43 α -domain, Ag43 β -domain and *gapA* respectively. mRNA levels of Ag43 were normalized to the reference gene *gapA* and calculated using the comparative *C*_T method.

Refolding of outer membrane β -barrel proteins into detergent micelle and protease digestions

To initiate refolding, the purified Ag43 β -domain held in 8 M urea was diluted 10 times into 0.5% LDAO buffer to a final

concentration of 5 μM at 37°C. To test the effect of OsmY on refolding, the buffer contained 25 μM OsmY, 25 μM BSA or neither. Folding progress was followed by removing aliquots of this folding reaction at various time intervals and adding them into SDS-reducing sample buffer. Folding status was determined by comparing the migration on SDS-PAGE with and without heating at 95°C for 5 min. Many folded OMPs, including Ag43, are known to migrate more rapidly on SDS-PAGE than unfolded versions (Owen *et al.*, 1996) when analyzed by SDS-PAGE and western blotting using anti-serum raised against Ag43's β -domain. Folded status was further determined by proteolysis. In these experiments, purified Ag43 β -domain, EspP β -domain, OmpA, OmpC and OmpT were refolded in 0.5% LDAO for 20 min, respectively, after preincubation with 25 μM of known chaperones or control proteins, including BSA, lysozyme, OsmY, Spy, SurA, Skp or the chaperone-active protease-inactive DegP variant DegP-S210A. Following various intervals of folding, proteinase K or trypsin was added to the reaction mixture. For the Ag43 β -barrel folding reaction, an Ag43: proteinase K ratio of 2000:1 was used, and for the other OMPs, a 1000:1 protein: proteinase K ratio was used. Equal volumes of sample were removed from the reaction mixture into 10 mM phenylmethylsulfonyl fluoride (PMSF) to stop protease activity, then analyzed by SDS-PAGE and western blotting using antiserum raised against the corresponding OMPs.

Protein expression and purification

Expression and purification of OsmY was performed as reported previously (Lennon *et al.*, 2015). His-tagged SurA and various OMPs (EspP Δ 5, OmpA, OmpC and OmpT) lacking signal sequences were expressed and purified as previously described (Roman-Hernandez *et al.*, 2014; Hussain and Bernstein, 2018). Skp was purchased directly from MyBiosource.com. Ag43 β -domain was cloned into pET21a with a C-terminal His6 tag. The recombinant proteins were overexpressed in *E. coli* BL21(DE3) strain and isolated from inclusion bodies using buffer A (8M urea, 25 mM Tris, 150 mM NaCl, pH 8.0). The proteins were purified on a 5 ml HisTrap HP column (Amersham Biosciences) equilibrated with buffer A. After washing with buffer A containing 50 mM imidazole, the proteins were eluted with buffer A containing 500 mM imidazole. The *degP* gene was cloned into pET28b with a C-terminal His6 tag. The QuikChange site directed mutagenesis kit (Stratagene) was then used to introduce the S210A mutation into *degP*. Recombinant DegP S210A was overexpressed in *E. coli* BL21(DE3) and isolated from cell lysates using buffer B (PBS containing 20 mM imidazole). The protein was purified on a Ni-NTA (Qiagen) column equilibrated with buffer B. After washing with buffer B containing 400 mM NaCl and 850 mM NaCl, the protein was eluted with buffer B containing 250–500 mM imidazole. Elution fractions containing DegP S210A were pooled and buffer exchanged into PBS using PD-10 Sephadex desalting columns (GE Healthcare). The purified protein was then concentrated using Amicon 30 kD centrifugal filter units (Millipore) and the concentration was determined using the DC Protein Assay (Bio-Rad).

Acknowledgements

We thank Patricia Clark for advice and Ke Wan in the Bardwell lab for protein purification. JCAB is a Howard Hughes Medical Institute Investigator. This work was also supported by Wacker Chemie AG, and the Intramural Research Program of the National Institute of Diabetes and Digestive and Kidney Diseases. The authors have no conflicts of interest.

Author contributions

ZY and JCAB designed the research. ZY, SH and XW performed the experiments. ZY, HDB and JCAB analyzed data. ZY, HDB and JCAB wrote the manuscript with input from all authors.

Data availability statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

References

- Albenne, C. and Ieva, R. (2017) Job contenders: roles of the beta-barrel assembly machinery and the translocation and assembly module in autotransporter secretion. *Molecular Microbiology*, **106**, 505–517.
- Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., *et al.* (2006) Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. *Molecular Systems Biology*, **2**, 2006.0008.
- Babu, M., Bundalovic-Torma, C., Calmettes, C., Phanse, S., Zhang, Q., Jiang, Y., *et al.* (2018) Global landscape of cell envelope protein complexes in *Escherichia coli*. *Nature Biotechnology*, **36**, 103–112.
- Bader, M.W., Navarre, W.W., Shiau, W., Nikaido, H., Frye, J.G., McClelland, M., *et al.* (2003) Regulation of *Salmonella typhimurium* virulence gene expression by cationic antimicrobial peptides. *Molecular Microbiology*, **50**, 219–230.
- Bamert, R.S., Lundquist, K., Hwang, H., Webb, C.T., Shiota, T., Stubenrauch, C.J., *et al.* (2017) Structural basis for substrate selection by the translocation and assembly module of the beta-barrel assembly machinery. *Molecular Microbiology*, **106**, 142–156.
- Barnard, T.J., Dautin, N., Lukacik, P., Bernstein, H.D. and Buchanan, S.K. (2007) Autotransporter structure reveals intra-barrel cleavage followed by conformational changes. *Nature structural & molecular biology*, **14**, 1214–1220.
- Bernstein, H.D. (2015) Looks can be deceiving: recent insights into the mechanism of protein secretion by the autotransporter pathway. *Molecular Microbiology*, **97**, 205–215.
- Bernstein, H.D. (2019) Type V secretion in gram-negative bacteria. *EcoSal Plus*, **8**(2). doi: 10.1128/ecosalplus.ESP-0031-2018.
- Bokinsky, G., Peralta-Yahya, P.P., George, A., Holmes, B.M., Steen, E.J., Dietrich, J., *et al.* (2011) Synthesis of three advanced biofuels from ionic liquid-pretreated switchgrass

- using engineered *Escherichia coli*. *Proceedings of the National Academy of Sciences of the United States of America*, **108**, 19949–19954.
- Braselmann, E., Chaney, J.L., Champion, M.M. and Clark, P.L. (2016) DegP chaperone suppresses toxic inner membrane translocation intermediates. *PLoS ONE*, **11**, e0162922.
- Burmam, B.M., Wang, C. and Hiller, S. (2013) Conformation and dynamics of the periplasmic membrane-chaperone complexes OmpX-Skp and tOmpA-Skp. *Nature Structural & Molecular Biology*, **20**, 1265–1272.
- Celik, N., Webb, C.T., Leyton, D.L., Holt, K.E., Heinz, E., Gorrell, R., *et al.* (2012) A bioinformatic strategy for the detection, classification and analysis of bacterial autotransporters. *PLoS ONE*, **7**, e43245.
- Cheng, C.M., Tzou, S.C., Zhuang, Y.H., Huang, C.C., Kao, C.H., Liao, K.W., *et al.* (2014) Functional production of a soluble and secreted single-chain antibody by a bacterial secretion system. *PLoS ONE*, **9**, e97367.
- Cho, S.H., Szewczyk, J., Pesavento, C., Zietek, M., Banzhaf, M., Roszczenko, P., *et al.* (2014) Detecting envelope stress by monitoring beta-barrel assembly. *Cell*, **159**, 1652–1664.
- Datsenko, K.A. and Wanner, B.L. (2000) One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proceedings of the National Academy of Sciences of the United States of America*, **97**, 6640–6645.
- Dautin, N. and Bernstein, H.D. (2007) Protein secretion in gram-negative bacteria via the autotransporter pathway. *Annual Review of Microbiology*, **61**, 89–112.
- Denoncin, K., Schwalm, J., Vertommen, D., Silhavy, T.J. and Collet, J.F. (2012) Dissecting the *Escherichia coli* periplasmic chaperone network using differential proteomics. *Proteomics*, **12**, 1391–1401.
- Diderichsen, B. (1980) flu, a metastable gene controlling surface properties of *Escherichia coli*. *Journal of Bacteriology*, **141**, 858–867.
- Dong, T. and Schellhorn, H.E. (2009) Global effect of RpoS on gene expression in pathogenic *Escherichia coli* O157:H7 strain EDL933. *BMC Genomics*, **10**, 349.
- Fan, E., Chauhan, N., Udath, D.B., Leo, J.C. and Linke, D. (2016) Type V secretion systems in bacteria. *Microbiology Spectrum*, **4**(1). doi: 10.1128/microbiolspec.VMBF-0009-2015
- Goemans, C., Denoncin, K. and Collet, J.F. (2014) Folding mechanisms of periplasmic proteins. *Biochimica et Biophysica Acta*, **1843**, 1517–1528.
- Gupta, S., Adlakha, N. and Yazdani, S.S. (2013) Efficient extracellular secretion of an endoglucanase and a beta-glucosidase in *E. coli*. *Protein Expression and Purification*, **88**, 20–25.
- Henderson, I.R. and Nataro, J.P. (2001) Virulence functions of autotransporter proteins. *Infection and Immunity*, **69**, 1231–1243.
- Henderson, I.R., Navarro-Garcia, F., Desvaux, M., Fernandez, R.C. and Ala'Aldeen, D. (2004) Type V protein secretion pathway: the autotransporter story. *Microbiology and Molecular Biology Reviews*, **68**, 692–744.
- Heras, B., Totsika, M., Peters, K.M., Paxman, J.J., Gee, C.L., Jarrott, R.J., *et al.* (2014) The antigen 43 structure reveals a molecular Velcro-like mechanism of autotransporter-mediated bacterial clumping. *Proceedings of the National Academy of Sciences of the United States of America*, **111**, 457–462.
- Hong, W., Jiao, W., Hu, J., Zhang, J., Liu, C., Fu, X., *et al.* (2005) Periplasmic protein HdeA exhibits chaperone-like activity exclusively within stomach pH range by transforming into disordered conformation. *Journal of Biological Chemistry*, **280**, 27029–27034.
- Hussain, S. and Bernstein, H.D. (2018) The Bam complex catalyzes efficient insertion of bacterial outer membrane proteins into membrane vesicles of variable lipid composition. *Journal of Biological Chemistry*, **293**, 2959–2973.
- Ieva, R. and Bernstein, H.D. (2009) Interaction of an autotransporter passenger domain with BamA during its translocation across the bacterial outer membrane. *Proceedings of the National Academy of Sciences of the United States of America*, **106**, 19120–19125.
- Ieva, R., Skillman, K.M. and Bernstein, H.D. (2008) Incorporation of a polypeptide segment into the β -domain pore during the assembly of a bacterial autotransporter. *Molecular microbiology*, **67**, 188–201.
- Ieva, R., Tian, P., Peterson, J.H. and Bernstein, H.D. (2011) Sequential and spatially restricted interactions of assembly factors with an autotransporter beta domain. *Proceedings of the National Academy of Sciences of the United States of America*, **108**, E383–E391.
- Jain, S. and Goldberg, M.B. (2007) Requirement for YaeT in the outer membrane assembly of autotransporter proteins. *Journal of Bacteriology*, **189**, 5393–5398.
- Kang'ethe, W. and Bernstein, H.D. (2013) Charge-dependent secretion of an intrinsically disordered protein via the autotransporter pathway. *Proceedings of the National Academy of Sciences of the United States of America*, **110**, E4246–E4255.
- Kern, R., Malki, A., Abdallah, J., Tagourti, J. and Richarme, G. (2007) *Escherichia coli* HdeB is an acid stress chaperone. *Journal of Bacteriology*, **189**, 603–610.
- Kim, Y.E., Hipp, M.S., Bracher, A., Hayer-Hartl, M. and Hartl, F.U. (2013) Molecular chaperone functions in protein folding and proteostasis. *Annual Review of Biochemistry*, **82**, 323–355.
- Konovalova, A., Kahne, D.E. and Silhavy, T.J. (2017) Outer Membrane Biogenesis. *Annual Review of Microbiology*, **71**, 539–556.
- Konovalova, A., Perlman, D.H., Cowles, C.E. and Silhavy, T.J. (2014) Transmembrane domain of surface-exposed outer membrane lipoprotein RcsF is threaded through the lumen of beta-barrel proteins. *Proceedings of the National Academy of Sciences of the United States of America*, **111**, E4350–E4358.
- Kotzsch, A., Vernet, E., Hammarstrom, M., Berthelsen, J., Weigelt, J., Graslund, S. and Sundstrom, M. (2011) A secretory system for bacterial production of high-profile protein targets. *Protein Science*, **20**, 597–609.
- Lazar, S.W. and Kolter, R. (1996) SurA assists the folding of *Escherichia coli* outer membrane proteins. *Journal of Bacteriology*, **178**, 1770–1773.
- Lennon, C.W., Thamsen, M., Friman, E.T., Cacciaglia, A., Sachsenhauser, V., Sorgenfrei, F.A., *et al.* (2015) Folding optimization in vivo uncovers new chaperones. *Journal of Molecular Biology*, **427**, 2983–2994.

- Leyton, D.L., Rossiter, A.E. and Henderson, I.R. (2012) From self sufficiency to dependence: mechanisms and factors important for autotransporter biogenesis. *Nature Reviews Microbiology*, **10**, 213–225.
- Mee-Ngan, Y.F. and Bernstein, H.D. (2013) Mutations in the *Escherichia coli* ribosomal protein L22 selectively suppress the expression of a secreted bacterial virulence factor. *Journal of bacteriology*, **195**, 2991–2999.
- Mendez, J., Cascales, D., Garcia-Torrico, A.I. and Guijarro, J.A. (2018) Temperature-dependent gene expression in *Yersinia ruckeri*: tracking specific genes by bioluminescence during in vivo colonization. *Frontiers in Microbiology*, **9**, 1098.
- Miller, S.I. and Salama, N.R. (2018) The gram-negative bacterial periplasm: size matters. *PLoS Biology*, **16**, e2004935.
- Missiakas, D., Betton, J.M. and Raina, S. (1996) New components of protein folding in extracytoplasmic compartments of *Escherichia coli* SurA, FkpA and Skp/OmpH. *Molecular Microbiology*, **21**, 871–884.
- Noinaj, N., Gumbart, J.C. and Buchanan, S.K. (2017) The beta-barrel assembly machinery in motion. *Nature Reviews Microbiology*, **15**, 197–204.
- Oomen, C.J., van Ulsen, P., Van Gelder, P., Feijen, M., Tommassen, J. and Gros, P. (2004) Structure of the translocator domain of a bacterial autotransporter. *The EMBO journal*, **23**, 1257–1266.
- Owen, P., Meehan, M., de Loughry-Doherty, H. and Henderson, I. (1996) Phase-variable outer membrane proteins in *Escherichia coli*. *FEMS Immunology and Medical Microbiology*, **16**, 63–76.
- Pavlova, O., Peterson, J.H., Ieva, R. and Bernstein, H.D. (2013) Mechanistic link between beta barrel assembly and the initiation of autotransporter secretion. *Proceedings of the National Academy of Sciences of the United States of America*, **110**, E938–E947.
- Piazza, I., Kochanowski, K., Cappelletti, V., Fuhrer, T., Noor, E., Sauer, U., et al. (2018) A map of protein-metabolite interactions reveals principles of chemical communication. *Cell*, **172**, 358–372.e23.
- Pohlner, J., Halter, R., Beyreuther, K. and Meyer, T.F. (1987) Gene structure and extracellular secretion of *Neisseria gonorrhoeae* IgA protease. *Nature*, **325**, 458–462.
- Qian, Z.G., Xia, X.X., Choi, J.H. and Lee, S.Y. (2008) Proteome-based identification of fusion partner for high-level extracellular production of recombinant proteins in *Escherichia coli*. *Biotechnology and Bioengineering*, **101**, 587–601.
- Quan, S., Hiniker, A., Collet, J.F. and Bardwell, J.C. (2013) Isolation of bacteria envelope proteins. *Methods in Molecular Biology*, **966**, 359–366.
- Quan, S., Koldewey, P., Tapley, T., Kirsch, N., Ruane, K.M., Pfizenmaier, J., et al. (2011) Genetic selection designed to stabilize proteins uncovers a chaperone called Spy. *Nature Structural & Molecular Biology*, **18**, 262–269.
- Rizzitello, A.E., Harper, J.R. and Silhavy, T.J. (2001) Genetic evidence for parallel pathways of chaperone activity in the periplasm of *Escherichia coli*. *Journal of Bacteriology*, **183**, 6794–6800.
- Roman-Hernandez, G., Peterson, J.H. and Bernstein, H.D. (2014) Reconstitution of bacterial autotransporter assembly using purified components. *Elife*, **3**, e04234.
- Ruiz, N., Kahne, D. and Silhavy, T.J. (2006) Advances in understanding bacterial outer-membrane biogenesis. *Nature Reviews Microbiology*, **4**, 57–66.
- Ruiz-Perez, F., Henderson, I.R., Leyton, D.L., Rossiter, A.E., Zhang, Y. and Nataro, J.P. (2009) Roles of periplasmic chaperone proteins in the biogenesis of serine protease autotransporters of Enterobacteriaceae. *Journal of Bacteriology*, **191**, 6571–6583.
- Schiffirin, B., Calabrese, A.N., Devine, P.W.A., Harris, S.A., Ashcroft, A.E., Brockwell, D.J., et al. (2016) Skp is a multivalent chaperone of outer-membrane proteins. *Nature Structural & Molecular Biology*, **23**, 786–793.
- Selkig, J., Mosbahi, K., Webb, C.T., Belousoff, M.J., Perry, A.J., Wells, T.J., et al. (2012) Discovery of an archetypal protein transport system in bacterial outer membranes. *Nature Structural & Molecular Biology*, **19**, 506–510, S501.
- Sklar, J.G., Wu, T., Kahne, D. and Silhavy, T.J. (2007) Defining the roles of the periplasmic chaperones SurA, Skp, and DegP in *Escherichia coli*. *Genes & Development*, **21**, 2473–2484.
- Spiess, C., Beil, A. and Ehrmann, M. (1999) A temperature-dependent switch from chaperone to protease in a widely conserved heat shock protein. *Cell*, **97**, 339–347.
- Stull, F., Betton, J.M. and Bardwell, J.C.A. (2018) Periplasmic chaperones and prolyl isomerases. *EcoSal Plus*, **8**. doi: 10.1128/ecosalplus.ESP-0005-2018.
- van den Berg, B. (2010) Crystal structure of a full-length autotransporter. *Journal of molecular biology*, **396**, 627–633.
- van der Woude, M.W. and Henderson, I.R. (2008) Regulation and function of Ag43 (flu). *Annual Review of Microbiology*, **62**, 153–169.
- Van Wielink, J.E. and Duine, J.A. (1990) How big is the periplasmic space? *Trends in Biochemical Sciences*, **15**, 136–137.
- Vertommen, D., Ruiz, N., Leverrier, P., Silhavy, T.J. and Collet, J.F. (2009) Characterization of the role of the *Escherichia coli* periplasmic chaperone SurA using differential proteomics. *Proteomics*, **9**, 2432–2443.
- Vo, J.L., Martinez Ortiz, G.C., Subedi, P., Keerthikumar, S., Mathivanan, S., Paxman, J.J. and Heras, B. (2017) Autotransporter adhesins in *Escherichia coli* pathogenesis. *Proteomics*, **17**. doi: 10.1002/pmic.201600431.
- Walton, T.A., Sandoval, C.M., Fowler, C.A., Pardi, A. and Sousa, M.C. (2009) The cavity-chaperone Skp protects its substrate from aggregation but allows independent folding of substrate domains. *Proceedings of the National Academy of Sciences of the United States of America*, **106**, 1772–1777.
- Wells, T.J., Totsika, M. and Schembri, M.A. (2010) Autotransporters of *Escherichia coli*: a sequence-based characterization. *Microbiology*, **156**, 2459–2469.
- Wickner, W., Driessen, A.J. and Hartl, F.U. (1991) The enzymology of protein translocation across the *Escherichia coli* plasma membrane. *Annual Review of Biochemistry*, **60**, 101–124.
- Ye, Y., Gao, J., Jiao, R., Li, H., Wu, Q., Zhang, J. and Zhong, X. (2015) The membrane proteins involved in virulence of *Cronobacter sakazakii* virulent G362 and attenuated L3101 isolates. *Frontiers in Microbiology*, **6**, 1238.

- Yeats, C. and Bateman, A. (2003) The BON domain: a putative membrane-binding domain. *Trends in Biochemical Sciences*, **28**, 352–355.
- Yim, H.H. and Villarejo, M. (1992) osmY, a new hyperosmotically inducible gene, encodes a periplasmic protein in *Escherichia coli*. *Journal of Bacteriology*, **174**, 3637–3644.
- Yuan, X., Johnson, M.D., Zhang, J., Lo, A.W., Schembri, M.A., Wijeyewickrema, L.C., *et al.* (2018) Molecular basis for the folding of beta-helical autotransporter passenger domains. *Nature Communications*, **9**, 1395.
- Zhai, Y., Zhang, K., Huo, Y., Zhu, Y., Zhou, Q., Lu, J., *et al.* (2011) Autotransporter passenger domain secretion requires

a hydrophobic cavity at the extracellular entrance of the beta-domain pore. *The Biochemical Journal*, **435**, 577–587.

Zheng, Z., Chen, T., Zhao, M., Wang, Z. and Zhao, X. (2012) Engineering *Escherichia coli* for succinate production from hemicellulose via consolidated bioprocessing. *Microbial Cell Factories*, **11**, 37.

Supporting Information

Additional supporting information may be found online in the Supporting Information section at the end of the article.