

## ORIGINAL ARTICLE

# Expanding the genotypic spectrum of Jalili syndrome: Novel *CNNM4* variants and uniparental isodisomy in a north American patient cohort

Lev Prasov<sup>1,2,3</sup> | Ehsan Ullah<sup>1</sup> | Amy E. Turriff<sup>1</sup> | Blake M. Warner<sup>4</sup> |  
Julie Conley<sup>5</sup> | Paul R. Mark<sup>6</sup> | Robert B. Hufnagel<sup>1</sup> | Laryssa A. Huryn<sup>1</sup> 

<sup>1</sup>Ophthalmic Genetics and Visual Function Branch, National Eye Institute, National Institutes of Health, Bethesda, Maryland

<sup>2</sup>Department of Ophthalmology and Visual Sciences, W.K. Kellogg Eye Center, University of Michigan, Ann Arbor, Michigan

<sup>3</sup>Department of Human Genetics, University of Michigan, Ann Arbor, Michigan

<sup>4</sup>National Institute of Dental and Craniofacial Research, National Institutes of Health, Bethesda, Maryland

<sup>5</sup>Section of Pediatric Ophthalmology, Helen DeVos Children's Hospital, Grand Rapids, Michigan

<sup>6</sup>Spectrum Health Division of Medical Genetics, Grand Rapids, Michigan

## Correspondence

Laryssa A. Huryn, Ophthalmic Genetics and Visual Function Branch, National Eye Institute, National Institutes of Health, Building 10, Rm 10D45; 10 Center Dr., Bethesda, MD 20892. Email: laryssa.huryn@nih.gov

## Funding information

NEI Intramural Research Grant, Grant/Award Number: NEI K12-EY022299

## Abstract

Jalili syndrome is a rare multisystem disorder with the most prominent features consisting of cone-rod dystrophy and amelogenesis imperfecta. Few cases have been reported in the Americas. Here we describe a case series of patients with Jalili syndrome examined at the National Eye Institute's Ophthalmic Genetics clinic between 2016 and 2018. Three unrelated sporadic cases were systematically evaluated for ocular phenotype and determined to have cone-rod dystrophy with bull's eye maculopathy, photophobia, and nystagmus. All patients had amelogenesis imperfecta. Two of these patients had Guatemalan ancestry and the same novel homozygous *CNNM4* variant (p.Arg236Trp c.706C > T) without evidence of consanguinity. This variant met likely pathogenic criteria by the American College of Medical Genetics guidelines. An additional patient had a homozygous deleterious variant in *CNNM4* (c.279delC p.Phe93Leufs\*31), which resulted from paternal uniparental isodisomy for chromosome 2p22-2q37. This individual had additional syndromic features including developmental delay and spastic diplegia, likely related to mutations at other loci. Our work highlights the genotypic variability of Jalili syndrome and expands the genotypic spectrum of this condition by describing the first series of patients seen in the United States.

## KEYWORDS

amelogenesis, *CNNM4*, cone-rod dystrophy, Jalili syndrome, retinal degeneration, uniparental isodisomy

## 1 | INTRODUCTION

Jalili syndrome is a rare multisystem disorder characterized by a childhood cone-rod retinal degeneration and amelogenesis imperfecta (Jalili, 2010). It was initially described in a large extended family in Gaza, as an autosomal recessive condition in which all affected family members had photophobia, nystagmus, and achromatopsia along with abnormal tooth enamel (Jalili & Smith, 1988). Since that time, most cases have

been described in large consanguineous families on multiple continents, including one family in Newfoundland (Doucette et al., 2013) and one family in Guatemala (Parry et al., 2009), with few reported sporadic cases (Daneshmandpour, Darvish, Pashazadeh, & Emamalzadeh, 2019).

Though variants in many genes can lead to amelogenesis imperfecta or cone-rod dystrophy, the constellation of these two findings in the absence of other syndromic features is highly suggestive of one gene, *CNNM4*, as the underlying cause (Daneshmandpour et al., 2019;

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.


© 2020 The Authors. *American Journal of Medical Genetics Part A* published by Wiley Periodicals, Inc.

Parry et al., 2009; Polok et al., 2009). *CNNM4* encodes a transmembrane protein with a critical role in magnesium homeostasis and is expressed in photoreceptors and in the dental enamel (Gomez Garcia, Oyenarte, & Martinez-Cruz, 2011; Parry et al., 2009; Polok et al., 2009). Here, we describe a series of sporadic cases with Jalili syndrome in a cohort of patients originating in North and Central America. These patients carry novel variants in *CNNM4* and further highlight the variable phenotypic and genotypic spectrum of the condition.

## 2 | METHODS

This study was carried out according to the standards of the Common Rule of the United States Federal Government (46CFR45) under protocols approved by the Institutional Review Board at the National

Institutes of Health and informed consent was obtained for all participants. Patients were evaluated in the Ophthalmic Genetics and Visual Function Branch Clinic between 2016 and 2018. All patients underwent ophthalmic clinical evaluation, including best corrected visual acuity (BCVA) using the Early Treatment of Diabetic Retinopathy Study (ETDRS) chart and guidelines, refraction, color vision testing by monocular Farnsworth D15, slit-lamp biomicroscopy, dilated eye exam, Goldmann visual field testing, fundus color, and autofluorescence (FAF) imaging (Topcon, Tokyo, Japan; Optos, Dunfermline, Scotland), optical coherence tomography (Cirrus HD-OCT, Carl Zeiss Meditec, Dublin, CA; Bioptigen Inc., Research Triangle Park, NC), electroretinography (ERG) (LKC, Gaithersburg, MD), and medical evaluation by a clinical geneticist. ERGs were recorded using Burian-Allen electrodes and according to the International Society for Clinical Electrophysiology of Vision standards. Ocular, medical, and dental



**FIGURE 1** Ocular features of Jalili syndrome cases. (a-c) Multimodal imaging of the right eye of patient 1: SD-OCT (a) demonstrating ellipsoid zone disruption in the fovea; ring of hyper- hypo-autofluorescence on wide-field Optos autofluorescence (b) and bull's eye maculopathy on 50° color fundus photo (c) are all consistent with the cone-rod dystrophy. (d-f) Multimodal imaging of patient 2's left eye showing loss of foveal ellipsoid zone on OCT (d), with hypofluorescent sector of inferior retina on wide-field Optos autofluorescence (e), corresponding to the areas of atrophy and retinal pigment migration on color fundus photo (f). (g-i) Multimodal imaging of patient 3's right eye showing macular changes on Bioptigen OCT (g), with corresponding bull's eye hyper- hypo-autofluorescence pattern on wide-field Optos autofluorescence (h), and diffuse granular pigment changes and bull's eye maculopathy on wide-field Optos color photograph (i). All patients share bull's eye maculopathy, with loss of ellipsoid and interdigitation zone as evidenced by the hyporeflective dark area under the fovea on OCT. However, patient 2 also notably has a large hypoautofluorescent area of atrophy and pigment clumping in the periphery (e,f) [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

health records were also reviewed, including panoramic and bite wing X-rays when available. Patients without molecular results underwent clinical genetic testing with an inherited retinal dystrophy panel (Molecular Vision Laboratory [MVL]; Hillsboro, OR; Clinical Laboratory Improvement Amendments identifier, 38D2059762). *CNNM4* variants were evaluated for pathogenicity using established American College of Medical Genetics (ACMG) criteria (Richards et al., 2015).

### 3 | RESULTS

#### 3.1 | Case reports


##### 3.1.1 | Patient 1

Patient 1 is a 15-year-old girl born from Guatemalan parents who presented with poor central vision, nystagmus, and light sensitivity. Examination was notable for Snellen BCVA at distance of 20/250 in the right eye and 20/200 in the left eye. Auto-refraction showed a myopic refraction with astigmatism:  $-2.75 + 2.25 \times 103$  and  $-3.00 + 3.00 \times 076$  in the right and left eye, respectively. She had high frequency, low amplitude horizontal nystagmus and was orthophoric. Anterior segment exam was unremarkable, and fundus exam was notable for bull's eye maculopathy and granular pigment changes in the periphery. Spectral domain optical coherence tomography (SD-OCT) revealed loss of the ellipsoid zone in the fovea, and FAF demonstrated a bull's eye pattern of hyper- and hypo-autofluorescence (Figure 1). Farnsworth D15 color vision testing revealed multiple axis errors (achromatic pattern). Loss of the I1e isopter was noted on the Goldmann visual field testing in each eye, with preservation of I4e and V4e isopters. Scotopic electroretinography showed diminished amplitudes (~50%) and mildly delayed implicit times, while photopic bright flash and flicker ERG

responses were unrecordable. The dental examination and dental radiographs (bitewings, anterior and posterior periapical radiographs, and panoramic radiograph) revealed an entirely restored dentition with crowns on all molar teeth and composite restorations on the anterior teeth (Figure 2). The pulp chambers appear slightly enlarged on some teeth (20, 23, 26, 29), while others exhibit appearance within normal limits for this age (Figure 2, Figure S1). Clinically, the enamel is thin or absent on the non-restored surfaces of the teeth. Periapical and bitewing radiographs reveal isolated residual enamel on nonrestored and under restored surfaces, respectively (Figure S1, arrows). There were otherwise no systemic abnormalities noted. A 581 gene panel from Molecular Vision Lab (MVL Panel v1) revealed a homozygous *CNNM4* c.706C > T (p.Arg236Trp) variant with one copy inherited from each parent.

##### 3.1.2 | Patient 2

Patient 2 is a 16-year-old boy from Guatemala, who presented with poor central vision and light sensitivity. Examination was notable for Snellen BCVA at distance of 20/200 in the right eye and 20/160 in the left eye, with a manifest refraction of  $+1.00 + 4.50 \times 109$  and  $-0.25 + 4.25 \times 069$ , respectively. He had mild end-gaze nystagmus and was orthophoric. Anterior segment exam was unremarkable, and fundus exam was notable for bull's eye maculopathy, perivasculature, and segmental pigment deposition in the inferior retina, and mild vascular attenuation (Figure 1a-hyphen;d). OCT revealed loss of the ellipsoid zone in the fovea, and FAF imaging revealed a hypofluorescent sector corresponding to the areas of retinal atrophy and pigment deposition. Color vision testing by Farnsworth D15 revealed multiple axis errors. Goldmann visual fields demonstrated superior constriction, and loss of the I1e isopter centrally, consistent with the retinal changes inferiorly (Figure 1). Scotopic ERG showed mildly diminished amplitudes and



**FIGURE 2** Dental features of Jalili syndrome cases. (a,b) External photos of mouth of patient 1 (a) and patient 3 (b) showing tooth decay and crowns on multiple teeth. (c) Panoramic X-ray of patient 1 showing crowns on all adult molar teeth [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



delayed implicit times, while photopic bright flash and flicker ERG were extinguished. Dental examination revealed crowns on multiple teeth and the absence of enamel. There were otherwise no systemic abnormalities. Molecular Vision Lab NGS Retinal Dystrophy SmartPanel v11 (281 genes) revealed a homozygous *CNNM4* c.706C > T (p.Arg236Trp) variant with one copy inherited from each parent.

### 3.1.3 | Patient 3

Patient 3 is a 3-year-old boy born from a Puerto Rican father and Caucasian mother who presented to ophthalmology clinic with light sensitivity and multiple congenital anomalies. On examination, binocular visual acuity was 2.4cy/cm by Teller acuity cards (approximately 20/360 Snellen equivalent). Cycloplegic refraction showed myopia with astigmatism in both eyes:  $-3.25 + 2.25 \times 090$  and  $-3.25 + 2.50 \times 095$ , in the right and left eye, respectively. Anterior segment exam was unremarkable. Fundus exam was notable for diffuse granular pigment changes and bull's eye maculopathy, with typical corresponding bull's eye pattern on FAF (Figure 1). Bioptogen OCT revealed loss of the ellipsoid zone at the fovea (Figure 1). Dental examination revealed crowns on multiple molar teeth and yellow/opaque appearance of the anterior teeth with thinning and complete absence of enamel on some teeth (Figure 2). Systemic evaluation was notable for spastic paraplegia, developmental delay, and fatty liver. The patient was dependent on tracheostomy for respiration and gastrostomy tube for feeding. Clinical whole exome sequencing revealed homozygosity for a frameshift variant in *CNNM4*: c.279delC (p.Phe93Leufs\*31). Segregation analysis revealed that this variant was present in only the patient's father and not mother. Subsequent analysis examining other single nucleotide polymorphisms (SNPs) revealed paternal uniparental isodisomy (UPD) on chromosome 2p22-2q37. With the exception of homozygosity for the *CNNM4* variant, no additional variants within the UPD region were directly linked to the phenotypic features of the patient.

## 3.2 | Genetic analysis

We evaluated pathogenicity of *CNNM4* variants in each patient using established ACMG criteria. The *CNNM4* c.279delC (p.Phe93Leufs\*31) creates a frameshift early in the protein, which would be predicted to result in nonsense mediated decay of RNA or generation of a non-functional protein. The *CNNM4* c.706C > T (p.Arg236Trp) variant creates a nonconservative substitution in the protein in a highly conserved residue (GERP score 4.99) located in the DUF21 domain of the protein. The missense variant p.Arg236Trp was predicted to be damaging by 14 different in-silico prediction tools (SIFT, PolyPhen2, PROVEAN, DANN, DEOGEN2, EIGEN, FATHMM-MKL, M-CAP, MVP, MutationAssess, MutationTaster, REVEL, MetaSVM, MetaLR) whereas it was predicted benign by only one in-silico tool (PrimateAI) (Kopanos et al., 2019). Also, the missense variant p.Arg236Trp had a

high CADD score (Rentzsch, Witten, Cooper, Shendure, & Kircher, 2019) of 32, while the gnomAD missense Z score (2.37) indicates that *CNNM4* gene is moderately constrained against missense changes. The p.Arg236Trp variant was found to be absent in population databases comprised healthy (gnomAD, <https://gnomad.broadinstitute.org>) as well as diseased individuals (HGMD, <http://www.hgmd.cf.ac.uk>; ClinVar, <https://www.ncbi.nlm.nih.gov/clinvar/>). Notably, limited haplotype analysis on genetic panel data did not reveal additional homozygous variants in *CNNM4* or surrounding genes *ALMS1* and *MERTK* on chromosome 2, suggesting this is a founder mutation in the Guatemalan population rather than a result of consanguinity or previously unknown relatedness between families of patients 1 and 2. A previous variant in this same amino acid, *CNNM4* c.707G > A (p.Arg236Gln), segregated with disease in a large family with Jalili syndrome, with a 0.5<sup>17</sup> probability of cosegregation by chance (Polok et al., 2009). This variant was also absent from the gnomAD database. Based on 2015 ACMG/AMP variant interpretation criteria, *CNNM4* p.Phe93Leufs\*31 is considered pathogenic and p.Arg236Gln is considered likely pathogenic (Richards et al., 2015).

## 4 | DISCUSSION

The phenotypic spectrum of Jalili syndrome typically comprises cone-rod dystrophy and amelogenesis imperfecta, but there is significant variability in the severity of the retinal degeneration, even within families with the same mutation (Daneshmandpour et al., 2019; Gerth-Kahlert et al., 2015; Jalili, 2010). Consistent with this observation, we see a difference in structural and functional measures among two patients of similar age and demographic background carrying the same *CNNM4* p.Arg236Trp variant. While both have significant cone dysfunction and a bull's eye maculopathy, Patient 2 has a significant peripheral sector of retinal atrophy and pigment deposition with a corresponding visual field defect, while Patient 1 has disease mostly limited to the posterior pole. This suggests that factors beyond the genotype influence the disease severity. Both probands were of Guatemalan descent, denied consanguinity, and yet carried the same homozygous variant. This suggests a founder effect, though the ancestor may be more distant as neither patient carries homozygous variants in neighboring retinopathy genes *ALMS1* nor *MERTK*, which would constitute a ~39 Mb haplotype block. Likewise, the previously reported Guatemalan family carried two different mutations (Parry et al., 2009), suggesting that there are at least 3 alleles present in this population.

We report the first case of uniparental isodisomy associated with Jalili syndrome. This patient had additional features including spastic paraplegia, fatty liver, and developmental delay, while both his father and mother were phenotypically normal. Though the patient carried a large stretch of UPD on chromosome 2p22-2q37, complete UPD of chromosome 2 can be tolerated (Ou et al., 2013; Zhang et al., 2019). Uniparental isodisomy causes disease by two predominant mechanisms, loss of heterozygosity uncovering rare deleterious variants and through disruption of imprinting. In our case, a frameshift variant in *CNNM4*, which was paternally inherited, was uncovered and led to

the Jalili syndrome phenotype. Exome failed to reveal any rare coding variants that were thought to cause the nonretinal and nondental phenotypes. Likewise, no imprinting disorders with overlapping features have been described for these regions on chromosome 2. It remains possible that homozygosity for noncoding variants explain the other neurological phenotypes.

Our work describes the detailed phenotypic features of this condition in an underreported patient population, identifies a novel mis-sense variant shared by two of the patients, and highlights an unusual case of UPD leading to the disorder. Future efforts will be necessary to identify the full spectrum of UPD-associated disorders on chromosome 2 and identify the precise cause of the additional features in this patient. Additionally, mechanistic studies exploring the structure and function of the DUF21 domain of *CNNM4* will be necessary to identify the function of this critical portion of the protein.

### ACKNOWLEDGEMENTS

The authors are grateful to the individuals and families for participating in the study; to Denise Cunningham, Mike Arango and the ophthalmic photographers and technicians at the National Institutes of Health Ophthalmic Genetics Clinic. Dr. Lev Prasov is supported in part by NEI K12-EY022299.

### WEB RESOURCES

gnomAD Database: <http://gnomad.broadinstitute.org/>

NCBI Human Reference Genome Build 37.1: <http://www.ncbi.nlm.nih.gov/genome/assembly/2928/>.

Clinvar: <https://www.ncbi.nlm.nih.gov/clinvar/>

HGMD: <http://www.hgmd.cf.ac.uk>

### DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

### ORCID

Laryssa A. Hury  <https://orcid.org/0000-0002-0309-9419>

### REFERENCES

- Daneshmandpour, Y., Darvish, H., Pashazadeh, F., & Emamalizadeh, B. (2019). Features, genetics and their correlation in Jalili syndrome: A systematic review. *Journal of Medical Genetics*, 56(6), 358–369.
- Doucette, L., Green, J., Black, C., Schwartzentruber, J., Johnson, G. J., Galutira, D., & Young, T. L. (2013). Molecular genetics of achromatopsia in Newfoundland reveal genetic heterogeneity, founder effects and the first cases of Jalili syndrome in North America. *Ophthalmic Genetics*, 34(3), 119–129.

- Gerth-Kahlert, C., Seebauer, B., Dold, S., Hanson, J. V., Wildberger, H., Sporri, A., ... Berger, W. (2015). Intra-familial phenotype variability in patients with Jalili syndrome. *Eye*, 29(5), 712–716.
- Gomez Garcia, I., Oyenarte, I., & Martinez-Cruz, L. A. (2011). Purification, crystallization and preliminary crystallographic analysis of the CBS pair of the human metal transporter *CNNM4*. *Acta Crystallographica Section F, Structural Biology and Crystallization Communications*, 67(3), 349–353.
- Jalili, I. K. (2010). Cone-rod dystrophy and amelogenesis imperfecta (Jalili syndrome): Phenotypes and environs. *Eye*, 24(11), 1659–1668.
- Jalili, I. K., & Smith, N. J. (1988). A progressive cone-rod dystrophy and amelogenesis imperfecta: A new syndrome. *Journal of Medical Genetics*, 25(11), 738–740.
- Kopanos, C., Tsiolkas, V., Kouris, A., Chapple, C. E., Albarca Aguilera, M., Meyer, R., & Massouras, A. (2019). VarSome: The human genomic variant search engine. *Bioinformatics*, 35(11), 1978–1980.
- Ou, X., Liu, C., Chen, S., Yu, J., Zhang, Y., Liu, S., & Sun, H. (2013). Complete paternal uniparental isodisomy for chromosome 2 revealed in a parentage testing case. *Transfusion*, 53(6), 1266–1269.
- Parry, D. A., Mighell, A. J., El-Sayed, W., Shore, R. C., Jalili, I. K., Dollfus, H., ... Inglehearn, C. F. (2009). Mutations in *CNNM4* cause Jalili syndrome, consisting of autosomal-recessive cone-rod dystrophy and amelogenesis imperfecta. *American Journal of Human Genetics*, 84(2), 266–273.
- Polok, B., Escher, P., Ambresin, A., Chouery, E., Bolay, S., Meunier, I., ... Schorderet, D. F. (2009). Mutations in *CNNM4* cause recessive cone-rod dystrophy with amelogenesis imperfecta. *American Journal of Human Genetics*, 84(2), 259–265.
- Rentzsch, P., Witten, D., Cooper, G. M., Shendure, J., & Kircher, M. (2019). CADD: Predicting the deleteriousness of variants throughout the human genome. *Nucleic Acids Research*, 47(D1), D886–D894.
- Richards, S., Aziz, N., Bale, S., Bick, D., Das, S., Gastier-Foster, J., ... Committee, A. L. Q. A. (2015). Standards and guidelines for the interpretation of sequence variants: A joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genetics in Medicine: Official Journal of the American College of Medical Genetics*, 17(5), 405–424.
- Zhang, X., Ding, Z., He, R., Qi, J., Zhang, Z., & Cui, B. (2019). Complete paternal Uniparental Disomy of chromosome 2 in an Asian female identified by short tandem repeats and whole genome sequencing. *Cytogenetic and Genome Research*, 157(4), 197–202.

### SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Prasov L, Ullah E, Turriff AE, et al. Expanding the genotypic spectrum of Jalili syndrome: Novel *CNNM4* variants and uniparental isodisomy in a north American patient cohort. *Am J Med Genet Part A*. 2020;182A:493–497. <https://doi.org/10.1002/ajmg.a.61484>