

Abstract

Whilst the prevalence of obesity has rapidly increased in the low- and middle-income countries of the Middle East and North Africa (MENA) and Latin America and the Caribbean (LAC) regions, child undernutrition remains a public health challenge. We examined region-specific sociodemographic determinants of this double burden of malnutrition, specifically, the co-occurrence of child stunting and overweight, using Demographic and Health Survey and Multiple Indicator Cluster Survey data (2003-2016) from 11 countries in the MENA (n=118,585) and 13 countries in the LAC (n=77,824) regions. We used multiple logistic regressions to model region-specific associations of maternal education, and household wealth with child nutritional outcomes (6-59 months). The prevalence of stunting, overweight, and their co-occurrence was 24%, 10%, 4.3% in children in the MENA region, respectively, and 19%, 5%, and 0.5% in children in the LAC region, respectively. In both regions, higher maternal education and household wealth were significantly associated with lower odds of stunting and higher odds of overweight. As compared to the poorest wealth quintiles, decreased odds of co-occurring stunting and overweight were observed among children from the 2nd, 3rd and 4th wealth quintiles in the LAC region. In the MENA region, this association was only statistically significant for the 2nd wealth quintile. In both regions, double burden was not statistically significantly associated with maternal education. The social patterning of co-occurring stunting and overweight in children varied across the two regions, indicating potential differences in the underlying etiology of the double burden across regions, and stages of the nutrition transition.

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1111/mcn.12923](https://doi.org/10.1111/mcn.12923)

Keywords: double burden, LAC, MENA, stunting, overweight, children

Author Manuscript

Background

A global nutrition transition is under way, characterized by a rapid shift in diet composition and behavioral trends towards increased intakes of foods high in energy in the form of saturated fats and sugars, and decreased intakes of nutrient- and fiber- rich foods (B. M. Popkin, 2004). This transition has translated into a rise in overweight across all age groups globally (Abarca-Gómez et al., 2017). Overweight and obesity prevalence are particularly high in the Middle East and North Africa (MENA) and Latin America and the Caribbean (LAC) regions, whilst stunting remains a public health challenge in subpopulations in these regions (Black et al., 2013; Farrag, Cheskin, & Farag, 2017; Ng et al., 2014; Rivera, Pedraza, Martorell, & Gil, 2014).

The co-occurrence of under- and over-nutrition within populations, households and even individuals has been described in these contexts, and termed as the double burden of malnutrition (B. M. Popkin, 2004). Population-level double burden refers to the prevalence of undernutrition and overweight in the same country or community. Whereas household-level double burden, has been defined as the co-existence of undernutrition in children and overweight in adults within the same household (Black et al., 2013; Gubert, Spaniol, Segall-Corrêa, & Pérez-Escamilla, 2017; Jones, Acharya, & Galway, 2016; Wojcicki, 2014). Both population- level and household level double burdens have been relatively well documented in the LAC region, with findings highlighting that stunting, anemia and overweight are prevalent in women and children in these countries (Atalah, Amigo, & Bustos, 2014; Conde & Monteiro, 2014; Freire, Silva-Jaramillo, Ramírez-Luzuriaga, Belmont, & Waters, 2014; Gubert et al., 2017; Kroker-Lobos, Pedroza-

Tobías, Pedraza, & Rivera, 2014; Ramirez-Zea, Kroker-Lobos, Close-Fernandez, & Kanter, 2014; Rivera et al., 2014; Sarmiento et al., 2014; Severi & Moratorio, 2014). The rapid increases in overweight and obesity in the LAC region have been paralleled by similar surges in the MENA region (Abarca-Gómez et al., 2017; Ng et al., 2014). Although one analysis found height-for-age z-score (HAZ) to be correlated with BMI-for-age z-score (BMI-Z) among children under the age of five (El Taguri et al., 2009), few studies have described double burdens in the MENA region.

Much of the literature has described the household- level double burden of maternal overweight and childhood stunting (El Kishawi, Soo, Abed, & Muda, 2016; Garrett & Ruel, 2003; Kosaka & Umezaki, 2017). Factors associated with this double-burden include: 1) urbanicity: with higher odds of double-burden in urban and peri-urban areas as compared to rural areas (Jones et al., 2016); 2) economic status (income and wealth): with some positive associations found between higher income and double burden whilst some studies show no association (Bassett, Romaguera, Giménez, Lobo, & Samman, 2014; Kosaka & Umezaki, 2017; Saibul et al., 2009); and, 3) maternal or head of household education: where the associations are mixed (Kosaka & Umezaki, 2017).

Few studies have thus far focused on individual-level double burdens such as overweight and anemia coinciding in the same individual (women or children) (Eckhardt, Torheim, Monterrubio, Barquera, & Ruel, 2008; Gartner et al., 2014; Pinhas-Hamiel et al., 2003), and the co-existence of stunting and overweight in individual children (Bates, Gjonca, & Leone, 2017; Fernald &

Neufeld, 2007; Mamabolo, Alberts, Steyn, Delemarre-van de Waal, & Levitt, 2005; Barry M Popkin, Richards, & Montiero, 1996; Said-Mohamed, Bernard, Ndzana, & Pasquet, 2012; Varela-Silva et al., 2012; Wang et al., 2009).

Bates et al. (2017) recently reported prevalences of stunted-overweight in under-5-year-old children from 79 low- and middle-income countries, and estimated that approximately ten million children are concurrently stunted and overweight in these countries (Bates et al., 2017). The authors argue that this is a neglected phenomenon that requires further attention and that more research into the determinants of co-occurring stunting and overweight is needed (Bates et al., 2017).

Household wealth and maternal education are well documented key determinants of childhood nutritional status; both stunting and overweight (Black et al., 2013; Hong & Mishra, 2006; Keino, Plasqui, Ettyang, & van den Borne, 2014; Vollmer et al., 2014). Lower wealth and maternal education are both associated with higher prevalence of stunting, whereas the associations of these factors with childhood overweight are not consistent (Keino et al., 2014; Kosaka & Umezaki, 2017; Tzioumis, Kay, Bentley, & Adair, 2016). Differentials exist by level of economic development and stage of the nutrition transition; in low-income countries, most studies point to positive associations between household wealth and maternal education and childhood overweight, whereas in high-income countries, overweight appears to be more common in low socioeconomic and education groups, with mixed results in middle-income countries that have faced rapid changes in food environments and parallel increases in the

prevalence of overweight (Martorell, Khan, Hughes, & Grummer-Strawn, 2000; Subramanian, Perkins, Ozaltin, & Davey Smith, 2011). Some studies also suggest that there may be interactions between wealth and education in overweight differentials (Aitsi-Selmi et al., 2012). Household wealth and maternal education are thus likely to also explain variability in the double burden of stunting and overweight in children, yet few studies have explored these (Fernald & Neufeld, 2007).

The aim of the present analysis is to describe the double burden of stunting and overweight in under-5-year-old children at both the population and individual level in low and middle income countries (LMIC) of the MENA and LAC regions. We also examine the associations between household wealth and maternal education, and stunting, overweight and their co-occurrence in children across the two regions.

Methods

Data sources

Publicly available data from the Multiple Indicator Cluster Surveys (MICS available- <http://mics.unicef.org/> Accessed 2018 June 1) and the Demographic and Health Surveys (DHS available- <https://dhsprogram.com/Data/>. Accessed 2018 June 1) were used. Both MICS and DHS are household survey initiatives that collect comparable, nationally representative data from multiple countries including demographic, economic, health and nutrition data for women and children. The standardized designs of these surveys allow for cross-national comparisons of

indicators. Both MICS and DHS surveys use a multistage stratified cluster sampling design where each primary unit has a defined probability of selection.

Sample definition

We included the most recent publicly available datasets from countries of the MENA and LAC regions, that collected anthropometric data for under-5 children (length/height, weight), and household data on assets for the construction of a wealth index. Eight MICS and four DHS datasets from the MENA region and seven MICS and six DHS datasets from the LAC region were available and met inclusion criteria. We used the anthropometry module from the questionnaire for under-5 children, and the education and household characteristics modules from the household questionnaire. Data on child anthropometric variables, age, gender, maternal education, household assets and urbanicity were extracted and coded to harmonize categorization of variables across datasets. The datasets from each region were independently pooled into 2 regional (MENA and LAC) datasets to produce of region-specific estimates with high precision. We excluded all infants under the age of 6 months from this analysis. Table 1 lists the datasets used and sample size of the datasets across the different countries for 6-59-month-old children.

Measures

Dependent variables

To analyze the nutritional status of children aged 6-59 months, indicators for stunting and overweight were derived according to WHO Child Growth standards (WHO, 2015). Z-scores for

height-for-age (HAZ) and weight-for-height (WHZ) were calculated using the Stata command *zscore06*. Stunting was defined as HAZ < 2 SD below the reference median value, and overweight as WHZ > 2 SD above the median. Outliers were defined as any index value with a z-score > 6 SD or < 6 SD away from the median and were therefore excluded from the analysis. Child-level double burden was defined as the co-occurrence of stunting and overweight in a single individual, according to the above classification assuming these conditions are independent from one another. If any of the measurements used to calculate the standardized z-scores were missing, the outcome variable was not generated for that particular individual.

Independent variables

The independent variables of interest were maternal education and wealth, the primary social determinants of child malnutrition. Maternal education was standardized across all datasets as a categorical variable (no education/primary education/secondary education or more). Wealth index is a measure of household's cumulative living standards and is calculated using Principal Component Analysis (PCA) by DHS and MICS. The standard wealth index is based on household ownership of durable assets (such as a television and a refrigerator), materials used for housing construction (such as the type of floor) and variables such as household's main source of drinking water and sanitation facilities (Filmer & Pritchett, 2001; Rutstein & Johnson, 2004). We re-calculated the wealth index by excluding hygiene and sanitation variables, in order to explore the individual associations between these variables and nutritional outcomes. The resulting index captures the ownership of assets (namely, radio, television, refrigerator, bicycle, motorcycle, and

car), type of flooring, and access to electricity. This set of factors reflects information that was uniformly collected in all countries. The first principal component was used to predict a unidimensional index, which was then categorized into five quintiles.

Hygiene was represented by a dichotomous indicator measuring the source of water (piped or otherwise) while sanitation was represented by a dichotomous indicator measuring the type of toilet (piped sewage/septic tank or otherwise). A dichotomous variable was created for urban-rural dwelling. Child age was categorized into 3 categories (age 6-11 months, 12 to 23 months and 24 to 59 months) and treated as a discrete variable.

Statistical analyses

Analyses were conducted on Stata 13.0 (StataCorp, College Station, TX, USA). The prevalence of single and double burdens was estimated for each country. A total prevalence estimate (unweighted by country size) was also generated for each region. We calculated the expected prevalence of double burden within each dataset by multiplying the prevalence of overweight by the prevalence of stunting in children. The difference between observed and expected prevalences of the double burden was calculated for each country.

Regional pooled datasets were then used to examine the correlates of stunting, overweight and child-level double burden. Bivariate analyses were used to describe the outcomes of interest across the selected social determinants and the χ^2 test was used to detect statistically significant

associations between wealth or education and nutritional status. Bivariate logistic regression models were used to calculate unadjusted odds ratios (ORs) of the association between the various socio-demographic determinants and each of the three outcomes: child stunting, overweight, and double burden of stunting and overweight. Multivariable logistic regression models of the correlates of stunting, overweight and child-level double burden included variables with a strong theoretical basis: wealth, maternal education, hygiene, sanitation, urbanicity, age and gender. All models were adjusted for country fixed effects and within country clustering effect.

Interactions between maternal education and wealth index were tested and data were stratified by maternal education level to further examine significant interactions between maternal education and wealth (Aitsi-Selmi, Bell, Shipley, & Marmot, 2014). All country-level estimates presented were weighted using in-country sampling weights as provided by MICS and DHS. A p-value of 0.05 was used to indicate statistical significance. All data were publicly available, therefore ethical review was not required.

Results

The analytical sample included 118,585 children aged 6-59 months in the MENA region and 77,824 children in the LAC region (Table 1).

Table 2 presents the prevalence of stunting, overweight and co-occurring stunting and overweight among 6-59-month-old children in both regions, alongside GDP per capita;

Supplementary Table 1 presents the prevalence of maternal education levels by country. The prevalence of stunting in the MENA region varied from 7.4% in Palestine to 49.1% in Yemen, and in the LAC region from 5.7% in Paraguay to 48.2% in Guatemala. Total (unweighted by country population size) regional prevalence estimates of stunting were slightly lower in the LAC region as compared to the MENA region. The prevalence of overweight in children in the MENA region was lowest in Yemen (2.1%) and highest in Syria (18.2%), whereas in the LAC region, it ranged from 3.2% in Guatemala and Haiti to 12.6% in Paraguay. Overall, total overweight prevalence was substantially lower in the LAC region (5.5%) as compared to the MENA region (10.0%). Consequently, the prevalence of the double burden was very low at 0.5% in the LAC region, whereas it was 4.3% in the MENA region. The prevalence of the double burden reached 1.9% in Barbados and 10.7% in Syria (numbers from Syria are from data collected in 2006 and pre-date the current ongoing conflict).

The differences between observed and expected prevalence of stunted-overweight children ranged from 0.1 to 5.5% in the MENA region with a mean of 1.9%, while for the LAC region the difference ranged from -1.0 to 1.2% with a mean of -0.5% (Table 2). The observed prevalence of the double burden exceeded the expected prevalence in 5 of the 13 LAC countries, and in all of the 11 MENA countries studied.

Supplementary Table 2 presents bivariate analyses from aggregated LAC and MENA datasets showing associations between hypothesized correlates and stunting, overweight and child-level double burden.

Multivariable models then assessed differentials in stunting, overweight and co-occurring stunting and overweight by wealth quintiles (Figure 1). Across both datasets, children in the lowest quintiles had the highest odds of being stunted. In the MENA region, children belonging to the richest subgroup had lower odds to be stunted as compared to children living in the poorest subgroup (0.63; 95%CI 0.58-0.69) (Figure 1A). For the LAC region, the association was even stronger (0.32; 95%CI 0.28-0.37) (Figure 1D).

Wealth inequalities were also observed in overweight. The correlation between overweight and wealth was more pronounced in the LAC region as compared to the MENA region. The richest subgroup in the MENA region was found to be 1.15 times more likely to be overweight as compared to the poorest subgroup (Figure 1B) whereas their LAC counterparts were 1.72 times more likely to be overweight as compared to the poorest subgroup (Figure 1E). Co-occurring stunting and overweight among children was correlated with wealth in both regions. As compared to the poorest quintiles, lower odds of the stunted-overweight double burden were observed among children from the 2nd, 3rd and 4th quintiles in the LAC region (Figure 1F). In the MENA region, this association was only statistically significant for the 2nd wealth quintile (Figure 1C).

Differentials in child nutritional status were also investigated by maternal education for both regions (Figure 2). Across both regions, multivariable analysis showed that higher maternal education was associated with a statistically significantly lower odds of stunting (Figures 2A and 2D). This association was stronger in the LAC region (0.39; 95%CI 0.35-0.44 for secondary

education). The odds of child overweight increased with increased maternal education level in both regions. Secondary maternal education level was more strongly associated with child overweight in the LAC region (1.42; 95%CI 1.16-1.75) (Figure 2E) than in the MENA region (1.14; 95%CI 1.04-1.23) (Figure 2B). Maternal education was not statistically significantly associated with the co-occurrence of stunting and overweight in either region (Figures 2C and 2F).

To examine whether the association between wealth and child nutritional status outcomes varied according to maternal education, tests for interaction between wealth and maternal education were found to be significant for the MENA but not the LAC region. Data were stratified by maternal education level (none, primary, secondary plus) for both regions for comparative purposes. Within these strata, we examined the association of wealth with stunting, overweight and the double burden while adjusting for sex, age, urbanicity, hygiene and country effect (Figure 3). In the MENA region, the association between wealth and stunting was attenuated with increasing maternal education; in children of women with no education, odds of stunting were 0.5 (95%CI:0.4-0.6) in the richest compared to the poorest quintile; whereas in children of women with secondary education or higher, these odds were 0.7 (95%CI: 0.6-0.8) (Figures 3A, 3B, 3C); this was not the case in the LAC region (Figures 3D, 3E, 3F). Odds of overweight increased with increasing wealth quintiles among mothers educated to the primary level in both the MENA and LAC regions (Figures 3H, 3K). For mothers with higher education levels in the

LAC region, the association between overweight and wealth followed a similar pattern (Figures 3J, 3K, 3L) while there was no association in the MENA region (Figures 3G, 3H, 3I).

In the MENA region, stunted-overweight children followed a similar pattern to overweight children (Figures, 3M, 3N, 3O). However, the number of children with double burden in the LAC region was too small to allow us to explore these associations (Figures, 3P, 3Q, 3R).

Discussion

We present here a comparative analysis of the associations of socio-demographic characteristics with the double burden of malnutrition across two global regions where the nutrition transition has been documented to be under way (Abarca-Gómez et al., 2017). Whereas the literature generally describes prevalence estimates and studies burden associations within countries, this is the first paper to systematically compare regional patterns in child stunting, overweight and the double burden, and assess their determinants in the LAC and MENA regions.

Our findings confirm that in these regions, chronic undernutrition (stunting) is consistently rooted in poverty, with the odds of stunting significantly decreasing with increasing wealth and maternal education, consistent with global literature (Hong, 2007; Keino et al., 2014; D Makoka, 2013; Semba et al., 2008; Tzioumis et al., 2016; Vollmer et al., 2014). Our results also highlight the differential association of maternal education on stunting across the two regions. In the MENA region particularly, the effect of wealth on stunting is attenuated by maternal education (Figure 3), adding to the global evidence on the particular importance of maternal education for

the prevention of child stunting in this region (Abuya, Ciera, & Kimani-Murage, 2012; D. Makoka & Masibo, 2015; Negash, Whiting, Henry, Belachew, & Hailemariam, 2015; Semba et al., 2008; Wachs, Creed-Kanashiro, Cueto, & Jacoby, 2005).

While stunting is generally on the decline in these regions, obesity has steadily increased in adults and is increasingly reported amongst children (Abarca-Gómez et al., 2017; Black et al., 2013; Ng et al., 2014). We find the prevalence of overweight among under-5-year-old children to be higher in the MENA region as compared to LAC region. The findings also highlight stronger socioeconomic differentials in child overweight in the LAC region vis-à-vis the MENA region. This finding of generally high levels of overweight in the MENA region aligns with other evidence from the region highlighting elevated prevalence of obesity in children and women, and may be explained by the particularly high penetration of globalized food markets in the MENA region, and weaker policies to promote breastfeeding and regulate unhealthy foods, including to infants and children (Abarca-Gómez et al., 2017; Akik, Ghattas, Filteau, & Knai, 2017; Colchero, Rivera-Dommarco, Popkin, & Ng, 2017; Fraser, 2013). Additional cultural factors such as notions of beauty and health, and social restrictions on physical activity may contribute to these differences across regions (Obermeyer, Bott, & Sassine, 2015; Sharara, Akik, Ghattas, & Obermeyer, 2018). It may be that the co-occurrence of stunting and overweight which was reported in LAC populations a decade ago has stimulated some action at the policy level to reduce both these burdens (Hodge, Verstraeten, & Ochoa-Avilés, 2016; Hoey & Pelletier, 2011; Huicho et al., 2016).

The prevalence of co-occurring stunting and overweight among children is consequently low in the LAC region and may not have allowed sufficient statistical power for us to adequately investigate social determinants; whereas the numbers appear to represent a greater burden in the MENA region.

It has been argued that double burdens may not be an independent phenomenon, but rather a by-product of the differences in pace at which chronic malnutrition is disappearing and overnutrition is appearing (Dieffenbach & Stein, 2012). These studies, focusing on the intra-household stunted child-overweight mother double burden have shown that in most countries, the computed observed prevalence of stunted-child, overweight-mother pairs does not surpass the expected prevalence (Dieffenbach & Stein, 2012; Rivera et al., 2014). In the case of the present study, we observed a difference of around 2% between the expected and observed prevalence of stunted-overweight children in the MENA region (Table 2), while the difference was close to 0 in the LAC region. These findings suggest that in the MENA region, the stunted-overweight child is a true phenomenon rather than a statistical artifact. Potential explanations for the existence of the stunted-overweight child include poor quality diets with limiting micronutrients, but sufficient caloric density (Fernald & Neufeld, 2007; Mamabolo et al., 2005; Barry M Popkin et al., 1996; Said-Mohamed et al., 2012; Varela-Silva et al., 2012; Wang et al., 2009), in-utero epigenetic alterations (Barker, 2006; Kitsiou-Tzeli & Tzetis, 2017), or poor early nutrition that alters the physiology of stunted children to accumulate fat instead of lean mass (Freire et al., 2014; Kroker-Lobos et al., 2014; Ramirez-Zea et al., 2014; Tzioumis et al., 2016).

Our data further suggest that the co-occurrence of stunting and overweight among children is driven differentially in each region by the underlying burdens. In the LAC region, the direction of the associations of wealth and maternal education with the double burden follow those of stunting. Whereas in the MENA region, these associations follow those of overweight, even after stratifying for maternal education. This may reflect different etiologies underlying the double burden, the more advanced nutrition transition of the MENA region as compared to the LAC region, or that these analyses are limited by the small samples of stunted-overweight children.

In the present analysis, children with the double burden were included in both the stunted only and the overweight only categories, although it has been suggested that the stunted-overweight double burden should be explored using mutually exclusive categories in order to better understand its etiology. However, when these categories were generated, the results remained unchanged for all the analyses presented (data not shown).

The advantages of using DHS and MICS datasets include the comparability across regions of surveys having used standardized methodologies, on representative samples. Our findings are limited, however, by the cross-sectional nature of the data, and our inability to examine temporal trends. Although we did not aim to capture changes in patterns and trends over time, it would be worth conducting such analysis on longitudinal datasets to better elucidate the effects of the nutrition transition on the stunted-overweight double burden. And though DHS and MICS are highly relied upon surveys for the estimation of child malnutrition globally, it is important to note that any systematic error that may have occurred in height measurement could exaggerate

bias in the estimation of double burden prevalence. It should also be noted that although country-level analyses utilized sampling weights provided by MICS and DHS, the pooled analyses do not adjust for population size of each country, as this would require country-level data on population size for under-5-year-old children. Another limitation of this analysis is the variability in the waves of MICS and DHS surveys included. Although we captured the most recent available surveys for both regions, for countries such as Syria and Yemen, these data pre-date ongoing conflicts and are unlikely to represent the current nutritional situation of children, which has been catastrophically affected (El Bcheraoui, Jumaan, Collison, Daoud, & Mokdad, 2018; Eshaq, Fothan, Jensen, Khan, & AlAmodi, 2017; Meiqari, Hoetjes, Baxter, & Lenglet, 2018).

Conclusion

The systematic comparison of stunting, overweight and stunted-overweight burdens across two rapidly developing regions has highlighted that despite being on a similar development trajectory, nutritional burdens vary greatly and are driven differentially within and across the LAC and MENA regions. This highlights the need for regional or country-level focus on these determinants, rather than global analyses which may be divorced from local contexts and make decision-making and specific policy formulation/recommendations more challenging.

The high prevalence of stunted-overweight children in the MENA region is of public health importance and requires further research on the pathways leading to this double burden, as well as its short- and long-term impacts on health and development of these children. Policies and programs aiming to promote breastfeeding, and regulate the marketing and sales of unhealthy foods to infants and children are needed, and there may be lessons to derive on this from the LAC region.

Targeting population nutritional intervention programs to both stunting and overweight is a challenge and should take into account the possible coexistence of under- and over-nutrition in the same household or individual. This is particularly relevant at a time when there have been shifts from in-kind food assistance to cash-based programs in many countries of these two regions.

Key Messages

- In the MENA region, the prevalence of stunted-overweight children was higher than expected.
- Wealth drives the co-occurrence of stunting and overweight among children in different directions in the MENA and LAC regions.
- Targeting population nutritional intervention programs to both stunting and overweight is a challenge and it will be important to account for the possible coexistence of under- and over-nutrition in the same child.

References

- Abarca-Gómez, L., Abdeen, Z. A., Hamid, Z. A., Abu-Rmeileh, N. M., Acosta-Cazares, B., Acuin, C., . . . Aguilar-Salinas, C. A. (2017). Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128·9 million children, adolescents, and adults. *The Lancet*.
- Abuya, B. A., Ciera, J., & Kimani-Murage, E. (2012). Effect of mother's education on child's nutritional status in the slums of Nairobi. *BMC Pediatr*, *12*(1), 80.
- Aitsi-Selmi, A., Bell, R., Shipley, M. J., & Marmot, M. G. (2014). Education modifies the association of wealth with obesity in women in middle-income but not low-income countries: an interaction study using seven national datasets, 2005-2010. *PLoS One*, *9*(3), e90403.
- Aitsi-Selmi, A., Chandola, T., Friel, S., Nouraei, R., Shipley, M. J., & Marmot, M. G. (2012). Interaction between education and household wealth on the risk of obesity in women in Egypt. *PLoS One*, *7*(6), e39507.
- Akik, C., Ghattas, H., Filteau, S., & Knai, C. (2017). Barriers to breastfeeding in Lebanon: A policy analysis. *Journal of public health policy*, *38*(3), 314-326.
- Atalah, E., Amigo, H., & Bustos, P. (2014). Does Chile's nutritional situation constitute a double burden?–. *The American journal of clinical nutrition*, *100*(6), 1623S-1627S.
- Barker, D. J. (2006). Adult consequences of fetal growth restriction. *Clinical obstetrics and gynecology*, *49*(2), 270-283.
- Bassett, M., Romaguera, D., Giménez, M., Lobo, M., & Samman, N. (2014). Prevalencia y determinantes de la doble carga de malnutrición en hogares en La Puna y Quebrada de Humahuaca, Jujuy, Argentina. *Nutricion hospitalaria*, *29*(2), 322-330.

- Bates, K., Gjonca, A., & Leone, T. (2017). Double burden or double counting of child malnutrition? The methodological and theoretical implications of stunting/overweight in low and middle income countries. *J Epidemiol Community Health*, *71*(8), 779-785. doi:10.1136/jech-2017-209008
- Black, R. E., Victora, C. G., Walker, S. P., Bhutta, Z. A., Christian, P., De Onis, M., . . . Martorell, R. (2013). Maternal and child undernutrition and overweight in low-income and middle-income countries. *The Lancet*, *382*(9890), 427-451.
- Colchero, M. A., Rivera-Dommarco, J., Popkin, B. M., & Ng, S. W. (2017). In Mexico, evidence of sustained consumer response two years after implementing a sugar-sweetened beverage tax. *Health Affairs*, *36*(3), 564-571.
- Conde, W. L., & Monteiro, C. A. (2014). Nutrition transition and double burden of undernutrition and excess of weight in Brazil—. *The American journal of clinical nutrition*, *100*(6), 1617S-1622S.
- Dieffenbach, S., & Stein, A. D. (2012). Stunted child/overweight mother pairs represent a statistical artifact, not a distinct entity. *J Nutr*, *142*(4), 771-773. doi:10.3945/jn.111.153387
- Eckhardt, C. L., Torheim, L., Monterrubio, E., Barquera, S., & Ruel, M. T. (2008). The overlap of overweight and anaemia among women in three countries undergoing the nutrition transition. *Eur J Clin Nutr*, *62*(2), 238.
- El Bcheraoui, C., Jumaan, A. O., Collison, M. L., Daoud, F., & Mokdad, A. H. (2018). Health in Yemen: losing ground in war time. *Global Health*, *14*(1), 42. doi:10.1186/s12992-018-0354-9
- El Kishawi, R. R., Soo, K. L., Abed, Y. A., & Muda, W. A. M. W. (2016). Prevalence and Associated Factors for Dual Form of Malnutrition in Mother-Child Pairs at the Same Household in the Gaza Strip-Palestine. *PLoS One*, *11*(3), e0151494.
- El Taguri, A., Besmar, F., Abdel Monem, A., Betimal, I., Ricour, C., & Rolland-Cachera, M. F. (2009). Stunting is a major risk factor for overweight: results from national surveys in 5 Arab countries. *East Mediterr Health J*, *15*(3), 549-562.
- Eshaq, A. M., Fothan, A. M., Jensen, E. C., Khan, T. A., & AlAmodi, A. A. (2017). Malnutrition in Yemen: an invisible crisis. *The Lancet*, *389*(10064), 31-32. doi:10.1016/S0140-6736(16)32592-2
- Farrag, N. S., Cheskin, L. J., & Farag, M. K. (2017). A systematic review of childhood obesity in the Middle East and North Africa (MENA) region: Health impact and management. *Advances in pediatric research*, *4*, 6. doi:10.12715/apr.2017.4.6
- Fernald, L. C., & Neufeld, L. M. (2007). Overweight with concurrent stunting in very young children from rural Mexico: prevalence and associated factors. *Eur J Clin Nutr*, *61*(5), 623-632. doi:10.1038/sj.ejcn.1602558
- Filmer, D., & Pritchett, L. H. (2001). Estimating wealth effects without expenditure data—or tears: an application to educational enrollments in states of India. *Demography*, *38*(1), 115-132.
- Fraser, B. (2013). Latin American countries crack down on junk food. *The Lancet*, *382*(9890), 385-386. doi:10.1016/S0140-6736(13)61657-8
- Freire, W. B., Silva-Jaramillo, K. M., Ramírez-Luzuriaga, M. J., Belmont, P., & Waters, W. F. (2014). The double burden of undernutrition and excess body weight in Ecuador—. *The American journal of clinical nutrition*, *100*(6), 1636S-1643S.
- Garrett, J. L., & Ruel, M. T. (2003). Stunted child-overweight mother pairs: An emerging policy concern. *Washington, DC: International Food Policy Research Institute (IFPRI)*, 13-18.

- Gartner, A., El Ati, J., Traissac, P., Bour, A., Berger, J., Landais, E., . . . Delpuech, F. (2014). A double burden of overall or central adiposity and anemia or iron deficiency is prevalent but with little socioeconomic patterning among Moroccan and Tunisian urban women. *J Nutr*, *144*(1), 87-97. doi:10.3945/jn.113.178285
- Gubert, M. B., Spaniol, A. M., Segall-Corrêa, A. M., & Pérez-Escamilla, R. (2017). Understanding the double burden of malnutrition in food insecure households in Brazil. *Maternal & Child Nutrition*, *13*(3), e12347. doi:doi:10.1111/mcn.12347
- Hodge, J., Verstraeten, R., & Ochoa-Avilés, A. (2016). Malnutrition's new frontier: The challenge of obesity. *Nourishing Millions*, 81-88.
- Hoey, L., & Pelletier, D. L. (2011). Bolivia's multisectoral Zero Malnutrition Program: insights on commitment, collaboration, and capacities. *Food and Nutrition Bulletin*, *32*(2_suppl2), S70-S81.
- Hong, R. (2007). Effect of economic inequality on chronic childhood undernutrition in Ghana. *Public Health Nutr*, *10*(4), 371-378.
- Hong, R., & Mishra, V. (2006). Effect of wealth inequality on chronic under-nutrition in Cambodian children. *Journal of Health, Population and Nutrition*, 89-99.
- Huicho, L., Segura, E. R., Huayanay-Espinoza, C. A., de Guzman, J. N., Restrepo-Méndez, M. C., Tam, Y., . . . Group, P. C. C. S. W. (2016). Child health and nutrition in Peru within an antipoverty political agenda: a Countdown to 2015 country case study. *The lancet global health*, *4*(6), e414-e426.
- Jones, A. D., Acharya, Y., & Galway, L. P. (2016). Urbanicity gradients are associated with the household- and individual-level double burden of malnutrition in Sub-Saharan Africa. *J Nutr*, jn226654.
- Keino, S., Plasqui, G., Etyang, G., & van den Borne, B. (2014). Determinants of stunting and overweight among young children and adolescents in sub-Saharan Africa. *Food and Nutrition Bulletin*, *35*(2), 167-178.
- Kitsiou-Tzeli, S., & Tzetzis, M. (2017). Maternal epigenetics and fetal and neonatal growth. *Curr Opin Endocrinol Diabetes Obes*, *24*(1), 43-46. doi:10.1097/med.0000000000000305
- Kosaka, S., & Umezaki, M. (2017). A systematic review of the prevalence and predictors of the double burden of malnutrition within households. *British Journal of Nutrition*, 1-10.
- Kroker-Lobos, M. F., Pedroza-Tobías, A., Pedraza, L. S., & Rivera, J. A. (2014). The double burden of undernutrition and excess body weight in Mexico—. *The American journal of clinical nutrition*, *100*(6), 1652S-1658S.
- Makoka, D. (2013). The impact of maternal education on child nutrition: evidence from Malawi, Tanzania, and Zimbabwe. *Calverton, Maryland, USA: ICF International*.
- Makoka, D., & Masibo, P. K. (2015). Is there a threshold level of maternal education sufficient to reduce child undernutrition? Evidence from Malawi, Tanzania and Zimbabwe. *BMC Pediatr*, *15*, 96. doi:10.1186/s12887-015-0406-8
- Mamabolo, R. L., Alberts, M., Steyn, N. P., Delemarre-van de Waal, H. A., & Levitt, N. S. (2005). Prevalence and determinants of stunting and overweight in 3-year-old black South African children residing in the Central Region of Limpopo Province, South Africa. *Public Health Nutr*, *8*(5), 501-508.
- Martorell, R., Khan, L. K., Hughes, M. L., & Grummer-Strawn, L. M. (2000). Obesity in women from developing countries. *Eur J Clin Nutr*, *54*(3), 247.

- Meiqari, L., Hoetjes, M., Baxter, L., & Lenglet, A. (2018). Impact of war on child health in northern Syria: the experience of Medecins Sans Frontieres. *Eur J Pediatr*, *177*(3), 371-380. doi:10.1007/s00431-017-3057-y
- Negash, C., Whiting, S. J., Henry, C. J., Belachew, T., & Hailemariam, T. G. (2015). Association between maternal and child nutritional status in Hula, rural Southern Ethiopia: a cross sectional study. *PLoS One*, *10*(11), e0142301.
- Ng, M., Fleming, T., Robinson, M., Thomson, B., Graetz, N., Margono, C., . . . Gakidou, E. (2014). Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*, *384*(9945), 766-781. doi:10.1016/s0140-6736(14)60460-8
- Obermeyer, C. M., Bott, S., & Sassine, A. J. (2015). Arab adolescents: health, gender, and social context. *Journal of Adolescent Health*, *57*(3), 252-262.
- Pinhas-Hamiel, O., Newfield, R. S., Koren, I., Agmon, A., Lilos, P., & Phillip, M. (2003). Greater prevalence of iron deficiency in overweight and obese children and adolescents. *Int J Obes Relat Metab Disord*, *27*(3), 416-418. doi:10.1038/sj.ijo.0802224
- Popkin, B. M. (2004). The nutrition transition: an overview of world patterns of change. *Nutr Rev*, *62*(7 Pt 2), S140-143.
- Popkin, B. M., Richards, M. K., & Montiero, C. A. (1996). Stunting is associated with overweight in children of four nations that are undergoing the nutrition transition. *J Nutr*, *126*(12), 3009.
- Ramirez-Zea, M., Kroker-Lobos, M. F., Close-Fernandez, R., & Kanter, R. (2014). The double burden of malnutrition in indigenous and nonindigenous Guatemalan populations-. *The American journal of clinical nutrition*, *100*(6), 1644S-1651S.
- Rivera, J. A., Pedraza, L. S., Martorell, R., & Gil, Á. (2014). Introduction to the double burden of undernutrition and excess weight in Latin America. *The American journal of clinical nutrition*, *100*(6), 1613S-1616S.
- Rutstein, S. O., & Johnson, K. (2004). *The DHS wealth index: ORC Macro, MEASURE DHS*.
- Saibul, N., Shariff, Z. M., Lin, K. G., Kandiah, M., Ghani, N. A., & Rahman, H. A. (2009). Food variety score is associated with dual burden of malnutrition in Orang Asli (Malaysian indigenous peoples) households: implications for health promotion. *Asia Pacific journal of clinical nutrition*, *18*(3), 412-422.
- Said-Mohamed, R., Bernard, J. Y., Ndzana, A. C., & Pasquet, P. (2012). Is overweight in stunted preschool children in Cameroon related to reductions in fat oxidation, resting energy expenditure and physical activity? *PLoS One*, *7*(6), e39007. doi:10.1371/journal.pone.0039007
- Sarmiento, O. L., Parra, D. C., González, S. A., González-Casanova, I., Forero, A. Y., & Garcia, J. (2014). The dual burden of malnutrition in Colombia-. *The American journal of clinical nutrition*, *100*(6), 1628S-1635S.
- Semba, R. D., de Pee, S., Sun, K., Sari, M., Akhter, N., & Bloem, M. W. (2008). Effect of parental formal education on risk of child stunting in Indonesia and Bangladesh: a cross-sectional study. *The Lancet*, *371*(9609), 322-328.
- Severi, C., & Moratorio, X. (2014). Double burden of undernutrition and obesity in Uruguay-. *The American journal of clinical nutrition*, *100*(6), 1659S-1662S.

- Sharara, E., Akik, C., Ghattas, H., & Obermeyer, C. M. (2018). Physical inactivity, gender and culture in Arab countries: a systematic assessment of the literature. *BMC public health*, *18*(1), 639.
- Subramanian, S. V., Perkins, J. M., Ozaltin, E., & Davey Smith, G. (2011). Weight of nations: a socioeconomic analysis of women in low- to middle-income countries. *Am J Clin Nutr*, *93*(2), 413-421. doi:10.3945/ajcn.110.004820
- Tzioumis, E., Kay, M. C., Bentley, M. E., & Adair, L. S. (2016). Prevalence and trends in the childhood dual burden of malnutrition in low-and middle-income countries, 1990–2012. *Public Health Nutr*, *19*(8), 1375-1388.
- Varela-Silva, M. I., Dickinson, F., Wilson, H., Azcorra, H., Griffiths, P. L., & Bogin, B. (2012). The nutritional dual-burden in developing countries--how is it assessed and what are the health implications? *Coll Antropol*, *36*(1), 39-45.
- Vollmer, S., Harttgen, K., Subramanyam, M. A., Finlay, J., Klasen, S., & Subramanian, S. (2014). Association between economic growth and early childhood undernutrition: evidence from 121 Demographic and Health Surveys from 36 low-income and middle-income countries. *The lancet global health*, *2*(4), e225-e234.
- Wachs, T. D., Creed-Kanashiro, H., Cueto, S., & Jacoby, E. (2005). Maternal education and intelligence predict offspring diet and nutritional status. *J Nutr*, *135*(9), 2179-2186. doi:10.1093/jn/135.9.2179
- Wang, X., Hojer, B., Guo, S., Luo, S., Zhou, W., & Wang, Y. (2009). Stunting and 'overweight' in the WHO Child Growth Standards - malnutrition among children in a poor area of China. *Public Health Nutr*, *12*(11), 1991-1998. doi:10.1017/s1368980009990796
- WHO. (2015). Retrieved from <http://www.who.int/childgrowth/standards/en/>
- Wojcicki, J. M. (2014). The double burden household in sub-Saharan Africa: maternal overweight and obesity and childhood undernutrition from the year 2000: results from World Health Organization Data (WHO) and Demographic Health Surveys (DHS). *BMC public health*, *14*(1), 1124.

Legends to figures and tables

Figure 1. Odds of child stunting, overweight and the stunted-overweight double burden, by wealth quintiles in the MENA and LAC regions (OR; 95%CI).

Multivariable regression models adjusted for child sex and age, maternal education, urbanicity, hygiene, country fixed effects and within country clustering effect.

Figure 2. Odds of child stunting, overweight and the stunted-overweight double burden, by maternal education in the MENA and LAC regions (OR; 95%CI).


Multivariable regression models adjusted for child sex and age, wealth quintiles, urbanicity, hygiene, country fixed effects and within country clustering effect.

Figure 3. Odds of stunting, overweight and the stunted-overweight double burden, by wealth quintiles in children with mothers with varying educational status (none, primary, and secondary or more school attainment) (OR; 95%CI).


Multivariable regression models adjusted for child sex and age, urbanicity, hygiene, country fixed effects and within country clustering effect.

Table 1. Datasets included from the Middle East and North Africa (MENA) and Latin American and Caribbean (LAC) regions


Table 2. Prevalence of stunting, overweight, stunted-overweight 6-59-month-old children in the MENA and LAC regions.



MCN_12923_F1.jpg



MCN_12923_F2.tif



MCN_12923_F3.tif