

Should we let fever run its course in the early stages of COVID-19?

Alexandre A. Steiner, PharmD, PhD

Associate Professor, Departamento de Imunologia, Instituto de Ciencias Biomedicas, Universidade de Sao Paulo, Sao Paulo, SP, Brazil

Address for correspondence: Av. Prof. Lineu Prestes 1730, Ed. Biomedicas IV, Sao Paulo, SP 05508-000, Brazil. Tel.: +55-11-3091-7378; e-mail: asteiner@usp.br

No competing interests: The author has no conflict of interest, financial or otherwise, to declare.

Keywords: coronavirus, antipyretic, acetaminophen, ibuprofen, aspirin, immunity.

Abstract

Antipyretic drugs are routinely taken by most people at the first sign of a fever, but the symptomatic relief achieved with these drugs comes with the price of putting out a host defense response known to aid the immune system in the clearance of pathogens. This may be a steep price to pay when lives are at stake amid the turmoil of the COVID-19 pandemic. Here, experimental and clinical data are discussed in light of the notion that fever is particularly adaptive in the early, less severe stages of an infection, when the benefits of a raised body temperature to immunity far exceed its costs to the host. It is, therefore, reasonable to advise that allowing fever to run its natural course for a few days after the onset of symptoms in COVID-19 may reduce the risk of long-lasting infection and, consequently, the risk of complications.

This article is a preprint and has not been peer reviewed. It reports new medical research or thought that has yet to be evaluated and so should not be used to guide clinical practice.

Copyright © 2020 by Alexandre A. Steiner. Posted on Annals of Family Medicine COVID-19 Collection courtesy of Alexandre A. Steiner.

Antipyretic drugs are readily available as over-the-counter medication, being taken routinely by most people at the first sign of a fever. But the symptomatic relief achieved with these drugs comes with the price of putting out a host defense response that has evolved for millions of years to aid the immune system in the clearance of pathogens^{1,2}. While it may be ok to pay this price in unthreatening infections such as a regular cold or a mild flu, the situation may be different when lives are at stake amid the uncertainty of the COVID-19 pandemic.

Animal studies provide substantial support for a role of fever in survival, with it being particularly adaptive in the early, less severe stages of an infection, when the benefits of a raised body temperature to immunity far exceed its costs to the host³. Albeit less compelling, clinical evidence is also consistent with this notion. In a prospective trial in trauma patients, an aggressive antipyretic protocol strongly tended to increase the risk of acquiring an infection and developing complications, which prompted interruption of the trial after the first interim analysis of the data⁴. Furthermore, two randomized, double-blind, placebo-controlled trials in rhinovirus-infected volunteers have shown that the use of aspirin, acetaminophen or ibuprofen is associated with increased or prolonged viral shedding^{5,6}. In one of those trials⁵, aspirin and acetaminophen were further shown to suppress the neutralizing antibody response and, paradoxically, worsen nasal symptoms. These drugs are not identical with regard to their effects on local inflammation, but have the suppression of fever as a common denominator. And although fever was thought to be occasional in the rhinovirus-infected volunteers, infrequent measures of body temperature (no more than four times a day) are likely to have underestimated the prevalence of low-grade fever in those

This article is a preprint and has not been peer reviewed. It reports new medical research or thought that has yet to be evaluated and so should not be used to guide clinical practice.

Copyright © 2020 by Alexandre A. Steiner. Posted on Annals of Family Medicine COVID-19 Collection courtesy of Alexandre A. Steiner.

trials. Indeed, many of the placebo-treated volunteers reported chilliness⁶, a behavior known to be linked with development of fever.

Therefore, it is plausible to advise that suppression of fever early in the course of COVID-19 may increase the risk of long-lasting infection and, consequently, the risk of complications. It is also plausible to expect that antipyresis can weaken the development of antibody-dependent herd immunity, which, together with an enhancement in viral shedding, can favor the spread of the disease in the population. These arguments call further attention the recent warning to avoid ibuprofen in COVID-19. The warning was based on the effects exerted by the drug at the site of infection, be that the suppression of inflammation⁷ or the facilitation of ACE2-dependent targeting of host cells by the SARS-CoV-2 virus⁸. However, it cannot be ignored that the impacts of ibuprofen on COVID-19 may also be related to antipyresis, in which case recommending the use of alternate antipyretic drugs such as acetaminophen and aspirin may not suit the patients' needs in terms of immunity.

The positive association between age and the risk of complications in COVID-19⁹ is another aspect that can be looked upon from the perspective of fever as a host-defense response. Humans and experimental animals are known to mount less pronounced fevers as they age, with fever being completely absent in 20%-30% of elderly patients harboring a serious infections¹⁰. While the mechanisms underlying the suppression of fever in the elderly are still poorly understood, an experimental study provides evidence that this phenomenon may be restricted to the autonomic mechanisms of fever, with the behavioral mechanisms being spared to the extent that old rats regain the ability to develop fever if given the chance to move to a warmer environment¹¹. If the same thermoeffector pattern occurs in humans, it

This article is a preprint and has not been peer reviewed. It reports new medical research or thought that has yet to be evaluated and so should not be used to guide clinical practice.

Copyright © 2020 by Alexandre A. Steiner. Posted on Annals of Family Medicine COVID-19 Collection courtesy of Alexandre A. Steiner.

may open a therapeutic opportunity. More specifically, if an elderly person with respiratory symptoms complains of chilliness, perhaps it would be more adequate to provide him or her with warmth than to seek symptom relief with drugs.

The question then arises as to why so many people are afraid of fever. It has been documented that fever phobia is based on unrealistic concerns and misconceptions that have persisted in our culture despite decades of research on the true nature of fever¹². The perception that body temperature will reach dangerous levels if fever is not treated is unsubstantiated by scientific evidence. On the contrary, the available evidence indicates that body temperature is precisely regulated during fever, and that endogenous antipyretic mechanisms are at work to prevent excessive rises in body temperature. Accordingly, the body temperature of infected patients usually stays around 38.0-39.5°C, rarely reaching 40-41°C and never exceeding this ceiling. Within this physiological range, the temperature rise per se has never been shown to be harmful to neurons or other cellular phenotypes. In young children, febrile seizure is a concern, but it must be considered that only a very small fraction of the children are predisposed to this condition, that febrile seizures are usually self-limiting and benign, and, what is more, that the preemptive use of antipyretic drugs may be ineffective at preventing the seizures¹³.

Recognizing the value of fever in the early stages of an infection does not imply that fever has no costs, but the available evidence indicates that the costs of fever outweigh its benefits only at more advanced and severe stages of infection³, typically when patients are hospitalized because of cardiovascular or respiratory complications^{14,15}. In such states of compromised physiological fitness (sepsis), the energetic cost of fever may be too much for

This article is a preprint and has not been peer reviewed. It reports new medical research or thought that has yet to be evaluated and so should not be used to guide clinical practice.

Copyright © 2020 by Alexandre A. Steiner. Posted on Annals of Family Medicine COVID-19 Collection courtesy of Alexandre A. Steiner.

the host to bear, and, additionally, fever may heighten the maladaptive inflammation underlying the complications. Interestingly, it often happens that fever is replaced with a self-limiting, spontaneous form of hypothermia in severe sepsis¹⁶, which, at least in experimental animals, appears to be launched by the host as an alternate defense strategy aimed at tolerating the pathogen while preserving vital bodily functions³. This dichotomy in the thermal adaptation of critically ill patients may underlie the controversial results obtained in some trials on the use of antipyresis in this subset of patients^{4,14,17}. More refined strategies of patient stratification, perhaps involving temperature trajectories¹⁸, may help to solve this matter. But regardless of how complex the situation may be with critically ill patients, the situation is unquestionably less complex when an infection such as COVID-19 is at its inception, and competing demands are not at play to offset the value of fever to immunity.

Taken together, the lines of evidence discussed herein call attention to the value of fever as a host defense strategy and its possible implications to COVID-19. Prioritizing symptom relief over host immunity in the early stages of a potentially life-threatening infection may not be working in our favor in the COVID-19 pandemic. In this moment of uncertainty and crisis, it may be proper to break cultural habits and advise against the use of antipyretic drugs for at least a few days after the onset of symptoms. This recommendation is based on lessons learned from studies on the pathophysiology of other infectious diseases, and does not eliminate the need for placebo-controlled studies specifically tailored to COVID-19. Such studies are likely to be quite challenging for requiring patient enrollment at the earliest sign of infection, as well as adherence to fever management protocols in an outpatient setting, but those challenges certainly do not undermine their importance.

This article is a preprint and has not been peer reviewed. It reports new medical research or thought that has yet to be evaluated and so should not be used to guide clinical practice.

Copyright © 2020 by Alexandre A. Steiner. Posted on Annals of Family Medicine COVID-19 Collection courtesy of Alexandre A. Steiner.

Acknowledgments

The author's research reviewed herein was supported by Fundacao de Amparo a Pesquisa do Estado de Sao Paulo (FAPESP; grant Nos. 12/03831-8, 16/04921-1 & 18/03418-0).

References

1. Evans SS, Repasky EA, Fisher DT. Fever and the thermal regulation of immunity: the immune system feels the heat. *Nat Rev Immunol*. 2015;15(6):335-349.
2. Hasday JD, Thompson C, Singh IS. Fever, immunity, and molecular adaptations. *Compr Physiol*. 2014;4(1):109-148.
3. Steiner AA, Romanovsky AA. Energy trade-offs in host defense: immunology meets physiology. *Trends Endocrinol Metab*. 2019;30(12):875-878.
4. Schulman CI, Namias N, Doherty J, et al. The effect of antipyretic therapy upon outcomes in critically ill patients: a randomized, prospective study. *Surg Infect*. 2005;6(4):369-375.
5. Graham NM, Burrell CJ, Douglas RM, Debelle P, Davies L. Adverse effects of aspirin, acetaminophen, and ibuprofen on immune function, viral shedding, and clinical status in rhinovirus-infected volunteers. *J Infect Dis*. 1990;162(6):1277-1282.
6. Stanley ED, Jackson GG, Panusarn C, Rubenis M, Dirda V. Increased virus shedding with aspirin treatment of rhinovirus infection. *JAMA*. 1975;231(12):1248-1251.
7. Day M. Covid-19: ibuprofen should not be used for managing symptoms, say doctors and scientists. *BMJ*. 2020;368:m1086.
8. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *Lancet Respir Med*. 2020.
9. Verity R, Okell LC, Dorigatti I, et al. Estimates of the severity of coronavirus disease 2019: a model-based analysis. *The Lancet Infectious Diseases*. 2020;10.1016/s1473-3099(20)30243-7.
10. Norman DC. Fever in the elderly. *Clin Infect Dis*. 2000;31(1):148-151.
11. Florez-Duquet M, Peloso E, Satinoff E. Fever and behavioral thermoregulation in young and old rats. *Am J Physiol Regul Integr Comp Physiol*. 2001;280(5):R1457-1461.
12. Crocetti M, Moghbeli N, Serwint J. Fever phobia revisited: have parental misconceptions about fever changed in 20 years? *Pediatrics*. 2001;107(6):1241-1246.
13. Rosenbloom E, Finkelstein Y, Adams-Webber T, Kozer E. Do antipyretics prevent the recurrence of febrile seizures in children? A systematic review of randomized controlled trials and meta-analysis. *Eur J Paediatr Neurol*. 2013;17(6):585-588.
14. Schortgen F, Clabault K, Katsahian S, et al. Fever control using external cooling in septic shock: a randomized controlled trial. *Am J Respir Crit Care Med*. 2012;185(10):1088-1095.
15. Netzer G, Dowdy DW, Harrington T, et al. Fever is associated with delayed ventilator liberation in acute lung injury. *Ann Am Thorac Soc*. 2013;10(6):608-615.
16. Fonseca MT, Rodrigues AC, Cezar LC, Fujita A, Soriano FG, Steiner AA. Spontaneous hypothermia in human sepsis is a transient, self-limiting, and nonterminal response. *J Appl Physiol (1985)*. 2016;120(12):1394-1401.

This article is a preprint and has not been peer reviewed. It reports new medical research or thought that has yet to be evaluated and so should not be used to guide clinical practice.

Copyright © 2020 by Alexandre A. Steiner. Posted on Annals of Family Medicine COVID-19 Collection courtesy of Alexandre A. Steiner.

17. Young P, Saxena M, Bellomo R, et al. Acetaminophen for fever in critically ill patients with suspected infection. *N Engl J Med*. 2015;373(23):2215-2224.
18. Bhavani SV, Carey KA, Gilbert ER, Afshar M, Verhoef PA, Churpek MM. Identifying novel sepsis subphenotypes using temperature trajectories. *Am J Respir Crit Care Med*. 2019;200(3):327-335.

This article is a preprint and has not been peer reviewed. It reports new medical research or thought that has yet to be evaluated and so should not be used to guide clinical practice.

Copyright © 2020 by Alexandre A. Steiner. Posted on Annals of Family Medicine COVID-19 Collection courtesy of Alexandre A. Steiner.