

Guided tissue regeneration combined with bone allograft in infrabony defects: Clinical outcomes and assessment of prognostic factors

Jad Majzoub*, BDS, Shayan Barootchi*, DMD, Lorenzo Tavelli*, DDS, Chin-Wei Wang*, DDS, DMSc, Hsun-Liang Chan*, DDS, MS, and Hom-Lay Wang*, DDS, MSD, PhD

* Department of Periodontics and Oral Medicine, University of Michigan School of Dentistry, Ann Arbor, MI, USA

Corresponding author:

Hom-Lay Wang, DDS, MSD, PhD

Department of Periodontics and Oral Medicine, University of Michigan School of Dentistry

1011 North University Avenue

Ann Arbor, Michigan 48109-1078, USA.

TEL: (734) 763-3325; FAX: (734) 936-0374

E-mail address: homlay@umich.edu

Conflict of interest

The authors do not have any financial interests, either directly or indirectly, in the products or information listed in the paper.

Word count: 3485

Tables and figures: 2 figures, 4 tables

Running title: Clinical outcomes of guided tissue regeneration

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/JPER.19-0336](#).

This article is protected by copyright. All rights reserved.

One sentence summary: This study showed using guided tissue regeneration with bone graft to treat periodontal infrabony defects improves tooth retention and clinical outcomes.

Keywords: Guided Tissue Regeneration, survival analysis, periodontitis, smoking

Author contribution:

Majzoub Jad, contributed to the conception and design of the work, drafted the work and revised it critically for important intellectual content, agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Barootchi Shayan, contributed to the analysis of data for the work, drafted the work and revised it critically for important intellectual content, agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Tavelli Lorenzo, contributed to the conception and design of the work, drafted the work and revised it critically for important intellectual content, agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Hsun-Liang Chan, contributed to the interpretation of data for the work revised the work critically for important intellectual content, agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Wang Hom-Lay, contributed to the conception and design of the work; the acquisition, analysis, and interpretation of data for the work, critical revision of the work for important intellectual content, final approval of the version to be published, agrees to be accountable for all aspects of the work in

ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Abstract

Background and Aim: Clinical data on the outcomes of guided tissue regeneration (GTR) is scarce. The aim of this retrospective cohort study was to evaluate the outcomes after GTR, their stability and the survival of the treated periodontal infrabony defects.

Materials and methods: Infrabony defects treated with GTR using a bioabsorbable membrane and a bone graft substitute with at least 1-year follow-up were included. Survival and regression analyses were conducted to evaluate the outcomes, their stability, and the retention of the teeth. The effect of recorded variables on clinical attachment gain (CAL) and tooth survival were assessed via Cox Proportional-Hazards Models and Multivariate Generalized Linear Models.

Results: 175 treated defects were selected from a total of 641 charts. The average follow-up was 5.75 ± 4.6 years. At baseline, the mean CAL was 9.56 ± 1.93 mm with a mean pocket depth (PD) of 8.41 ± 1.42 mm. At the 1-year post-surgical recall, 3.55 ± 1.85 mm of CAL gain and 3.87 ± 1.87 mm PD reduction were observed ($p < 0.05$). The 5- and 10-year survival rates of the treated teeth were 85.0% and 72.7%, respectively. Baseline PD, smoking, and membrane exposure were significantly related to CAL gain, whereas baseline CAL, age, frequency in maintenance visits significantly affected tooth survival.

Conclusion: Within the limitations of this study, data suggests GTR is a good option for the treatment of infrabony defects since it can improve both tooth retention rate and overall clinical outcomes.

1.Introduction

Periodontitis is a complex multifactorial disease that often leads to the formation of deep infrabony defects¹. Their presence has been shown to increase the risk for the progression of periodontitis by more than 10 times². The treatment of infrabony defects poses clinical challenges, making its presence one of the influential factors when determining the complexity of periodontal disease³. In order to address this concern, several approaches have been proposed for their treatment, including scaling and root planing, open flap debridement, resective surgeries and various periodontal regenerative therapies⁴.

Among the proposed regeneration procedures, evidence supports the use of guided tissue regeneration (GTR) that employed barrier membranes, and many studies have reported significant clinical attachment level (CAL) gain, pocket depth (PD) reduction and bone fill with the usage of membranes⁵⁻⁹. Additionally, histologic evidence has confirmed the regeneration of new cementum, periodontal ligament and alveolar bone^{10,11}. Moreover, the superiority of GTR outcomes over traditional flap surgeries in the treatment of infrabony defects has been well established^{6,9,12}. Cortellini et al. in a 20-year follow-up of a randomized clinical trial, observed that treated sites with GTR showed better long-term stability of their clinical outcomes compared to sites that were treated with a modified Widman flap procedure. Interestingly, the authors found the differences between regenerative therapy and flap surgery more pronounced in the second decade of the observation¹³.

Nevertheless, patient related factors, defect morphology and surgical techniques have all been reported to significantly impact the overall predictability of the GTR procedure^{8,14}. In addition, most of the studies only assessed the short-term outcomes of GTR and have been performed on a very carefully selected pool of patients^{13,15,16}.

Therefore, the aim of the present study was to investigate the clinical outcomes of GTR and assess potential prognostic factors that can affect the clinical outcomes and tooth survival.

2.Materials and methods

2.1. Study design

The current investigation was designed according to the principles presented in the Helsinki Declaration of 1975, as revised in 2000 for biomedical research involving human patients. The study was approved by the Institutional Review Board for Human Studies, School of Dentistry, University of

Michigan, Ann Arbor, MI, (HUM00154408) to be conducted at the Department of Periodontology within the same institution.

This retrospective cohort study selected all patients that had undergone treatment for infrabony defects with GTR at the School of Dentistry, University of Michigan, Ann Arbor, MI. All paper files and digital charts of patients treated with GTR were carefully scanned and analyzed by two independent and pre-calibrated investigators (JM, SB). At every stage, after examining the gathered data, in case of a disagreement, discussion was held by the two reviewers. If resolution was not possible, a third investigator (LT) was consulted to reach a consensus. This study was conducted by obtaining anonymized data and all personally identifiable patient information was removed; hence there was no need for informed consent.

2.2. Inclusion criteria

Patients that met the following inclusion criteria were included in this study:

- (1) A patient who had previously received GTR therapy for at least one infrabony defect with a probing depth (PD) of ≥ 6 mm¹⁷.
- (2) Prior to the GTR procedure, all individuals must have previously received a comprehensive periodontal treatment (including oral hygiene instructions, scaling/root planing, prophylaxis, etc.).
- (3) GTR procedures must have included utilization of a bioabsorbable membrane in combination with a bone graft substitute.
- (4) Patient records must have had at least 1-year follow-up after GTR treatment.
- (5) Patients charts should have contained complete clinical data including radiographs at baseline (pre-surgical stage) and at least 1-year after the surgical procedure.

2.3. Exclusion criteria

Patients that had the following conditions were excluded from this study:

- (1) Patients without a post-surgical follow-up data reaching 12 months.

(2) The use of barrier membranes for procedures other than GTR (sinus lift, guided bone regeneration, socket augmentation, etc.).

(3) GTR procedures in furcation defects.

(4) The sole use of bone graft without utilizing a barrier membrane, or the utilization of a membrane without placement of a bone graft.

(5) Placement of a non-resorbable membrane.

(6) A medically compromised patients or those taking medications that are known to interfere with the normal healing response process (e.g., bisphosphonates, anti-cancer therapy, etc.).

2.4. Data collection and Classification

The following information were obtained for all qualified individuals. **1)** patient related factors (such as age, gender, etc. ; **2)** medical history (including documentation of smoking, diabetes, other systemic or local diseases); **3)** location of the treated defect (mandible/maxilla); **4)** related clinical parameters such as: PD, CAL, gingival recession (REC); **5)** flap design (envelope/papilla preservation); **6)** occurrence of post-surgical complications (such as membrane exposure); **7)** follow-up time (until tooth extraction or last maintenance appointment); **8)** frequency of maintenance appointments throughout the entire follow-up; and **9)** patient radiographs.

2.5. Study outcomes: The study outcomes of the present project were three folds:

2.5.1. Clinical outcomes of GTR

- a. To assess the outcomes of the treatment, changes in the clinical parameters (PD, CAL, REC) were compared from baseline to the 1-year post-op/follow-up.
- b. Additionally, the influence of other recorded variables was assessed on the CAL results.
- c. Stability of the treatment results throughout the entire follow-up was assessed for all sites as previously performed in other studies^{18,19}. Briefly, stability was defined for a treated site that maintained the 1-year post-surgical CAL throughout time, presenting with less than 2 mm of change. This was assessed with the Kaplan-Meier estimator.

2.5.2. Survival: The survival of a treated tooth was assessed according to the Kaplan-Meier analysis. Additionally, the effect of the recorded variables on the treated sites was assessed for potential effect on tooth retention/survival.

2.5.3 Assessment of the radiographic defect angle Extracted baseline radiographs were used to measure the radiographic defect angle of the treated site^{20, 21}, by using a digital software†. The angle outlined by the bony defect wall and the root surface of the corresponding tooth was measured by using the criteria described by Bjorn et al. (1969)²² and Schei et al. (1959)²³. Next, the effect of the width of the angle on tooth survival and post-surgical CAL gain and tooth survival was evaluated. All radiographic analyses were performed by two individual and calibrated examiners (JM, SB). In case of uncertainty or a substantial difference among the measurements, a third reviewer (LT) was consulted for reassessing the radiographs.

2.5.4 Data management and Statistical analysis

The extracted data were entered into pre-fabricated spread sheets by the same investigators. All analyses were performed by an author with expertise in biostatistics (SB) using Rstudio ‡, the survminer²⁴, survival^{25, 26}, and ggplot2²⁷ packages. Descriptive statistics were used for presenting the retrieved data at baseline as means \pm standard deviations, and range. The treated defects served as the unit of analysis. The changes in clinical parameters from baseline to the 1-year outcome were assessed with dependent t-test. Mixed-effects Uni- and Multi-level Regression analyses were performed to identify prognostic factors for CAL gain. Kaplan-Meier survival probabilities were calculated and subsequently the curves were plotted. Multi-variate Cox Proportional Hazzard models were used for assessing correlations between independent variables and tooth loss, accounting for the fact that an individual may have attributed to multiple treated infrabony defects (shared frailty was accounted for by including random effects). Step-wise regression analyses were performed using likelihood ratio tests, with variables presenting a p value of <0.05 in the initial univariate analysis. Exponentiated regression coefficients (Hazzard ratios) and corresponding 95% confidence intervals (CI) were produced, and a p value threshold of 0.05 was set for the statistical significance.

3. Results

3.1. Study population

A total of 641 patient charts were retrieved and screened as a result of the initial search. Subsequently, 513 were excluded due to the following reasons: guided bone regeneration (121) or socket augmentation procedures (61), regenerative approaches performed with a non-resorbable membrane (94), with the use of membranes alone (6), or those treated solely with a bone graft material (16). Additionally, files with less than 1-year of follow-up (81), sinus augmentation procedures (28) and regeneration attempts in furcation defects (106) were also excluded, a flowchart diagram has been presented in supplementary file Data S1 in online *Journal of Periodontology* to visualize the screening process.

As a result, a total of 128 patients (63 males and 65 females) with a mean age of 51.7 ± 13.88 (16 - 85 years), with 175 GTR-treated infrabony defects were included. In all GTR procedures, the combination of a collagen membrane with an allograft bone material was used. The utilized bone grafts were all allogenic in nature either Freeze-dried bone allograft § (in 38 defects), or Solvent-dehydrated bone allograft || (in 137 defects). The utilized membranes were of two types, both collagenous in nature BioMend ¶ in 128 defects, and Bio-Gide # in 47 cases. The mean follow-up for the selected cases was 5.75 ± 4.6 years. The average maintenance visits for patients was $2.2 (\pm 1.1)$ times per year. Table 1 presents details on the characteristics of the included patients.

3.2. Clinical outcomes of GTR

At baseline, 60.8% of sites presented with BOP, a mean PD of 8.41 ± 1.42 mm, REC of 1.21 ± 1.11 mm, and CAL of 9.56 ± 1.93 mm. At the 1-year post-surgical recall, the BOP dropped to 23.8%, a PD reduction of 3.87 ± 1.87 mm, an increase in recession of 0.35 ± 1.16 mm, and 3.55 ± 1.85 mm of CAL gain were observed, all of which were statistically significant.

An access flap with intrasulcular incision^{13, 28} was used in most of the surgical cases; however, a papilla preservation design was performed in 20 of the procedures (without statistically significant differences between the two). Vertical incisions were used in 38 surgeries to allow better access to the defect and the occurrence of a membrane exposure was noted in 22 of the cases.

Table 2 depicts the results of the regression models, exploring the effect of different variables on the gained CAL at the 1-year post-op. Results from univariate analyses demonstrated that smoking (-1.01 (95% CI $[-1.85, -0.16]$, $p=0.01$)), post-op membrane exposure (-0.26 (95% CI $[-$

0.50, -0.01], $p=0.02$), and wider initial radiographic angle (-0.02 (95% CI [-0.05, -0.002], $p=0.03$)), were significantly related to a lower CAL, while an increase in initial PD (0.55 (95% CI [0.26, 0.85], $p<0.001$)), and CAL (0.34 (95% CI [0.14, 0.55], $p=0.001$)), were significant predictors for higher CAL gains. The negative effect of smoking on CAL gain was observed (CAL gain in smoker was 2.97 ± 2.06 mm vs. 3.98 ± 1.90 mm in non-smokers).

Results from the multivariate analysis including the stated statistically significant variables demonstrated that, smoking (-0.91 (95% CI [-1.73, -0.07], $p=0.03$)) and membrane exposure (-1.18 (95% CI [-2.28, -0.06], $p=0.03$)), were associated with lower gains in CAL, while initial PD (0.57 (95% CI [0.16, 0.97], $p=0.006$)), was positively correlated to higher CAL gains. Moreover, factors such as initial radiographic angle, and initial CAL did not prove to be statistical predictors from the model.

Lastly, gender, age, diabetes, placement of a vertical releasing incisions during the surgery, and endodontic treatment, did not seem to impact the CAL gains ($p>0.05$). Regarding the stability of the gained attachment levels, the Kaplan-meier analysis showed that $70.4 \pm 5.85\%$ and $54.9 \pm 7.26\%$ of the treated sites remained stable at 5 and 10 years, respectively (supplementary file Data S2 in online *Journal of Periodontology*).

3.3. Survival analysis

Throughout the follow-up period 30 teeth in 27 patients were lost. The 5- and 10-year survival rates were 85.0% and 72.7%, respectively (76 treated teeth at the 5-year timepoint, and 21 at the 10-year timepoint). Figure 1 demonstrates the survival curves of the treated teeth, and the life table presented in Table 3. descriptively presents the number of followed and extracted teeth at every timepoint.

Univariate analyses demonstrated that smoking significantly affected the survival of the treated teeth (amounting to 21 of the total extractions). Visual representation comparing the survival curves of smokers versus non-smokers is presented in Figure 2. Frequency in maintenance visits was also found to have a significant impact on tooth survival (0.45 (95% CI [0.29, 0.67], $p<0.0001$)). The average maintenance visits/year for the teeth that were extracted was 1.43 ± 0.90 while the mean number of visits/years for the ones that remained was 2.39 ± 1.07 visit/year ($p<0.01$).

Age, endodontic treatment and membrane exposure were also significant factors associated with tooth loss. Nevertheless, factors such as gender, diabetes, flap design, presence of vertical incisions did not seem to be statistically correlated in the survival analysis.

When the significant factors from the univariate models were evaluated in a multi-variate cox proportional hazard model, it was shown that maintenance (0.34 (95% CI [0.20, 0.58], $p < 0.001$)), initial CAL (1.53 (95% CI [1.03, 2.28], $p = 0.03$)), and age (1.08 (95% CI [1.03, 1.14], $p = 0.002$)), were the most significant predictors affecting tooth survival (Table 4).

4. Discussion

4.1. Clinical outcomes after GTR

Results from this study showed 3.55 ± 1.85 mm gain of CAL and 3.87 ± 1.87 mm PD reduction 1 year following the GTR procedure. The gain in CAL observed at the 1-year recall are in line with previous studies. In fact, Kher et al in a study evaluating the effect GTR using human allograft combined with a collagen membranes on infrabony defects found a mean CAL gain of 3.54 ± 0.36 mm one year following the surgery²⁹. Sculean et al, observed an average CAL gain of 4.07 ± 1.3 mm, 1 year after the GTR procedure using bovine bone xenograft in combination with a collagen membrane³⁰, and 4.1 ± 0.9 mm using a composite bovine bone xenograft combined with a collagen membrane³¹.

When other treatment approaches were attempted for the treatment of infrabony defects, Nibali and colleagues found that minimally invasive non-surgical therapy achieved 3.0 mm CAL gain 1 year after treatment. However, the study included patients with a lower initial PD compared to the present study (7 mm vs 9.6 mm)³². Whether deeper pockets such as the ones included in our study could be addressed with non-surgical therapies remains unknown.

Regarding the stability, it was observed that approximately 70.4 % and 54.9% of the treated sites remained stable after 5 and 10 years, respectively. In another retrospective study, with similar definition of stability, the authors reported a higher estimate of approximately 80% and 70% after 5 and 10 years, respectively¹⁸. It can be speculated that their superior results in terms of stability of the CALs might be attributed to a more meticulous patient selection, the private practice setting, and the expertise of the surgeon¹³. In another publication, evaluating the 10-year results of a randomized clinical trial, Nickles and colleagues showed that 3 out of the 18 defects that remained 10

years after the GTR procedure were unstable. However, it should be noted that the mentioned study had a relative limited sample size compared to the present study.

4.2. The effect of GTR on tooth survival

One of the goals of this study was to evaluate the survival of teeth treated with GTR.

We found that the 5- and 10-year survival rates of the treated teeth were 85.0% and 72.7%, respectively. Higher survival rates have been reported in the literature. In a 5-year study assessing the outcomes of GTR, only 2 out of 50 teeth were lost, none of which were attributed to the loss of the periodontal structure. This study included infrabony defects in a population which were 12% smokers and 86% attending at least one SPT visit per year³³. Another study presenting the 10-year results after treatment of 38 patients with EMD, GTR, a combination of EMD and GTR, and OFD, showed that none of the teeth treated with GTR or any other procedure were lost. Patients enrolled in this study were all non-smokers and attending a regular maintenance program consisting of four visits per year, including oral hygiene reinforcement and professional tooth cleaning³⁴. While, the population enrolled in the current study consisted of 32% smokers and with patients showing variability in compliance with maintenance appointments. Our results showed that both smoking and maintenance were significantly associated with tooth loss. In fact, from the 30 teeth that were lost throughout the entire follow-up, 21 had been in patients that were smokers. Furthermore, the extracted teeth belonged to individuals presenting with significantly less maintenance appointments/year (1.43 ± 0.90) than the ones that had retained their teeth (2.40 ± 1.07).

4.3. Predictors for regenerative outcomes

In the pursuit of achieving improvement in results, predictability of clinical outcomes and higher survival rates for teeth treated with GTR, the identification of factors related to the patient, defect and surgical technique that potentially affect the result is imperative. In the present study, it was observed that higher levels of PD at baseline showed better results in terms of CAL gain. This is in line with several previous studies^{17, 35}. Teeth that presented with higher CAL at baseline had an increased risk of failure. A smoking habit was another factor that we found having a significantly negative effect on CAL gain. We observed that on average smokers had 1 mm less CAL gain when

compared to non-smokers. Tonetti et al. were the first to assess the effect of smoking on periodontal regeneration following GTR in a retrospective study³⁶. They found that smokers obtained significantly less CAL than non-smokers 1-year following GTR surgery in deep infrabony defects (2.1 ± 1.2 mm versus 5.2 ± 1.9 mm)³⁶. Later on, Stavropoulos and colleagues when identifying factors influencing GTR treatment outcomes found that patients that were smokers gain 1 mm less in CAL than nonsmokers (3.2 ± 1.4 in smokers and 4.3 ± 1.3 in non-smokers), and that smokers had 7 times less chance of obtaining 4 mm CAL as compared to non-smokers³⁷. Finally, Nickles et al. when comparing clinical outcomes of teeth with infrabony defects 10 years after OFD and GTR with a bioabsorbable barrier found that current smoking negatively impacts CAL gain³⁸.

Another factor that affected the 1-year post-surgical outcomes was membrane exposure. This complication occurred in 22 cases and was associated with less favorable CAL gains. In line with these results, a systematic review and meta-analysis, reported that the mean CAL gain was statistically higher in the group that had not experienced membrane exposure compared to the ones which had¹⁴.

Periodontal maintenance was shown to be of utmost importance, as one of the very significant factors associated with tooth survival. Patients who had their teeth extracted presented on average with 1 less maintenance appointment per year than those who did not. In line with the literature, Weigel and colleagues in a 4-year study evaluating patients treated with GTR, showed that the number of recall visits highly affected the long-term outcomes³⁹. Similarly, Cortellini et al. when evaluating 175 infrabony defects treated with GTR highlighted the importance of maintenance appointments by showing that patients who did not attend regular maintenance visits were more prone to tooth loss¹⁸. Indeed, periodontal maintenance has been shown to highly impact the outcome of regenerative therapy^{19,40}.

We also found that that endodontic treatment did not affect CAL gain. This is in agreement with a study that evaluated GTR outcomes of non-vital, endodontically treated teeth⁴¹.

Finally, age was shown to be one of the most significant predictors affecting tooth survival. In line with these results, a study identifying prognostic factors relating to tooth survival including more than two hundred thousand subjects, found that the risk of losing ≥ 4 teeth was associated with an increase in age⁴². Other studies have also highlighted upon the significance of age as a predictor of tooth loss⁴³⁻⁴⁵.

4.4. Limitations

Among the limitations of our research are the retrospective nature of the study design and the lack of a control group. In addition, the infrabony component of the defect was not taken into consideration when evaluating the radiographic defect angle⁴⁶. Our study includes a large number of smoking patients. Nonetheless, this allowed us to focus on the impact of smoking on GTR outcomes. Finally, although the change in bleeding on probing provides a general idea about the oral hygiene of the patients^{47, 48}, full-mouth bleeding and local plaque scores were not available to evaluate the influence of oral hygiene status on the outcomes of GTR.

4.5. Conclusion

Within the limitations of the present study, the following conclusions can be obtained: GTR with the use of a collagen membrane and bone graft is a viable treatment for the management of teeth with an infrabony defects. The treatment outcomes following the GTR procedure are significantly influenced by factors such as frequency of maintenance visits, smoking, occurrence of a post-surgical membrane exposure, and initial defect characteristics (pocket depth and clinical attachment level).

Conflict of interest and source of funding

The authors do not have any financial interests neither directly nor indirectly in the companies whose materials were evaluated in this study. This manuscript was partially supported by the University of Michigan Periodontal Graduate Student Research Fund.

Footnotes

† Image J, U. S. National Institutes of Health, Bethesda, Maryland, USA

‡ Rstudio Version 1.1.383, RStudio, Inc., Boston, Massachusetts, USA

§ Salvin Dental Specialties, Charlotte, NC, USA

|| Puros, Zimmer Biomet, Carlsbad, CA, USA

¶ Zimmer Biomet, Carlsbad, CA, USA

Geistlich Pharma, Wolhusen, Switzerland

References

1. Waerhaug J. The infrabony pocket and its relationship to trauma from occlusion and subgingival plaque. *J Periodontol* 1979;50:355-365.
2. Rams TE, Listgarten MA, Slots J. Radiographic alveolar bone morphology and progressive periodontitis. *J Periodontol* 2018;89:424-430.
3. Papapanou PN, Sanz M, Buduneli N, et al. Periodontitis: Consensus report of workgroup 2 of the 2017 World Workshop on the Classification of Periodontal and Peri-Implant Diseases and Conditions. *J Clin Periodontol* 2018;45 Suppl 20:S162-S170.
4. Cortellini P, Tonetti MS. Clinical concepts for regenerative therapy in intrabony defects. *Periodontol* 2000 2015;68:282-307.
5. Needleman IG, Worthington HV, Giedrys-Leeper E, Tucker RJ. Guided tissue regeneration for periodontal infra-bony defects. *Cochrane Database Syst Rev* 2006:CD001724.
6. Parrish LC, Miyamoto T, Fong N, Mattson JS, Cerutis DR. Non-bioabsorbable vs. bioabsorbable membrane: assessment of their clinical efficacy in guided tissue regeneration technique. A systematic review. *J Oral Sci* 2009;51:383-400.
7. Stoecklin-Wasmer C, Rutjes AW, da Costa BR, Salvi GE, Juni P, Sculean A. Absorbable collagen membranes for periodontal regeneration: a systematic review. *J Dent Res* 2013;92:773-781.
8. Kao RT, Nares S, Reynolds MA. Periodontal regeneration - intrabony defects: a systematic review from the AAP Regeneration Workshop. *J Periodontol* 2015;86:S77-104.
9. Tu YK, Needleman I, Chambrone L, Lu HK, Faggion CM, Jr. A Bayesian network meta-analysis on comparisons of enamel matrix derivatives, guided tissue regeneration and their combination therapies. *J Clin Periodontol* 2012;39:303-314.
10. Nyman S, Lindhe J, Karring T, Rylander H. New attachment following surgical treatment of human periodontal disease. *J Clin Periodontol* 1982;9:290-296.
11. Cortellini P, Pini Prato G, Tonetti MS. Periodontal regeneration of human infrabony defects. II. Re-entry procedures and bone measures. *J Periodontol* 1993;64:261-268.

12. Murphy KG, Gunsolley JC. Guided tissue regeneration for the treatment of periodontal intrabony and furcation defects. A systematic review. *Ann Periodontol* 2003;8:266-302.
13. Cortellini P, Buti J, Pini Prato G, Tonetti MS. Periodontal regeneration compared with access flap surgery in human intra-bony defects 20-year follow-up of a randomized clinical trial: tooth retention, periodontitis recurrence and costs. *J Clin Periodontol* 2017;44:58-66.
14. Machtei EE. The effect of membrane exposure on the outcome of regenerative procedures in humans: a meta-analysis. *J Periodontol* 2001;72:512-516.
15. Falk H, Laurell L, Ravald N, Teiwik A, Persson R. Guided tissue regeneration therapy of 203 consecutively treated intrabony defects using a bioabsorbable matrix barrier. Clinical and radiographic findings. *J Periodontol* 1997;68:571-581.
16. Gorski B, Jalowski S, Gorska R, Zaremba M. Treatment of intrabony defects with modified perforated membranes in aggressive periodontitis: subtraction radiography outcomes, prognostic variables, and patient morbidity. *Clin Oral Investig* 2019;23:3005-3020.
17. Trombelli L, Kim CK, Zimmerman GJ, Wikesjo UM. Retrospective analysis of factors related to clinical outcome of guided tissue regeneration procedures in intrabony defects. *J Clin Periodontol* 1997;24:366-371.
18. Cortellini P, Tonetti MS. Long-term tooth survival following regenerative treatment of intrabony defects. *J Periodontol* 2004;75:672-678.
19. Silvestri M, Rasperini G, Milani S. 120 infrabony defects treated with regenerative therapy: long-term results. *J Periodontol* 2011;82:668-675.
20. Steffensen B, Webert HP. Relationship between the radiographic periodontal defect angle and healing after treatment. *J Periodontol* 1989;60:248-254.
21. Tonetti MS, Pini-Prato G, Cortellini P. Periodontal regeneration of human intrabony defects. IV. Determinants of healing response. *J Periodontol* 1993;64:934-940.
22. Bjorn H, Halling A, Thyberg H. Radiographic assessment of marginal bone loss. *Odontol Revy* 1969;20:165-179.
23. Schei O, Waerhaug, J. , Lovdal, A., Arno, A. Alveolar Bone Loss as Related to Oral Hygiene and Age. *J Periodontol* 1959;30:7-16.
24. Kassambara A, Kosinski M. survminer: Drawing Survival Curves using 'ggplo2'. R package version 0.4.2. 2018.
25. Therneau T. *_A Package for Survival Analysis in S_*. version 2.38. 2015.
26. Terry M, Therneau T, Grambsch PM. *_Modeling Survival Data: Extending the Cox Model_*. New York: Springer; 2000.
27. Wickham H. *ggplot2: Elegant Graphics for Data Analysis*. New York; 2009.

28. Cortellini P, Pini Prato G, Tonetti MS. Periodontal regeneration of human intrabony defects with titanium reinforced membranes. A controlled clinical trial. *J Periodontol* 1995;66:797-803.
29. Kher VK, Bhongade ML, Shori TD, Kolte AP, Dharamthok SB, Shirao TS. A comparative evaluation of the effectiveness of guided tissue regeneration by using a collagen membrane with or without decalcified freeze-dried bone allograft in the treatment of infrabony defects: A clinical and radiographic study. *J Indian Soc Periodontol* 2013;17:484-489.
30. Sculean A, Berakdar M, Chiantella GC, Donos N, Arweiler NB, Brex M. Healing of intrabony defects following treatment with a bovine-derived xenograft and collagen membrane. A controlled clinical study. *J Clin Periodontol* 2003;30:73-80.
31. Sculean A, Pietruska M, Schwarz F, Willershausen B, Arweiler NB, Auschill TM. Healing of human intrabony defects following regenerative periodontal therapy with an enamel matrix protein derivative alone or combined with a bioactive glass. A controlled clinical study. *J Clin Periodontol* 2005;32:111-117.
32. Nibali L, Yeh YC, Pometti D, Tu YK. Long-term stability of intrabony defects treated with minimally invasive non-surgical therapy. *J Clin Periodontol* 2018;45:1458-1464.
33. Eickholz P, Krigar DM, Kim TS, Reitmeir P, Rawlinson A. Stability of clinical and radiographic results after guided tissue regeneration in infrabony defects. *J Periodontol* 2007;78:37-46.
34. Sculean A, Kiss A, Miliauskaitė A, Schwarz F, Arweiler NB, Hannig M. Ten-year results following treatment of intra-bony defects with enamel matrix proteins and guided tissue regeneration. *J Clin Periodontol* 2008;35:817-824.
35. Tonetti MS, Cortellini P, Lang NP, et al. Clinical outcomes following treatment of human intrabony defects with GTR/bone replacement material or access flap alone. A multicenter randomized controlled clinical trial. *J Clin Periodontol* 2004;31:770-776.
36. Tonetti MS, Pini-Prato G, Cortellini P. Effect of cigarette smoking on periodontal healing following GTR in infrabony defects. A preliminary retrospective study. *J Clin Periodontol* 1995;22:229-234.
37. Stavropoulos A, Mardas N, Herrero F, Karring T. Smoking affects the outcome of guided tissue regeneration with bioresorbable membranes: a retrospective analysis of intrabony defects. *J Clin Periodontol* 2004;31:945-950.
38. Nickles K, Ratka-Kruger P, Neukrantz E, Raetzke P, Eickholz P. Open flap debridement and guided tissue regeneration after 10 years in infrabony defects. *J Clin Periodontol* 2009;36:976-983.
39. Weigel C, Bragger U, Hammerle CH, Mombelli A, Lang NP. Maintenance of new attachment 1 and 4 years following guided tissue regeneration (GTR). *J Clin Periodontol* 1995;22:661-669.

40. Pretzl B, Kim TS, Steinbrenner H, Dorfer C, Himmer K, Eickholz P. Guided tissue regeneration with bioabsorbable barriers III 10-year results in infrabony defects. *J Clin Periodontol* 2009;36:349-356.
41. Cortellini P, Tonetti MS. Evaluation of the effect of tooth vitality on regenerative outcomes in infrabony defects. *J Clin Periodontol* 2001;28:672-679.
42. Kim YT, Choi JK, Kim DH, Jeong SN, Lee JH. Association between health status and tooth loss in Korean adults: longitudinal results from the National Health Insurance Service-Health Examinee Cohort, 2002-2015. *J Periodontal Implant Sci* 2019;49:158-170.
43. Helal O, Gostemeyer G, Krois J, Fawzy El Sayed K, Graetz C, Schwendicke F. Predictors for tooth loss in periodontitis patients: Systematic review and meta-analysis. *J Clin Periodontol* 2019;46:699-712.
44. Nilsson H, Berglund JS, Renvert S. Longitudinal evaluation of periodontitis and tooth loss among older adults. *J Clin Periodontol* 2019. doi: 10.1111/jcpe.13167
45. Graetz C, Schutzhold S, Plaumann A, et al. Prognostic factors for the loss of molars--an 18-years retrospective cohort study. *J Clin Periodontol* 2015;42:943-950.
46. Kim CK, Choi SH, Kim TS, Kaltschmitt J, Eickholz P. The infrabony defect and its determinants. *J Periodontal Res* 2006;41:498-502.
47. Checchi L, Montevercchi M, Checchi V, Zappulla F. The relationship between bleeding on probing and subgingival deposits. An endoscopical evaluation. *Open Dent J* 2009;3:154-160.
48. Matuliene G, Pjetursson BE, Salvi GE, et al. Influence of residual pockets on progression of periodontitis and tooth loss: results after 11 years of maintenance. *J Clin Periodontol* 2008;35:685-695.

Supplementary Data S1. Flowchart diagram displaying the patient file screening process leading to the final inclusion of the selected patients.

Supplementary Data S2. Kaplan-Meier analysis displaying the stability results throughout the entire follow-up. Stability was defined for a treated site that maintained the 1-year post-surgical CAL, presenting less than 2 mm of change. Every event displays a site that showed CAL loss ≥ 2 mm. The grayish hue represents the upper and lower limit of the 95% confidence bands.

Tables:

Table 1. characteristics of the included patients/defects at baseline

Characteristic		Frequency
Males (N, %)		63, 49.2%
Age		51.7 ± 13.9
Diabetics (N, %)		11, 8.6%
Smoker (N, %)		41, 32.0 %
2017 World workshop ³		
	Stage 3 grade A Periodontitis (N, %)	17,13.3 %
	Stage 3 grade B Periodontitis (N, %)	46, 35.9%
	Stage 3 grade C Periodontitis (N, %)	27, 21.1%
	Stage 4 grade A Periodontitis (N, %)	10, 7.8%
	Stage 4 grade B Periodontitis (N, %)	13, 10.2%
	Stage 4 grade C Periodontitis (N, %)	15, 11.7%
Maxilla (N, %)		66, 37.7%
Mandible (N, %)		109, 62.3%
Endodontically treated (N, %)		14, 8%
Clinical attachment level [CAL (mm)]		9.56 ± 1.93 mm
Pocket depth [PD (mm)]		8.41 ± 1.42 mm
Recession [REC (mm)]		1.21 ± 1.11 mm
Initial radiographic angle		36.86° ± 15.3°
Data are expressed as mean ± standard deviation; or N, numbers, percentages		

Table 2. Results of the regression models evaluating the effect of different variables on the clinical attachment levels of the treated defects at the 1-year recall.

Variable	Univariate analysis			Multivariate analysis			
	Estimate	95% CI	p-value	Estimate	Std. Error	95% CI	p-value
Gender (Male)	-0.83	-1.67, 0.001	0.05				
Age	0.01	-0.01, 0.04	0.26				
Smoking	-1.01	-1.85, -0.16	0.01	-0.91	0.41	-1.73, -0.07	0.03
Diabetes	-0.18	-1.43, 1.07	0.77				
Membrane exposure	-0.26	-0.50, -0.01	0.02	-1.18	0.55	-2.28, -0.06	0.03
Endodontic treatment	-0.31	-1.91, 1.27	0.69				
Initial PD	0.55	0.26, 0.85	<0.001	0.57	0.20	0.16, 0.97	0.006
Initial CAL	0.34	0.14, 0.55	0.001	0.14	0.14	-0.13, 0.42	0.3
Flap design	0.55	-0.58, 1.69	0.33				
Vertical incision	-0.03	-1.04, 0.97	0.94				
Initial radiographic angle	-0.02	-0.05, -0.002	0.03	-0.01	0.01	-0.03, 0.01	0.26
<i>Bold signifies statistical significance; CI, confidence intervals</i>							

Table 3. Life table descriptively summarizing the survival probabilities according to the Kaplan-Meier analysis. Survival estimations for the 5- and 10- year timepoints have been lightly shaded in grey.

Time (months)	Number at risk	Number of event(s)	Survival	Standard Error	95% Confidence Interval	
					Lower bound	Upper bound
0	175	0	0.1	-	-	-
18	159	1	0.994	0.00627	0.981	1.000
19	157	1	0.987	0.00887	0.970	1.000
22	152	1	0.981	0.01093	0.960	1.000
28	138	1	0.974	0.01296	0.949	1.000
30	131	1	0.966	0.01484	0.938	0.996
31	129	1	0.959	0.01651	0.927	0.992
32	125	1	0.951	0.01807	0.916	0.987
40	117	1	0.943	0.01966	0.905	0.982
41	114	1	0.935	0.02116	0.894	0.977
42	113	1	0.927	0.02253	0.883	0.972
44	108	1	0.918	0.02390	0.872	0.966
47	104	2	0.900	0.02650	0.850	0.954
49	101	1	0.891	0.02769	0.839	0.947
55	91	2	0.872	0.03035	0.814	0.933
58	81	1	0.861	0.03183	0.801	0.926
60	76	1	0.850	0.03337	0.787	0.918
71	63	1	0.836	0.03546	0.770	0.909
78	61	1	0.822	0.03743	0.752	0.899
82	58	2	0.794	0.04116	0.717	0.879
90	47	2	0.760	0.04583	0.676	0.856
105	23	1	0.727	0.05447	0.628	0.842
125	19	1	0.689	0.06365	0.575	0.826


150	18	1	0.651	0.07069	0.526	0.805
151	17	1	0.612	0.07619	0.480	0.782
170	14	1	0.569	0.08236	0.428	0.755
207	9	1	0.506	0.09438	0.351	0.729

Table 4. Results of the multilevel cox proportional hazard models evaluating the effect of different variables on the survival of the treated teeth.

Variable	Univariate analysis			Multivariate analysis			
	HR	95% CI	p-value	HR	Std. Error	95% CI	p-value
Gender (Male)	0.89	0.43, 1.85	0.77				
Age	1.03	1.001, 1.063	0.04	1.08	0.03	1.03, 1.14	0.002
Smoking	3.36	1.52, 7.43	0.002	1.75	0.50	0.65, 4.71	0.26
Diabetes	0.98	0.23, 4.15	0.98				
Membrane exposure	3.31	1.51, 7.26	0.003	1.76	0.52	0.63, 4.90	0.28
Maintenance/year	0.45	0.29, 0.67	<0.001	0.34	0.26	0.20, 0.58	<0.001
Endodontic treatment	2.76	1.12, 6.82	0.03	1.35	0.71	0.34, 5.42	0.67
Initial PD	1.27	1.04, 1.56	0.019	0.89	0.22	0.57, 1.38	0.60
Initial CAL	1.28	1.07, 1.54	0.008	1.53	0.20	1.03, 2.28	0.03
Flap design	0.30	0.04, 2.26	0.245				
Vertical incision	0.998	0.38, 2.63	0.99				
Initial radiographic angle	1.015	0.99, 1.05	0.25				


Figures:

Figure 1. Kaplan-Meier survival curve for the entire follow-up period. Each event represents a tooth loss. The grayish hue represents the upper and lower limit of the 95% confidence bands.



Authoi

Figure 2. Kaplan-Meier survival curves displaying the comparison between smokers and non-smokers. Event = Tooth loss. Blue and red hues represent the upper and lower limit of the 95% confidence bands.



Author N