

# Ecology eclipses phylogeny as a major driver of nematode parasite community structure in a graminivorous primate

India Schneider-Crease<sup>1,2,3,4</sup>, Jacinta C. Beehner<sup>3,5,6</sup>, Thore J. Bergman<sup>3,6,7</sup>, Megan A. Gomery<sup>3</sup>, Lia Koklic<sup>1</sup>, Amy Lu<sup>2,3</sup>, Noah Snyder-Mackler<sup>1,3,4,8,9</sup>

<sup>1</sup>Department of Psychology, University of Washington, Seattle, Washington, 98195

<sup>2</sup>Department of Anthropology, Stony Brook University, Stony Brook, New York, 11794

<sup>3</sup>Simien Mountains Gelada Research Project, Sankaber, Ethiopia

<sup>4</sup>Center for Evolution and Medicine, Arizona State University, Tempe, Arizona, 85281

<sup>5</sup>Department of Anthropology, University of Michigan, Ann Arbor, Michigan, 48109

<sup>6</sup>Department of Psychology, University of Michigan, Ann Arbor, Michigan, 48109

<sup>7</sup>Department of Ecology and Evolution, University of Michigan, Ann Arbor, Michigan, 48109

<sup>8</sup>Center for Studies in Demography and Ecology, University of Washington, Seattle, Washington, 98195

<sup>9</sup>Washington National Primate Research Center, University of Washington, Seattle, Washington, 98195

## Data Accessibility

Data and code are available at <https://doi.org/10.5281/zenodo.3827516>.

## Acknowledgements

We thank the Ethiopian Wildlife Conservation Authority, the Amhara National Regional Parks State Development and Protection Authority, and all park officials who have permitted and aided our research. We thank all past and present field crew of the Simien Mountains Gelada Research Project for their dedication to over twelve years of data collection, and we thank Kenneth L. Chiou for intellectual support. Long-term gelada research was supported by the National Science Foundation (IOS-1255974, BCS-0715179, BCS-1723237, BCS-1723228), the National Institutes of Health (5R00AG051764-04), the Leakey Foundation, and the National Geographic

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/1365-2435.13603](https://doi.org/10.1111/1365-2435.13603)

This article is protected by copyright. All rights reserved

Society (Gr. #8989-11, Gr. #8100-06). ISC acknowledges funding from the National Science Foundation Directorate of Social and Behavioral Sciences Fellowship Program and the Arizona State University Center for Evolution and Medicine. Finally, we thank the reviewers and editors of this journal for their indispensable comments. The authors declare no conflict of interest.

### **Author Contributions**

ISC conceptualized the study and designed the methodology with input from AL, JCB, NSM, and TJB. ISC and MAG collected data, and AL, JCB, NSM, and TJB facilitated data collection. ISC and LK analyzed data with contributions from NSM. ISC wrote the manuscript with important intellectual contributions from AL, JCB, NSM, and TJB.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31

DR INDIA A SCHNEIDER-CREASE (Orcid ID : 0000-0002-2699-5304)

Article type : Research Article

**Corresponding Author Email ID: indiaaugusta@gmail.com**

Section: Community Ecology

Editor: Dr Seth Barribeau

**Abstract**

- Understanding the relative strength of ecology and phylogeny in shaping parasite communities can inform parasite control and wildlife conservation initiatives while contributing to the study of host species evolution.
- We tested the relative strengths of phylogeny and ecology in driving parasite community structure in a host whose ecology diverges significantly from that of its closest phylogenetic relatives.
- We characterized the gastrointestinal (GI) parasite community of wild geladas (*Theropithecus gelada*), primates that are closely related to baboons but specialized to graminivory in the Ethiopian Highlands.
- Geladas exhibited very constrained GI parasite communities: only two genera (*Oesophagostomum* and *Trichostrongylus*) were identified across 303 samples. This is far below the diversity reported for baboons (*Papio* spp.) and at the low end of the range of domestic grazers (e.g., *Bos taurus*, *Ovis aries*) inhabiting the same region and ecological niche.
- Using deep amplicon sequencing, we identified 15 amplicon sequence variants (ASVs) within the two genera, seven of which matched to *Oesophagostomum* sp., seven to *Trichostrongylus* sp., and one to *T. vitrinus*.

- 32 • Population was an important predictor of ASV richness. Geladas in the most ecologically  
33 disturbed area of the national park exhibited ~4x higher ASV richness than geladas at a  
34 less disturbed location within the park.
- 35 • In this system, ecology was a stronger predictor of parasite community structure than  
36 phylogeny, with geladas sharing more elements of their parasite communities with other  
37 grazers in the same area than with closely related sister taxa.

38

39 **Keywords:** parasite community structure; habitat-sharing; parasite ecology; parasite evolution;  
40 primate parasite ecology; cercopithecines; gastrointestinal parasites; nemabiome

41

## 42 **Introduction**

43

44 Wild animals are almost invariably infected with at least one parasite throughout their lives  
45 (Haque 2007). The immunological and reproductive costs of these infections have driven  
46 physiological and behavioral adaptations in hosts, ultimately centering parasites as major drivers  
47 of host evolution (Anderson & May 1979, Thomas et al., 1996, Ezenwa et al., 2016). The  
48 structure of parasite communities within a host species or population can thus be understood both  
49 as an evolutionary product and as a dynamic network to which changes can affect host health and  
50 survival. Understanding the major drivers of parasite community structure can accordingly offer  
51 insights into host evolution and into contemporary changes in susceptibility and disease (Patz et  
52 al., 2000).

53

54 The parasite community of a host population is shaped by a number of factors related to host  
55 ecology and phylogeny (Arneberg 2002, Lindenfors et al., 2007). Habitat-sharing may drive the  
56 structure of host parasite communities by increasing the likelihood that host species in a given  
57 geographic area encounter infective parasitic stages shed by other host species in that area. This  
58 is particularly salient in the case of generalist parasites, which are able to infect multiple host  
59 species upon exposure (Zaffaroni et al., 1999, Woolhouse et al. 2001, Ezenwa 2003,  
60 VanderWaal et al., 2014). Ecological changes that result in newly sympatric host communities  
61 allow parasites to accumulate adaptations to new hosts, a phenomenon that underlies the  
62 emergence of novel infectious diseases in wildlife, domestic animals, and humans (Mayer 2000,

63 Daszak et al., 2001). Animals living in habitats in which multiple generalist parasites are  
64 endemic are thus expected to share parasite species and other elements of parasite community  
65 structure such as species richness and prevalence (Poulin 1995, Ezenwa 2003, VanderWaal et al.,  
66 2014).

67  
68 Animals that share a recent common ancestor are also expected to share elements of parasite  
69 community structure, relative to more distantly related animals, even if they live in ecologically  
70 divergent habitats (Poulin 1995, Beer et al., 2019). Certain parasites evolve with their hosts,  
71 accumulating adaptations that facilitate continued infection. During speciation, populations that  
72 evolve to be distinct species are expected to retain the parasite communities of their most recent  
73 ancestor, which then embark on their own evolutionary trajectories (Poulin 1995). Thus,  
74 phylogeny is expected to be a powerful driver of parasite community structure across hosts,  
75 driven by co-evolutionary patterns of hosts and their parasites (Poulin 1995). Supporting this  
76 expectation, the inclusion of phylogeny in analyses of parasite species richness has revealed a  
77 consistently strong effect of shared ancestry across terrestrial mammals (Morand & Poulin 1998)  
78 and within certain mammalian clades (Nunn et al., 2003).

79  
80 The similarity of parasite community structures between two species is thus a consequence of  
81 their phylogenetic proximity and shared ecology, but little is known about the relative strength of  
82 these factors within particular systems. To investigate the interplay between phylogeny and  
83 ecology in shaping the dynamics of parasite-host systems, we turned to a host species whose  
84 ecology differs substantially from its sister taxa and for which predictions about parasite  
85 community based on ecology or phylogeny diverge sharply. Geladas (*Theropithecus gelada*) are  
86 primates that diverged from other members of the Papionini tribe (baboons (*Papio* spp.) and  
87 mangabeys (*Lophocebus* sp. and *Rungwecebus* sp.)) approximately 4-7 mya (Zinner et al., 2018)  
88 and were long considered to be a species of baboon. However, where baboon species are highly  
89 omnivorous and consume fruits, leaves, and meat (Swedell 2011), geladas specialize on  
90 graminoid leaves and supplement with underground plant parts (Fashing et al., 2014, Jarvey et  
91 al., 2018). While baboon species are widely dispersed across the African continent, geladas are  
92 found only in the Ethiopian highlands. On these high-altitude plateaus, geladas share their niche  
93 with domestic grazers (i.e., sheep (*Ovis aries*), cows (*Bos taurus*), donkeys (*Equis asinus*), and

94 horses (*Equus equus*) and less frequently with wild grazers (klipspringer (*Oreotragus oreotragus*),  
95 bushbuck (*Tragelaphus sylvaticus*)) and closely related omnivorous primates (i.e., baboons).  
96 Geladas thus provide a useful model system in which to examine the relative power of  
97 phylogeny and ecology in shaping parasite communities.

98  
99 We first characterized the gastrointestinal nematode parasite community of gelada populations in  
100 the Simien Mountains National Park (SMNP), Ethiopia. We focused on nematode parasites  
101 (Order Nematoda) because many species are generalists, meaning that they possess the capacity  
102 to infect a multitude of hosts across phylogenetic divisions (Zaffaroni et al., 2000, Walker &  
103 Morgan 2014). In addition, many nematode parasite species are transmitted through the fecal-  
104 oral route, increasing the likelihood of exposure for animals that inhabit the same area (Anderson  
105 1988). This means that any observed divergences from expectations based on phylogeny or  
106 niche-sharing will reflect the biological forces shaping parasite communities apart from species-  
107 specificity of certain parasites or differences in exposure related to transmission route.

108  
109 We evaluated the role of habitat-sharing in shaping parasite communities by comparing the  
110 gelada parasite community structure to that reported for domestic grazers in the same region of  
111 Ethiopia where our population of geladas is located (Amhara). We then performed a parallel  
112 analysis to evaluate the role of phylogeny in shaping parasite communities, comparing the gelada  
113 parasite community to that reported for baboons across Africa (*Papio* spp.). To assess on a finer  
114 scale the relative roles of micro-habitat and demography on the gelada parasite community, we  
115 characterized parasite genetic population structure using a recently developed deep amplicon  
116 sequencing approach.

## 117 118 **Materials and Methods**

### 119 120 **Study Sites and Populations**

121  
122 All samples were collected from wild geladas inhabiting the Simien Mountain National Park  
123 (SMNP), Ethiopia (13.1833°N, 38.0667°E). The SMNP is located in the North Gondar Zone of  
124 the Amhara region, covers 13,600 hectares, and is characterized by Afro-montane and Afro-

125 Alpine habitats. The park faces intense anthropogenic pressure from villages within its  
126 boundaries and at its peripheries as well as from high tourist presence with low infrastructural  
127 development. The SMNP is home to the largest remaining population of geladas, which number  
128 approximately ~10,000 across the park (Beehner & Bergman, pers. comm.).

129  
130 Geladas are gaminivorous and terrestrial primates endemic to the Ethiopian Highlands. The  
131 most basic unit of their multi-tiered social system is a reproductive unit, which comprises one  
132 dominant (“leader”) male, ~2-15 related adult females and dependent offspring. The reproductive  
133 unit can also include one or more subordinate (“follower”) males, which are typically previous  
134 leaders (Snyder-Mackler et al., 2012). Male offspring disperse upon reaching maturity, joining  
135 all-male (“bachelor”) groups and eventually becoming leader males of non-natal groups by  
136 overthrowing current leaders. Single reproductive units associate to form “bands” that tend to  
137 forage, travel, and sleep together.

138  
139 The samples analyzed for this study come from three areas within the SMNP (Fig 1): Sankaber,  
140 Chenek, and Limalimo. Sankaber (~3250 masl) is home to the Simien Mountains Gelada  
141 Research Project (SMGRP) field site, a small park ranger village, and a tourist campsite.  
142 Limalimo (~3000 masl) sits at the park’s western boundary and has the closest proximity to large  
143 villages. Chenek (~3600 masl) has a park ranger village and campsite, and serves both as the  
144 ultimate destination for many of the park’s tourists and as a transportation hub for commercial  
145 traffic crossing the park. The linear distance between Sankaber and Limalimo is approximately  
146 16 km; however, since geladas travel along the plateaus of the park, the actual traveling distance  
147 between the two sites is approximately 40 km. Similarly, the linear distance between Chenek and  
148 both Sankaber and Limalimo is 17 km, but geladas would need to travel 21 km along the plateau  
149 to get from Chenek to Sankaber, and 60 km to get from Chenek to Limalimo. Geladas have  
150 notably small day ranges, and the home ranges of units at Chenek and Limalimo fall well outside  
151 those of units at Sankaber (Snyder-Mackler et al., 2012). Thus, the geladas sampled at each of  
152 these sites are likely to belong to separate populations.

153  
154 **Figure 1.** Simien Mountains National Park, Ethiopia. National park boundaries indicated with  
155 dark green, sampling sites indicated with white circles.

156

157 In Sankaber, we collected samples from known individuals under long-term study by the  
158 SMGRP. In Chenek and Limalimo, we habituated individuals over two to three days prior to  
159 sample collection, and recorded descriptions that included age category (i.e., adult, juvenile,  
160 infant), sex, and distinguishing features (e.g., obvious injuries, facial scars, hair patterns). We  
161 cross-checked descriptions upon sample collection to minimize the possibility that individuals  
162 were erroneously sampled more than once.

163

### 164 **Gastrointestinal nematode identification**

165

166 We characterized the gelada gastrointestinal nematode community using both traditional  
167 microscopy and a recently developed deep amplicon sequencing approach (Avramenko et al.,  
168 2015, Avramenko et al., 2017, Pafčo et al., 2018). We used this combination of approaches to  
169 ensure the most rigorous approach possible: microscopy is limited to differentiating between egg  
170 types but can guide the selection of primers for deep amplicon sequencing; and deep amplicon  
171 sequencing provides higher resolution on the level of genetic diversity and genera present.

172

173 **Microscopy.** We non-invasively collected 49 fresh fecal samples from 43 individually  
174 identifiable geladas under long-term study by the SMGRP. A bolus weighing ~1g was taken  
175 from the center of each fecal sample and placed in 5mL tube with ~4mL of 10% buffered  
176 formalin. We performed modified Wisconsin sugar flotations at 1 egg per gram sensitivity at  
177 Duke University. Briefly, samples were placed into 15mL tubes with 10mL of water. Tubes were  
178 spun at 1500 rpm for 10 minutes in a swing-bucket centrifuge, after which the supernatant was  
179 discarded and Sheather's Sugar Flotation Solution added. Pellets were broken up by mixing with  
180 wooden applicators, and additional Flotation Solution was added in order to form a positive  
181 meniscus. Cover slips were placed on the tubes, which were centrifuged at 1500 rpm for 10  
182 minutes. Cover slips were then placed on labeled slides and read using a compound microscope.

183

184 **Amplicon sequencing: DNA extraction and library preparation.** We collected 396 fecal samples  
185 from 160 individually identifiable geladas in the SMNP (9 individuals in Limalimo, 92 in  
186 Sankaber, and 59 in Chenek). Of these fecal samples, 74 were cultured in the field and processed



187 with the Baermann technique (Appendix S1 in Supporting Information), with the resulting larvae  
188 stored in RNAlater. An additional 322 fecal samples were stored in RNAlater without culturing  
189 (Appendix S2). In the laboratory, larval DNA was extracted using the MoBio DNEasy Blood and  
190 Tissue Kit and fecal DNA was extracted with the Powerlyser Powersoil kit as per the  
191 manufacturer protocols, and DNA concentration was measured using a Qubit Fluorometer  
192 (ThermoFisher Scientific). Samples were processed for next generation sequencing following  
193 Avramenko et al. (2015) with primers for the Internal Transcription Spacer 2 (ITS-2), a region  
194 commonly used for Class V nematode identification, designed by Avramenko et al. (2015) and  
195 Pafčo et al. (2018). Briefly, 2ng of each DNA sample was added to 10ul NEBNext buffer and 1ul  
196 of each ITS-2 nematode primer (NC1\_ITS-2: ACG TCT GGT TCA GGG TTG TT; NC2\_ITS-2:  
197 ATG CTT AAG TTC AGC GGG TA). We performed the first PCR under the following  
198 conditions: 95°C for 3 minutes (1x), 98°C for 20 seconds, 62°C for 15 seconds, and 72°C for 15  
199 seconds (25x), and then 72°C for 2 minutes (1x). After a 1x bead-based purification (AMPure  
200 XP Magnetic Beads, Beckman Coulter, Inc), we performed a second PCR to add unique dual  
201 molecular indexes using the following conditions: 98°C for 45 seconds (1x), 98°C for 20  
202 seconds, 63°C for 20 seconds, and 72°C for two minutes (7x). After another 1x bead cleanup,  
203 barcoded samples were pooled in equimolar quantities and sequenced on an Illumina Novaseq  
204 SP flowcell using paired-end 250 bp reads. To increase diversity in the flowcell, we sequenced  
205 this pool along with a different, non-ITS-2 library pool and 10% PhiX. The full protocol can be  
206 accessed at <https://smack-lab.com/protocols/>.

207  
208 Amplicon sequencing: Sequence clustering. We used QIIME2 to align the sequencing reads,  
209 generate amplicon sequence variants (ASVs), and match ASVs with taxonomic data available  
210 through the NCBI Nucleotide database (<https://doi.org/10.5281/zenodo.3827516>). Our initial  
211 filtering steps included removing matches with an e-value of less than  $10^{-100}$  and a percent  
212 identity under 85% (Figure S3), based on a bimodal e-value and percent identity distributions  
213 across the dataset. This filtering rendered a dataset with percent identity matches between 92.04  
214 and 100%, and all potential matches for each sequence were within the same genus. The best  
215 taxonomic match for each ASV was then determined using the reported percent identity matches.  
216 If no match had a species percent identity above 98%, the sequence was assigned at the genus-  
217 level (all genera matches had percent identity matches of above 92.04% in this dataset). To

218 reduce noise caused by minor differences in nucleotide bases and low frequency ASVs, we  
219 filtered our dataset to include only those ASVs that comprised at least 1% of the total abundance  
220 of at least 10% of samples in any of our sampling sites. We imposed this threshold on each site  
221 independently to ensure that we were able to capture any site-based differences in ASV  
222 abundance that would otherwise be eliminated by analyzing all of the samples together (given  
223 the difference in sample sizes collected at each site). This filtering step eliminated 1,128 ASVs  
224 from our dataset, rendering a final taxonomic set of 15 ASVs (Table S4). We then removed all  
225 samples that failed to amplify ( $n = 30$ ) or were not sequenced deeply enough ( $< 1,000$  reads;  $n =$   
226  $61$ ), rendering a final dataset of 305 samples. All downstream analyses were done in R using the  
227 phyloseq package (McMurder & Holmes 2013) and the DESEQ2 package (Love et al., 2014).

228  
229

### 230 **Gelada nematode parasite community: composition, richness and prevalence**

231

232 Because we were able to resolve certain ASVs only to the genus-level, we report parasite  
233 richness at the genus level (“genus richness”). We calculated genus richness by summing the  
234 number of genera represented in our final dataset of 305 gelada fecal samples. We also  
235 calculated the ASV richness by summing the number of ASVs represented across the dataset.  
236 We then calculated the prevalence of each taxon and ASV as the number of samples in which it  
237 appeared divided by the total number of samples.

238

### 239 **Cross-species comparisons: composition and richness**

240

241 We first obtained average genus richness estimates for (1) domestic grazing species in Ethiopia  
242 that share their habitat with geladas (*O. aries*, *B. taurus*, *E. asinus*, *E. equis*) and (2) baboon  
243 species (*Papio* spp.) across Africa. Data on grazer parasites were gathered through a Google  
244 Scholar search using the terms ‘gastrointestinal parasites’, ‘helminths’, ‘nematodes’, and  
245 ‘Amhara, Ethiopia’ paired with Latin and common names for each target species, and baboon  
246 data were gathered from the Global Mammal Parasite Database (gmpd.org). Because some  
247 papers included in these analyses report parasite species identification based on morphology  
248 alone, we calculated richness on the genus level for both ruminant and *Papio* studies. Where

249 studies reported strongyles to their genus or species without morphological examination of  
250 cultured larvae, we reduced all strongyle-species to a ‘strongyle’ category. Since our  
251 morphological and molecular analyses only targeted nematodes (and our molecular approach  
252 only targeted Class V nematodes), we further restricted these datasets to taxa in the nematode  
253 phylum. We then qualitatively compared the composition and richness of the gelada parasite  
254 community to the composition and richness reported for grazers in Amhara, Ethiopia and Papio  
255 spp. across Africa.

256

### 257 **Within-species drivers of ASV richness and abundance: habitat and demography**

258

259 To identify the drivers of gelada parasite community structure on a finer scale, we assessed the  
260 importance of sampling site and demographic predictors (i.e., age, sex) on two metrics of  
261 parasite community: richness (the number of unique ASVs in each fecal sample) and relative  
262 abundance (the number of reads of each ASV normalized by the total number of reads in the  
263 sample). Because robust ages (estimated based on validated morphological cues or known by  
264 date of birth) are only known for Sankaber geladas, models analyzing demographic predictors  
265 include only samples collected at Sankaber ( $n = 237$ ). This dataset includes 146 females and 91  
266 males, of which 215 were adults, 5 were infants, and 17 were juveniles. The overall dataset used  
267 for site-based analyses contained all 305 samples, with 9 samples from Limalimo, 237 samples  
268 from Sankaber, and 59 from Chenek. Of these, 175 were females, 112 were males, and 18 were  
269 individuals of unknown sex. This dataset included 252 adults, 7 infants, 33 juveniles, and 13  
270 individuals of unknown age category. We worked to reduce the likelihood of collecting repeated  
271 samples from individuals at Limalimo and Chenek by cross-checking individual descriptions  
272 upon sample collection.

273

274 First, we evaluated the impact of sampling site, age, and sex on ASV richness. We modeled the  
275 log-transformed ASV richness of each sample as a function of sampling site and included  
276 individual ID as a random effect (‘lmer’ function in the ‘lme4’ package; Pinheiro et al., 2018,  
277 with default parameters). We then modeled the log-transformed ASV richness of each sample as  
278 a function of individual age (in years), sex, and the interaction between the two, with individual  
279 ID as a random effect. We performed the same analyses on other measures of sample alpha

280 diversity (i.e., Shannon, Simpson, Inverse Simpson, and Chao1 indices of diversity). We also  
281 ensured that our approach was equally likely to pick up similar ASV compositions in both  
282 cultured and uncultured samples by performing an analysis that modeled log-transformed ASV  
283 richness as a function of type of sample (0/1; uncultured/cultured) while controlling for  
284 individual ID as a random effect.

285  
286 Second, we evaluated the impact of sampling site, age, and sex on relative ASV abundance. We  
287 modeled the number of reads of each ASV, normalized with a scaling factor (DESEQ 2  
288 package), as a function of site and with individual ID as a random effect. We fitted a GLMM  
289 with a negative binomial distribution and a term for zero-inflation using the glmmTMB package  
290 (Bolker 2014). To address the variation in sample sizes between sites, we performed a secondary  
291 analysis that excluded samples from Limalimo (n=9). For this analysis, we subsampled 60  
292 samples from Sankaber without replacement 10,000 times and examined the distribution of  
293 coefficients to assess the role of sampling site in predicting abundance of ASVs.

294  
295 We then modeled normalized ASV abundance as a function of individual age (in years) and sex  
296 across the Sankaber-only dataset, with individual ID as a random effect. We again used the  
297 glmmTMB package, fitting a GLMM with a negative binomial distribution. We adjusted all p-  
298 values with a Bonferroni correction to account for multiple comparisons.

299

## 300 **Results**

301

### 302 **Microscopy**

303

304 Strongyle-type eggs were virtually the sole egg type observed with microscopy across 48 fecal  
305 samples from individuals across age-sex groups. At least one strongyle-type egg was observed in  
306 94.2% (17/18) of samples from females and in 96.8% (29/30) of samples from males. Similarly,  
307 at least one strongyle-type egg was observed in 90.9 % (11/12) of samples from juveniles and in  
308 97.3% (36/37) of samples from adults. A single instance of a possible non-strongylid egg type  
309 (ascarid-type morphology) was recorded during the validation stages, with a prevalence of 2.7%  
310 of the formalin-fixed sample set (1/37) and 0.3% of all analyzed samples (1/347).

311

312 **Amplicon sequencing**

313

314 With the deep amplicon approach, we confidently identified 15 ASVs mapping to two genera  
315 and one species (*Oesophagostomum* sp., *Trichostrongylus* sp., and *Trichostrongylus vitrinus*)  
316 (Table S4, Figure S5). All sequences are available on NCBI (BioProject Accession:  
317 PRJNA609008). The application of our approach to cultured samples (versus uncultured  
318 samples) did not have an impact on the richness of ASV communities ( $t = -1.618$ ,  $p = 0.1$ ).

319

320 **Gelada nematode parasite community: composition, richness and prevalence**

321

322 Our examination of the gelada gastrointestinal nematode community revealed a high degree of  
323 homogeneity. Across the entire SMNP gelada population, only two genera were ever observed  
324 (*Trichostrongylus* and *Oesophagostomum*), and both genera were found in almost all samples:  
325 *Oesophagostomum* appeared in 97.7% of samples (298/305 samples), while *Trichostrongylus*  
326 appeared in 98.7% of samples (301/305 samples).

327

328 At the individual sequence level, the ASV community was more complex: 15 ASVs observed  
329 across the sample set. Seven of these ASVs matched to *Oesophagostomum* sp., seven to  
330 *Trichostrongylus* sp., and one to *T. vitrinus*. The low relative identity match (~94%) of the ASV  
331 that mapped to *T. vitrinus* may suggest that this ASV represents a strain belonging to a species  
332 closely related to *T. vitrinus*. ASV prevalence ranged from 0% to 100% within each of the three  
333 sites (Table S6).

334

335 **Cross-species comparisons: composition and richness**

336

337 Our dataset of grazers in the Amhara region of Ethiopia contained 10 independent studies on  
338 domestic sheep, cows, donkeys, horses, and mules. Nematode genus richness ranged from 2.0-  
339 7.0, with an average of 4.8 (Table S7). Our dataset of *Papio* species (*P. anubis*, *P. cynocephalus*,  
340 *P. papio*, *P. ursinus*) included 30 independent studies. Nematode parasite genus richness ranged  
341 from 4.0-8.2, with an average of 6.3 (Table S8). Comprised of only two genera, gelada nematode

342 parasite genus richness fell within the distribution of grazer genus richness and below the range  
343 of the distributions for *Papio* species (Fig 2).

344

345 **Figure 2:** Nematode parasite richness by genus for 5 grazing species in Amhara, Ethiopia (light  
346 blue) and 5 *Papio* species (purple). The parasite genus richness of geladas is indicated by the  
347 dashed line.

348

### 349 **Within-species drivers of ASV richness and abundance: habitat and demography**

350

351 Sampling site was a strong predictor of ASV richness (Fig 3). Chenek samples had  
352 approximately 39% more ASVs than those from Sankaber (estimate: 0.39;  $p < 0.001$ ). No strong  
353 difference was observed between Limalimo and either Sankaber or Chenek, possibly due to the  
354 small sample size at Limalimo. Sampling site had a similar effect on other metrics of alpha  
355 diversity (Figure S9). No demographic predictors were strongly associated with ASV richness  
356 (Table S10).

357

358 **Figure 3.** (A) The highest richness of parasite amplicon sequence variants (ASVs) was observed  
359 at Chenek (in green). (B) Relative ASV abundance differed at each site, here represented with a  
360 network based on the Jaccard index of dissimilarity (using the 'igraph' package; Csardi &  
361 Nepusz 2006). Nodes represent the 15 unique ASVs, colors reflect the relative proportion of  
362 ASVs at each site, and node size reflects the relative abundance of each ASV.

363

364 The effect of sampling site on normalized relative abundance varied by ASV. Subsampling  
365 demonstrated that only three ASVs showed a trend towards differential relative abundance by  
366 site. Two *Trichostrongylus* ASVs and one *Oesophagostomum* ASV trended towards higher  
367 relative abundances in Sankaber across 10,000 random subsample iterations (Table S11). No  
368 demographic predictors were associated with changes in normalized abundance for any ASV  
369 (Table S12).

370

## 371 **Discussion**

372

373 Wild geladas in the SMNP exhibited a highly constrained parasite community, below the range  
374 of diversity reported for *Papio* spp. and even at the low end of the range reported for other  
375 grazers. Geladas were infected by only two nematode genera, *Trichostrongylus* and  
376 *Oesophagostomum*, both of which are generalist parasites that infect a broad array of herbivores  
377 as well as humans. Infections were ubiquitous, occurring in geladas at all three sampling sites  
378 and of all ages and sexes, but ASV diversity was highest at the Chenek site (Fig 3A). Within  
379 these two genera, we identified 15 ASVs: seven from *Oesophagostomum* and eight from  
380 *Trichostrongylus*.

381

### 382 **Cross-species comparisons of nematode parasite communities: composition and richness**

383

384 The nematode parasite community of geladas fell within the distribution of nematode genus  
385 richness observed in grazing species across the Amhara region of Ethiopia, where our study  
386 gelada population is located, and fell below the distribution observed in *Papio* species across  
387 Africa. Strongyles, including *Oesophagostomum* and *Trichostrongylus*, were found across  
388 geladas, grazers, and *Papio* sp., but geladas lacked any of the other parasites found in either  
389 group. In addition to strongyles, grazers were commonly observed to harbor infections with  
390 *Ascaris* sp., *Dictyocaulus* sp., *Parascaris* sp., and *Trichuris* sp. In *Papio* species, the most  
391 common non-strongyle parasites were *Ascaris* sp., *Enterobius* sp., *Physaloptera* sp., *Ternidens*  
392 sp., *Streptopharagus* sp., and *Trichuris* sp. Geladas were infected with only the most general of  
393 gastrointestinal nematode parasites.

394

395 The absence of eggs characteristic of *Ascaris* sp. or *Trichuris* sp. in gelada samples analyzed  
396 with microscopy (with one possible exception) is notable, particularly because these parasites are  
397 found in both grazers and *Papio* species. One possible avenue of explanation for the unusually  
398 low nematode parasite diversity in geladas is their evolutionary history. The extinct sister taxa to  
399 geladas ranged across much of Africa, Europe, and India, and all possessed morphological  
400 adaptations that suggest similar levels of terrestriality and graminivory (Krentz 1993). Thus,  
401 geladas have likely been terrestrial grazers since their origin, a characteristic that likely was  
402 accompanied by high levels of exposure to soil-transmitted helminth parasites. If the costs  
403 associated with infection were high, resulting in damages to reproductive success or survival,

404 parasitism may have exacted sufficient pressure for selection of immunological mechanisms to  
405 mitigate those costs in this particular system.

406

407 **Within-species drivers of ASV richness and abundance: habitat and demography**

408

409 While both *Oesophagostomum* sp. and *Trichostrongylus* sp. were observed across sampling sites,  
410 ages, and sexes, certain ASVs appeared in varying frequencies and abundances at different sites  
411 (Fig 3B). The overall homogeneity of the parasite communities of geladas across all three sites  
412 indicates a lack of parasite population structure at the genus level. This could be attributable to  
413 gelada dispersal patterns, in which males may travel extensively to reach new groups upon  
414 maturity. Because both genera found in geladas were also found in other ruminants in Ethiopia, it  
415 is also possible that multiple host species transport these parasites across the park and contribute  
416 to the homogeneity of the gelada parasite community.

417

418 Our ASV-level analysis revealed fine-grained parasite population structure. The higher ASV  
419 richness in Chenek individuals compared to Sankaber could be tied to higher overall host species  
420 diversity; Chenek is the only of our sampled sites where geladas overlap with *Walia ibex* (*Capra*  
421 *walia*) and Ethiopian wolves (*Canis simensis*). In Limalimo and Sankaber, geladas overlap only  
422 with klipspringer (*Oreotragus oreotragus*), bushbuck (*Tragelaphus sylvaticus*), and domestic  
423 ruminants, all of which are also found in Chenek. Another possible explanation hinges on the  
424 exposure of Chenek geladas to human trash and fecal waste. The Chenek campsite, unlike the  
425 park's other campsites, contains multiple open trash pits and is plagued by the extensive and  
426 open deposition of human excrement. The campsite is also a transportation hub, and local  
427 travelers regularly feed the geladas. As a result, geladas at the Chenek campsite display atypical  
428 behavior that includes raiding trash pits and spending time concentrated around human waste.  
429 The higher ASV richness of Chenek could thus be tied to exposure to strains of  
430 *Oesophagostomum* sp. and *Trichostrongylus* sp. in human waste.

431

432 While richness and abundance typically vary according to age- and sex- based differences in  
433 immune function and pathogen exposure, geladas were uniformly infected across demographic  
434 categories and neither ASV richness nor abundance was affected by age or sex. The lack of



435 demographic structure in ASV richness and abundance may be related to gelada ecology. As  
436 terrestrial grazers, geladas are likely to encounter infective stages of these parasites at the  
437 beginning of their lives. Even while dependent offspring are still nursing and not yet feeding on  
438 grass, they frequently dismount from their mothers and experiment with placing objects (e.g.,  
439 soil, feces, grasses) in their mouths. This may provide sufficient exposure to eggs and larvae for  
440 infections to establish, with continued exposure throughout their lives.

441  
442 The near-ubiquitous infection of geladas with species of both parasite genera, coupled with the  
443 lack of detectable demographic structure in ASV abundance, suggests that the gelada immune  
444 system may tolerate infections as opposed to working to resist or eliminate them. Resistance and  
445 tolerance are the two principal approaches available to hosts upon the establishment of a parasite  
446 infection (Best et al., 2008, Read et al., 2008, Råberg et al. 2009, Medzhitov et al., 2012, Kutzer  
447 & Armitage 2016). Resistance includes the reduction or elimination of infections, which are  
448 accompanied by collateral costs to the host, while tolerance includes the mitigation of the  
449 damage of infection while avoiding the costs of active defense (Read et al., 2008, Medzhitov et  
450 al., 2012, Råberg 2014).

451  
452 The evolution of resistance in a host-parasite system is expected when the costs of infection are  
453 higher than the costs of mounting an immune response to limit the infection, whereas the  
454 evolution of tolerance is expected when the costs of infection are not higher than the costs of the  
455 immune response (Råberg et al., 2009). In practice, resistance should reduce parasite prevalence  
456 while tolerance is expected increase it or have no effect (Råberg et al., 2009). The high  
457 prevalence of both *Oesophagostomum* sp. and *Trichostrongylus* infections fits the pattern  
458 expected in a system in which tolerance—as opposed to resistance—has evolved. To test the  
459 hypothesis that the gelada-nematode parasite system is characterized by tolerance-focused  
460 strategies, further work must be done to quantify infection intensity (a measure of parasitism that  
461 we did not gather here) and to assess the relationship between intensity and host fitness (Råberg  
462 et al. 2009, Jackson et al., 2014) or other correlates of health such as body condition (Hayward et  
463 al., 2014, Jackson et al., 2014, Råberg 2014) across this population.

464

465 In this system, habitat-sharing appears to be a stronger force in shaping the gastrointestinal  
466 nematode parasite community of geladas than phylogeny. The gelada gastrointestinal nematode  
467 parasite community of geladas diverged sharply in richness and composition from those of their  
468 sister taxa in the *Papio* genus, but fell within the distribution grazer parasite genus richness in the  
469 same area and was similarly composed. However, it should be noted that we collapsed any egg-  
470 based identification of strongyle-type egg to genus- or species-level to a ‘strongyle’ category for  
471 our analyses. In many of the grazer studies we included, this category likely represents more than  
472 one genus. Thus, it is possible-- and perhaps even likely-- that geladas fall even below the  
473 distribution of grazer nematode parasite genus richness. Our results emphasize the importance of  
474 a dynamic confluence of factors – including abiotic factors such as altitude and climate as well as  
475 biotic factors such as niche-sharing and host diversity – in shaping parasite communities.

476

#### 477 **References**

478

479 Anderson, R. C. (1988). Nematode transmission patterns. *Journal of Parasitology*, 30-45.

480

481 Anderson, R. M., & May, R. M. (1979). Population biology of infectious diseases: Part I. *Nature*,  
482 280(5721), 361.

483

484 Appleton, C. C., & Brain, C. (1995). Gastro-intestinal parasites of *Papio cynocephalus ursinus*  
485 living in the central Namib desert, Namibia. *African Journal of Ecology*, 33(3), 257-265.

486

487 Arneberg, P. (2002). Host population density and body mass as determinants of species richness  
488 in parasite communities: comparative analyses of directly transmitted nematodes of mammals.  
489 *Ecography*, 25(1), 88-94.

490

491 Avramenko, R. W., Redman, E. M., Lewis, R., Yazwinski, T. A., Wasmuth, J. D., & Gilleard, J.  
492 S. (2015). Exploring the gastrointestinal “nemabiome”: deep amplicon sequencing to quantify  
493 the species composition of parasitic nematode communities. *PLoS One*, 10(12), e0143559.

494

495 Avramenko, R. W., Redman, E. M., Lewis, R., Bichuette, M. A., Palmeira, B. M., Yazwinski, T.  
496 A., & Gilleard, J. S. (2017). The use of nemabiome metabarcoding to explore gastro-intestinal  
497 nematode species diversity and anthelmintic treatment effectiveness in beef calves. *International*  
498 *Journal for Parasitology*, 47(13), 893-902.  
499  
500 Beer, A., Ingram, T., & Randhawa, H. S. (2018). Role of ecology and phylogeny in determining  
501 tapeworm assemblages in skates (Rajiformes). *Journal of Helminthology*, 1-14.  
502  
503 Bezjian, M., Gillespie, T. R., Chapman, C. A., & Greiner, E. C. (2008). Coprologic evidence of  
504 gastrointestinal helminths of forest baboons, *Papio anubis*, in Kibale National Park, Uganda.  
505 *Journal of Wildlife Diseases*, 44(4), 878-887.  
506  
507 Bogale, B., Melaku, A., Chanie, M., Fentahun, T., & Berhanu, A. (2014). First report of helminth  
508 parasites of *Walia ibex* (*Capra walie*) at Simen Mountains National Park, Natural World  
509 Heritage Site, Northern Ethiopia. *International Journal of Animal and Veterinary Advances*, 6, 1-  
510 4.  
511  
512 Bogale, B., Chanie, M., Melaku, A., Fentahun, T., & Berhanu, A. (2014). Occurrence, intensity  
513 and parasite composition of gastrointestinal helminth parasites in *Walia ibex* (*Capra walie*) at  
514 Semien Mountains National Park, Natural World Heritage Site, northern Ethiopia. *Acta*  
515 *Parasitologica Globalis*, 5(1), 19-25.  
516  
517 Csardi G, Nepusz T (2006). The igraph software package for complex network  
518 research. *InterJournal, Complex Systems*, 1695. <http://igraph.org>.  
519  
520 Dagnachew, S., Amamute, A., & Temesgen, W. (2011). Epidemiology of gastrointestinal  
521 helminthiasis of small ruminants in selected sites of North Gondar zone, Northwest Ethiopia.  
522 *Ethiopian Veterinary Journal*, 15(2).  
523

524 Damiana, F. R., O'Riain, M. J., Vera, J. A., & Christopher, C. A. (2012). A coprological survey  
525 of protozoan and nematode parasites of free-ranging chacma baboons (*Papio ursinus*) in the  
526 southwestern Cape, South Africa. *African Journal of Wildlife Research*, 42(1), 35-45.  
527

528 Daszak, P., Cunningham, A. A., & Hyatt, A. D. (2001). Anthropogenic environmental change  
529 and the emergence of infectious diseases in wildlife. *Acta Tropica*, 78(2), 103-116.  
530

531 Ezenwa, V. O. (2003). Habitat overlap and gastrointestinal parasitism in sympatric African  
532 bovids. *Parasitology*, 126(4), 379-388.  
533

534 Ezenwa, V. O., Archie, E. A., Craft, M. E., Hawley, D. M., Martin, L. B., Moore, J., & White, L.  
535 (2016). Host behaviour–parasite feedback: an essential link between animal behaviour and  
536 disease ecology. *Proceedings of the Royal Society B: Biological Sciences*, 283(1828), 20153078.  
537

538 Fashing, P. J., Nguyen, N., Venkataraman, V. V., & Kerby, J. T. (2014). Gelada feeding ecology  
539 in an intact ecosystem at Guassa, Ethiopia: variability over time and implications for theropith  
540 and hominin dietary evolution. *American Journal of Physical Anthropology*, 155(1), 1-16.  
541

542 Hahn, N. E., Proulx, D., Muruthi, P. M., Alberts, S., & Altmann, J. (2003). Gastrointestinal  
543 parasites in free-ranging Kenyan baboons (*Papio cynocephalus* and *P. anubis*). *International*  
544 *Journal of Primatology*, 24(2), 271-279.  
545

546 Hayward, A. D., Nussey, D. H., Wilson, A. J., Berenos, C., Pilkington, J. G., Watt, K. A., ... &  
547 Graham, A. L. (2014). Natural selection on individual variation in tolerance of gastrointestinal  
548 nematode infection. *PLoS Biology*, 12(7), e1001917.  
549

550 Jackson, J. A., Hall, A. J., Friberg, I. M., Ralli, C., Lowe, A., Zawadzka, M., ... & Bradley, J. E.  
551 (2014). An immunological marker of tolerance to infection in wild rodents. *PLoS Biology*, 12(7),  
552 e1001901.  
553

554 Kemal, J., & Terefe, Y. (2013). Prevalence of gastrointestinal parasitism of cattle in Gedebano  
555 Gutazer Wolene district, Ethiopia. *Journal of Veterinary Medicine and Animal Health*, 5(12),  
556 365-370.

557

558 Kutzer, M. A., & Armitage, S. A. (2016). Maximising fitness in the face of parasites: a review of  
559 host tolerance. *Zoology*, 119(4), 281-289.

560

561 Lindenfors, P., Nunn, C. L., Jones, K. E., Cunningham, A. A., Sechrest, W., & Gittleman, J. L.  
562 (2007). Parasite species richness in carnivores: effects of host body mass, latitude, geographical  
563 range and population density. *Global Ecology and Biogeography*, 16(4), 496-509.

564

565 Medzhitov, R., Schneider, D. S., & Soares, M. P. (2012). Disease tolerance as a defense strategy.  
566 *Science*, 335(6071), 936-941.

567

568 Mezgebu, T., Tafess, K., & Tamiru, F. (2013). Prevalence of gastrointestinal parasites of horses  
569 and donkeys in and around Gondar Town, Ethiopia. *Open Journal of Veterinary Medicine*, 3(06),  
570 267.

571

572 Nunn, C. L., Altizer, S., Jones, K. E., & Sechrest, W. (2003). Comparative tests of parasite  
573 species richness in primates. *The American Naturalist*, 162(5), 597-614.

574

575 Pafčo, B., Čížková, D., Kreisinger, J., Hasegawa, H., Vallo, P., Shutt, K., ... & Modrý, D. (2018).  
576 Metabarcoding analysis of strongylid nematode diversity in two sympatric primate  
577 species. *Scientific Reports*, 8(1), 1-11.

578

579 Patz, J. A., Graczyk, T. K., Geller, N., & Vittor, A. Y. (2000). Effects of environmental change  
580 on emerging parasitic diseases. *International Journal for Parasitology*, 30(12-13), 1395-1405.

581

582 Poulin, R. (1995). Phylogeny, ecology, and the richness of parasite communities in vertebrates.  
583 *Ecological Monographs*, 65(3), 283-302.

584

585 Råberg, L. (2014). How to live with the enemy: Understanding tolerance to parasites. *PLoS*  
586 *Biology*, 12(11), e1001989.

587

588 Råberg, L., Graham, A. L., & Read, A. F. (2008). Decomposing health: tolerance and resistance  
589 to parasites in animals. *Philosophical Transactions of the Royal Society B: Biological Sciences*,  
590 364(1513), 37-49.

591

592 Read, A. F., Graham, A. L., & Råberg, L. (2008). Animal defenses against infectious agents: is  
593 damage control more important than pathogen control. *PLoS Biology*, 6(12), e1000004

594

595 Snyder-Mackler, N., Beehner, J. C., & Bergman, T. J. (2012). Defining higher levels in the  
596 multilevel societies of geladas (*Theropithecus gelada*). *International Journal of*  
597 *Primatology*, 33(5), 1054-1068.

598

599 VanderWaal, K., Omondi, G. P., & Obanda, V. (2014). Mixed-host aggregations and helminth  
600 parasite sharing in an East African wildlife–livestock system. *Veterinary Parasitology*, 205(1-2),  
601 224-232.

602

603 Walker, J. G., & Morgan, E. R. (2014). Generalists at the interface: nematode transmission  
604 between wild and domestic ungulates. *International Journal for Parasitology: Parasites and*  
605 *Wildlife*, 3(3), 242-250.

606

607 Woolhouse, M. J., L. H. Taylor, and D. T. Haydon. 2001. Population biology of multihost  
608 pathogens. *Science*, 292:1109–1112.

609

610 Workentine, M. L., Chen, R., Zhu, S., Gavriliuc, S., Shaw, N., de Rijke, J., ... & Gilleard, J. S.  
611 (2019). A database for ITS2 sequences from nematodes. *bioRxiv*, 689695.

612

613 Zaffaroni, E., Manfredi, M. T., Citterio, C., Sala, M., Piccolo, G., & Lanfranchi, P. (2000). Host  
614 specificity of abomasal nematodes in free ranging alpine ruminants. *Veterinary Parasitology*,  
615 90(3), 221-230.

616

617 Zinner, D., Atickem, A., Beehner, J. C., Bekele, A., Bergman, T. J., Burke, R., ... & Knauf, Y.


618 (2018). Phylogeography, mitochondrial DNA diversity, and demographic history of geladas

619 (*Theropithecus gelada*). *PloS One*, 13(8), e0202303.

620


621


Author Manuscript





Number of Studies





## ASV Abundance by Site

