


RESEARCH ARTICLE

Temporal trends of sulfur levels in soils of northwest Ohio (USA) between 2002 and 2014

Leandro Michalovicz¹  | Warren A. Dick¹ | Cassio A. Tormena² |
Marcelo M. L. Müller³ | Eduardo C. Cervi⁴

¹School of Environmental and Natural Resources, Ohio Agricultural Research and Development Center, Wooster, OH

²Graduate Program in Agronomy, The State University of Maringa, Maringa, PR, Brazil

³Graduate Program in Agronomy, The Mid-West State University, Guarapuava, PR, Brazil

⁴School for Environment and Sustainability, University of Michigan, Ann Arbor, MI

Correspondence

Leandro Michalovicz, School of Environmental and Natural Resources, Ohio Agricultural Research and Development Center, 1680 Madison Av., Wooster, OH 44691, USA
Email: leandromichalovicz@gmail.com

Funding information

The Ohio State University College of Food, Agricultural, and Environmental Sciences, 7637285

Abstract

Sulfur (S) is an essential nutrient for plant growth. Despite increasing reports of yield responses of crops to S fertilization, there is limited information about changes in the soil test concentrations of S. This study used a soil chemical analysis dataset from 2002 to 2014 to evaluate changes in soil S and other nutrient levels. The soil-test database comprised 8,428 topsoil samples (0–20 cm depth layer) collected from 143 farm fields located in the northwest (NW) Ohio counties of Defiance, Paulding, and Williams. Except for S, the database showed no significant changes in soil chemical properties from NW Ohio between 2002 and 2014. Soil sulfate (SO_4^{2-}) levels have linearly decreased by 63% from 2002 to 2014, reaching the range of concentration considered deficient for the main cereal crops. With no changes in soil organic matter (SOM) and pH, this result was attributed primarily to enactment of air quality regulations, since soil SO_4^{2-} decreases were directly correlated with the reductions of SO_2 emissions (–70%), SO_4^{2-} in rainwater (–66%) and deposited (–52%) in NW Ohio between the years of 2002–2013. Furthermore, combined increasing crop yields and insufficient compensation by fertilization had role on decreasing soil SO_4^{2-} levels. Current fertilization practices and wet deposition of S have not been sufficient to balance S removals from soil leading to the declines in the soil test S levels. It is imperative to paid more attention to practices that maintain soil S fertility levels to avoid yield penalties associated with soil S deficiencies.

KEYWORDS

atmospheric depositions, greenhouse gases, soil sulfur fertility, sulfur levels, temporal trends, wet deposition

1 | INTRODUCTION

Sulfur (S) is the ninth most abundant element on Earth, being naturally found in the form of pure sulfide and sulfate minerals (Khan & Mazid, 2011). Although considered a secondary macronutrient, S is the fourth most essential nutrient for plants (Franzen & Grant, 2008), performing several important roles in growth, development, and survival (Tripathi et al., 2014). Adequate soil levels of this nutrient are required in order to maintain satisfactory yields (Dick, Kost, & Chen, 2008).

Plants uptake S mainly in the sulfate form (SO_4^{2-}), but soil retention of this anion, however, changes according to both soil chemical and physical properties (Tabatai, 1987; Raji, 2008). Soil surface layers have a lower capacity to retain SO_4^{2-} due to the predominance of negative charges generated by soil organic matter (SOM) and higher pH values (Scherer, 2009), and due to the presence of other competitive anions like phosphates and carbonates (Eriksen, 2009; Sokolova & Alekseeva, 2008).

The organic pool makes up almost 95% of the total S in non-calcareous soils and the mineralization of SOM pool often is capable

of supplying much of the plant's requirement for S (Kovar & Grant, 2011). Therefore, any management practice that leads to decreases in both the amount of organic residue inputs and residual SOM will negatively affect S availability for crops (Blanco-Canqui et al., 2015; Kibet, Blanco-Canqui, & Jasa, 2016). Sulfur deficiencies are corrected by applying inorganic fertilizers that include elemental S, ammonium sulfate, simple superphosphate, FGD gypsum or phosphogypsum, and potassium and magnesium sulfates (Camberato & Casteel, 2017; Lucheta & Lambais, 2012).

Another critical source of soil S is atmospheric deposition (Aas et al., 2019). The S in the atmosphere is a result of energy production that comes from the burning of fossil fuels (Gautam et al., 2019). Gases containing S (e.g., sulfur dioxide—SO₂) that are generated by burning fossil fuels can return to the Earth's surface dissolved in rainwater or attached to solid particles (Eriksen, 2009). However, the adoption of strict regulations for emissions of greenhouse gases around the world has drastically reduced S atmospheric depositions (Haneklaus, Bloem, & Schnug, 2008; United States Environmental Protection Agency, 2020; Vieira-Filho, Lehmann, & Fornaro, 2015).

The United States approved its first federal regulation dealing with air quality control in 1955. This regulation, continuously improved until its current version and active since 1990, covers the control of acid deposition and the emission levels for 189 gases (United States Environmental Protection Agency, 2013). This decrease in S deposition has occurred at the same time as increased S uptake and extraction by plants that has greatly increased in the last 50 years. Not only is S removal due to higher plant yields that increased almost 200% for the most cultivated cereals (Food and Agriculture Organization of the United Nations, 2014) but also by the increases in the harvest indexes achieved by plant breeding (Koester, Skoneczka, Cary, Diers, & Ainsworth, 2014; Pan & Deng, 2007).

Historically, over the most recent decades until now, S has generally had soil concentrations above the critical deficiency limits, mainly due to inputs of atmospheric depositions (Kost, Chen, & Dick, 2008). As a result, S availability in soils has not been considered a limiting factor for plant growth and crop yields, resulting in S receiving less attention than other macronutrients such as N (Li et al., 2019). However, in the last decades, crops like soybean, wheat, and maize have shown positive yield and nutrition responses when supplying S under different pedoclimatic conditions in the USA (Rehm, 2005; Chen, Dick, & Kost, 2008; Chen, Dick, & Nelson, 2005; Sloan, Dowdy, Dolan, & Rehm, 1999) and other countries around the world (Broch, Pavinato, Possentti, Martin, & Del Quiqui, 2011; Pias, Tiecher, Cherubin, Mazurana, & Bayer, 2019; Singh, Meena, Bharati, & Gade, 2013; Tiecher et al., 2012; Tisdale, Reneau, & Platou, 1986). These results provide evidence that both current fertilization practices and atmospheric depositions have not been sufficient to maintain adequate soil S levels, and consequently, leading to an inability of crops to realize their maximum yield potential (Mikkelsen & Norton, 2013).

Despite increasing reports of crop's positive response to S fertilization, there is limited information about temporal changes in soil S levels. This research hypothesizes that the current yield level of crops,

the reduction of S atmospheric emissions and depositions, and the absence of compensation by the use of S fertilizers is leading to a gradual decrease in soil S levels over time. This study aimed to use a soil chemical analysis dataset from NW Ohio (USA) farms to evaluate changes in soil S and other nutrient levels from 2002 to 2014.

2 | MATERIAL AND METHODS

2.1 | Characterization of the study area

The State of Ohio is located in what is called the Eastern Cornbelt of the United States. It is divided into 88 counties and totals approximately 116,096 km² of total area (Figure 1). The NW region of Ohio is composed of 14 counties along with another 10 counties that are frequently reported as belonging to the NW region of the State (State of Ohio, 2010).

The climate in NW Ohio is classified as Dfa in the Köppen-Geiger scale (Köppen & Geiger, 1928) and is characterized by temperate temperatures that average between 3°C to 18°C in the three coldest months and above 10°C in the hottest month. There are well-defined winter and summer seasons and no dry periods. The data of annual accumulated rainfall and average temperatures to the NW Ohio during the period of study (Figure 2) were obtained from National Oceanic and Atmospheric Administration (NOAA) weather stations.

Chemical elements deposition data were taken from the National Atmospheric Deposition Program (NAPD) station located in Crawford and responsible for monitoring the NW Ohio State. Emissions data of S and N due primarily to coal burning were obtained from US Energy Information Administration (US EIA). Information about cereal crops




FIGURE 1 Location of NW Ohio counties in the State of Ohio, the United States of America

area and historical yields in Ohio were taken from the National Agricultural Statistics Service (NASS).

2.2 | Database description

The soil's database was originally made up of 9,080 soil chemical tests of soil samples (0–20 cm depth) of farms localized in the Ohio




FIGURE 2 Accumulated annual rainfall (circles) and average annual temperature (squares) in NW Ohio State between the years of 2002 and 2014. The dotted lines represent the average values of annual precipitation and rainfall in the last 100 years. Source: National Oceanic and Atmospheric Administration (2015)

counties of Defiance, Paulding and Williams totaling an area of 5,900 ha (Figure 1 and Table 1).

The soil chemical data (Figure 3) included cation exchange capacity (CEC) by the sum of exchangeable cations, pH 1:1 in H₂O (McLean, 1982), H + Al by SMP solution (Shoemaker, Mclean, & Pratt, 1961), soil organic matter (SOM) determined as the loss of mass by ignition at 360°C (Schulte & Hopkins, 1996), inorganic N extracted by 1 M KCl (Dahnke, 1990) and SO₄²⁻, P, Ca²⁺, Mg²⁺, K⁺ and Na⁺ extracted by Mehlich III (Mehlich, 1984). The determination of SO₄²⁻, was performed using the turbidimetric method according to Bartlett and Neller (1960). The soil chemical attributes which had values under or above one standard deviation (SD) when compared to the overall data average were considered outliers and removed from the statistical analysis.

The final database was composed by 8,428 soil chemical reports. The mean values and other descriptive statistics of chemical properties are shown in Figure 3.

2.3 | Statistical analysis

After the removal of the outliers, data from all soil chemical properties were separated by the respective years and submitted to the Shapiro–Wilk normality and Bartlett tests for variance homogeneity using the XLSTAT 2015 statistical package (Addinsoft, 2019). The P levels data did not show normal distribution and were transformed using the square root function. The data were then submitted to analysis of variance (ANOVA) and regression. Models were chosen based

TABLE 1 Description of the farms that provided samples which have composed the soil chemical analysis database

Farm	County	Number of samples	Area (ha)	Period (years)	Soil series ^a	Additional information
AD	Williams	900	809	2003–2013	Blount Loam, Glywood Loam, Pewamo silty clay Loam, Mermill Loam, Fulton Loam, Haskins Loam	Long-term no-till system, cover crops and wheat-soybean-wheat succession
BW	Williams	2,010	2,225	2000–2013	Blount Loam, Glywood Loam, Pewamo silty clay Loam, Haskins Loam	Milk production, maize (silage), soybean, alfalfa and wheat. Cattle manure applied to the soil every 3 years
KH	Defiance, Paulding	1,393	1,012	2002–2013	Latty silty clay, Hoytville silty clay Loam, Fulton Loam, Nappanee silty clay Loam	Long-term no-till system and cultivation of wheat-soybean succession.
MP	Williams	620	809	2002–2013	Blount Loam, Glywood Loam, Pewamo silty clay Loam	Long-term no-till system and cultivation of wheat-soybean succession.
RC	Williams	236	303	2005–2013	Blount Loam, Glywood Loam, Pewamo silty clay Loam	Soil tillage before corn seeding.
RF	Defiance	2,746	1,618	2000–2014	Rensselaer Loam, Martinsville Loam, Whitaker silt Loam, Blount Loam, Hoytville silty clay Loam	Long-term no-till system and cultivation of wheat-soybean succession.
SM	Defiance, Williams	1,015	809	2003–2013	Blount Loam, Glywood Loam, Pewamo silty clay Loam, Mermill Loam, Fulton Loam, Haskins Loam, Hoytville SCL, Nappanee SCL, Kibbie Loam, Colwood Loam	No-till or reduced soil tilling. Cultivation of maize and soybean in succession.

^aSoil series classified according to Soil Survey Staff (1999).




FIGURE 3 Descriptive statistics of the variables in the soil (0.0–0.20 m depth soil layer) data set. The central rectangle of the boxplots spans the first to the third input quartile. The thin line inside the rectangle is the median, the bold line is the mean, and the horizontal lines to the left and right of the rectangle extend to the minimum and maximum values, respectively. The solid circles represent the minimum and maximum outliers. † Cation exchange capacity; ‡ soil organic matter; § inorganic N levels ($\text{NO}_3^- + \text{NH}_4^+$)

upon statistically significant ($p < .05$) values and the highest coefficients of determination (R^2). The correlation between independent variables was analyzed by the Pearson linear correlation ($p < .05$).

3 | RESULTS AND DISCUSSION

Except for SO_4^{2-} , the database showed no significant differences in soil chemical properties from NW Ohio for the years of 2002–2014 ($p < .05$). Concentrations of SO_4^{2-} , however, significantly declined from 27.3 ± 6.1 to $10.0 \pm 1.7 \text{ mg kg}^{-1}$ (Figure 4). The concentrations were constant between 2002 and 2006 (average of 26.6 mg kg^{-1}), but then sharply decreased up to 2014, with an average concentration that was 52% lower than values observed in the first 4 years (2002–2006).

Soil SO_4^{2-} levels observed in 2014 (average of $10.1 \pm 1.7 \text{ mg kg}^{-1}$) were in the range of concentrations (3.8 to 8.4 mg kg^{-1}) considered deficient for the main cereal crops (Blair, Chinoim, Lefroy, Anderson, & Crocker, 1991; Chen et al., 2008; Horneck,




FIGURE 4 Average sulfate (SO_4^{2-}) levels in topsoils (0–20 cm) from NW Ohio for the years of 2002–2014. Bars indicate twice the standard deviation from the mean for each year. Statistical significance at $p < .01$ is denoted by two asterisks (**)

Sullivan, Owen, & Hart, 2011). If this trend is maintained, soil SO_4^{2-} concentrations will reach values considered restrictive for plant growth. Kost et al. (2008) evaluated 1,473 soil samples representing 443 of the 475 soil series in Ohio, and concluded that for a crop requiring 15 kg S ha^{-1} , most soils (62.5%) were classified as variably deficient, indicating the existence of potential for crop's response to S supply. Camberato and Casteel (2017) summarized soil tests from northern and southern Indiana and concluded that the percentage of samples with soil SO_4^{2-} levels lower than 8 mg kg^{-1} has increased from less than 5% to about 70% in the period of 2005–2017.

Reduced availability of S has also the potential to compromise the uptake and assimilation of N by plants, given that S is a fundamental component of essential amino acids (Hawkesford & Kok, 2006; Salvagiotti & Miralles, 2008). On average, for each kg of crop's S shortfall, 15 kg of N cannot be taken up by plants and, therefore, is subject to loss by leaching and/or volatilization (Haneklaus et al., 2008). Chen et al. (2008) verified interactions between N and S for the nutrition and yield of corn in Ohio soils and concluded that S addition increased yields even at the lower rates of N fertilizer, suggesting that N fertilizer use can be more efficiently utilized when combined with S sources. This can improve profitability in addition to reducing water contamination by sulfates and nitrates from fertilizers (Bindraban, Dimkpa, Nagarajan, Roy, & Rabbinge, 2015; Divito, Echeverria, Andrade, & Sadras, 2015).

The occurrence of soil SO_4^{2-} fluctuations throughout the year is strongly dependent on variations in soil texture, the balance between atmospheric inputs, fertilizer addition, leaching, plant uptake, and microbial activity (Eriksen, 2009). Considering the organic pool, it is expected that soil levels of SO_4^{2-} will be lower over winter due to low mineralization rates associated with reduced soil temperature, moisture and microbial activity (Edwards, 1998). In NW Ohio, farmers usually grow a single crop each year (generally corn and soybean in rotation). Soil testing is generally performed in late autumn or early

spring and represents a one-time picture of nutrient availability that is then used to make decisions regarding fertilization practices. As soils warm in the next spring and summer, more S is mineralized. This increases its availability for uptake and may mitigate some of the expected yield limitations associated with S nutrition.

Considering the agricultural soils evaluated, all under aerobic conditions, SOM and pH changes would be expected to have a key role in controlling soil SO_4^{2-} levels and its availability to crops (Lucheta & Lambais, 2012). The studied areas did not have significant changes in SOM levels between 2002 and 2014 (Figure 5), with an average value of 3.1%, and minimum and maximum values of 1.9 and 4.3% (Figure 3), respectively. The adoption of a long-term no-till system in all the evaluated farms (Table 1), with practices as growing cover crops, maintenance of crops straw on soil surface, and the absence of plowing kept SOM levels stable over the years (Blanco-Canqui et al., 2015; Kibet et al., 2016). Any practices that change SOM can affect SO_4^{2-} levels (Lu et al., 2016), since more than 95% of soil S is in the organic pool, and the mineralization process, which changes reduced S forms into SO_4^{2-} by oxidation, depends on the chemical (pH) and microbiological interaction with SOM (Dick et al., 2008; Eriksen, 2009).

The studied areas also did not have significant changes in pH values between 2002–2014 (Figure 5), with an average value of 6.56, and minimum and maximum values of 5.8 and 7.5 (Figure 3), respectively. Soil SO_4^{2-} adsorption has an inversely proportional relationship with pH (Fuentes-Lara et al., 2019), reaching its maximum at pH 3.0 and minimum at pH 6.5 (Scherer, 2009). Soils with slight acidity (i.e., pH values close to neutral), characteristic of surface layers (0–20 cm) in many agricultural soils including most of the soils in this study, have a predominance of negative charges (CEC), favoring the adsorption of cations instead of anions. However, soils with strongly acid conditions, more commonly found in subsurface profile layers, favor the retention of SO_4^{2-} by its adsorption on Fe and Al oxides as well on the edges of clay particles (Tabatai, 1987).




FIGURE 5 Average pH values and soil organic matter (SOM) levels in topsoils (0–20 cm) from NW Ohio for the years of 2002–2014. Bars indicate the standard deviation from the mean for each year

Given the pedoclimatic characteristics of the studied areas (Figures 2 and 3), neither reduced S-compounds nor S-minerals should have relevance in affecting soil SO_4^{2-} levels. Sulphur inputs by weathering of parent material is difficult to distinguish from other sources, such as mineralization, and do not provide more than 1 kg of S $\text{ha}^{-1} \text{yr}^{-1}$, mainly because of its constant and slow release process (Haneklaus, Bloem, & Schnug, 2000). Reduced sulfur compounds as sulfides (S^{2-}), elemental sulfur (S^0), and sulfites (SO_3^{2-}) are found, in small amounts, mostly in strongly acid and/or in reduced soils (Fuentes-Lara et al., 2019). Other sulfur minerals, as Ca and Mg sulfates, are significant for incipient soils and/or drier regions of the world, since in long-term agricultural soils or humid areas these minerals are leached by rainfall and rarely found (Dick et al., 2008).

The effects of anthropogenic SO_2 emissions results in increased S deposition that can cause acid rain and concurrent acidification of terrestrial and aquatic ecosystems worldwide (Lehmann, 2008). Despite these negative environmental effects, atmospheric S depositions also has been having an essential role in balancing soil S levels over the years. However, emissions of gases containing S, either from combustion of coal or other fossil fuels, systematically decreased in the United States since 1950. This change was especially evident after the 1990s decade when the US Environmental Protection Agency (U.S. EPA) started to adopt more rigid protocols to control the emission of greenhouse gases (United States Environmental Protection Agency, 2013).

The SO_2 emissions by coal-burning from the Ohio electric power plants increased 46% between 1990 and 2001 whereas, in the period of 2002–2012, the emissions decreased by 70% (Figure 6; United States Energy Information Administration, 2014). Similarly, to what was observed for soil SO_4^{2-} levels (Figure 4), the emissions of SO_2 were higher between 2002 and 2007, and then were significantly reduced after 2008, showing a linear decrease trend up to 2013 with averages 48% lower compared to the 2002–2007 period (Figure 6).




FIGURE 6 SO_2 emissions by coal burning for the production of electric power in NW Ohio between 2002 and 2013. Statistical significance ($p < .01$) is denoted with two asterisks (**). Data from United States Energy Information Administration (2015)

As a direct consequence of the reduction in SO_2 emissions, both wet depositions and S concentrations in rainwaters have been decreasing (Figure 7a,b). The major reductions in S wet depositions has been recorded in the States of Maryland, New York, West Virginia, Virginia, Pennsylvania and the region of the Ohio River Valley (United States Environmental Protection Agency, 1998). Data from National Atmospheric Deposition Program (2014) indicate decreases of 66% in the concentration of SO_4^{2-} in rainwaters (Figure 7a) and 52% in the amount of SO_4^{2-} deposited in NW Ohio between the years of 2002–2013 (Figure 7b).

Soil SO_4^{2-} levels were positively correlated with both the concentrations of SO_4^{2-} in rainwater ($r = 0.89$, $p < .05$) and the amount of SO_4^{2-} deposited on soils ($r = 0.91$, $p < .05$). However, there was no correlation between these variables and the annual precipitation volume (Figure 2). Once SO_4^{2-} concentration in rainwater decreases,




FIGURE 7 Sulfate (SO_4^{2-}) concentrations in rainwater (a) and depositions (b) in NW Ohio soils between the years of 2002 and 2013. Statistical significance ($p < .01$) is denoted with two asterisks (**). Bars indicate the standard deviation from the mean for each year. Source: Data from the National Atmospheric Deposition Program station (National Atmospheric Deposition Program, 2014), situated in the Crawford County (Lat. 40.55 Long. -82.99), NW Ohio

changes in the absolute amount of precipitation become more relevant in the final account of the SO_4^{2-} deposited on soil. Nevertheless, the variation in the average volume of precipitation in the evaluated period (Figure 2) did not correlate with the reduction of soil SO_4^{2-} levels. Besides the effects of precipitation volume on S depositions, the volume of rainwater that moves through the soil is important due to the potential of SO_4^{2-} leaching (Edwards, 1998; Scherer, 2009).

Further to the rainfall effects, the use of irrigation is another important aspect to considerate in the balance of soil SO_4^{2-} , especially in arid regions and/or for fruit and horticulture growing (United States Geographic Survey, 2018). Depending on the volume of water used to irrigate and the potential evapotranspiration (PE), irrigation can either increase or decrease soil SO_4^{2-} levels (Kivi & Bailey, 2017). If the applied irrigation volume is higher than PE, than there will be SO_4^{2-} leaching potential. Otherwise, if the irrigation volume is lower than PE, a positive balance of SO_4^{2-} will occur (Haneklaus et al., 2000). In the NW Ohio region, and especially in the evaluated farms, the use of irrigation to enhance production of grain crops is almost absent (Figure 2). In the three counties where the soil samples came from, the sum of the total irrigated area was about 356 and 570 ha in 2010 and 2015 years, respectively (United States Geographic Survey, 2018), and were mostly sprinkler and microirrigation system types which are not suitable for grain and forage crops.

Increases in nutrient export, mostly due to higher crop yields and harvest indices (HI), are also directly associated with decreases in soil S concentrations. In the past 50 years, major crops in Ohio such as soybean, maize, and wheat (United States Department of Agriculture, 2014a, 2014b) had average yield increases of 104, 258 and 166%, respectively (Figure 8). In the same period, the HI was increased 50% for soybeans (Koester et al., 2014) resulting in higher S export from soil since more of the total plant's biomass is being directed to harvested reproductive structures, in this case, grains.




FIGURE 8 Average yields of maize, soybean and wheat in NW Ohio between the years of 1950 and 2013. For the utilization of the equations, the x value should be accounted sequentially by considering 1950 = 1 and 2013 = 64. Statistical significance ($p < .01$) is denoted with two asterisks (**). Source: United States Department of Agriculture (2014a, 2014b)

Combining the NW Ohio current average grain yields (Figure 8) with estimates of S removal by crops like soybean (3.25 kg of S Mg⁻¹ of grains; Hitsuda et al., 2008), corn (1.30 kg of S Mg⁻¹ of grains, Lamond, 1997), and wheat (1.50 kg of S Mg⁻¹ of grains, Györi, 2005), it can be concluded that the average S export from soil has been higher than S depositions (Figure 7b). This contributes directly to the reduction of soil S levels over time.




FIGURE 9 Agricultural consumption of the main sources of S fertilizers in the USA between the years of 1990 and 2011. Statistical significance ($p < .01$) is denoted with two asterisks (**). Source: Economic Research Service, United States Department of Agriculture (2013)

Crop rotation also affects soil S concentration. Crops within the *Poaceae* family (wheat and maize) removed much more S from soil than crops within the *Fabaceae* family (soybean). Other crops, like species of *Brassica napus* and its cultivars, also have a high demand and capacity for soil S removal (up to 35 kg of S Mg⁻¹ of grain, Mašauskiene & Mašauskas, 2012) mainly from soil subsurface layers (Franzen & Grant, 2008). Thus, both grain yield and growing crops with higher demand for S uptake and extraction, like oilseed rape crops (*B. napus* L.), can predispose the following crop to more severe S deficiencies (Mašauskiene & Mašauskas, 2012).

Sulfur reductions from wet depositions and increases in nutrient removal by crops create soil S deficits that can be compensated by the use of fertilizers. However, the use of S fertilizers in the USA has kept relatively constant since the beginning of the 1990s decade. The use of gypsum (CaSO₄·2H₂O) and elemental S in 2011 was similar to that observed in 1990 (Figure 9). An exception to this trend is the application of fertilizers containing ammonium sulfate. However, utilization of this source is minor when compared to other options for N fertilization, and it is not considered a major fertilizer source of S (Figure 9).

As observed for S, the emissions of N compounds (i.e., NO, NO₂) from coal-burning plants in the State of Ohio decreased 38% between 1990 and 2001, and 77% between 2002 and 2012 (United States Energy Information Administration, 2014). Consequently, a linear decrease of N concentrations in the rainwaters and wet depositions was also noticed (Table 2). However, in contrast to S, soil N concentrations in the NW Ohio were constant in the evaluated period, keeping the values close to the average of 39.0 g kg⁻¹ (Table 2). The main

TABLE 2 Soil N concentrations and wet depositions, N concentration in rainwaters, and emission of N compounds by electric power plants in the State of Ohio between 2002–2012

Year	N ^a g kg ⁻¹	SD ^b	[N] in rainwaters mg L ⁻¹	N depositions ^c kg ha ⁻¹	NO emissions ^d x10 ⁶ mg yr ⁻¹
2002	40.0	5.33	2.24	20.7	3.47
2003	40.3	5.37	2.17	16.1	3.31
2004	38.9	5.69	2.39	18.4	2.51
2005	39.4	5.29	2.10	21.2	2.38
2006	37.5	5.58	1.69	17.9	2.24
2007	38.9	5.69	1.77	18.2	2.27
2008	38.3	5.40	1.75	17.7	2.22
2009	38.6	6.03	1.43	15.8	1.10
2010	39.8	5.46	1.51	16.8	1.22
2011	38.7	5.19	1.38	10.9	1.21
2012	39.4	5.62	1.40	10.8	0.91
Regression	n.s	—	[N] = 207** - 0.10**·year	N = 1587* - 0.80*·year	NO = 506 - 0.25**·year
R ²	—	—	.86	.60	.91

^aAverage concentration of inorganic N (NO₃⁻ + NH₄⁺) in the soils of NW Ohio, calculated from the database.

^bStandard deviation.

^cDepositions of inorganic N (NO₃⁻ + NH₄⁺) obtained in the National Atmospheric Deposition Program obtained located in Crawford County (Lat. 40.55 Long. -82.99), NW Ohio.

^dEmission of N compounds by the coal burning in the electric power plants in the State of Ohio (United States Energy Information Administration, 2014).

* $p < .05$; ** $p < .01$.

difference between N compared to S is that between 1990 and 2011 the consumption of N fertilizers in North America increased by 47%. More specifically, the consumption of urea increased 67%, reaching 5.52×10^6 tons in 2011 (Economic Research Service, United States Department of Agriculture, 2013).

Differences in temporal trends for both soil S and N emissions in NW Ohio indicate that as soon as the SO_2 and NO emissions began to decrease, a cumulative deficit began for both elements. In this scenario, the use of fertilizers has become even more critical. Increases in the application of N fertilizers were an adequate and fast response to this condition, and efficiently balanced soil N concentrations, even considering the increases in the crop's yield and HI.

4 | CONCLUSIONS

In this detailed study of soil SO_4^{2-} levels in NW Ohio, the reduction of 70% in SO_2 emissions and 52% in SO_4^{2-} deposition, combined with increasing crop yields and insufficient compensation by fertilization, has led to a decreasing of 63% in soil SO_4^{2-} concentrations between 2002–2014. With this trend established, it is predicted that S soil concentrations will increasingly fall below critical levels needed to support optimum crops yields.

To overcome the S deficiencies in soil, several management options may be adopted including (i) adopting practices to increase soil organic matter levels and subsequent rates of S mineralization, and (ii) replace and replenish the S in soil lost by crop removal using S sources like organic and inorganic fertilizers or various types of industrial by-products.

ACKNOWLEDGMENTS

The authors thank Nester Ag Consultants (Bryan, OH) for providing the soil database. This research was also supported by state and federal funds appropriated to The Ohio State University and The Ohio Agricultural Research and Development Center, Wooster, OH, USA.

CONFLICT OF INTEREST

The authors declare no potential conflict of interest.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author (leandromichalovicz@gmail.com) upon reasonable request.

ORCID

Leandro Michalovicz  <https://orcid.org/0000-0003-1692-4024>

REFERENCES

- Aas, W., Mortier, A., Bowersox, V., Cherian, R., Faluvegi, G., Fagerli, H., ... Xu, X. (2019). Global and regional trends of atmospheric sulfur. *Scientific Reports*, 9, 953. <https://doi.org/10.1038/s41598-018-37304-0>
- Addinsoft. (2019). XLStat v. 2018.1: *Data Analysis and statistics software for Microsoft Excel*. Retrieved from <http://www.xlstat.com/en/download.html>
- Bartlett, F. D., & Neller, J. R. (1960). Turbidimetric determination of sulphate in soil extracts. *Soil Science*, 90(3), 201–204.
- Bindraban Prem S., Dimkpa Christian, Nagarajan Latha, Roy Amit, Rabbinge Rudy (2015). Revisiting fertilisers and fertilisation strategies for improved nutrient uptake by plants. *Biology and Fertility of Soils*, 51, (8), 897–911. <http://dx.doi.org/10.1007/s00374-015-1039-7>.
- Blair, G. J., Chinoim, N., Lefroy, R. D. B., Anderson, G. C., & Crocker, G. J. (1991). A soil sulfur test for pastures and crops. *Australian Journal of Soil Research*, 29, 619–626. <https://doi.org/10.1071/SR9910619>
- Blanco-Canqui, H., Shaver, T. M., Lindquist, J. L., Shapiro, C. A., Elmore, R. W., Francis, C. A., & Hergert, G. W. (2015). Cover crops and ecosystem services: Insights from studies in temperate soils. *Agronomy Journal*, 107(6), 2449–2474. <https://doi.org/10.2134/agronj15.0086>
- Broch, D. L., Pavinato, P. S., Possenti, J. C., Martin, T. N., & Del Quiqui, E. M. (2011). Soybean grain yield in cerrado region influenced by sulphur sources. *Revista Ciência Agrônômica*, 42, 791–796. <https://doi.org/10.1590/S1806-66902011000300027>
- Camberato, J., & Casteel, S. (2017). Sulfur deficiency. In *Soil Fertility Update* (Vol. 1, pp. 1–6). West Lafayette, IN: Purdue University. Retrieved from. <https://ag.purdue.edu/agry/Documents/Sulfur%20deficiency%202017.pdf>
- Chen, L., Dick, W. A., & Kost, D. (2008). Flue gas desulfurization products as sulfur sources for corn. *Soil Science Society of America Journal*, 72, 1464–1470. <https://doi.org/10.2136/sssaj2007.0221>
- Chen, L., Dick, W. A., & Nelson, S. (2005). Flue gas desulfurization products as sulfur sources for alfalfa and soybean. *Agronomy Journal*, 97, 265–271. <https://doi.org/10.2134/agronj2005.0265>
- Dahnke, W. C. (1990). Testing soils for available nitrogen. In R. L. Westerman (Ed.), *Soil testing and plant analysis Soil Science Society of America Book Series* (Vol. 3, pp. 120–140). Madison: ASA. <https://doi.org/10.2136/sssabookser3.3ed.c6>
- Dick, W. A., Kost, D., & Chen, L. (2008). Availability of sulfur to crops from soil and other sources. In J. Jez (Ed.), *Sulfur: A missing link between soils, crops and nutrition* (pp. 59–82). Madison: ASA. <https://doi.org/10.2134/agronmonogr50.c5>
- Divito, G. A., Echeverría, H. E., Andrade, F. H., & Sadras, V. O. (2015). Diagnosis of S deficiency in soybean crops: Performance of S and N:S determinations in leaf, shoot and seed. *Field Crops Research*, 180, 167–175. <https://doi.org/10.1016/j.fcr.2015.06.006>
- Economic Research Service, United States Department of Agriculture. (2013). Fertilizer use and price: U.S. consumption of selected secondary, micronutrients, and natural organic materials, 1986–2011. Retrieved from <http://https://www.ers.usda.gov/webdocs/DataFiles/50341/fertilizeruse.xls?v=7832.7>
- Edwards, P. J. (1998). *Sulfur cycling, retention, and mobility in soils: A review (Gen. Tech. Rep. NE-250)* (p. 18). Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Research Station.
- Eriksen, J. (2009). *Soil sulfur cycling in temperate agricultural systems*. In D. L. Sparks (Ed.), *Advances in Agronomy* (pp. 55–89). London: Academic Press. [https://doi.org/10.1016/S0065-2113\(09\)01002-5](https://doi.org/10.1016/S0065-2113(09)01002-5)
- Food and Agriculture Organization of the United Nations. (2014). World cereals, total Yield 1961–2014. Retrieved from <http://faostat3.fao.org/compare/E>
- Franzen, D., & Grant, C. A. (2008). Sulfur response based on crop, source, and landscape position. In J. Jez (Ed.), *Sulfur: A missing link between soils, crops and nutrition* (pp. 105–116). Madison: ASA. <https://doi.org/10.2134/agronmonogr50.c7>
- Fuentes-Lara, L. O., Medrano-Macias, J., Pérez-Labrada, F., Rivas-Martínez, E. N., García-Enciso, E. L., González-Morales, S., ... Benavides-Mendoza, A. (2019). From elemental sulfur to hydrogen sulfide in agricultural soils and plants. *Molecules*, 24(12), 2282. <https://doi.org/10.3390/molecules24122282>
- Gautam, K., Kumar, S., & Lokhandwala, S. (2019). Energy-aware intelligence in megacities. In *Current Developments in Biotechnology and*

- Bioengineering* (pp. 211–238). Amsterdam: Elsevier. <https://doi.org/10.1016/B978-0-444-64083-3.00011-7>
- Györi, Z. (2005). Sulphur content of winter wheat grain in long term field experiments. *Communications in Soil Science and Plant Analysis*, 36 (1–3), 373–382. <https://doi.org/10.1081/CSS-200043098>
- Haneklaus, S., Bloem, E., & Schnug, E. (2000). Sulphur in Agroecosystems. *Folia Universitatis Agriculturae Stetinensis*, 81, 17–23. <https://doi.org/10.2134/agronmonogr50.c4>
- Haneklaus, S., Bloem, E., & Schnug, E. (2008). History of sulfur deficiency in crops. In J. JEZ (Ed.), *Sulfur: A missing link between soils, crops and nutrition* (pp. 45–58). Madison: ASA. <https://doi.org/10.2134/agronmonogr50.c4>
- Hawkesford, M. J., & Kok, L. J. (2006). Managing sulphur metabolism in plants. *Plant, Cell and Environment*, 29, 382–395. <https://doi.org/10.1111/j.1365-3040.2005.01470.x>
- Hitsuda, K., Toriyama, K., Subbarao, G.V., & Ito, O. (2008). Sulfur Management for Soybean Production. Joseph Jez *Sulfur: A missing link between soils, crops, and nutrition*, (117–142). Madison, WI: ASA, CSSA and SSSA. <https://doi.org/10.2134/agronmonogr50.c8>
- Horneck, D. A., Sullivan, D. M., Owen, J. S., & Hart, J. M. (2011). *Soil test interpretation guide* (p. 12). Corvallis, OR: Oregon State University Extension Service Publications and Multimedia.
- Khan, T. A., & Mazid, M. (2011). Nutritional significance of sulphur in pulse cropping system. *Biology and Medicine*, 3(2), 114–133. https://biomedonline.com/Articles/MAASCON-1/Vol3_2_114-133.pdf
- Kibet, L. C., Blanco-Canqui, H., & Jasa, P. (2016). Long-term tillage impacts on soil organic matter components and related properties on a Typic Argiudoll. *Soil and Tillage Research*, 155, 78–84. <https://doi.org/10.1016/j.still.2015.05.006>
- Kivi, S. R., & Bailey, R. T. (2017). Modeling sulfur cycling and sulfate reactive transport in an agricultural groundwater system. *Agricultural Water Management*, 185, 78–92. <https://doi.org/10.1016/j.agwat.2017.02.002>
- Koester, R. P., Skoneczka, J. A., Cary, T. R., Diers, B. W., & Ainsworth, E. A. (2014). Historical gains in soybean (*Glycine max* Merr.) seed yield are driven by linear increases in light interception, energy conversion, and partitioning efficiencies. *Journal of Experimental Botany*, 65, 3311–3321. <https://doi.org/10.1093/jxb/eru187>
- Köppen, W., & Geiger, R. (1928). *Klimate der Erde*. Gotha: Verlag Justus Perthes.
- Kost, D., Chen, L., & Dick, W. A. (2008). Predicting plant sulfur deficiency in soils: Results from Ohio. *Biology and Fertility of Soils*, 44, 1091–1098. <https://doi.org/10.1007/s00374-008-0298-y>
- Kovar, J. L., & Grant, C. A. (2011). *Nutrient cycling in soils: Sulfur* (Paper 1383). Lincoln, NE: USDA-ARS/UNL Faculty. <https://digitalcommons.unl.edu/usdaarsfacpub/1383>
- Lamond, R. E. (1997). *Sulphur in Kansas: Plant, soil and fertilizer considerations* (p. 4). Manhattan, KS: Kansas State University Agricultural Experiment Station and Cooperative Extension Service. <https://bookstore.ksre.ksu.edu/pubs/MF2264.pdf>
- Lehmann, J. (2008). Atmospheric SO emissions since the late 1800 change organic sulfur forms in humic substance extracts of soils. *Environmental Science & Technology*, 42, 3550–3555. <https://doi.org/10.1021/es702315g>
- Li, N., Yang, Y., Wang, L., Zhou, C., Jing, J., Sun, X., & Tian, X. (2019). Combined effects of nitrogen and sulfur fertilization on maize growth, physiological traits, N and S uptake, and their diagnosis. *Field Crops Research*, 242, 107593. <https://doi.org/10.1016/j.fcr.2019.107593>
- Lu, Q., Bai, J., Fang, H., Wang, J., Zhao, Q., & Jia, J. (2016). Spatial and seasonal distributions of soil sulfur in two marsh wetlands with different flooding frequencies of the Yellow River Delta, China. *Ecological Engineering*, 96, 63–71. <https://doi.org/10.1016/j.ecoleng.2015.10.033>
- Lucheta, A. R., & Lambais, M. R. (2012). Sulfur in Agriculture. *Revista Brasileira de Ciência do Solo*, 36, 1369–1379. <https://doi.org/10.1590/S0100-06832012000500001>
- Mašauskiene, A., & Mašauskas, V. (2012). Soil Sulphur problems and management. *Christine Jakobsson Sustainable Agriculture* (p. 113–116). Uppsala, SE: Baltic University Press, Uppsala University. <http://uu.diva-portal.org/smash/get/diva2:603516/FULLTEXT01>
- McLean, E. O. (1982). *Soil pH and lime requirement*. In Page, et al. (Eds.), *Methods of soil analysis, Part 2: Chemical and Microbiological Properties. Agronomy Monogr.* 9 (2nd ed., pp. 199–223). Madison, WI: ASA. <https://doi.org/10.2134/agronmonogr9.2.2ed.c12>
- Mehlich A. (1984). Mehlich 3 soil test extractant: A modification of Mehlich 2 extractant. *Communications in Soil Science and Plant Analysis*, 15(12), 1409–1416. <http://dx.doi.org/10.1080/00103628409367568>
- Mikkelsen, R., & Norton, R. (2013). Soil and fertilizer sulfur. *Better Crops*, 97(1), 7–10. [http://www.ipni.net/publication/bettercrops.nsf/0/583E6D994D59373685257B18005ACC3F/\\$FILE/BC-2013-1.pdf](http://www.ipni.net/publication/bettercrops.nsf/0/583E6D994D59373685257B18005ACC3F/$FILE/BC-2013-1.pdf)
- National Atmospheric Deposition Program. (2014). Total sulfur depositions in Ohio. Retrieved from <http://nadp.slh.wisc.edu/datalib/AIRMoN/AIRMoN-ALL.csv>
- National Oceanic and Atmospheric Administration. (2015). Climatological data publications. Retrieved from <http://www1.ncdc.noaa.gov/pub/orders/IPS/IPS-07269548-4AEE-442E-9AE1-4F126310B2B4.pdf>
- Pan, X. H., & Deng, Q. H. (2007). Review on crop harvest index. *Acta Agriculturae Universitatis Jiangxiensis*, 29, 1–5.
- Pias, O. H. d. C., Tiecher, T., Cherubin, M. R., Mazurana, M.I., & Bayer, C. (2019). Crop yield responses to sulfur fertilization in Brazilian no-till soils: A systematic review. *Revista Brasileira de Ciência do Solo*, 43, 1–21. <https://doi.org/10.1590/18069657rbc20180078>
- Raij, B., van. (2008). *Gesso na Agricultura*, Campinas, BR: IAC.
- Rehm, G. W. (2005). Sulfur management for corn grown with conservation tillage. *Soil Science Society of American Journal*, 69, 709–717. <https://doi.org/10.2136/sssaj2004.0151>
- Salvagiotti, F., & Miralles, D. J. (2008). Radiation interception, biomass production and grain yield as affected by the interaction of nitrogen and sulfur fertilization in wheat. *European Journal of Agronomy*, 28, 282–290. <https://doi.org/10.1016/j.eja.2007.08.002>
- Scherer, H. W. (2009). Sulfur in soils. *Journal of Plant Nutrition and Soil Science*, 172, 326–335. <https://doi.org/10.1002/jpln.200900037>
- Schulte, E. E., & Hopkins, B. G. (1996). Estimation of soil organic matter by weight loss-on-ignition. In F. R. Magdoff, M. A. Tabatabai, & E. A. Hanlon, Jr. (Eds.), *Soil organic matter: Analysis and interpretation (Special Publication No. 46)* (pp. 21–32). Madison: ASA. <https://doi.org/10.2136/sssaspecpub46.c3>
- Shoemaker H. E., McLean E. O., Pratt P. F. (1961). Buffer Methods for Determining Lime Requirement of Soils With Appreciable Amounts of Extractable Aluminum. *Soil Science Society of America Journal*, 25(4), 274–277. <http://dx.doi.org/10.2136/sssaj1961.03615995002500040014x>
- Singh A. K., Manibhushan, Meena M. K., Upadhyaya A. (2012). Effect of Sulphur and Zinc on Rice Performance and Nutrient Dynamics in Plants and Soil of Indo Gangetic Plains. *Journal of Agricultural Science*, 4(11). <http://dx.doi.org/10.5539/jas.v4n11p162>
- Sloan, J. J., Dowdy, R. H., Dolan, M. S., & Rehm, G. W. (1999). Plant and soil responses to field-applied flue gas desulfurization residue. *Fuel*, 78, 169–174. <https://doi.org/10.1016/S0016-2361>
- Soil Survey Staff. (1999). *Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys* (2nd ed. ed. USDA–NRCS Agric. Handbook 436). Washington, DC: U.S. Government Printing Office.
- Sokolova, T. A., & Alekseeva, S. A. (2008). Adsorption of sulfate ions by soils (a review). *Eurasian Soil Science*, 41, 140–148. <https://doi.org/10.1134/S106422930802004X>
- State of Ohio. (2010). Census 2010. Retrieved from <https://www.census.gov/prod/cen2010/cph-1-37.pdf>
- Tabatabai M. Ali (1987). Physicochemical fate of sulfate in soils. *JAPCA*, 37(1), 34–38. <http://dx.doi.org/10.1080/08940630.1987.10466197>
- Tiecher, T., Santos, D. R., Rasche, J. W. A., Brunetto, G., Mallmann, F. J. K., & Piccin, R. (2012). Crop responses and sulfur

- availability in soils with different contents of clay and organic matter submitted to sulfate fertilization. *Bragantia*, 71, 518–527. <https://doi.org/10.1590/S0006-87052013005000010>.
- Tisdale, S. L., Reneau, R. B., & Platou, J. S. (1986). Atlas of sulfur deficiencies. In M. A. Tabatabai (Ed.), *Sulfur in agriculture* (pp. 295–322). Madison: ASA. <https://doi.org/10.2134/agronmonogr27.c10>.
- Tripathi R. D., Tripathi P., Dwivedi S., Kumar A., Mishra A., Chauhan P. S., Norton G. J., Nautiyal C. S. (2014). Roles for root iron plaque in sequestration and uptake of heavy metals and metalloids in aquatic and wetland plants. *Metallomics*, 6(10), 1789–1800. <http://dx.doi.org/10.1039/c4mt00111g>.
- United States Department of Agriculture. (2014a). *National agricultural statistics service (NASS)*. Retrieved from <https://quickstats.nass.usda.gov/>
- United States Department of Agriculture. (2014b). State agriculture overview, Ohio. Retrieved from https://www.nass.usda.gov/Quick_Stats/Ag_Overview/stateOverview.php?state=OHIO
- United States Energy Information Administration. (2014). *Electric power industry emissions estimates, 1990 through 2014, Ohio*. Retrieved from: <http://www.eia.gov/electricity/state/ohio/xls/sept07OH.xls>
- United States Environmental Protection Agency (1998). Atmospheric deposition of sulfur and nitrogen compounds. In *National air quality and emissions trends report*, Washington, D.C.: U.S. EPA. Retrieved from. https://www.epa.gov/sites/production/files/2017-11/documents/trends_report_1998.pdf
- United States Environmental Protection Agency. (2020). History of the Clean Air Act. Retrieved from <https://www.epa.gov/clean-air-act-overview>
- United States Environmental Protection Agency. (2013). Global Mitigation of Non-CO₂ Greenhouse Gases: 2010-2030. *Office of Atmospheric Programs, Washington, DC.*–410p. https://www.epa.gov/sites/production/files/2016-06/documents/mac_report_2013.pdf.
- United States Geographic Survey. (2018). National water information system: Water use data for Ohio. Retrieved from https://waterdata.usgs.gov/oh/nwis/water_use?format=html_table&rdb_compression=file&wu_area=County&wu_year=1985%2C1990%2C1995%2C2000%2C2005%2C2010%2C2015&wu_county=033%2C039%2C125%2C171&wu_category=ALL%2CIT%2CIC%2CIG&wu_county_nms=Crawford%2BCounty%252CDefiance%2BCounty%252CPaulding%2BCounty%252CWilliams%2BCounty&wu_category_nms=--ALL%2BCategories--%252CIrrigation%252C%2BTotals%252CIrrigation%252C%2BCrop%252CIrrigation%252C%2BGolf%2BCourses
- Vieira-Filho, M. S., Lehmann, C., & Fornaro, A. (2015). Influence of local sources and topography on air quality and rainwater composition in Cubatão and São Paulo, Brazil. *Atmospheric Environment*, 101, 200–208. <https://doi.org/10.1016/j.atmosenv.2014.11.025>

How to cite this article: Michalovicz L, Dick WA, Tormena CA, Müller MML, Cervi EC. Temporal trends of sulfur levels in soils of northwest Ohio (USA) between 2002 and 2014. *Land Degrad Dev*. 2021;32:573–582. <https://doi.org/10.1002/ldr.3745>