



# Interpersonal Trauma Exposure and Interpersonal Problems in Adolescent Posttraumatic Stress Disorder

Kathryn A. Hughesdon <sup>1</sup>, Julian D. Ford <sup>2</sup>, Ernestine C. Briggs,<sup>3</sup> Julia S. Seng,<sup>4</sup> Alison L. Miller,<sup>5</sup> and Sarah A. Stoddard<sup>4</sup>

<sup>1</sup>School of Nursing, Eastern Michigan University, Ypsilanti, Michigan, USA

<sup>2</sup>Department of Psychology, University of Connecticut, Farmington, Connecticut, USA

<sup>3</sup>School of Medicine, Duke University, Durham, North Carolina, USA

<sup>4</sup>School of Nursing, University of Michigan, Ann Arbor, Michigan, USA

<sup>5</sup>School of Public Health, University of Michigan, Ann Arbor, Michigan, USA

Traumatic experiences have been differentiated as interpersonal (i.e., the direct result of actions by other people) or noninterpersonal (i.e., other life-threatening events, such as severe accidents). Interpersonal trauma exposure generally has been shown to be associated with more severe posttraumatic stress disorder (PTSD) symptoms than noninterpersonal trauma exposure. Interpersonal problems also tend to be associated with trauma exposure and PTSD symptoms, but it is unclear whether a mediating association exists between trauma type, interpersonal problems, and PTSD symptoms. A clinical sample of 4,275 adolescents (age range: 12–18 years) from the National Child Traumatic Stress Network Core Data Set were classified as having experienced interpersonal trauma, noninterpersonal trauma, or both. Interpersonal problems were operationalized by social problem behaviors (e.g., immature and dependent behaviors) and aggressive behaviors on the Child Behavior Checklist. The results of path analyses showed that cumulative interpersonal trauma exposure was both directly and indirectly associated with PTSD symptoms via social problem behaviors but not aggressive behaviors, total effect  $\beta = .20$ , 95% CI [.17, .23]. In a second model, path analyses showed that cumulative interpersonal trauma exposure was associated directly and indirectly via PTSD symptoms with social problem behaviors, total effect  $\beta = .15$ , 95% CI [.11, .18], and aggressive behaviors, total effect  $\beta = .13$ , 95% CI [.09, .17]. These findings suggest that during adolescence, interpersonal problems play an important role in the association between interpersonal trauma exposure and PTSD symptoms.

Traumatic experiences during childhood have been associated with negative mental health outcomes throughout the lifespan (Dube et al., 2006). Such experiences have been differentiated as interpersonal (i.e., the direct result of actions by other people) and noninterpersonal (i.e., other life-threatening events,

such as severe accidents, disasters, or illness; Charak et al., 2018; Dunn et al., 2017; Haldane & Nickerson, 2016; Jaffe et al., 2019; Woodward et al., 2015). The results of previous research suggest that experiencing interpersonal traumatic events may lead to an increased risk of developing posttraumatic stress disorder (PTSD) and complex PTSD (CPTSD; i.e., symptoms of PTSD plus symptoms of emotional, cognitive, and interpersonal difficulties; World Health Organization, 2020) compared to experiencing noninterpersonal traumatic events (Alisic et al., 2014; Brewin et al., 2017; Charuvastra & Cloitre, 2008; Frost et al., 2020; Kerig et al., 2009). In a meta-analysis that aimed to determine the incidence of PTSD in trauma-exposed children and adolescents who were 2–18 years old, Alisic and colleagues (2014) found that approximately one in four youth developed PTSD after experiencing an interpersonal traumatic event, whereas approximately one in 10 youth developed PTSD after experiencing a noninterpersonal traumatic event. Similarly, in a sample of adolescents involved in the juvenile justice system, Kerig and colleagues (2009) found that among both male and female youth, more severe PTSD symptoms were associated with the number of past interpersonal trauma types reported but not associated with noninterpersonal trauma types.

Kathryn Hughesdon acknowledges predoctoral support from the Rita and Alex Hillman Foundation Hillman Scholars in Nursing Innovation. This project was developed in part under grant number 2U79SM054284 from the Center for Mental Health Services (CMHS), Substance Abuse and Mental Health Services Administration (SAMHSA), U.S. Department of Health and Human Services (HHS). The views, policies, and opinions expressed are those of the authors and do not necessarily reflect those of the funding sources listed above.

The authors would like to thank Robert C. Lee and Alison Cooke for their statistical contributions to this project, as well as the 56 sites within the National Child Traumatic Stress Network that have contributed data to the Core Data Set and the children and families who have contributed to our growing understanding of child traumatic stress.

Correspondence concerning this article should be addressed to Kathryn Hughesdon, School of Nursing, Eastern Michigan University, 318F Porter, Ypsilanti, MI, 48197, USA. Email: kabramos@emich.edu

© 2021 International Society for Traumatic Stress Studies. View this article online at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com/doi/10.1002/jts.22687)  
DOI: 10.1002/jts.22687

The timing of interpersonal and noninterpersonal trauma exposure may play a role in the differential risk of adverse sequelae of the two types of traumatic experiences. In a study of low-income, African American adults, Dunn et al. (2017) found that exposure to maltreatment in early childhood, considered to be an interpersonal trauma, was associated with more severe depression and PTSD symptoms than exposure at later developmental periods, after controlling for exposure to noninterpersonal or other types of interpersonal trauma concurrently. Moreover, exposure to other types of interpersonal trauma (e.g., witnessing an attack) in middle childhood was shown to be associated with ratings of depressive symptoms that were twice as high as ratings for the same symptoms in adults who first experienced interpersonal trauma in adulthood, again after controlling for exposure to noninterpersonal trauma (Dunn et al., 2017). These findings support the assertion that exposure to interpersonal trauma in childhood may have a particularly adverse impact on PTSD symptoms.

A potential factor that could play a role in the association between trauma type and PTSD symptoms is interpersonal problems, or problematic interpersonal relationships and difficulties relating to others that create a negative impact on one's ability to form healthy and rewarding relationships, such as being too aggressive or having trouble with affection or sympathy (Horowitz et al., 1993, 2000). Findings from previous studies suggest that disruptions in social interactions and relationships after traumatic experiences can impact the development of PTSD (Brewin et al., 2000; Hyman et al., 2003; Trickey et al., 2012), possibly through shaping posttraumatic cognitions (Charuvastra & Cloitre, 2008). In a study of adult survivors of intimate partner violence and victims of motor vehicle accidents, social support from family and friends but not from a close other was associated with lower levels of posttraumatic cognitions and PTSD symptoms among participants in both the former interpersonal trauma cohort and the latter noninterpersonal trauma cohort (Woodward et al., 2015). Sperry and Widom (2013) found that a lack of social support explained the association between childhood trauma exposure and anxiety and depression in adulthood. However, although social support and interpersonal problems may be associated, they are not synonymous, and social support most often has been investigated as a moderator of the association between trauma exposure and PTSD symptoms but not as a potential mediator. In contrast to social support, interpersonal problems that result from or are exacerbated by trauma exposure could be a link between trauma exposure and PTSD symptoms. However, no studies of which we are aware have examined interpersonal problems as a mediator of the association between trauma exposure and PTSD symptoms.

Although findings from previous research provide evidence that interpersonal problems may occur after trauma exposure and before the development of PTSD symptoms, there could also be a bidirectional relation between interpersonal problems and PTSD. For example, in a sample of adolescents who had a history of childhood sexual abuse, higher levels of PTSD

symptom severity were associated with poorer social functioning (McLean et al., 2013). Moreover, PTSD has been associated with problems related to trust, closeness, and peer relationships (Beck et al., 2008; McFarlane & Bookless, 2001). Thus, PTSD could also lead to or exacerbate interpersonal problems following trauma exposure.

In addition, to our knowledge, no studies have focused on whether interpersonal problems are differentially associated with exposure to interpersonal versus noninterpersonal trauma. Although there is evidence that the cumulative effect of exposure to multiple types of trauma, including both interpersonal and noninterpersonal traumatic events, is associated with a wide variety of emotional, behavioral, and cognitive problems in childhood (Hebert et al., 2018; Hodges et al., 2013; Solva et al., 2020), there also is evidence that cumulative exposure to interpersonal trauma may have more debilitating effects than cumulative exposure to noninterpersonal trauma (Baker et al., 2020). Given that interpersonal traumatic events are often stigmatizing and can lead to a sense of betrayal, distrust, and insecurity in relationships (Charuvastra & Cloitre, 2008; Freyd, 1994; Punamaki et al., 2005), exposure to interpersonal trauma may have a greater impact on interpersonal problems than exposure to noninterpersonal trauma. Children who are exposed to multiple types of interpersonal trauma have been described as polyvictims (Finkelhor et al., 2007), and polyvictimization has been shown to be associated with PTSD, depression, and substance abuse symptoms and interpersonal problems among adolescents when compared to less-extensive trauma exposure (Ford, Elhai, et al., 2010).

Examining the association between trauma type, interpersonal problems, and PTSD symptoms in adolescence is important given the relevance of interpersonal relationships during this developmental period. In adolescence, positive relationships with peers can provide a sense of belonging and feelings of closeness, as adolescents become increasingly independent and develop their own identity (Graber et al., 2016). Positive interpersonal relationships in adolescence have been associated with lower depression rates (Lahey & Cronin, 2008) and higher ratings of self-esteem (Cast & Burke, 2002), whereas interpersonal problems in adolescence have been associated with higher levels of stress (Segrin, 2001; Shahar et al., 2004), generalized anxiety disorder (Borkovec et al., 2002; Eng & Heimberg, 2006), and depression (Petty et al., 2004; Vittengl et al., 2003). Thus, interpersonal relationships and experiences may play an important role in mental health outcomes for trauma-exposed adolescents.

The goal of the present study was to address gaps in the research by identifying associations between interpersonal versus noninterpersonal trauma exposure, interpersonal problems, and PTSD symptoms in a clinical sample of trauma-exposed adolescents. We hypothesized that (a) cumulative interpersonal trauma exposure would be associated with PTSD symptoms and with interpersonal problems, (b) interpersonal problems would mediate the association between interpersonal trauma and PTSD symptoms, and (c) noninterpersonal trauma would

not be associated with interpersonal problems. Given the potential bidirectional nature of interpersonal problems and PTSD symptoms, in a separate analysis, we hypothesized that (d) PTSD symptoms would mediate the association between interpersonal trauma and interpersonal problems.

## Method

### Participants and Procedure

The present study included a secondary analysis of data collected as part of a large clinical multisite dataset, the National Child Traumatic Stress Network (NCTSN) Core Data Set (CDS). The dataset consists of clinic-referred children 0–21 years of age ( $N = 14,088$ ) with trauma histories who were seen in a variety of treatment settings, including hospitals, community-based clinics, and outpatient and residential treatment settings throughout the United States. Data were collected from 56 NCTSN sites between 2004 and 2010. For the present analysis, we used baseline data collected before individuals started any trauma treatment. The CDS is part of a quality improvement effort by the NCTSN. Additional details on the CDS are reported in previous studies (Briggs et al., 2012; Greeson et al., 2014; Steinberg et al., 2014). Clinicians were licensed providers or provisionally licensed providers operating under the supervision of a licensed provider. All staff members were trained on the assessment measures and CDS data collection protocols. Deidentified data for the present study were obtained and analyzed at the UCLA–Duke University National Center for Child Traumatic Stress (NCCTS). Institutional Review Board (IRB) approval was obtained from the University of Michigan. For the present study, we restricted the sample to a subset of the NCTSN CDS that included adolescents ages 12–18 years who had experienced at least one traumatic event ( $N = 4,275$ ). This age range was chosen because previous research on adolescence has typically encompassed youth who are 12–18 years old (Jaworska & MacQueen, 2015). In addition, the Child Behavior Checklist, a measure used in the present study, is intended for use in children aged 6–18 years (Achenbach & Rescorla, 2001).

The mean participant was 14.8 years ( $SD = 1.7$ , range: 12–18). The sample was 60.4% female. The distribution of race/ethnicity backgrounds was 34.7% White/Non-Hispanic, 34.5% Hispanic, 24.7% Black/Non-Hispanic, and 6.1% “other”/Non-Hispanic. In addition, 60.0% of the sample had public insurance.

### Measures

#### Childhood Trauma Exposure

Childhood trauma exposure was assessed using the CDS General Trauma Information Form. This form was adapted from the trauma exposure section of the UCLA PTSD Reaction Index (RI; Steinberg et al., 2004). Clinicians use the form to report which of 20 different trauma types a child may have

experienced based on interviews with the child, caregiver, and collateral individuals (e.g., caseworkers; Briggs et al., 2012). For each trauma type, response options include “yes,” “no,” “suspected,” or “unknown.” For the present study, only answers reported as “yes” were analyzed. Trauma exposure types were categorized as interpersonal or noninterpersonal for analyses. Interpersonal traumas included: sexual maltreatment or abuse; sexual assault or rape; physical maltreatment or abuse; physical assault; emotional abuse or psychological maltreatment; neglect; domestic violence; war, terrorism, or political violence inside the United States; war, terrorism, or political violence outside the United States; kidnapping; forced displacement; impaired caregiver; extreme interpersonal violence not reported elsewhere; community violence not reported elsewhere; and school violence not reported elsewhere. This categorization is consistent with previous studies that have compared exposure to interpersonal and noninterpersonal trauma exposure (e.g., Alisic et al., 2014; Kerig et al., 2009; Vibhakar et al., 2019). A summary score was created to reflect the cumulative number of interpersonal trauma types adolescents experienced (range: 1–15).

Noninterpersonal traumas included illness or medical trauma, serious injury accident, natural disaster, and traumatic loss or bereavement. This categorization is consistent with previous studies that have compared exposure to interpersonal versus noninterpersonal traumatic events (e.g., Alisic et al., 2014; Kerig et al., 2009; Vibhakar et al., 2019). A summary score was created to reflect the cumulative number noninterpersonal trauma types adolescents experienced (range: 1–4).

#### Interpersonal Problems

Interpersonal problems in adolescence include problematic interpersonal behaviors and interactions with others (e.g., Johnson et al., 2002). The Social Problem Behaviors and Aggressive Behaviors subscales of the Child Behavior Checklist (CBCL; Achenbach & Rescorla, 2001) were used to operationalize interpersonal problems. On the CBCL, the parent or caregiver who spends the most time with the child rates items, each of which begins with the stem “For each item that describes your child now or within the past 6 months...,” using response options ranging from 0 (*not true*) to 2 (*very true or often true*). For subscale analyses, a continuous  $t$  score was used for each scale to determine clinical significance. Per Achenbach and Rescorla (2001),  $t$  scores less than 67 are considered normal,  $t$  scores of 67–70 are considered borderline clinical, and  $t$  scores higher than 70 are considered to be in the clinical range. The CBCL has been normed, and several studies have documented its robust psychometric properties, including validity and reliability (e.g., Achenbach & Rescorla, 2001; Dang et al., 2017).

The 11-item CBCL Social Problems subscale was used to assess immature, dependent, and socially incompetent behaviors. Sample subscale items include, “Doesn’t get along with other kids,” “Complains of loneliness,” and “Gets teased a lot” (Achenbach & Rescorla, 2001). In the present sample,  $t$  scores

for the Social Problems subscale ranged from 50 to 91, and Cronbach's alpha was .88.

Aggressive behaviors (i.e., aggressive behaviors that create problematic and poor relationships) were measured using the 18-item CBCL Aggressive Behavior subscale. Sample subscale items are, "Argues a lot," "Disobedient at home," "Cruelty, bullying, or meanness to others" (Achenbach & Rescorla, 2001). In the present sample, *t* scores for the Aggressive Behaviors subscale ranged from 50 to 98, and Cronbach's alpha was .93.

### PTSD Symptoms

The UCLA PTSD Reaction Index (RI; Steinberg et al., 2004) was used to assess PTSD symptoms based on the criteria in the fourth edition (text rev.) of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*; American Psychiatric Association [APA], 2000). For the present study, 20 PTSD symptom items and two associated feature items (i.e., trauma-related guilt, fear of trauma recurrence) were used. Responses were collected via clinical interviews and adolescent self-report. The items correlate with the *DSM-IV* intrusion, avoidance, and arousal criteria while also assessing fear of recurrence and trauma-related guilt (Steinberg et al., 2004). Response options range from 0 (*none*) to 4 (*most*). Several studies have demonstrated evidence of the measure's validity and reliability in children and adolescents (e.g., Steinberg et al., 2004, 2013). For the current analyses, PTSD symptoms were treated as a continuous variable, with higher scores indicating higher levels of PTSD symptom severity. Youth frequently do not meet the criteria for a full PTSD diagnosis but still have severe symptoms that can interfere with functioning (Newman, 2002). Thus, research on adolescent PTSD is best served by continuous ratings of PTSD symptoms rather than diagnostic status alone (Kerig et al., 2009). In the present sample, Cronbach's alpha was .95.

### Demographic Characteristics

Sociodemographic characteristics used as control variables included race and ethnicity (Black/Non-Hispanic, White/Non-Hispanic, "other"/Non-Hispanic, and Hispanic), gender (male, female, other, unknown), and public insurance status (yes, no).

### Data Analysis

First, descriptive statistics and bivariate correlations among all study variables were examined using SAS (Version 9.4; SAS Institute Inc., 2015). For each model, the inclusion of sociodemographic characteristics was determined based on theory and support from previous studies (e.g., Greeson et al., 2014; Jones, 2007; Mercado-Crespo & Mbah, 2013; Valois et al., 2002; Winer & Thompson, 2013). Given the limitations of the cross-sectional data and an inability to determine whether the interpersonal problems occurred before PTSD symptom onset, path analyses were used to test two models representing the hypothesized associations. The first model tested the role of interpersonal problems as a mediator between trauma type and PTSD symptoms. The second model was used to examine

PTSD symptoms as a mediator between trauma type and interpersonal problems. Path analyses were completed in *Mplus* (Version 8.0; Muthén & Muthén, 2017). Both models were estimated using full information maximum likelihood estimation with 500 bootstrap samples to provide conservative estimates of 95% confidence intervals that were robust to nonnormality. The reported results were obtained from bias-corrected bootstrapped models. Standardized 95% confidence intervals were evaluated to determine the significance of path coefficients; confidence intervals that did not include 0 were considered statistically significant. Total and specific indirect associations were also evaluated using the same criteria.

Model fit was evaluated using the root mean square error of approximation (RMSEA), comparative fit index (CFI), Tucker-Lewis Index (TLI), and standardized root mean square residual (SRMR). For the RMSEA, values less than .06 were used to indicate good model fit. For the CFI and TLI, values above .95 were used to indicate good model fit. For the SRMR, values less than .08 were used to indicate good model fit (Hu & Bentler, 1999; West et al., 2012). Data were missing in low proportions (less than 10%). Of the full sample, 346 participants were removed due to missingness on the predictor variables ( $n = 119$ ) or missingness on all outcome variables ( $n = 227$ ). No other patterns of missingness were observed.

## Results

### Descriptive Statistics

Descriptive statistics and correlations are depicted in Table 1. Participants experienced a mean of 3.15 ( $SD = 2.31$ ) types of interpersonal traumatic events and 0.90 ( $SD = 0.82$ ) noninterpersonal traumatic events. Female participants had significantly higher levels of PTSD symptoms than their male counterparts,  $\beta_s = .21-.22$ . Youth who were Black,  $\beta = -.08$ ; Hispanic,  $\beta_s = -.05-.06$ ; or reported their race/ethnicity as "other"  $\beta_s = -.04-.05$ , had fewer social problem behaviors than youth who were White/Non-Hispanic. Youth who reported public insurance had higher levels of social problem behaviors,  $\beta_s = .04-.05$ , than those who did not. Female youth had lower rates of aggressive behaviors,  $\beta_s = -.05-.07$ , than male youth. Participants who were Hispanic had lower rates of aggressive behaviors,  $\beta = -.06$ , than youth who were White/Non-Hispanic.

### Model 1

The first path model fit the data well,  $RMSEA = .011$ ,  $CFI = .999$ ,  $TLI = .995$ ,  $SRMR = .007$  (see Figure 1 and Supplementary Table S1). Social problem behaviors and aggressive behaviors were significantly correlated with each other in the model,  $r = .61$ ,  $p < .001$ . Cumulative interpersonal trauma exposure was associated with more severe social problem behaviors, aggressive behaviors, and PTSD symptoms,  $\beta_s = .13-.17$ ,  $p < .001$ . Social problem behavior severity was associated with


**Table 1**  
*Descriptive Statistics and Correlation Table for Variables*

	2	3	4	5	6	7	8	9	10	M	SD	Skewness	Kurtosis	Frequency
1. PTSD symptoms	.13*	.20*	.24*	.07*	.24*	-.02	-.01	.01	.00	25.92	15.03	.31	-.69	
2. Aggressive behavior	—	.62*	.13*	.03	-.03	.00	-.05*	.01	.06*	63.50	10.99	.77	.07	
3. Social problems		—	.17*	.06*	.03	-.07*	-.01	-.01	.06*	61.56	9.15	.58	-.15	
4. Interpersonal trauma			—	.18*	.14*	-.09*	-.05	.07*	.16*	3.15	2.31	.83	.37	
5. Noninterpersonal trauma				—	-.01	-.03	.01	.04*	.08*	0.90	0.82	.82	.54	
6. Female <sup>a</sup>					—	-.01	-.03	.02	.01	—	—	—	—	60.4%
7. Black <sup>a</sup>						—	-.40*	-.14*	.08*	—	—	—	—	24.7%
8. Hispanic <sup>a</sup>							—	-.18*	-.03	—	—	—	—	34.5%
9. "Other" race								—	.00	—	—	—	—	6.1%
10. Public insurance									—	—	—	—	—	60.0%

*Note.* N = 4,275. Mean values and standard deviations are reported for continuous variables; frequencies are reported for categorical variables. PTSD = posttraumatic stress disorder.  
<sup>a</sup>White race, non-Hispanic ethnicity, and male gender were reference groups.  
 \*p < .05. \*\*p < .01.

**Figure 1**

Model 1: Trauma Exposure, Interpersonal Problems, and Posttraumatic Stress Disorder (PTSD) Symptoms as an Outcome Variable



Note. Standardized parameter estimates (i.e., beta values) are included. Solid lines represent significant direct pathways; dotted lines represent nonsignificant pathways. Only covariates significant at the  $p < .05$  level are included in the figure. White race, Non-Hispanic ethnicity, and male gender were used as the reference groups. RMSEA = .011, CFI = .999, TLI = .995, SRMR = .007.

higher levels of PTSD symptoms,  $\beta = .16$ ,  $p < .001$ . Exposure to interpersonal trauma was directly associated with PTSD symptoms,  $\beta = .17$ ,  $p < .001$ , and indirectly associated with PTSD symptoms through social problem behaviors (i.e., partial mediation),  $\beta = .02$ ,  $p < .001$ . Cumulative noninterpersonal trauma exposure was associated with more severe PTSD symptoms,  $\beta = .03$ ,  $p = .045$ . There were no associations between cumulative noninterpersonal trauma exposure and social problem behaviors or aggressive behaviors. The model explained 19% of the variance in PTSD symptom severity.

## Model 2

The second model also fit the data well, RMSEA = .006, CFI = 1.0, TLI = .998, SRMR = .006 (see Figure 2 and Supplementary Table S2). Social problem behaviors and aggressive behaviors were significantly correlated with each other in the model,  $r = .61$ ;  $p < .001$ . Cumulative interpersonal trauma exposure was associated with more severe PTSD symptoms, aggression, and social problem behaviors,  $\beta = .11$ –.20,  $p < .001$ . More severe PTSD symptoms were associated with increased aggression and social problem behaviors,  $\beta_s = .12$  and  $.18$ ,  $p < .001$ , respectively. Interpersonal trauma exposure was directly


associated with social problem behaviors,  $\beta = .11$ ,  $p < .001$ , and aggressive behaviors,  $\beta = .11$ ,  $p < .001$ , and indirectly associated (i.e., partial mediation) with social problem behaviors,  $\beta = .04$ ,  $p < .001$ , and aggressive behaviors,  $\beta = .02$ ,  $p < .001$ , via PTSD symptoms. Cumulative noninterpersonal trauma exposure was associated with more severe PTSD symptoms,  $\beta = .03$ ,  $p = .034$ , as well as with social problem behaviors indirectly via PTSD symptoms,  $\beta = .01$ ,  $p = .039$ . The model explained 21% of the variance in interpersonal problems.

## Discussion

The present findings support the hypothesis that cumulative interpersonal trauma exposure (i.e., polyvictimization) is associated with adolescent PTSD symptoms and interpersonal problems and that these variables also are partial mediators when the other variable is modeled as an outcome. These findings are consistent with previous research that has shown an association between exposure to interpersonal trauma, interpersonal problems, and PTSD (McLean et al., 2013; Beck et al., 2008) and that disruptions in social interactions can impact the development of PTSD (Brewin et al., 2000; Hyman et al., 2003; Trickey et al., 2012). However, this was the first study to our knowledge

**Figure 2**

Model 2: Trauma Exposure, Interpersonal Problems, and Posttraumatic Stress Disorder (PTSD) Symptoms as a Mediating Variable



Note. Standardized parameter estimates (i.e., beta values) are included. Solid lines represent significant direct pathways; dotted lines represent nonsignificant pathways. Only covariates significant at the  $p < .05$  level are included in the figure. White race, Non-Hispanic ethnicity, and male gender were used as the reference groups. RMSEA = .006, CFI = 1.0, TLI = .998, SRMR = .006.

that examined the role of interpersonal problems as a mediator between the type of trauma exposure and PTSD symptoms as well as PTSD symptoms as a mediator between trauma type and interpersonal problems. Interpersonal trauma exposure may involve psychological factors that can make it difficult for an individual to engage in positive social interactions, such as feelings of betrayal (Freyd, 1994), a sense of distrust and stigma (Punamaki et al., 2005), and associated maladaptive cognitions, such as self-blame, feelings of incompetence, and beliefs that the world is dangerous (Alisic et al., 2014; Tolin & Foa, 2006). Such psychological effects of interpersonal trauma warrant investigation as potential contributors in the pathways that were identified in the current study.

Although the present findings showed that exposure to interpersonal trauma was associated with interpersonal problems and PTSD symptoms in both models, it is important to note that when PTSD symptoms were tested as a mediator, noninterpersonal trauma exposure was associated indirectly with social problem behaviors through PTSD symptoms. Although noninterpersonal trauma exposure has been shown to be associated with a lower risk of adverse outcomes than exposure to interpersonal trauma (Alisic et al., 2014; Musicaro et al., 2020; Vibhakar et al., 2019), children who experience noninterpersonal traumatic events are at risk for multiple adverse outcomes, including full and partial PTSD and CPTSD symptoms (Elliott et al., 2021). These symptoms, in turn, can affect their interpersonal (Bolton et al., 2004), cognitive (Malarbi et al., 2020), be-

havioral (Price et al., 2013), and affective functioning (Briggs-Gowan et al., 2010).

We examined two models in an attempt to obtain a more nuanced understanding of the association between trauma exposure, interpersonal problems, and PTSD symptoms. Both models fit the data well and explained similar levels of variance. Due to the cross-sectional nature of the present study, it is not possible to draw definitive conclusions about the directionality of some of the results. Specifically, the association between PTSD symptoms and interpersonal problems could be bidirectional. Future research using longitudinal data would help to better describe these relations.

The first model examined the association between trauma type and interpersonal problems, with PTSD symptoms as a mediator. In the first model, aggressive behaviors did not explain significant variance in PTSD symptoms after accounting for the association between social problem behaviors with PTSD symptoms. Aggression is often associated with poor peer relationships (Card et al., 2008; Laible et al., 2000; Leary et al., 2006) and PTSD (Marsee, 2008; Silvern & Griese, 2012), but these studies did not account for social problem behaviors. In this model, social problem behaviors partially mediated the association between interpersonal trauma exposure and PTSD symptoms. Thus, social problem behaviors may be a more robust risk factor for PTSD symptoms than aggressive behaviors.

The second model examined the alternative possibility that PTSD symptoms might mediate the relation between

interpersonal trauma exposure and interpersonal problems. The results indicated that PTSD symptoms mediated the association between interpersonal trauma exposure and both social problem behaviors and aggressive behaviors. The two models indicate that there is an association between interpersonal problems and PTSD symptoms (e.g., avoidance, social detachment, hyperarousal) after exposure to interpersonal trauma, although the directionality of these associations is still uncertain. It is possible that interpersonal problems lead to PTSD symptoms, interpersonal problems result from PTSD symptoms (e.g., reactive aggression; Ford, Fraleigh, et al., 2010; Hecker et al., 2015), or both adverse outcomes may occur concurrently but be independent of each other. Prospective studies are needed to determine which of the potential mediation pathways identified in the present study best describe the associations among interpersonal trauma exposure, interpersonal problems, and PTSD symptoms as well as other posttraumatic outcomes.

It is important to note that the models controlled for significant sociodemographic differences. Female youth had significantly higher levels of PTSD symptoms than male youth, which is consistent with previous findings in children and adolescents (Alisic et al., 2014). Adolescent girls tend to report more close friends and more intimacy among friends than adolescent boys (Hall, 2011); thus, a disruption in positive peer relationships could impact female adolescents more than their male counterparts. Youth on public insurance had higher levels of social problem behaviors, which could be related to the high levels of mistrust that can occur in disadvantaged neighborhoods (Ross et al., 2001). Youth on public insurance also had higher levels of aggressive behaviors. Previous researchers have indicated that poverty predicts violence, possibly due to feelings of frustration and injustice (Valois et al., 2002). Youth who were Black, Hispanic, or reported “other” as their race/ethnicity had fewer social problem behaviors compared to non-Hispanic/White youth, and youth who were Hispanic had lower aggressive behaviors compared to non-Hispanic/White youth, which is similar to previous findings from this dataset (Greeson et al., 2014). Future research is needed to examine what role, if any, these sociodemographic differences play in the association between exposure to interpersonal trauma, interpersonal problems, and PTSD.

Several limitations should be noted. Although this study utilized validated measures with a large sample of youth, the data were cross-sectional, so causal relations cannot be inferred. Additional details about the nature and complexity of traumatic experiences, interpersonal problems, and PTSD symptoms were not available for this study but could serve as the basis for future research in this area. Similarly, the extent of the youths’ interpersonal problems before trauma exposure was unknown, so it was not possible to determine the impact of preexisting problems. Although the data were collected before youth started trauma treatment, youth could have had other mental health treatments previously that impacted the baseline data. Symptoms of PTSD were assessed based on the *DSM-IV* rather than *DSM-5* criteria, so the current *DSM-5* category of negative al-

terations in cognitions and mood was not fully captured in this study (APA, 2013). Assessments of social problems and aggression were based on parent report and did not specify key distinctions, such as peer versus family problems and reactive versus proactive aggression. Finally, the sample comprised treatment-seeking youth; therefore, the results are only generalizable to that population.

Preliminary clinical implications include the importance of assessing social problem behaviors with youth exposed to interpersonal trauma, particularly those exposed to several types of interpersonal traumatic events (i.e., polyvictims; Finkelhor et al., 2007). Teaching and practicing communication skills as well as facilitating prosocial connections with peers could be important in PTSD treatment for interpersonal trauma-exposed youth to not only mobilize positive social support but also to reduce interpersonal problems (e.g., Ford et al., 2012).

In conclusion, the present study was the first to our knowledge that demonstrated partial mediation by social problem behaviors in the association between interpersonal trauma exposure and PTSD symptoms as well as partial mediation by PTSD symptoms in the association between interpersonal trauma exposure and interpersonal problems. Interpersonal relationships become more diverse and potentially problematic as children enter adolescence. Positive relationships can provide a sense of belonging and feelings of closeness, whereas negative relationships are associated with mental health problems and high levels of stress (Eng & Heimberg, 2006; Graber et al., 2016; Segrin, 2001; Shahar et al., 2004). Therefore, interpersonal trauma exposure and interpersonal problems are a particularly relevant focus for research and treatment with adolescents who have experienced trauma.

### Open Practices Statement

The secondary analysis study reported in this article was not formally preregistered. Neither the data nor the materials have been made available on a permanent third-party archive.

### References

- Achenbach, T.M., & Rescorla, L.A. (2001). *Manual for the ASEBA School Age-Forms & Profiles*. University of Vermont, Research Center for Children.
- Alisic, E., Zalta, A. K., Van Wesel, F., Larsen, S. E., Hafstad, G. S., Hassanpour, K., & Smid, G. E. (2014). Rates of post-traumatic stress disorder in trauma-exposed children and adolescents: Meta-analysis. *The British Journal of Psychiatry*, 204(5), 335–340. <https://doi.org/10.1192/bjp.bp.113.131227>
- American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders* (4th ed., text rev.). Author.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Author. <https://doi.org/10.1176/appi.books.9780890425596>
- Baker, D. E., Hill, M., Chamberlain, K., Hurd, L., Karlsson, M., Zielinski, M., Calvert, M., & Bridges, A.J. (2021). Interpersonal vs. non-interpersonal cumulative traumas and psychiatric symptoms in treatment-seeking incarcerated women. *Journal of Trauma & Dissociation*, 22(3), 249–264. <https://doi.org/10.1080/15299732.2020.1760172>



- Beck, J. G., Grant, D. M., Clapp, J. D., & Palyo, S. A. (2008). Understanding the interpersonal impact of trauma: Contributions of PTSD and depression. *Journal of Anxiety Disorders, 23*(4), 443–450. <https://doi.org/10.1016/j.janxdis.2008.09.001>
- Bolton, D., Hill, J., O’Ryan, D., Udwin, O., Boyle, S., & Yule, W. (2004). Long-term effects of psychological trauma on psychosocial functioning. *Journal of Child Psychology and Psychiatry, 45*(5), 1007–1014. <https://doi.org/10.1111/j.1469-7610.2004.t01-1-00292.x>
- Borkovec, T. D., Newman, M. G., Pincus, A. L., & Lytle, R. (2002). A component analysis of cognitive behavioral therapy for generalized anxiety disorder and the role of interpersonal problems. *Journal of Consulting and Clinical Psychology, 70*(2), 288–298. <https://doi.org/10.1037/0022-006X.70.2.288>
- Brewin, C. R., Andrews, B., & Valentine, J. D. (2000). Meta-analysis of risk factors for posttraumatic stress disorder in trauma-exposed adults. *Journal of Consulting and Clinical Psychology, 68*(5), 748–766. <https://doi.org/10.1037/0022-006X.68.5.748>
- Brewin, C. R., Cloitre, M., Hyland, P., Shevlin, M., Maercker, A., Bryant, R. A., Hamayun, A., Jones, L.M., Kagee, A., Rousseau, C., Somasundaram, D., Suzuki, Y., Wessely, S., Ommeren, M., & Reed, G. M. (2017). A review of current evidence regarding the *ICD-11* proposals for diagnosing PTSD and complex PTSD. *Clinical Psychology Review, 58*, 1–15. <https://doi.org/10.1016/j.cpr.2017.09.001>
- Briggs, E. C., Greeson, J. K., Layne, C. M., Fairbank, J. A., Knoverek, A. M., & Pynoos, R. S. (2012). Trauma exposure, psychosocial functioning, and treatment needs of youth in residential care: Preliminary findings from the NCTSN Core Data Set. *Journal of Child & Adolescent Trauma, 5*(1), 1–15. <https://doi.org/10.1080/19361521.2012.646413>
- Briggs-Gowan, M. J., Carter, A. S., Clark, R., Augustyn, M., McCarthy, K. J., & Ford, J. D. (2010). Exposure to potentially traumatic events in early childhood: differential links to emergent psychopathology. *Journal of Child Psychology and Psychiatry and Allied Disciplines, 51*(10), 1132–1140. <https://doi.org/10.1111/j.1469-7610.2010.02256.x>
- Card, N. A., Stucky, B. D., Sawalani, G. M., & Little, T. D. (2008). Direct and indirect aggression during childhood and adolescence: A meta-analytic review of gender differences, intercorrelations, and relations to maladjustment. *Child Development, 79*(5), 1185–1229. <https://doi.org/10.1111/j.1467-8624.2008.01184.x>
- Cast, A. D., & Burke, P. J. (2002). A theory of self-esteem. *Social Forces, 80*(3), 1041–1068. <https://doi.org/10.1353/sof.2002.0003>
- Charak, R., Ford, J. D., Modrowski, C. A., & Kerig, P. K. (2018). PTSD and dissociation symptoms as mediators of the relationship between polyvictimization and psychosocial and behavioral problems among justice-involved adolescents. *Journal of Trauma & Dissociation, 19*(3), 325–346. <https://doi.org/10.1080/15299732.2018.1441354>
- Charuvastra, A., & Cloitre, M. (2008). Social bonds and posttraumatic stress disorder. *Annual Review of Psychology, 59*(1), 301–328. <https://doi.org/10.1146/annurev.psych.58.110405.085650>
- Dang, H. M., Nguyen, H., & Weiss, B. (2017). Incremental validity of the child behavior checklist (CBCL) and the strengths and difficulties questionnaire (SDQ) in Vietnam. *Asian Journal of Psychiatry, 29*, 96–100. <https://doi.org/10.1016/j.ajp.2017.04.023>
- Dube, S. R., Miller, J. W., Brown, D. W., Giles, W. H., Felitti, V. J., Dong, M., & Anda, R. F. (2006). Adverse childhood experiences and the association with ever using alcohol and initiating alcohol use during adolescence. *Journal of Adolescent Health, 38*(4), 444e1–444.e10. <https://doi.org/10.1016/j.jadohealth.2005.06.006>
- Dunn, E. C., Nishimi, K., Powers, A., & Bradley, B. (2017). Is developmental timing of trauma exposure associated with depressive and post-traumatic stress disorder symptoms in adulthood? *Journal of Psychiatric Research, 84*, 119–127. <https://doi.org/10.1016/j.jpsychires.2016.09.004>
- Elliott, R., McKinnon, A., Dixon, C., Boyle, A., Murphy, F., Dahm, T., Travers-Hill, E., Mul, C., Archibald, S., Smith, P., Dalglish, T., Meiser-Stedman, R., & Hitchcock, C. (2021). Prevalence and predictive value of *ICD-11* posttraumatic stress disorder and complex PTSD diagnoses in children and adolescents exposed to a single-event trauma. *Journal of Child Psychology and Psychiatry, 62*(3), 270–276. <https://doi.org/10.1111/jcpp.13240>
- Eng, W., & Heimberg, R. G. (2006). Interpersonal correlates of generalized anxiety disorder: Self versus other perception. *Journal of Anxiety Disorders, 20*(3), 380–387. <https://doi.org/10.1016/j.janxdis.2005.02.005>
- Finkelhor, D., Ormrod, R. K., & Turner, H. A. (2007). Poly-victimization: A neglected component in child victimization. *Child Abuse & Neglect, 31*(1), 7–26. <https://doi.org/10.1016/j.chiabu.2006.06.008>
- Ford, J. D., Elhai, J. D., Connor, D. F., & Frueh, B. C. (2010). Poly-victimization and risk of posttraumatic, depressive, and substance use disorders and involvement in delinquency in a national sample of adolescents. *Journal of Adolescent Health, 46*(6), 545–552. <https://doi.org/10.1016/j.jadohealth.2009.11.212>
- Ford, J. D., Fraleigh, L. A., & Connor, D. F. (2010). Child abuse and aggression among seriously emotionally disturbed children. *Journal of Clinical Child and Adolescent Psychology, 39*(1), 25–34. <https://doi.org/10.1080/15374410903401104>
- Ford, J. D., Steinberg, K. L., Hawke, J., Levine, J., & Zhang, W. (2012). Randomized trial comparison of emotion regulation and relational psychotherapies for PTSD with girls involved in delinquency. *Journal of Clinical Child and Adolescent Psychology, 41*(1), 27–37. <https://doi.org/10.1080/15374416.2012.632343>
- Freyd, J. J. (1994). Betrayal trauma: Traumatic amnesia as an adaptive response to childhood abuse. *Ethics & Behavior, 4*(4), 307–329. [https://doi.org/10.1207/s15327019eb0404\\_1](https://doi.org/10.1207/s15327019eb0404_1)
- Frost, R., Hyland, P., Shevlin, M., & Murphy, J. (2020). Distinguishing complex PTSD from borderline personality disorder among individuals with a history of sexual trauma: A latent class analysis. *European Journal of Trauma and Dissociation, 4*(1), 10080. <https://doi.org/10.1016/j.ejtd.2018.08.004>
- Graber, R., Turner, R., & Madill, A. (2016). Best friends and better coping: Facilitating psychological resilience through boys’ and girls’ closest friendships. *British Journal of Psychology, 107*(2), 338–358. <https://doi.org/10.1111/bjop.12135>
- Greeson, J. K., Briggs, E. C., Layne, C. M., Belcher, H. M., Ostrowski, S. A., Kim, S., Lee, R.C., Vivrette, R. L., Pynoos, R.S., & Fairbank, J. A. (2014). Traumatic childhood experiences in the 21st century: Broadening and building on the Ace Studies with data from the National Child Traumatic Stress Network. *Journal of Interpersonal Violence, 29*(3), 536–556. <https://doi.org/10.1177/0886260513505217>
- Haldane, J., & Nickerson, A. (2016). The impact of interpersonal and noninterpersonal trauma on psychological symptoms in refugees: The moderating role of gender and trauma type. *Journal of Traumatic Stress, 29*(5), 457–465. <https://doi.org/10.1002/jts.22132>
- Hall, J. A. (2011). Sex differences in friendship expectations: A meta-analysis. *Journal of Social and Personal Relationships, 28*(6), 723–747. <https://doi.org/10.1177/0265407510386192>
- Hebert, M., Langevin, R., & Oussaid, E. (2018). Cumulative childhood trauma, emotion regulation, dissociation, and behavior problems in school-aged sexual abuse victims. *Journal of Affective Disorders, 225*, 306–312. <https://doi.org/10.1016/j.jad.2017.08.044>
- Hecker, T., Fetz, S., Ainamani, H., & Elbert, T. (2015). The cycle of violence: Associations between exposure to violence, trauma-related symptoms, and

- aggression—findings from Congolese refugees in Uganda. *Journal of Traumatic Stress*, 28(5), 448–455. <https://doi.org/10.1002/jts.22046>
- Hodges, M., Godbout, N., Briere, J., Lanktree, C., Gilbert, A., & Kletzka, N. T. (2013). Cumulative trauma and symptom complexity in children: A path analysis. *Child Abuse & Neglect*, 37(11), 891–898. <https://doi.org/10.1016/j.chiabu.2013.04.001>
- Horowitz, L. M., Alden, L. E., Wiggins, J. S., & Pincus, A. L. (2000). *Inventory of Interpersonal Problems manual*. The Psychological Corporation.
- Horowitz, L. M., Rosenberg, S. E., & Bartholomew, K. (1993). Interpersonal problems, attachment styles, and outcome in brief dynamic psychotherapy. *Journal of Consulting and Clinical Psychology*, 61(4), 549–560. <https://doi.org/10.1037/0022-006X.61.4.549>
- Hu, L., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling: A Multidisciplinary Journal*, 6(1), 1–55. <https://doi.org/10.1080/10705519909540118>
- Hyman, S. M., Gold, S. N., & Cott, M. A. (2003). Forms of social support that moderate PTSD in childhood sexual abuse survivors. *Journal of Family Violence*, 18(5), 295–300. <https://doi.org/10.1023/A:1025117311660>
- Jaffe, A. E., DiLillo, D., Gratz, K. L., & Messman-Moore, T. L. (2019). Risk for revictimization following interpersonal and noninterpersonal trauma: Clarifying the role of Posttraumatic Stress symptoms and trauma-related cognitions. *Journal of Traumatic Stress*, 32(1), 42–55. <https://doi.org/10.1002/jts.22372>
- Jaworska, N., & MacQueen, G. (2015). Adolescence as a unique developmental period. *Journal of Psychiatry & Neuroscience: Journal of Psychiatry and Neuroscience*, 40(5), 291–293. <https://doi.org/10.1503/jpn.150268>
- Johnson, J. G., Cohen, P., Gould, M. S., Kasen, S., Brown, J., & Brook, J. S. (2002). Childhood adversities, interpersonal difficulties, and risk for suicide attempts during late adolescence and early adulthood. *Archives of General Psychiatry*, 59(8), 741–749. <https://doi.org/10.1001/archpsyc.59.8.741>
- Jones, J. M. (2007). Exposure to chronic community violence: Resilience in African American children. *Journal of Black Psychology*, 33(2), 125–149. <https://doi.org/10.1177/0095798407299511>
- Kerig, P. K., Ward, R. M., Vanderzee, K. L., & Moeddel, M. A. (2009). Post-traumatic stress as a mediator of the relationship between trauma and mental health problems among juvenile delinquents. *Journal of Youth and Adolescence*, 38(9), 1214–1225. <https://doi.org/10.1007/s10964-008-9332-5>
- Laible, D. J., Carlo, G., & Raffaelli, M. (2000). The differential relations of parent and peer attachment to adolescent adjustment. *Journal of Youth and Adolescence*, 29(1), 45–59. <https://doi.org/10.1023/A:1005169004882>
- Lakey, B., & Cronin, A. (2008). Low social support and major depression: Research, theory and methodological issues. In K. Dobson & D. Dozois (Eds.), *Risk factors in depression* (pp. 385–408). Elsevier. <https://doi.org/10.1016/B978-0-08-045078-0.00017-4>
- Leary, M. R., Twenge, J. M., & Quinlivan, E. (2006). Interpersonal rejection as a determinant of anger and aggression. *Personality and Social Psychology Review*, 10(2), 111–132. [https://doi.org/10.1207/s15327957pspr1002\\_2](https://doi.org/10.1207/s15327957pspr1002_2)
- Malarbi, S. T., Muscara, F., Barnett, P. L. J., Palmer, C. S., & Stargatt, R. (2020). Post-traumatic stress symptoms and cognition in children exposed to motor vehicle accident trauma. *Child Neuropsychology*, 26(4), 560–575. <https://doi.org/10.1080/09297049.2019.1700223>
- Marsee, M. A. (2008). Reactive aggression and posttraumatic stress in adolescents affected by Hurricane Katrina. *Journal of Clinical Child & Adolescent Psychology*, 37(3), 519–529. <https://doi.org/10.1080/15374410802148152>
- McLean, C. P., Rosenbach, S. B., Capaldi, S., & Foa, E. B. (2013). Social and academic functioning in adolescents with child sexual abuse-related PTSD. *Child Abuse & Neglect*, 37(9), 675–678. <https://doi.org/10.1016/j.chiabu.2013.03.010>
- McFarlane, A. C., & Bookless, C. (2001). The effect of PTSD on interpersonal relationships: Issues for emergency service workers. *Sexual and Relationship Therapy*, 16(3), 261–267. <https://doi.org/10.1080/14681990124457>
- Mercado-Crespo, M. C., & Mbah, A. K. (2013). Race and ethnicity, substance use, and physical aggression among U.S. high school students. *Journal of Interpersonal Violence*, 28(7), 1367–1384. <https://doi.org/10.1177/0886260512468234>
- Musicaro, R. M., Ford, J., Suvak, M. K., Sposato, A., & Andersen, S. (2020). Sluggish cognitive tempo and exposure to interpersonal trauma in children. *Anxiety Stress Coping*, 33(1), 100–114. <https://doi.org/10.1080/10615806.2019.1695124>
- Muthén, L. K., & Muthén, B. O. (2017). *Mplus user's guide* (8th ed.). Muthén & Muthén.
- Newman, E. (2002). Assessment of PTSD and trauma exposure in adolescents. *Journal of Aggression, Maltreatment & Trauma*, 6(1), 59–77. [https://doi.org/10.1300/J146v06n01\\_04](https://doi.org/10.1300/J146v06n01_04)
- Petty, S. C., Sachs-Ericsson, N., & Joiner Jr, T. E. (2004). Interpersonal functioning deficits: Temporary or stable characteristics of depressed individuals? *Journal of Affective Disorders*, 81(2), 115–122. [https://doi.org/10.1016/S0165-0327\(03\)00158-7](https://doi.org/10.1016/S0165-0327(03)00158-7)
- Price, M., Higa-McMillan, C., Kim, S., & Frueh, B. C. (2013). Trauma experience in children and adolescents: An assessment of the effects of trauma type and role of interpersonal proximity. *Journal of Anxiety Disorders*, 27(7), 652–660. <https://doi.org/10.1016/j.janxdis.2013.07.009>
- Punamäki, R. L., Komproe, I. H., Qouta, S., Elmasri, M., & de Jong, J. T. (2005). The role of peritraumatic dissociation and gender in the association between trauma and mental health in a Palestinian community sample. *American Journal of Psychiatry*, 162(3), 545–551. <https://doi.org/10.1176/appi.ajp.162.3.545>
- Ross, C. E., Mirowsky, J., & Pribesh, S. (2001). Powerlessness and the amplification of threat: Neighborhood disadvantage, disorder, and mistrust. *American Sociological Review*, 66(4), 568–591. <https://doi.org/10.2307/3088923>
- SAS Institute Inc. (2015). *SAS/STAT® 14.1 user's guide*. SAS Institute Inc.
- Segrin, C. (2001). *Interpersonal processes in psychological problems*. Guilford Press.
- Shahar, G., Joiner Jr, T. E., Zuroff, D. C., & Blatt, S. J. (2004). Personality, interpersonal behavior, and depression: Co-existence of stress-specific moderating and mediating effects. *Personality and Individual Differences*, 36(7), 1583–1596. <https://doi.org/10.1016/j.paid.2003.06.006>
- Silvern, L., & Griese, B. (2012). Multiple types of child maltreatment, post-traumatic stress, dissociative symptoms, and reactive aggression among adolescent criminal offenders. *Journal of Child & Adolescent Trauma*, 5(2), 88–101. <https://doi.org/10.1080/19361521.2012.671799>
- Solva, K., Haselgruber, A., & Lueger-Schuster, B. (2020). The relationship between cumulative traumatic experiences and ICD-11 post-traumatic symptoms in children and adolescents in foster care: The mediating effect of sense of coherence. *Child Abuse Neglect*, 101, 104388. <https://doi.org/10.1016/j.chiabu.2020.104388>
- Sperry, D. M., & Widom, C. S. (2013). Child abuse and neglect, social support, and psychopathology in adulthood: A prospective investigation. *Child Abuse & Neglect*, 37(6), 415–425. <https://doi.org/10.1016/j.chiabu.2013.02.006>
- Steinberg A. M., Brymer M. J., Decker K. B., & Pynoos R. S. (2004). The University of California at Los Angeles Post-Traumatic Stress Disorder Reaction Index. *Current Psychiatry Reports*, 6(2), 96–100. <https://doi.org/10.1007/s11920-004-0048-2>

- Steinberg, A. M., Brymer, M. J., Kim, S., Briggs, E. C., Ippen, C. G., Ostrowski, S. A., Gully, K.J., & Pynoos, R. S. (2013). Psychometric properties of the UCLA PTSD Reaction Index: Part I. *Journal of Traumatic Stress, 26*(1), 1–9. <https://doi.org/10.1002/jts.21780>
- Steinberg, A. M., Pynoos, R. S., Briggs, E. C., Gerrity, E. T., Layne, C. M., Vivrette, R. L., Beyerlein, B., & Fairbank, J. A. (2014). The National Child Traumatic Stress Network Core Data Set: Emerging findings, future directions, and implications for theory, research, practice, and policy. *Psychological Trauma: Theory, Research, Practice, and Policy, 6*(S1), S50–S57. <https://doi.org/10.1037/a0037798>
- Tolin, D. F., & Foa, E. B. (2006). Sex differences in trauma and posttraumatic stress disorder: A quantitative review of 25 years of research. *Psychological Bulletin, 132*(1), 37–85. <https://doi.org/10.1037/1942-9681.132.1.37>
- Trickey, D., Siddaway, A. P., Meiser-Stedman, R., Serpell, L., & Field, A. P. (2012). A meta-analysis of risk factors for post-traumatic stress disorder in children and adolescents. *Clinical Psychology Review, 32*(2), 122–138. <https://doi.org/10.1016/j.cpr.2011.12.001>
- Valois, R. F., MacDonald, J. M., Bretous, L., Fischer, M. A., & Drane, J. W. (2002). Risk factors and behaviors associated with adolescent violence and aggression. *American Journal of Health Behavior, 26*(6), 454–464. <https://doi.org/10.5993/AJHB.26.6.6>
- Vibhakar, V., Allen, L. R., Gee, B., & Meiser-Stedman, R. (2019). A systematic review and meta-analysis on the prevalence of depression in children and adolescents after exposure to trauma. *Journal of Affective Disorders, 255*, 77–89. <https://doi.org/10.1016/j.jad.2019.05.005>
- Vittengl, J. R., Clark, L. A., & Jarrett, R. B. (2003). Interpersonal problems, personality pathology, and social adjustment after cognitive therapy for depression. *Psychological Assessment, 15*(1), 29–40. <https://doi.org/10.1037/1040-3590.15.1.29>
- West, S. G., Taylor, A. B., & Wu, W. (2012). Model fit and model selection in structural equation modeling. In R. H. Hoyle (Ed.), *Handbook of structural equation modeling* (pp. 209–231). Guilford Press.
- Winer, A. C., & Thompson, R. (2013). How poverty and depression impact a child's social and emotional competence. *Policy Brief, Center for Poverty Research, 1*(10), 1–2.
- Woodward, M. J., Eddinger, J., Henschel, A. V., Dodson, T. S., Tran, H. N., & Beck, J. G. (2015). Social support, posttraumatic cognitions, and PTSD: The influence of family, friends, and a close other in an interpersonal and noninterpersonal trauma group. *Journal of Anxiety Disorders, 35*, 60–67. <https://doi.org/10.1016/j.janxdis.2015.09.002>
- World Health Organization. (2020, September). *ICD-11 for mortality and morbidity statistics*. <https://icd.who.int/browse11/l-m/en>