

Repeat RNA Toxicity Drives Ribosomal RNA Processing Defects in SCA2

Geena Skariah,^{1*} and Roger Lee Albin^{1,2}

¹Department of Neurology, University of Michigan, Ann Arbor, Michigan, USA

²Neurology Service and GRECC, VAAHS, Ann Arbor, Michigan, USA

Spinocerebellar ataxia type 2 (SCA2) is an autosomal dominant ataxia with no effective treatments. It is the second most prevalent form of dominant ataxia, accounting for ~15% of cases worldwide.^{1,2} Clinical features of SCA2 include gait ataxia, tremor, abnormal eye movements, peripheral neuropathy, and other multisystem features. SCA2 neuropathology is characterized by olivopontocerebellar atrophy with marked loss of Purkinje cells and inferior olivary, pontocerebellar, and substantia nigra neuron loss.³⁻⁶

SCA2 belongs to a growing class of nucleotide repeat expansion disorders arising from unstable repeats in the genome.^{7,8} In this issue of *Movement Disorders*, Li et al⁹ add to an expanding body of literature indicating that repeat expansion disorders are often characterized by plural molecular pathogenic mechanisms.

SCA2 is caused by expansion of CAG trinucleotide repeats in the first exon of the *ATXN2* gene, encoding a polyglutamine (polyQ) repeat containing protein. In individuals with SCA2, repeat numbers are in the range of 37 to 39 repeats, as opposed to a normal range of 13 to 23 repeats.¹⁰ Intermediate repeat lengths of 27 to 33 CAG repeats are associated with increased risk for amyotrophic lateral sclerosis (ALS).¹¹ The exact function of *ATXN2* protein is not yet known, but it is implicated in RNA processing and translation, stress granule assembly, endoplasmic reticular (ER) calcium response, and cytoskeletal reorganization.¹² The expanded CAG repeat in *ATXN2* encodes an abnormal stretch of polyglutamine (polyQ) in the N-terminal region. In multiple polyQ disorders, these canonically translated repeat harboring proteins show gain of function toxicities separate from their normal cellular functions.¹³ These pathogenic protein species may exert their effects via multiple proximate mechanisms.¹⁴ In the case of the polyQ disorder

Huntington's disease (HD), haploinsufficiency may also contribute to neurodegeneration.⁷

A second general molecular mechanism in nucleotide expansion disorders is the phenomenon of repeat-associated non-AUG (RAN) translation, which gives rise to toxic repeat peptides because of aberrant translation from expanded repeat regions. RAN translation, however, from the expanded SCA2 transcript appears to be minimal and may not be a significant pathogenic process in SCA2.¹⁵

Li et al⁹ evaluated a third potential pathogenic molecular mechanism involving repeat containing RNA transcripts. In some nucleotide expansion disorders, such as type 1 myotonic dystrophy, expanded repeats exert neurotoxicity at the RNA level by disrupting mRNA processing. One of the mechanisms of RNA mediated toxicity is sequestration of key RNA binding proteins (RBPs), making them unavailable for their normal cellular functions.¹⁶ RNA toxicity is often accompanied by formation of RNA foci. The authors find that expanded sense *ATXN2* RNA transcripts cause neurotoxicity and disrupt ribosomal RNA processing. They present evidence that overexpression of an expanded *ATXN2* transcript, not undergoing RAN translation, shows increased caspase 3/7 activity in human neuroblastoma cells and shows increased toxicity in nuclear condensation assays performed with primary mouse cortical neurons. The authors go on to show that the mutant transcripts bearing either 58 or 104 CAG triplets form more RNA foci when transfected into cells. RNA foci were also detected in cerebellar Purkinje cells of SCA2 transgenic mice and in one of five postmortem human patient brains.

To examine the possibility of RBP sequestration, the authors performed in vitro pull down of the mutant *ATXN2* transcript followed by mass spectrometry. They identified a number of RBPs with a predominant

© 2021 International Parkinson and Movement Disorder Society

*Correspondence to: Dr. G. Skariah, Department of Neurology, University of Michigan, Ann Arbor, MI, USA; E-mail: geskaria@med.umich.edu

Relevant conflicts of interest/financial disclosures: Nothing to report.

Received: 19 August 2021; Revised: 26 August 2021; Accepted: 26 August 2021

Published online in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/mds.28795




FIG. 1. Multiple modes of SCA2 pathogenesis. Normal CAG repeat number in the ATXN2 gene generates a WT protein that participates in a host of cellular functions. The intermediate repeat length is associated with ALS, whereas the expanded CAG repeats in the ATXN2 gene causes toxicity by RNA binding protein sequestration, repeat-associated non-AUG translation or via the ATXN2-antisense (AS) transcript. Each of these aberrant processes cause dysregulation of downstream molecular pathways and could collectively contribute to neurodegeneration. [Color figure can be viewed at wileyonlinelibrary.com]

nuclear and nucleolar localization pattern, suggesting the nucleus as the possible site for aberrant interactions. Several of the sequestered RBPs are critical for maturation of the small rRNA subunit. The authors selected transducing β -like protein 3 (TBL3) for further analysis because of its known interaction with the expanded huntingtin transcript, implicating a possible common disease mechanism in HD and SCA2. With a series of elegant biochemical studies, the authors demonstrate that TBL3 binds to the aberrant hairpin structure formed by continuous CAG repeats in the expanded repeat ATXN2 transcript.

The yeast homolog of TBL3 plays a role in 35S rRNA processing and 18S rRNA biogenesis.¹⁷ Based on these findings, Li et al⁹ tested the levels of 45S pre-rRNA (human equivalent of yeast 35S pre-rRNA) and the ratios of mature 18S and 28S rRNA after TBL3 knock-down in HEK293T cells. They observed an increase in 45S pre-rRNA levels and a decrease in the ratio of mature 18S and 28S rRNA, suggesting defective rRNA processing and maturation, respectively, in the absence of TBL3. Similar trends, albeit without statistical significance, were observed in postmortem HD and SCA2 brains. Defects in ribosome biogenesis are reported in HD, accompanied by nucleolar aggregates.^{18,19} This

study opens up lines of investigation into the possible role of ribosomal abnormalities in another neurodegenerative disorder.²⁰ Taken together, the report by Li et al,⁹ along with their earlier finding of a toxic anti-sense ATXN2 transcript, presents evidence for combinatorial protein-RNA driven pathology in SCA2 (Fig. 1).

Neurodegenerative disorders such as SCA2 typically exhibit age-related penetrance. The observation that ribosome biogenesis may be affected in SCA2 is intriguing and with implications for what might happen in aging brains, characterized by decreased global protein translation.^{21,22} Many age associated neurodegenerative disorders are because of dysfunctions in core components of the translation machinery.²³ Ribosomal proteins, as well as rRNA levels, are affected during healthy aging²⁴ and could conceivably change ribosomal assembly as well as mRNA translation dynamics. These age associated changes could influence ribosomal and proteasomal subunit stoichiometry, altering global protein homeostasis.²⁵ These age-related changes might make neurons more susceptible to the toxic effects of expanded repeat transcripts.

Another characteristic aspect of neurodegenerative disorders, such as SCA2, is degeneration of specific neuronal subtypes. Single-cell transcriptomic analyses of

the aging brain are beginning to unravel tissue and cell type specificities for components of the translation machinery, such as ribosomal proteins,^{26,27} suggesting distinct post-transcriptional outcomes based on neuronal cell type. This might explain the susceptibility of neuronal subtypes in specific disease contexts. In addition, high-energy neuronal subtypes, such as dopaminergic neurons, are particularly susceptible to mitochondrial dysfunction, an emerging factor in multiple neurodegenerative disorders.^{28,29}

Moreover, the most affected transcripts in the aging human cortex are involved in synaptic function.³⁰ Because activity-mediated dynamic regulation of mRNA translation at synapses is a key factor for neuronal function and survival,³¹ sequestration of crucial molecules, such as TBL3 as reported here, could be particularly impactful for synaptic mRNA translation. Ultimately leading to neuronal death, the initial manifestations would be synaptic dysfunction and degeneration, which are widely believed to be common features of many neurodegenerative disorders.

Within the context of an aging neuronal environment, the accumulation of expanded repeat containing RNA and proteins could trigger cascades of events that affect both arms of the peptide life cycle—synthesis and degradation—precipitating neurodegeneration.³² ■

Acknowledgments: The authors would like to acknowledge Dr. Indranil Malik for assistance with the figure and Dr. Peter Todd for helpful suggestions. G.S. receives support from the Claude D. Pepper Older Adults Independence Center (AG024824), Michigan Alzheimer Disease Center (AG053760), and National Institutes of Health grants to Dr. Todd (R01NS086810, R01NS099280, and P50HD104463). R.L.A. receives support from P50NS123067 and the Parkinson's Foundation.

Data Availability Statement

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

References

- Pulst S-M, Nechiporuk A, Nechiporuk T, et al. Moderate expansion of a normally biallelic trinucleotide repeat in spinocerebellar ataxia type 2. *Nat Genet* 1996;14:269–276.
- Imbert G, Saudou F, Yvert G, et al. Cloning of the gene for spinocerebellar ataxia 2 reveals a locus with high sensitivity to expanded CAG/glutamine repeats. *Nat Genet* 1996;14:285–291.
- Filla A, De Michele G, Santoro L, et al. Spinocerebellar ataxia type 2 in southern Italy: a clinical and molecular study of 30 families. *J Neurol* 1999;246:467–471.
- Velázquez-Pérez L, Rodríguez-Labrada R, García-Rodríguez JC, Almaguer-Mederos LE, Cruz-Mariño T, Laffita-Mesa JM. A comprehensive review of spinocerebellar ataxia type 2 in Cuba. *Cerebellum* 2011;10:184–198.
- Estrada R, Galarraga J, Orozco G, Nodarse A, Auburger G. Spinocerebellar ataxia 2 (SCA2): morphometric analyses in 11 autopsies. *Acta Neuropathol* 1999;97:306–310.
- Schöls L, Reimold M, Seidel K, et al. No parkinsonism in SCA2 and SCA3 despite severe neurodegeneration of the dopaminergic substantia nigra. *Brain* 2015;138:3316–3326.
- Lieberman AP, Shakkottai VG, Albin RL. Polyglutamine Repeats in Neurodegenerative Diseases. *Annu Rev Pathol* 2019;14:1–27.
- van Eyk CL, Richards RI. Dynamic mutations: where are they now? *Adv Exp Med Biol* 2012;769:55–77.
- Li PP, Moulick R, Feng H, et al. RNA toxicity and perturbation of rRNA processing in spinocerebellar ataxia type 2. *Mov Disord* 2021. DOI: 10.1002/mds.28729
- Antenora A, Rinaldi C, Roca A, et al. The multiple faces of spinocerebellar ataxia type 2. *Ann Clin Transl Neurol* 2017;4:687–695.
- Elden AC, Kim H-J, Hart MP, et al. Ataxin-2 intermediate-length polyglutamine expansions are associated with increased risk for ALS. *Nature* 2010;466:1069–1075.
- Magaña JJ, Velázquez-Pérez L, Cisneros B. Spinocerebellar ataxia type 2: clinical presentation, molecular mechanisms, and therapeutic perspectives. *Mol Neurobiol* 2013;47:90–104.
- Malik I, Kelley CP, Wang ET, Todd PK. Molecular mechanisms underlying nucleotide repeat expansion disorders. *Nat Rev Mol Cell Biol* 2021;22:644.
- Jafar-Nejad P, Ward CS, Richman R, Orr HT, Zoghbi HY. Regional rescue of spinocerebellar ataxia type 1 phenotypes by 14-3-3epsilon haploinsufficiency in mice underscores complex pathogenicity in neurodegeneration. *Proc Natl Acad Sci U S A* 2011;108:2142–2147.
- Scoles DR, Ho MH, Dansithong W, et al. Repeat associated non-aug translation (RAN translation) dependent on sequence downstream of the ATXN2 CAG repeat. *PLoS One* 2015;10:e0128769.
- Miller JW, Urbinati CR, Teng-Umnuay P, et al. Recruitment of human muscleblind proteins to (CUG)(n) expansions associated with myotonic dystrophy. *EMBO J* 2000;19:4439–4448.
- Dragon F, Gallagher JE, Compagnone-Post PA, et al. A large nucleolar U3 ribonucleoprotein required for 18S ribosomal RNA biogenesis. *Nature* 2002;417:967–970.
- Tsoi H, Chan HY. Roles of the nucleolus in the CAG RNA-mediated toxicity. *Biochim Biophys Acta* 1842;2014:779–784.
- Lee J, Hwang YJ, Ryu H, Kowall NW, Ryu H. Nucleolar dysfunction in Huntington's disease. *Biochim Biophys Acta* 1842;2014:785–790.
- Hetman M, Slomnicki LP. Ribosomal biogenesis as an emerging target of neurodevelopmental pathologies. *J Neurochem* 2019;148:325–347.
- Tavernarakis N. Ageing and the regulation of protein synthesis: a balancing act? *Trends Cell Biol* 2008;18:228–235.
- Rattan SI. Synthesis, modifications, and turnover of proteins during aging. *Exp Gerontol* 1996;31:33–47.
- Kapur M, Ackerman SL. mRNA translation gone awry: translation fidelity and neurological disease. *Trends Genet* 2018;34:218–231.
- Curran SP, Ruvkun G. Lifespan regulation by evolutionarily conserved genes essential for viability. *PLoS Genet* 2007;3:e56.
- Kelmer Sacramento E, Kirkpatrick JM, Mazzetto M, et al. Reduced proteasome activity in the aging brain results in ribosome stoichiometry loss and aggregation. *Mol Syst Biol* 2020;16:e9596.
- Davie K, Janssens J, Koldere D, et al. A single-cell transcriptome atlas of the aging drosophila brain. *Cell* 2018;174:982–998.e920.
- Ximerakis M, Lipnick SL, Innes BT, et al. Single-cell transcriptomic profiling of the aging mouse brain. *Nat Neurosci* 2019;22:1696–1708.
- Müller-Nedebeck AC, van der Westhuizen FH, Köks S, Bardien S. Nuclear genes associated with mitochondrial DNA processes as contributors to Parkinson's disease risk. *Mov Disord* 2021;36:815–831.

-
29. Norat P, Soldozy S, Sokolowski JD, et al. Mitochondrial dysfunction in neurological disorders: exploring mitochondrial transplantation. *NPJ Regen Med* 2020;5:22.
 30. Dillman AA, Majounie E, Ding J, et al. Transcriptomic profiling of the human brain reveals that altered synaptic gene expression is associated with chronological aging. *Sci Rep* 2017;7:16890.
 31. Kapur M, Monaghan CE, Ackerman SL. Regulation of mRNA translation in neurons-A matter of life and death. *Neuron* 2017;96:616–637.
 32. Skariah G, Todd PK. Translational control in aging and neurodegeneration. *Wiley Interdiscip Rev RNA* 2021;12:e1628.