

Time to Come Inside? A Novel Approach to Anastomotic Biliary Strictures

SEE ARTICLE ON PAGE 88

Liver transplantation is the definitive management for patients with decompensated cirrhosis and/or liver malignancy. Two types of liver transplantation are traditionally performed: deceased donor liver transplantation (DDLT) and living donor liver transplantation (LDLT). LDLT involves a partial liver from a live donor and is a response to the limited resource of donor livers. In LDLT, the biliary anastomosis can be accomplished by a Roux-en-Y hepaticojejunostomy or a duct-to-duct biliary reconstruction. Biliary strictures, which typically occur at the anastomosis, are known complications of both methods but occur more commonly in LDLT. While the duct-to-duct biliary anastomosis provides normal enteric anatomy and ease of endoscopic access, the anastomosis is more peripheral and angulated, and thus management strategies and outcomes have generally been studied separately from DDLT.⁽¹⁾

Standard treatment of these strictures is performed by endoscopic retrograde cholangiopancreatography (ERCP) and traditionally includes biliary sphincterotomy, dilation, and placement of transpapillary plastic, and/or self-expandable metal stents (SEMSs)

either alone or in combination.⁽¹⁾ Biliary sphincterotomy is implemented for the treatment and palliation of a wide spectrum of biliary and papillary diseases. This intervention also helps facilitate subsequent biliary interventions and potentially decreases the risk of post-ERCP pancreatitis when performed prior to biliary stent placement.⁽²⁾ However, it is not without risk. While intraprocedural risks such as bleeding and perforation are well described, the creation of a sphincterotomy also disrupts the inherent sphincter function, potentially leading to increased duodenobiliary reflux and contributing to both stent occlusion and ascending cholangitis.^(1,3) Transpapillary stenting similarly disrupts the sphincter function. In patients with anastomotic biliary strictures, placement of the multiple plastic stents is common practice and leads to high rates of stricture resolution.⁽⁴⁾ However, the development of ascending cholangitis with this approach is significant, with a systematic review confirming an incidence of 19% in LDLT.⁽⁵⁾

In this issue of *Liver Transplantation*, Nishikawa et al.⁽⁶⁾ describe a single-center, observational, retrospective study investigating a novel method of stent placement inside the bile duct (inside-stents) for anastomotic biliary strictures following LDLT. The inside-stents are plastic stents that are placed above the sphincter of Oddi (ie, not transpapillary), obviating the need for sphincterotomy and theoretically decreasing the risk of duodenobiliary reflux.^(1,3) While transpapillary stents can be placed without sphincterotomy,⁽⁷⁾ the technique described in their manuscript theoretically preserves the function of the sphincter entirely, both while the stent is in place and following removal upon stricture resolution. These stents were predominantly self-made until 2017 when commercially available stents became available.⁽⁶⁾

Throughout the study period, inside-stents were successfully placed in 112/136 (82.4%) patients with anastomotic biliary strictures. A total of 19 patients were excluded, leaving 93 patients who underwent inside-stenting. Stent exchanges were performed every 6 months, as needed. Of the 93 patients, 88 (94.6%) had initial stricture resolution, and 70/88 (79.5%) had persistent resolution after a median follow-up of 7.5 years. A total of 17/18 (94.4%) patients with re-stenosis

Abbreviations: DDLT, deceased donor liver transplantation; ERCP, endoscopic retrograde cholangiopancreatography; LDLT, living donor liver transplantation; SEMS, self-expandable metal stent.

Address reprint requests to Allison R. Schulman, M.D., M.P.H., Division of Gastroenterology and Hepatology, Department of Internal Medicine, University of Michigan, 3912 Taubman Center, 1500 East Medical Center Drive, SPC 5362, Ann Arbor, MI 48109-5362. Telephone: 734-936-3444; FAX: 734-936-7392; E-mail: arschulm@med.umich.edu

Allison R. Schulman consults for Apollo Endosurgery, Olympus, Boston Scientific, and MicroTech. She received grants from GI Dynamics.

Received September 27, 2021; accepted September 27, 2021.

Copyright © 2021 American Association for the Study of Liver Diseases

View this article online at wileyonlinelibrary.com.

DOI 10.1002/lt.26322

had new inside-stents placed with eventual resolution. Only 5 (5.4%) patients developed cholangitis during the study period.⁽⁶⁾

The work by Nishikawa et al. reports a novel approach to the treatment of anastomotic biliary strictures with placement of inside-stents and nicely highlights several advantages of this technique. Acceptable rates of stricture resolution were achieved using this strategy, and stent exchanges were only required every 6 months. Typically, transpapillary plastic biliary stents have a maximum dwell time of only 3 months given their propensity to become occluded and cause ascending cholangitis. The longer dwell time and low cholangitis rates with inside-stents further support the possibility that nontranspapillary stenting without biliary sphincterotomy may be mitigating duodenobiliary reflux and reducing stent occlusion and cholangitis. Besides, this strategy may be cost saving due to fewer required ERCPs and fewer interventions necessary during each ERCP. The inside-stents also had a lower migration rate, allowing for safer stenting of smaller intrahepatic bile ducts where SEMs cannot be used routinely.⁽⁶⁾

The exciting results presented in this paper will certainly guide future work in this area. To demonstrate true superiority of inside-stents in the treatment of anastomotic biliary strictures, this approach should be compared with conventional technique in a prospective randomized controlled trial. A study like this could also control for the presence or absence of sphincterotomy in the development of ascending cholangitis or recurrent biliary obstruction in this patient population. Moreover, the commercialization of an inside-stent has precluded the need for self-made stents, which were used during most of the observation period.⁽⁶⁾ A commercialized stent would be more generalizable to the majority of practice settings and should be used in research moving forward.

In conclusion, Nishikawa et al. describe an exciting and emerging approach using inside-stents which obviate the need for a sphincterotomy or a

transpapillary stent. These authors demonstrate low rates of ascending cholangitis with acceptable rates of stricture resolution.⁽⁶⁾ Prospective studies comparing inside-stents using commercially available products with conventional management will be informative to determine whether this approach should be considered first-line in the management of anastomotic biliary strictures.

Jeremy P. Farida, M.D.¹

Allison R. Schulman, M.D., M.P.H.^{1,2}

¹**Division of Gastroenterology and Hepatology**

Department of Internal Medicine

²**Department of Surgery**

University of Michigan

Ann Arbor, MI

REFERENCES

- 1) Tsujino T, Isayama H, Kogure H, Sato T, Nakai Y, Koike K. Endoscopic management of biliary strictures after living donor liver transplantation. *Clin J Gastroenterol* 2017;10:297-311.
- 2) Kato S, Kuwatani M, Onodera M, Kudo T, Sano I, Katanuma A, et al. Risk of pancreatitis following biliary stenting with/without endoscopic sphincterotomy: a randomized controlled trial. *Clin Gastroenterol Hepatol* 2021 Aug 12. <https://doi.org/10.1016/j.cgh.2021.08.016>. [Epub ahead of print]
- 3) Weickert U, Venzke T, König J, Janssen J, Remberger K, Greiner L. Why do bilioduodenal plastic stents become occluded? A clinical and pathological investigation on 100 consecutive patients. *Endoscopy* 2001;33:785-790.
- 4) Hsieh T-H, Mekeel KL, Crowell MD, Nguyen CC, Das A, Aqel BA, et al. Endoscopic treatment of anastomotic biliary strictures after living donor liver transplantation: outcomes after maximal stent therapy. *Gastrointest Endosc* 2013;77:47-54.
- 5) Kao D, Zepeda-Gomez S, Tandon P, Bain VG. Managing the post-liver transplantation anastomotic biliary stricture: multiple plastic versus metal stents: a systematic review. *Gastrointest Endosc* 2013;77:679-691.
- 6) Nishikawa Y, Uza N, Hata K, Marui S, Kuwada T, Matsumori T, et al. Long-term outcomes of stent placement inside the bile duct for biliary strictures after living donor liver transplantation. *Liver Transpl* 2022;28:88-97.
- 7) Wilcox CM, Kim H, Ramesh J, Trevino J, Varadarajulu S. Biliary sphincterotomy is not required for bile duct stent placement. *Dig Endosc* 2014;26:87-92.