






Contrasting recruitment of skin-associated adipose depots during cold challenge of mouse and human

Ildiko Kasza¹, Jens-Peter Kühn² , Henry Völzke³, Diego Hernando^{4,5}, Yaohui G. Xu⁶, John W. Siebert⁷, Angela L. F. Gibson⁷, C. -L. Eric Yen⁸, David W. Nelson⁸, Ormond A. MacDougald⁹, Nicole E. Richardson^{10,11} , Dudley W. Lamming^{10,11} , Philip A. Kern¹²  and C. M. Alexander¹ 

¹McArdle Laboratory for Cancer Research, University of Wisconsin-Madison, Madison, Wisconsin, USA

²Institute and Polyclinic of Diagnostic and Interventional Radiology, Medical Faculty Carl Gustav Carus, Technical University Dresden, Germany

³Institute of Community Medicine, University of Greifswald, Germany

⁴Department of Radiology, University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin, USA

⁵Department of Medical Physics, University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin, USA

⁶Department of Dermatology, University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin, USA

⁷Department of Surgery, University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin, USA

⁸Department of Nutritional Sciences, University of Wisconsin-Madison, Madison, Wisconsin, USA

⁹Department of Molecular & Integrative Physiology, University of Michigan, Ann Arbor, Michigan, USA

¹⁰Department of Medicine, University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin, USA

¹¹William S. Middleton Memorial Veterans Hospital, Madison, Wisconsin, USA

¹²Department of Internal Medicine, University of Kentucky, Lexington, Kentucky, USA

Edited by: Kim Barrett & Yasuhiko Minokoshi

Key points

- Several distinct strategies produce and conserve heat to maintain the body temperature of mammals, each associated with unique physiologies, with consequences for wellness and disease susceptibility
- Highly regulated properties of skin offset the total requirement for heat production
- We hypothesize that the adipose component of skin is primarily responsible for modulating heat flux; here we evaluate the relative regulation of adipose depots in mouse and human, to test their recruitment to heat production and conservation
- We found that insulating mouse dermal white adipose tissue accumulates in response to environmentally and genetically induced cool stress; this layer is one of two adipose depots closely apposed to mouse skin, where the subcutaneous mammary gland fat pads are actively recruited to heat production
- In contrast, the body-wide adipose depot associated with human skin produces heat directly, potentially creating an alternative to the centrally regulated brown adipose tissue

Ildiko Kasza, PhD, studied biology at the Semmelweis Medical University, Budapest, Hungary, and prepared her PhD thesis on the role of the cellular cholesterol exporter, ABCA1, in cholesterol homeostasis in humans. She started her postdoctoral career with Dr CM Alexander at the University of Wisconsin, where she focused on the mechanisms underlying the profound tumour resistance of mice with a syndecan-1 mutation. She discovered a novel link between the cancer resistance phenotype and the depletion of a specific fat depot under the skin, the so-called dermal white adipose tissue, leading to systemic cold stress. More specifically, she is interested in mammalian skin as a key metabolic regulator, as well as its influence on human disease susceptibility.



This article was first published as a preprint. Kasza L, Kühn J-P, Völzke H, Hernando D, Xu YG, Siebert JW, Gibson ALF, Yen C-L E, Nelson DW, MacDougald OA, Richardson NE, Lamming DW, Kern PA, Alexander CM. 2021. Contrasting recruitment of skin-associated adipose depots during cold challenge of mouse and human. bioRxiv. <https://doi.org/10.1101/2020.09.16.300533v2>.

Abstract Mammalian skin impacts metabolic efficiency system-wide, controlling the rate of heat loss and consequent heat production. Here we compare the unique fat depots associated with mouse and human skin, to determine whether they have corresponding functions and regulation. For humans, we assay a skin-associated fat (SAF) body-wide depot to distinguish it from the subcutaneous fat pads characteristic of the abdomen and upper limbs. We show that the thickness of SAF is not related to general adiposity; it is much thicker (1.6-fold) in women than men, and highly subject-specific. We used molecular and cellular assays of β -adrenergic-induced lipolysis and found that dermal white adipose tissue (dWAT) in mice is resistant to lipolysis; in contrast, the body-wide human SAF depot becomes lipolytic, generating heat in response to β -adrenergic stimulation. In mice challenged to make more heat to maintain body temperature (either environmentally or genetically), there is a compensatory increase in thickness of dWAT: a corresponding β -adrenergic stimulation of human skin adipose (*in vivo* or in explant) depletes adipocyte lipid content. We summarize the regulation of skin-associated adipocytes by age, sex and adiposity, for both species. We conclude that the body-wide dWAT depot of mice shows unique regulation that enables it to be deployed for heat preservation; combined with the actively lipolytic subcutaneous mammary fat pads they enable thermal defence. The adipose tissue that covers human subjects produces heat directly, providing an alternative to the brown adipose tissues.

(Received 6 October 2020; accepted after revision 2 March 2021; first published online 16 March 2021)

Corresponding author C. M. Alexander: McArdle Laboratory for Cancer Research, University of Wisconsin-Madison, 1111 Highland Ave, Madison, WI 53705-2275. Email: cmalexander@wisc.edu

Introduction

Specializations of mammalian adipose depots determine their individual impact on human health (Manolopoulos *et al.* 2010; Karastergiou & Fried, 2013; Lee *et al.* 2013; Yoneshiro *et al.* 2013; Pinnick *et al.* 2014; Walker *et al.* 2014). In general, obesity is associated with metabolic abnormalities and the development of a variety of diseases, from diabetes to cancer, yet not all large deposits of adipose tissue types are linked to disease susceptibility (Smith *et al.* 2019). For example, intraperitoneal visceral white adipose tissue (vWAT) is linked to cardiometabolic risk, whereas accumulation of subcutaneous white adipose tissue (scWAT), especially gluteofemoral adipose tissue in the lower body, appears to protect subjects from these health risks (Manolopoulos *et al.* 2010; Karastergiou *et al.* 2012).

Studies of scWAT typically describe the gluteofemoral and/or abdominal depots; however, even high-resolution techniques such as magnetic resonance imaging (MRI) tend to overlook the body-wide fat layer underneath the skin. Our studies have drawn attention to this depot; indeed a preliminary calculation suggested that this body-wide depot comprised at least half of the fat for the average lean woman (Kasza *et al.* 2016).

The term 'subcutaneous white adipose tissue' is an umbrella term that has been used differently by previous studies, and is likely to include a heterogeneous set of depots and adipocyte cell types (Kelley *et al.* 2000; Driskell *et al.* 2014; Walker *et al.* 2014; Kruglikov & Scherer, 2016b;

Nicu *et al.* 2018; Zwick *et al.* 2018). For this study we therefore distinguish a body-wide depot (skin-associated fat, SAF) from subcutaneous fat pads (sc fat pads; Fig. 1A). Subcutaneous fat pads accumulate around the abdomen and upper limbs, particularly in overweight women with gynoid (pear-shaped) fat deposition (Wajchenberg, 2000; Karastergiou & Fried, 2017), and they have been quantified by MRI (Smith *et al.* 2001). However, the body-wide SAF depot has never been quantified.

We are focused on the thermal defence properties of mouse and human SAF. Thus, a layer of fat, applied body-wide to skin, is predicted to impact heat transfer properties dramatically. In turn, this affects the metabolic budget and energy expenditure of mammals; for example, mice housed at temperatures 10°C lower than their thermoneutral temperature show a 50% increase in energy expenditure (Tschop *et al.* 2012; Speakman, 2013; Kasza *et al.* 2014). A thermoneutral environment for mice is warmer than for humans (approximately 31°C compared with room temperature/24°C), and the effect of room temperature housing on the physiology of mice is considerable, suppressing a wide range of inflammatory reactions (Tian *et al.* 2016; Giles *et al.* 2017; Qiao *et al.* 2018).

The dramatic physiological cost of heat production has not gone unnoticed; a major research effort has been directed to devising a strategy for treating the epidemic rates of obesity in westernized populations, by stimulating thermogenic adipose depots (beige/brown) to 'burn' calories, using either direct cold exposure or β -adrenergic

agonists (Cannon & Nedergaard, 2009; Harms & Seale, 2013; Yoneshiro *et al.* 2013; Bartelt & Heeren, 2014; Elattar & Satyanarayana, 2015). The physiology that results from beige adipose activation is different from that arising from brown adipose depots (Keipert & Jastroch, 2014; Keipert *et al.* 2020), since each adipose depot is endocrine and highly specialized (Kershaw & Flier, 2004; Villarroya & Giralt, 2015; Wang *et al.* 2015). It is not known how the relative contribution of each mechanism is determined.

Our previous studies have shown that the SAF depot of mice (dermal white adipose tissue, dWAT) is highly regulated in a manner that suggests it could be part of an acclimation process that mitigates heat loss (Alexander *et al.* 2015). Thus dWAT thickens in mice housed at room temperature (sub-thermoneutral), and mice that are unable to develop thicker dWAT are chronically cold

stressed (Kasza *et al.* 2014). Importantly, several mouse strains with defects of skin lipogenic enzymes lose heat at high rates; interestingly, these strains are also resistant to diet-induced obesity (Sampath *et al.* 2009; Shih *et al.* 2009; Neess *et al.* 2013, 2020; Sampath & Ntambi, 2014; Kruse *et al.* 2017; Kasza *et al.* 2019).

Here, we aim to provide insight into the functional homologies between the skin-associated adipose depots in mice and humans, so that models aimed at testing the role of the regulated properties of skin in systemic metabolism can be built more accurately. We quantify not just skin thickness, but SAF, for human subjects, and find that it is surprisingly variable between individuals. We show that this fat can directly produce heat, suggesting a highly tailored strategy for maintaining body temperature that varies between individuals depending




Figure 1. The thickness of skin-associated fat is highly variable between human subjects, and independent of BMI
 A, cartoon illustrates the use of assay terminology. Ectopic subcutaneous fat pads are coloured yellow; the body-wide depot of skin-associated fat (SAF) coloured dark blue. B, assay of SAF measured using coronal, fat-only MRI views of lower leg, 30% of the distance between knee and ankle. C, results of SAF thickness assay for men ($n = 236$) and women ($n = 286$) as determined by MRI: these data are shown together (left-hand side,) and after dividing the subjects into groups; namely, lean ($BMI < 25$), overweight ($BMI 25-30$) and obese ($BMI > 30$), with the indicated group sizes (right-hand side). Data are represented as a geometric means with 95% confidence intervals. The significance of differences was analysed by Kruskal–Wallis tests corrected for multiple comparisons with a Dunn’s test. D and E, evaluation of correlation between BMI, age and SAF thickness, for men and women, indicated using Spearman’s correlation coefficients. [Colour figure can be viewed at wileyonlinelibrary.com]

upon the size of their SAF depots. We derive data from a genetically cool-stressed mouse model that shows that the accumulation of dWAT is a component of the UCP1-independent heat-generating response, which together with a demonstration that this depot is entirely insensitive to β -adrenergic agonists, supports our claim that dWAT is designed as a functional insulator within mouse skin. Overall, we show that the mouse skin-associated adipose depots comprise two distinct types of fat, one insulating (dWAT), and the other a lipolytic thermogenic depot (scWAT). In contrast, the human skin-associated adipose tissues are surprisingly homogeneous with respect to their responsiveness to thermogenic cues, regardless of depth from skin, or body site; indeed we could find no evidence for extensive colonization of human skin by a mouse dWAT equivalent.

Methods

Ethical approval

The animal studies were performed in strict accordance with the recommendations in the *Guide for the Care and Use of Laboratory Animals of the National Institutes of Health*. Experimental protocols were approved by the University of Wisconsin School of Medicine and Public Health Animal Care and Use Committee (IACUC approval#M005471). The number of mice used to perform this study was minimized, and every effort was made to reduce the chance of pain or suffering. The method of killing was CO₂ asphyxiation, as per the guidelines. All authors understand the ethical principles operating at *The Journal of Physiology* and confirm that this work complies with the animal ethics checklist. The human subjects submitted their informed consent for the Study of Health in Pomerania (SHIP) study as part of the Community Medicine Research Net of the University of Greifswald, Germany; the University of Greifswald is a member of the 'Centre of Knowledge Interchange' programme of Siemens AG, and this study was compliant with the *Declaration of Helsinki*, except for registration in a database. The SHIP study is a longitudinal study of health parameters of over 4000 adults (ages 20–79 years), representative of an area with a population of >200,000 (Volzke *et al.* 2011). All analysis of the SHIP data analysed here was HIPAA-compliant and performed after obtaining approval from the ethics committee of the University of Greifswald (Germany) and the University of Wisconsin Institutional Review Board (IRB approval#2019-0082). Human skin specimens were taken from discarded tissues during defect reconstruction and body sculpting procedures and submitted, anonymously, to the BioCore Biobank (IRB approval# 2016-0934-CP057).

Mice

BALB/cJ (#00651), C57BL/6J (#00664), Agouti lethal yellow (A^y/a; C57BL/6J;#002468) and UCP1-/C57BL/6J;#003124) mice were obtained from Jackson Labs and bred in-house. All animals were housed at constant temperatures (19–23°C) in 12 h light/dark cycles with free access to water and standard chow (Harlan Teklad Global Diet 2020X). For high-fat feeding, C57BL/6J mice were provided with Harlan Teklad diet#TD06415 (45% calories from fat); for calorie restriction, *ad libitum* consumption rates of chow were measured for singly housed mice; then mice were 25% calorie restricted using a 60% mix of AIN93M:40% calorie restriction (Bioserv cat#F05314) in control diet (AIN93M, Bioserv cat#F05312), pair-fed daily at 5 pm, for 5 weeks.

Histological analysis

Skin, brown adipose tissue (BAT), perigonadal white adipose tissue (vWAT) and mammary gland (iWAT) were dissected for histological processing as follows. Samples were paraformaldehyde-fixed (4%) overnight at 4°C and then paraffin-embedded for evaluation. Tissue sections were deparaffinized, rehydrated and stained with either standard haematoxylin and eosin (H&E) or Trichrome protocols for visualization. For assay of dWAT thickness, six images of H&E-stained, non-anagen fields of skin (equivalent to $\geq 4500 \mu\text{m}$ linear dWAT) were assayed by image analysis (dividing total area by length). For immunofluorescent analysis, tissues were processed for heat-induced epitope retrieval, epitopes were blocked in 10% goat serum for 3 h; samples were either incubated with Alexa-conjugated primary antibodies for 1 h, or incubated overnight with the primary antibody, washed and incubated for 1 h with secondary antibodies. Samples were visualized on a confocal microscope (Nikon A1RS Confocal Microscope). For fluorescent intensity assessment, at least three independent fields were obtained for each mouse sample, and signal was quantified using an open-source Fiji image processing package. In each case, the specific signal was normalized to the signal from lipid droplet-associated perilipin. For quantification of lipid droplet size, six independent fields were obtained and quantified using the open source Fiji image processing package (<https://loci.wisc.edu/software/fiji>). Antibodies and immunohistochemical reagents were as follows: anti-CD31 (#3528; RRID:AB_2160882), anti-CD31 (#77699; RRID:AB_2722705), anti-Fabp4 (#3544; RRID:AB_2278527), anti-FASN (#3180; RRID:AB_2100796), anti-hormone-sensitive lipase (HSL) (#4107; RRID:AB_2296900), anti-pS565 HSL (#4137; RRID:AB_2135498), anti-pS660 HSL (#4126; RRID:AB_490997), anti-pS 235/226 S6 (#2211;

RRID:AB_331679), all from Cell Signalling Technology; Alexa Fluor488 anti-perilipin (#NB110-40760AF488; RRID:AB_2167264) and AlexaFluor647 anti-perilipin (#NB110-40760AF647; RRID:AB_1849889) both from Novus Biologicals; anti-UCP1 (#ab10983, RRID:AB_2241462; Abcam); secondary reagents were AlexaFluor546 goat anti-rabbit (#A-11035; RRID:AB_143051), AlexaFluor546 goat anti-mouse (#A-11030; RRID:AB_144695) and AlexaFluor633 wheat germ agglutinin (#W21404) from Thermo Fisher Scientific. Specificity of antisera was confirmed by immunohistochemical evaluation of positive controls.

3T3L1 cell culture

Mouse 3T3-L1 preadipocytes were from the American Tissue Culture Collection (ATCC; RRID: CVCL_0123) and were maintained in Dulbecco's modified Eagle's medium supplemented with 4.5 g l⁻¹ of glucose (Life Technologies), 10% fetal bovine serum and 100 U ml⁻¹ penicillin and streptomycin. Briefly, cells were differentiated into adipocytes using MDI medium (100 mg ml⁻¹ 3-isobutyl-1-methylxanthine, 100 ng ml⁻¹ dexamethasone and 1 mg ml⁻¹ insulin, both from Sigma) for 4 days, followed by 1 mg ml⁻¹ insulin for an additional 4 days (Kasza *et al.* 2014), and fixed for immunofluorescent staining using 3% paraformaldehyde (20 min/4°C).

β -adrenergic induction

The pan β -adrenergic agonist, isoproterenol hydrochloride (ISO) was from Sigma; the β 3-adrenoreceptor agonists, CL 316243 disodium salt (CL) was obtained from Tocris Bioscience (UK) and mirabegron (Mira) was from Cayman Chemical. For CL administration *in vivo*, mice were acclimated to thermoneutrality for 3 days, injected with 1 mg kg⁻¹ CL, and killed for tissue collection 60 min later. For β -adrenergic stimulation *in vitro*, 3T3-L1 cells were treated with ISO (10 μ M) for 40 min at 37°C. For β -adrenergic stimulation *ex vivo*, human fat samples (approx. 0.5 cm in size) were treated with ISO (10 μ M) for 40 min at 37°C. For β -adrenergic stimulation *in vivo*, human subjects were administered Mira for 10 weeks (50 mg day⁻¹), with abdominal adipose biopsies before and after treatment, as described previously (Finlin *et al.* 2018).

FLIR imaging

To measure surface temperatures by infrared thermography, we used a hand-held FLIR T360 camera (FLIR Systems, Oregon). Pin drops in the software of the FLIR

camera were used to record surface temperatures and quantified using FLIR Tools Advanced Thermal Analysis and Reporting software; each photograph is internally and externally calibrated to show actual temperatures (Kasza *et al.* 2019).

SHIP study design and acquisition of MRI images of skin-associated fat

The SHIP study is a longitudinal study of health parameters for over 4000 adults (ages between 20 and 79 years), representative of an area with a population of >200,000 (Volzke *et al.* 2011). The assay of the thickness of SAF was performed using a chemical shift-encoded 3D gradient-echo sequence of the whole-body (1.5 T MR images, Avanto, Siemens Healthcare), with the following parameters: repetition time = 12 ms, echo time (s) = 2.38 ms, flip angle = 5°, spatial resolution 1.95 mm \times 1.95 mm \times 5 mm, acquisition matrix 0/256/128/0, pixel bandwidth 1955 Hz/Pixel of each series with complete imaging of both calves. MR sequences were postprocessed to separate confounder-corrected water- and fat-only images. Confounder-corrected fat-only images were used to measure SAF (Kuhn *et al.* 2017). Skin-associated fat was measured at the lateral side of both legs, 30% of the distance between tibia and ankle diaphysis.

Collection of human samples

Breast tissues were collected during breast reduction surgeries by the University of Wisconsin Carbone Cancer Center Translational Science Biocore Biobank. Human skin samples (with associated fat) were obtained from patients undergoing elective reconstructive surgeries at our institution. The de-identified samples were exempt from the regulation of the University of Wisconsin-Madison Human Subjects Committee Institutional Review Boards. Data on patient age, sex and type of surgery were collected with the tissues. Skin samples were processed within 3 h of surgery.

Statistical analysis

Analyses were conducted using GraphPrism8 software, and the statistical tests appropriate to each analysis are indicated in the figure legends. To test for normal or lognormal distribution of sample values we used the Anderson-Darling test, outliers were identified using ROUT method. Box-and-whisker graphs show median values with 5–95 percentile whiskers; other data are expressed as means \pm standard deviation, unless specifically stated.

Results

To distinguish only the skin-associated body-wide depot (SAF), we used chemical shift MRI of lower limb, with fat/water separation (Fig. 1B). Fat-only images allow quantitative evaluation of fat content and SAF thickness, compared with the more standard fat/water MRI images (data not shown). This methodology is focused specifically on the fat content of skin, rather than the thickness of skin, which can be measured by skin-fold calipers or ultrasound (Perez-Chirinos Buxade *et al.* 2018; Storchle *et al.* 2018). These other techniques are not designed to determine only the thickness of fat.

In a previous pilot study we found that SAF thickness was highly variable from subject to subject, and surprisingly, not related to body mass index (BMI) (or other indices of obesity such as waist/height) (Kasza *et al.* 2016). Here, we assessed MRI images of volunteers recorded as part of a large public health study (Study of Health in Pomerania, SHIP3), comprising 286 women and 236 men. We found that women had thicker SAF than men (9.75 ± 1.41 compared with 5.95 ± 1.27 mm; Fig. 1C). Indeed, the SAF thickness for the top decile of women was 15.0 mm, *versus* 6.5 mm in the bottom decile; this translates to a total predicted weight of 24.3 kg for women with the thickest SAF layer, *versus* 10.5 kg for women with the thinnest SAF, approximated using the surface area (1.8 m^2) of the average woman (Kasza *et al.* 2016). For men, this range was 9.6 mm to 3.0 mm thick (15.5–4.9 kg).

We assessed whether the thickness of SAF depended upon general adiposity, dividing the cohorts into three: lean (BMI <25), overweight (BMI 25–30) and obese (BMI >30; Fig. 1C). Skin-associated fat did not thicken significantly in obese or overweight men. For women, there was a significant, if minor, increase in SAF thickness for lean *versus* overweight women (9.00 ± 1.73 compared with 10.41 ± 2.49 mm); which did not increase further in the obese cohort. These conclusions were restated by correlation analysis (Fig. 1D), showing no relationship of BMI (or waist/hip ratio; data not shown) with skin SAF for males, and a weak relationship for females ($r = 0.300$, $P < 0.0001$).

We tested the distributions of values for SAF thickness for men and women, and found that for men, the values were normally distributed; for women, the values were not, best fitting a lognormal distribution, and tailing towards thicker SAF (Fig. 1C). This implies an independent factor that promotes the accumulation of SAF in a minority of women. Perhaps surprisingly, we noted no significant correlation of SAF thickness with age, for either men or women (55 men and 57 women over 70; Fig. 1E).

To assess similarities and differences between mouse and human skin-associated adipose depots, we first

describe their morphology and anatomy. Mouse dWAT is clearly demarcated by a muscle layer (*panniculus carnosus*), making it simple to distinguish from the subcutaneous depots of the mammary glands (females) or mammary fat pads (males) (Alexander *et al.* 2015). The most commonly studied scWAT depot in mice is the inguinal fat pad, or iWAT; this is immediately adjacent to the dWAT layer (proximity is illustrated in Fig. 8).

Overall, the thickness of dWAT varies from almost zero (young C57BL/6J males or rats) to 400 μm , typical of anagen stage dorsal skin for BALB/cJ mice, or obese mice (Fig. 2A, B, D, E). The average thickness for non-anagen stage dorsal skin from chow-fed adult BALB/cJ females housed at room temperature is 350 μm (Fig. 2D).

We noticed that dWAT accumulates steeply in C57BL/6J males between the ages of 20 and 30 weeks; this occurs prior to a phase of rapid continuous body weight gain (Fig. 2B). We note that this reaction coincides with the loss of beiging potential in iWAT (Berry *et al.* 2017), and precedes their mid-life obesogenesis, suggesting a general evolution of metabolic efficiency and heat production strategies during this period.

Using an immunohistochemical stain to outline the fat globule of adipocytes (anti-perilipin), we found that mouse dermal adipocytes contain approximately the same amount of fat as inguinal/subcutaneous WAT adipocytes but show only half the cross-sectional area compared with visceral adipocytes (Fig. 2C). Assuming the fat globule is approximately spherical, this translates to approximately 4-fold higher lipid load in each visceral adipocyte compared with the adipocytes in iWAT and dWAT depots.

We compared the thickness of dWAT in males and females from two strains, C57BL/6J and BALB/cJ mice, and found that BALB/cJ mice showed a dramatic sex dimorphism, where dWAT was almost four times thicker in females than males (Fig. 2D). This was not observed for mature male and female C57BL/6J mice, though mature females showed high variability compared with males.

Bearing in mind the lack of correlation between human SAF thickness and obesity, we tested whether obesity and dWAT thickness were correlated in mice. We found that dWAT thickness expanded by 4-fold after 2 weeks of high-fat feeding for C57BL/6J males, in parallel with vWAT depots (Fig. 2E and Kasza *et al.* 2016). Likewise, genetically obese mice (A^y/a) showed thick dWAT and increased dermal adipocyte size; diameters of dermal adipocytes (obese and lean) are summarized in Table 1.

Inflammation of obese adipose depots is linked to metabolic deterioration, and adipose depots from obese mice are differentially sensitive to invasion by inflammatory macrophages, assayed by the accumulation of crown-like structures (Grove *et al.* 2010). We investigated whether crown-like structures appeared in dWAT depots in skin from mice fed with high-fat




Figure 2. Determinants of rodent dWAT thickness: age, sex, hair cycle and obesity

A, hair cycle-associated expansion. Representative H&E-stained sections of skin of rats and mice in non-anagen and anagen (follicular development) phases. (a) dermis/epidermis; (b) dWAT; (c) panniculus carnosus; (d) hair follicle. Scale bars = 100 μm . **B**, maturation dependent accumulation. Assay of dWAT thickness during the growth and maturation of dorsal skin of male C57BL/6J mice; $n = 65$. **C**, dermal adipocytes match inguinal adipocyte size. Comparison of the sizes of fat globules in dermal (dWAT), inguinal (subcutaneous; iWAT) and visceral (vWAT) adipocytes from lean mice, as measured by immunofluorescent visualization using anti-perilipin (PL); $n = 6$. Scale bars = 100 μm . **D**, thickness of dWAT is highly sex- and strain-dependent. dWAT thickness was measured from H&E-stained sections of skin from C57BL/6J or BALB/cJ male and female mice >30 weeks old ($n = 6$). Data for panels **B** and **D** were analysed by unpaired two tailed t test; groups compared for panel **C** were analysed by one-way ANOVA followed by Tukey's multiple comparisons test. **E**, number and size of dermal adipocytes increases in obese mice. Representative H&E sections of dorsal skins from C57BL/6J male mice (7–11 weeks; $n = 6$), fed chow or a high-fat diet for 5 weeks, or from a genetically obese model (1-year-old C57BL/6J- Ag^y ; $n = 3$). Average diameters are shown in Table 1. Scale bars = 100 μm . [Colour figure can be viewed at wileyonlinelibrary.com]

Table 1. Summary comparison of key features of skin-associated fat depots for mouse and human

Parameter	Mouse dWAT	Human skin-associated fat (SAF)	Comments
Morphology	Segregated from nearest adjacent scWAT depots by muscle layer.	Not partitioned by muscle: partitioned instead into collagen-wrapped 'bubble wrap'.	Driskell <i>et al.</i> 2014; Alexander <i>et al.</i> 2015; Kruglikov & Scherer, 2016b
Thickness	0–400 μm Varies depending upon age, sex, strain, adiposity.	3–15 mm Highly variable from subject to subject.	Dimensions quoted for lean subjects.
Size of adipocytes (diameter)	40 \pm 1.7 μm /lean mice 59 \pm 5.5 μm / obese mice \approx 50% increase Adipocytes from iWAT and dWAT are similar size, and half the size of vWAT adipocytes.	56 \pm 1.1 μm /lean subjects 67 \pm 0.7 μm /obese subjects \approx 20% increase Subcutaneous and visceral adipocytes are similar sizes.	Combined data from human, for several body sites (breast, abdominal, visceral, thigh; lean, $n = 7$, obese $n = 8$) and for mice, either genetically obese (A ^y) or fed high-fat diet ($n = 9$, obese; $n = 26$, lean).
Reaction to obesity	Thickens; size of adipocytes increases. Resistant to macrophage infiltration (crown-like structures).	No change of thickness in men; marginal change in women (<10%). Susceptible to macrophage infiltration (Cappellano <i>et al.</i> 2018).	Size of human abdominal scWAT adipocytes increases in obese subjects (Verboven <i>et al.</i> 2018; Finlin <i>et al.</i> 2019). High-fat diet feeding leads to expansion of mouse dWAT adipocytes (Kasza <i>et al.</i> 2016; Zhang <i>et al.</i> 2019).
Reaction to calorie restriction	Becomes depleted in parallel with other adipose depots (in C57BL/6J males), although approx. half of dWAT is protected.	Not known. However, lean and obese men show the same thickness of SAF.	Little association of low or high body-wide adiposity with SAF thickness in humans.
Rate of accumulation and involution	Highly dynamic in response to hair growth, skin infection, environmental temperature.	Unknown	Alexander <i>et al.</i> 2015; Zwick <i>et al.</i> 2018
Male/female dimorphism	Thicker in females than males for BALB/cJ mice, but not C57BL/6J mice.	Thicker in females than males.	
Age-dependent maturation	Thickness increases steeply from 20 to 300 μm from 20–30 weeks of age for C57BL/6J males before, and during, a phase of rapid weight gain. Other mice do not show this regulation.	No change of thickness for SHIP study subjects (both male and female), between ages 20 and 79.	
Molecular reaction to β-adrenergic stimulants	No significant reaction of lipolytic hub enzyme (activating pS ⁵⁶⁶⁰ HSL and inhibitory pS ⁵⁶⁵ HSL), or mTOR activation (pS ^{235/236} S6); no significant lipid depletion assayed from lipid droplet size.	Majority of skin-associated adipocytes (scWAT fat pads or SAF) react (including HSL and S6 activation), and become lipid depleted, measured from adipocyte size after <i>ex vivo</i> stimulation.	HSL activation corresponds to active lipolysis, assayed by adipocyte shrinkage. Little functional heterogeneity of human depots.

(Continued)

Parameter	Mouse dWAT	Human skin-associated fat (SAF)	Comments
Functional reaction to cooling	Thickens in response to activated thermogenesis (room temperature housing), and in mice with highly induced non-canonical thermogenic processes (UCP1KO). Accumulation depends upon timing of cool exposure.	Lipolysis is activated in majority of skin-associated adipocytes, and SAF is therefore predicted to become depleted. Thermogenic activation of scWAT directly induces heat production.	Skin-associated adipocytes of human and mouse have opposite responses to sub-thermoneutral environments: mouse dWAT increases to oppose heat loss; human SAF becomes lipolytic and actively participates in heat generation.

diets (compared with either iWAT or gWAT) and found that this depot was relatively resistant to this type of inflammation. In extremely obese mice, some crown-like structures developed (32 week-old A^y mice; data not shown).

Vice versa, calorie-restricted mice showed depleted dWAT. We used a moderate, 25% calorie restriction protocol to assess the impact on both C57BL/6J and BALB/cJ mice; this protocol induced a significant loss of body weight within 2 weeks. After 4 weeks of calorie restriction, C57BL/6J mice lost 18% body weight, and BALB/cJ mice lost 22% body weight (Fig. 3A). dWAT mirrored the loss of body weight; both strains lost >100 μm thickness of dWAT (Fig. 3B), so the leanest calorie-restricted BALB/cJ mice showed negligible dWAT levels after 4 weeks.

In contrast, human subcutaneous adipose tissues comprise capsules of yellow fat attached to the dermis/epidermis; capsule size ranges from 5 to 15 mm, where the lipid-filled adipocytes are encapsulated by thick seams of collagen (Fig. 4A). Several publications have reviewed the topic of potential heterogeneity of human subcutaneous adipose tissue (Kelley *et al.* 2000; Smith *et al.* 2001; Sbarbati *et al.* 2010; Kruglikov & Scherer, 2016a; Zwick *et al.* 2018); in particular, deep layers of SAF may have different properties compared with the more superficial layers (Enevoldsen *et al.* 2001; Cappellano *et al.* 2018). We found little difference in the size of adipocytes from superficial (within 5 mm of dermis), mid- (5–15 mm) and deeper layers (more than 15 mm from dermis; $n = 7$) for samples from lean individuals (Fig. 4A, B); neither did we observe significant differences in adipocyte sizes from different body locations (for example breast *versus* lower body (thigh) *versus* upper body (abdominal); $n = 7, 4, 5$, respectively) (Fig. 4B, C). The adipocytes from visceral depots and breast and abdominal subcutaneous sites all underwent hypertrophy (Fig. 4C). In summary, human subcutaneous adipocytes




Figure 3. Calorie restriction depletes dWAT

A, body weight changes for C57BL/6J or BALB/cJ male mice (28 to 30 weeks old) after 25% calorie restriction for 4 weeks (red arrow indicates start of dietary regime). Data were analysed by two-way ANOVA followed by multiple comparisons test ($n = 6$). B, dWAT thickness was assessed from H&E sections of dorsal skin. Data were analysed by unpaired two-tailed t tests. Scale bars = 300 μm . [Colour figure can be viewed at wileyonlinelibrary.com]




Figure 4. Like mouse dermal adipocytes, human skin-associated adipocyte hypertrophy in obese subjects

A, gross anatomy and histology of human skin-associated fat (SAF). Gross appearance and trichrome-stained sections of SAF from breast of a lean subject, with detail insets shown for dermis, superficial, mid- and deep layers of adipocytes. Scale bars = 5 mm on gross view, 2 mm on Trichrome-stained low power view, and $100 \mu\text{m}$ on high power detail insets. **B**, size of adipocytes is unaffected by distance from skin. Human thigh tissue sections were stained with perlipin antibody ($n = 3$); representative images are shown, with quantitation of the size of adipocytes from superficial, mid- and deep layers. Scale bars = $100 \mu\text{m}$. **C**, comparison of SAF from breast and abdomen with visceral WAT. Representative images of PLIN-stained fat globules from lean and obese subjects (left) were quantified (right); scale bars = $100 \mu\text{m}$. **D**, adipocytes of mouse dWAT and human SAF respond similarly to obesogenesis. The relative size of adipocytes in human SAF and mouse dWAT are shown for lean and obese subjects (human lean, $n = 5$; human obese, $n = 7$; mouse lean, $n = 7$; mouse obese, $n = 5$). Data were analysed using a one-way ANOVA followed by Tukey's multiple comparisons test. [Colour figure can be viewed at wileyonlinelibrary.com]

of lean subjects were larger on average (56 μm diameter) than mouse dermal adipocytes (40 μm diameter), and adipocytes from both human and mouse undergo hypertrophy during obesogenesis (67 and 59 μm , respectively; Fig. 4D; Table 1).

Human and mouse subcutaneous adipose depots are typically ascribed a supportive role in the thermogenic response; they are activated by β -adrenergic agents to become lipolytic, to provide fatty acid fuels for heat production by BAT, or to participate in the beiging response (Wu *et al.* 2012; Mottillo *et al.* 2014; Chondronikola & Sidossis, 2019). However, the analysis of the transcriptome of mouse dermal adipocytes performed by Scherer and colleagues suggested that this population was unlike adipocytes in brown or beiging depots, resembling vWAT instead (Zhang *et al.* 2019); they showed that dWAT showed no lipid depletion upon treatment with a β -adrenergic agent (scored as a reduction from mono- to multi-locularity of the lipid body).

To further evaluate the molecular response of dermal white adipocytes to cold stress, we evaluated the regulatory modification of HSL in response to the β -adrenergic agonist CL *in vivo* using an immunohistochemical assay (Fig. 5). HSL is part of a complex that integrates input stimuli from kinases to coordinate lipolysis of brown adipose tissue (Fruhbeck *et al.* 2014; Ogasawara *et al.* 2015). We confirmed the accuracy of HSL (and FASN) stains using mouse adipocyte cultures treated with the pan β -adrenergic agonist, isoproterenol (data not shown). Tissue samples of BAT and iWAT showed a gain of signal for the PKA-dependent activating phosphorylation of S⁶⁶⁰ (pHSL stim), whereas vWAT showed a gain of signal for the AMPK-dependent inhibitory phosphorylation of S⁵⁶⁵ (pHSL inh; Fig. 5A, B). Positive staining for fatty acid synthase, FASN and fatty acid binding protein-4, FABP4, was suppressed in iWAT and BAT, and unchanged/negative for vWAT and dWAT (data not shown). This confirms that dWAT was non-responsive to β -adrenergic-induced heat production, measured by the same criteria used to assess the recruitment of brown and beige depots.

To corroborate this result, we tested another β -adrenergic/PKA-induced reaction; thus, activation of mTORC1 is known to be essential for β -adrenergic-induced lipolysis in adipocytes (Liu *et al.* 2016). S6 and S6-kinase are activated and phosphorylated downstream of the mTORC1 kinase and are markers of mTORC1 activation status. Signal for phospho-S^{240/244} S6 was induced in BAT and iWAT in CL-administered mice but showed no reaction in dWAT (Fig. 5A), despite the local activation of muscle and sebocytes by β -adrenergic stimulation. To confirm that these immunohistochemical assays reflect active lipolysis, we measured the size of adipocyte lipid stores in mice that were activated to produce heat, either by transfer to 4°C environment or

60 min after CL administration. We found lipid stores significantly depleted in iWAT, but not in dWAT (Fig. 5C). In summary, when BAT and iWAT become lipolytic in response to β -adrenergic stimulation, dWAT shows little molecular response, like vWAT and bone marrow adipocytes (Scheller *et al.* 2019).

We hypothesized that in mice chronically stimulated to produce more heat, dWAT would instead react to mitigate heat loss. Mice housed at room temperature (19°C) compared with thermoneutral housing (31°C) show thicker dWAT (1.6 \times ; Fig. 6A). To test this hypothesis using a genetic model, we evaluated *UCP1*^{-/-} mice. These mice have a lesion in the most efficient heat production mechanism (affecting the uncoupling of BAT mitochondria), and deploy instead beiging and other 'non-canonical' mechanisms to compensate (Keipert *et al.* 2020). As with environmentally cool-stressed mice, genetically cool-stressed mice also show thicker dWAT (Fig. 6B). This supports the claim that the accumulation of fat in dWAT is an adaptive response to environmental cooling.

What triggers this adaptive response? In particular, we noticed that mammals typically experience variation of environmental temperature over the circadian daily cycle. To begin to isolate environmental cues, we divided the housing cycle into a warm phase (thermoneutral, 31°C/88°F) and a cool phase (cool room temperature, 19°C/66°F). Although the length of time in each phase was constant for mice in cycling environments, only the mice housed cool during their sleep cycle developed thicker dWAT (Fig. 6C); the same cue given to awake mice did not induce dWAT thickening. This suggests that there is a component of circadian control of dWAT lipogenesis.

Turning to the molecular regulation of SAF in humans, we administered β -adrenergic agonists either *in vivo* or *ex vivo*, and assayed HSL activation and lipid store mobilization. Human abdominal, breast and thigh SAF administered the pan- β adrenergic agent isoproterenol *ex vivo* showed uniform activation of HSL and lipid store depletion (Fig. 7A, B). Likewise, abdominal adipose tissues from obese subjects administered the β 3-adrenergic agent mirabegron daily, also showed activated HSL, and higher UCP1 staining (data not shown and (Finlin *et al.* 2018)). Skin-associated fat in calf, thus the same body site imaged for the data of Fig. 1, also showed HSL activation after exposure to isoproterenol. The β -adrenergic response was therefore a consistent feature of human skin-associated adipose tissues, regardless of body site or distance from skin (superficial or deep).

Typically, lipolysis of subcutaneous fat is considered to have a supporting role for the delivery of fatty acids to dedicated, heat-generating organs such as brown adipose tissue or muscle. However, emerging literature suggests that a number of UCP1-independent biochemical




Figure 5. Mouse dWAT is entirely insensitive to β -adrenergic lipolysis

A, HSL and mTOR are not activated in dWAT after treatment with a β -adrenergic agonist. Fat depots (brown adipose tissue, BAT; iWAT and vWAT and dWAT) were harvested from mice, killed 60 min after administration of CL (control $n = 5$ and treated $n = 4$). Immunofluorescent assay of regulatory modifications of hormone-sensitive lipase (HSL), either pS⁶⁶⁰ (pHSL stim) or pS⁵⁶⁵ (pHSL inh), or mTOR-activated S6 (pS^{235/236}) were quantified in (B) as described in Methods. Insets shown for pS6-stained dWAT show sebaceous glands (a, unstimulated; c, stimulated) and the panniculus carnosus muscle layer (b, unstimulated; d, stimulated); scale bars = 20 μ m. C, adipose globules of dWAT are not depleted after CL administration. The size of lipid droplets (outlined by anti-perilipin staining) was assayed in dWAT and iWAT, 60 min after CL injection; scale bars = 20 μ m. Data were analysed by unpaired two-tailed *t* tests. [Colour figure can be viewed at wileyonlinelibrary.com]

pathways can be harnessed for the purpose of regulated heat production (Chouchani *et al.* 2019), where the substrates used to fuel oxidation are not yet clearly defined. Therefore, to directly evaluate whether β -adrenergic activation can induce heat production in SAF, we used infrared thermography (FLIR) to assay fat explants *ex vivo* and found a significant rise in temperature in response to isoproterenol treatment (Fig. 7C). We conclude that human skin-associated adipose tissue is an active participant in heat production, and not just a passive insulator.

Discussion

We compared the most superficial layers of adipose tissue associated with the skin of humans and mice and showed that they respond differently to the activation of thermogenesis; the dermal adipocytes associated with mouse skin are protected from lipolysis, where the SAF of human skin becomes lipolytic and produces heat autonomously (Table 1). Thus, the human SAF layer comprises a dynamically warming blanket, not only able to resist heat loss, but actively contributing

to the total heat budget (see scheme in Fig. 8). Our data show that human skin-associated adipocytes are remarkably homogeneous in their ability to recognize and respond to β -adrenergic effectors, whether from obese or lean subjects, and irrespective of body site (upper *versus* lower body, or limb *versus* abdominal depots). A speculative review has suggested that there could be a dWAT equivalent around human hair follicles (Kruglikov & Scherer, 2016a), but we could find no β -adrenergic-resistant adipocyte population in the array of human samples we evaluated. Note that we use the umbrella terminology of ‘skin-associated fat’ for humans, aware that there is an extensive literature documenting site-specific regulation and properties of subcutaneous adipose tissues.

This study is the first to quantify this specific human fat depot, though many MRI-based studies have reported segregated fat volumes that discriminate between, for example, visceral and subcutaneous abdominal fat (Wajchenberg, 2000; Smith *et al.* 2001). Specific ‘fat-only’ MRI data can be extracted from the data obtained for routine diagnostics, though those scans are typically processed to show a combination of fat and water signals.




Figure 5. Continued

Other techniques aimed at assessing body composition use ultrasound or calipers to measure skin-fold thickness (Perez-Chirinos Buxade *et al.* 2018; Storchle *et al.* 2018); however, although these techniques may show broadly comparable results, they do not measure fat specifically.

Other imaging strategies used for the study of thermogenesis rely on the detection of glucose uptake by

^{18}F -fluorodeoxyglucose (FDG)-PET imaging of activated fat depots in cool-exposed individuals. The SAF depot has not yet been noted by these studies (Chen *et al.* 2016). This could be due to 1) the relatively lower resolution of this technique and the disseminated nature of the depot, or 2) low glucose uptake by activated SAF. This is possible, since we do not yet understand the substrate requirements of




Figure 6. Mouse dWAT accumulates when mice are challenged to produce heat

A, dWAT thickens in response to sub-thermoneutral environmental housing. dWAT thickness was assayed for mice housed at thermoneutrality (31°C) or room temperature (19°C) for 3 weeks; scale bars = 100 µm. B, dWAT thickens in UCP1^{-/-} mice producing heat by non-canonical means. H&E-stained sections are shown for iWAT and dWAT from UCP1^{-/-} or control C57BL/6J mice (10 to 19 weeks old); quantified on the right-hand side. Scale bars = 50 µm (iWAT) or 100 µm (skin). C, circadian cues promote dWAT accumulation. Mice were housed individually with alternating circadian cycles of thermoneutral and room temperature housing for 3 weeks. The experimental paradigm is shown on the left, continuous warm (31°C, red line), continuous cool (19°C, blue line), or alternating daily cycles of warm sleeping/cool-waking phases (black) or cool sleeping/warm waking phases (grey). The thickness of dWAT is shown on the right ($n = 5$). Data were analysed by unpaired two-tailed t tests. [Colour figure can be viewed at wileyonlinelibrary.com]

heat production in non-BAT thermogenic tissues, but they probably include creatine and amino acids (Ikeda *et al.* 2017; Mills *et al.* 2018; Kazak *et al.* 2019).

Women show thicker SAF than men, by 60% on average; by calculation this depot is 10.5–24.3 kg, making it the largest individual adipose depot in the lean woman's body. The number of women with thick SAF is higher than that predicted from a normal distribution, and we speculate that the health and energy expenditure of this population may be significantly affected. Likewise, the females of the BALB/cJ mouse strain show much thicker dWAT than their male counterparts; this depot is thus more prominent in females of both species.

The dWAT depot of mice correlates strongly with relative adiposity elsewhere, even from mouse to mouse in a given population. Mouse dWAT increases in obese mice, whether induced by high-fat feeding, by age-induced obesity, or in mice made obese genetically, including *Agouti^v*, MitoNEET and *ob/ob* (Kasza *et al.* 2016,2019; Zhang *et al.* 2019; and this study). In adult mice, the

majority of dWAT thickening is enabled by adipocyte hypertrophy rather than hyperplasia (Zhang *et al.* 2019).

In obese mice, we found this depot to be relatively resistant to the development of 'crown-like structures'; these reflect the infiltration of inflammatory macrophages into depots containing distended adipocytes, typically highest in male mouse vWAT (Grove *et al.* 2010). For human subjects, subcutaneous adipose depots are significantly associated with inflammatory macrophage infiltration (Cappellano *et al.* 2018).

Vice versa, we showed that dWAT is depleted in mice subjected to calorie restriction. Trajkovski and colleagues showed that beige depots were activated in calorie-restricted mice (Fabbiano *et al.* 2016); this may support the hypothesis that dWAT thinning leads to higher demand for heat.

Given *ad libitum* calorie intake, we propose that mice deploy the combination of iWAT activation and dWAT thickening to meet heat production demands, notably when BAT activation is insufficient. Thus UCP1KO mice




Figure 7. Evaluation of β -adrenergic lipolytic response for human skin-associated fat depots

A, human skin-associated adipocytes all respond to β -adrenergic agonists. Immunofluorescent stains of pHSL-stim and pHSL-inh in sections of skin-associated fat from thigh, exposed to isoproterenol (Iso) *ex vivo*, counterstained with perilipin (PL) and DAPI (nuc; nuclei) ($n = 3$). Scale bars = 20 μm . B, human skin-associated adipocytes are depleted by thermogenesis. The average size of lipid droplets was quantified for the adipocytes visualized in A. C, human skin-associated fat makes heat upon treatment with β -adrenergic agonist. Infrared thermographic images of freshly collected fat capsules, pretreated or not with isoproterenol for 40 min, were quantified (right-hand side). Scale bar = 5 mm. Data were analysed using unpaired two-tailed *t* tests. [Colour figure can be viewed at wileyonlinelibrary.com]

deploy a range of strategies for heat conservation and production (Chouchani *et al.* 2019), to ensure their body temperature is resilient to change upon cool stress. As well as thicker dWAT (shown here), UCP1 KO mice show a change of vascular tone, so mice have a sustained vasoconstrictor response after minor cool stress (Kajimura *et al.* 2015). Fgf21 was shown to be an effector of iWAT activation in UCP1KO mice (Keipert *et al.* 2020); perhaps Fgf21 is also an effector of dermal adipocyte hypertrophy. We note that the dWAT depot, alongside other fat depots, becomes depleted when mice are challenged with an extreme cold stress (transfer to 4°C) (Zhang *et al.* 2019), which could compound their energetic stress.

This study has shown little correlation between SAF thickness in human subjects and obesity; indeed the thickness of this layer is highly individual-specific. The marginal increase in thickness between lean and overweight/obese individuals could be accounted for by the 20% increase in SAF adipocyte fat globule size (Table 1). We speculate that both environmental and genetic factors may control the thickness, with an impact on the subsequent energy budget. Specific genes are known to regulate the pattern of human fat deposition, notably, for example, lower body subcutaneous depots (such as gluteofemoral) and peritoneal visceral fat (Loh *et al.* 2015; Lu *et al.* 2016); there may be similar genes that regulate the thickness of SAF.

We also note that high variation in SAF volume may lead to a correspondingly high variation in the total heat-generating capacity of this depot. Heat production

in mammalian bodies is a zero-sum reaction, the demand for heat is fulfilled by various means, whether muscle exercise, activation of BAT or activation of non-canonical futile cycles in beige fat, muscle or heart. This leads us to speculate that our observations could go some way to explaining the paradoxical differences observed for individual lean subjects in their capacity for BAT activation (van Marken Lichtenbelt *et al.* 2009); BAT activation for any given individual may be inversely related to the total amount of skin-associated adipose tissue. It would also explain why obese subjects show little BAT activation, since these individuals typically have a large volume of subcutaneous fat pads.

We considered the possibility that the lipolytic hub, HSL, was not present to be activated in dWAT in response to β -adrenergic activation; however, activation of this enzyme has been observed during the lipid depletion associated with dWAT involution during catagen (Rivera-Gonzalez *et al.* 2016). The lack of HSL activation in dWAT stands in contrast to human SAF, where it is known that lipolysis is dependent upon HSL (Langin *et al.* 2005). Indeed, human scWAT has been shown to become lipolytic during the cool season (Kern *et al.* 2014; Finlin *et al.* 2017, 2018), when mast cells are implicated as the lipolytic initiator. Only temporary but repeated application of an ice pack to the thigh (30 min for 10 days) was enough to induce systemic activation of the β -adrenergic response. We would expect sustained lipolysis to deplete the SAF layer (Fig. 7B); it remains to be tested whether intermittent cool exposure increases or




Figure 8. Summary of inferred functionality of skin-associated adipose depots for human and mouse

In human (left-hand side), the skin-associated fat (SAF) depot is responsive to β -adrenergic demand and contributes heat towards body temperature homeostasis. For mouse (right-hand side), there is a bilaminar adipose depot, the superficial dWAT layer is non-responsive to β -adrenergic demand (increasing insulative properties), whereas the subjacent iWAT depots are responsive. Immunofluorescent stained sections (both human and mouse) are stained for endothelial cells (CD31), perilipin (PL) and DAPI (nuc; nuclei); scale bars = 100 μ m. [Colour figure can be viewed at wileyonlinelibrary.com]

decreases the SAF layer or affects an individual's capacity for non-canonical heat generation by subcutaneous fat activation.

Mouse dWAT is highly dynamic compared with other fat depots (Alexander *et al.* 2015); however, the time required for dWAT thickening in response to environmental cues is still in the order of days. We propose that dWAT thickening is a chronic adaptive response that mitigates heat loss when thermogenic demand is high. Acclimation to cold exposure reflects adaptive changes to heat production strategies; these are important, since they impact glucose homeostasis and insulin sensitivity (van der Lans *et al.* 2013; Hanssen *et al.* 2016; Yoneshiro *et al.* 2016). To narrow down the degree of exposure required to cue the thickening of dWAT in response to cooler environmental temperatures, we exposed mice to room temperature for 12 h out of each day, either during the waking period (night-time) or sleep period (daytime). Interestingly, we found that the temperature of the mice during their sleep phase cued the development of thicker, cold-adapted dWAT and we surmise that there is a component of circadian control of this process.

Note that this cycling protocol was applied to singly housed mice; the typical sleeping temperature for group-housed mice is predicted to be warm, since mice build nests and huddle together to sleep. We conclude that our warm-sleep cycling protocol may be mimicked by group-housed mice. This implicates behaviour as a regulator of dWAT thickness. The cycle of warm-sleeping and cool-waking time was designed to mimic a typical human circadian environmental temperature exposure, with the goal of identifying whether there are dominant temperature exposures that train the thermogenic response of each individual mammal. Our data to date suggest that chronic adaptations are designed to minimize the total thermogenic load. Therefore, intermittent exposure to cool temperatures will likely have more benefit than continuous chronic exposures, when homeostatic mechanisms are activated.

Scherer and colleagues defined aspects of mouse dermal adipocytes that make them unique, such as micro-environment, transcriptome and regulation. This group found that these adipocytes express distinguishing gene markers such as cathelicidin (CAMP1), collagen5 and CCL4 (Zhang *et al.* 2019). We have shown that the most distinctive property of dWAT, that distinguishes it from its nearest neighbour, iWAT, is its ability to resist depletion in response to β -adrenergic activation (Fig. 5; Table 1).

Despite the distinct properties of human SAF and mouse dWAT, there is a functional similarity between the heat production strategies of mouse and human skin-associated depots. Thus, directly underneath the dWAT heat transfer barrier lie 10 subcutaneous fat pads (mammary glands or male equivalent; Fig. 8) spread across the peritoneum, where the approximate surface

area equals 10 cm². These are highly responsive to β -adrenergic activation (including heterogeneous UCP1 induction), and we conclude that together, the dWAT and iWAT combination are functionally homologous to human SAF.

References

- Alexander CM, Kasza I, Yen CL, Reeder SB, Hernando D, Gallo RL, Jahoda CA, Horsley V & MacDougald OA (2015). Dermal white adipose tissue: a new component of the thermogenic response. *J Lipid Res* **56**, 2061–2069.
- Bartelt A & Heeren J (2014). Adipose tissue browning and metabolic health. *Nat Rev Endocrinol* **10**, 24–36.
- Berry DC, Jiang Y, Arpke RW, Close EL, Uchida A, Reading D, Berglund ED, Kyba M & Graff JM (2017). Cellular aging contributes to failure of cold-induced beige adipocyte formation in old mice and humans. *Cell Metab* **25**, 166–181.
- Cannon B & Nedergaard J (2009). Thermogenesis challenges the adipostat hypothesis for body-weight control. *Proc Nutr Soc* **68**, 401–407.
- Cappellano G, Morandi EM, Rainer J, Grubwieser P, Heinz K, Wolfram D, Bernhard D, Lobenwein S, Pierer G & Ploner C (2018). Human macrophages preferentially infiltrate the superficial adipose tissue. *Int J Mol Sci* **19**, 1404.
- Chen KY, Cypess AM, Laughlin MR, Haft CR, Hu HH, Bredella MA, Enerback S, Kinahan PE, Lichtenbelt W, Lin FI, Sunderland JJ, Virtanen KA & Wahl RL (2016). Brown adipose reporting criteria in imaging Studies (BARCIST 1.0): Recommendations for standardized FDG-PET/CT experiments in humans. *Cell Metab* **24**, 210–222.
- Chondronikola M & Sidossis LS (2019). Brown and beige fat: From molecules to physiology. *Biochim Biophys Acta Mol Cell Biol Lipids* **1864**, 91–103.
- Chouchani ET, Kazak L & Spiegelman BM (2019). New advances in adaptive thermogenesis: UCP1 and beyond. *Cell Metab* **29**, 27–37.
- Driskell RR, Jahoda CA, Chuong CM, Watt FM & Horsley V (2014). Defining dermal adipose tissue. *Exp Dermatol* **23**, 629–631.
- Elattar S & Satyanarayana A (2015). Can brown fat win the battle against white fat?. *J Cell Physiol* **230**, 2311–2317.
- Enevoldsen LH, Simonsen L, Stallknecht B, Galbo H & Bulow J (2001). In vivo human lipolytic activity in preperitoneal and subdivisions of subcutaneous abdominal adipose tissue. *Am J Physiol Endocrinol Metab* **281**, E1110–E1114.
- Fabbiano S, Suarez-Zamorano N, Rigo D, Veyrat-Durebex C, Stevanovic Dokic A, Colin DJ & Trajkovski M (2016). Caloric restriction leads to browning of white adipose tissue through type 2 immune signaling. *Cell Metab* **24**, 434–446.
- Finlin BS, Confides AL, Zhu B, Boulanger MC, Memetimin H, Taylor KW, Johnson ZR, Westgate PM, Dupont-Versteegden EE & Kern PA (2019). Adipose tissue mast cells promote human adipose beiging in response to cold. *Sci Rep* **9**, 8658.
- Finlin BS, Memetimin H, Confides AL, Kasza I, Zhu B, Vekaria HJ, Harfmann B, Jones KA, Johnson ZR, Westgate PM, Alexander CM, Sullivan PG, Dupont-Versteegden EE & Kern PA (2018). Human adipose beiging in response to cold and mirabegron. *JCI Insight* **3**.

- Finlin BS, Zhu B, Confides AL, Westgate PM, Harfmann BD, Dupont-Versteegden EE & Kern PA (2017). Mast cells promote seasonal white adipose being in humans. *Diabetes* **66**, 1237–1246.
- Fruhbeck G, Mendez-Gimenez L, Fernandez-Formoso JA, Fernandez S & Rodriguez A (2014). Regulation of adipocyte lipolysis. *Nutr Res Rev* **27**, 63–93.
- Giles DA, Moreno-Fernandez ME, Stankiewicz TE, Graspentner S, Cappelletti M, Wu D, Mukherjee R, Chan CC, Lawson MJ, Klarquist J, Sunderhauf A, Softic S, Kahn CR, Stemmer K, Iwakura Y, Aronow BJ, Karns R, Steinbrecher KA, Karp CL, Sheridan R, Shanmukhappa SK, Reynaud D, Haslam DB, Sina C, Rupp J, Hogan SP & Divanovic S (2017). Thermoneutral housing exacerbates nonalcoholic fatty liver disease in mice and allows for sex-independent disease modeling. *Nat Med* **23**, 829–838.
- Grove KL, Fried SK, Greenberg AS, Xiao XQ & Clegg DJ (2010). A microarray analysis of sexual dimorphism of adipose tissues in high-fat-diet-induced obese mice. *Int J Obes* **34**, 989–1000.
- Hanssen MJ, van der Lans AA, Brans B, Hoeks J, Jardon KM, Schaart G, Mottaghy FM, Schrauwen P & van Marken Lichtenbelt WD (2016). Short-term cold acclimation recruits brown adipose tissue in obese humans. *Diabetes* **65**, 1179–1189.
- Harms M & Seale P (2013). Brown and beige fat: development, function and therapeutic potential. *Nat Med* **19**, 1252–1263.
- Ikeda K, Kang Q, Yoneshiro T, Camporez JP, Maki H, Homma M, Shinoda K, Chen Y, Lu X, Maretich P, Tajima K, Ajuwon KM, Soga T & Kajimura S (2017). UCP1-independent signaling involving SERCA2b-mediated calcium cycling regulates beige fat thermogenesis and systemic glucose homeostasis. *Nat Med* **23**, 1454–1465.
- Kajimura S, Spiegelman BM & Seale P (2015). Brown and beige fat: physiological roles beyond heat generation. *Cell Metab* **22**, 546–559.
- Karastergiou K & Fried SK (2013). Multiple adipose depots increase cardiovascular risk via local and systemic effects. *Curr Atheroscler Rep* **15**, 361.
- Karastergiou K & Fried SK (2017). Cellular mechanisms driving sex differences in adipose tissue biology and body shape in humans and mouse models. *Adv Exp Med Biol* **1043**, 29–51.
- Karastergiou K, Smith SR, Greenberg AS & Fried SK (2012). Sex differences in human adipose tissues - the biology of pear shape. *Biol Sex Differ* **3**, 13.
- Kasza I, Adler D, Nelson DW, Eric Yen CL, Dumas S, Ntambi JM, MacDougald OA, Hernando D, Porter WP, Best FA & Alexander CM (2019). Evaporative cooling provides a major metabolic energy sink. *Mol Metab* **27**, 47–61.
- Kasza I, Hernando D, Roldan-Alzate A, Alexander CM & Reeder SB (2016). Thermogenic profiling using magnetic resonance imaging of dermal and other adipose tissues. *JCI Insight* **1**, e87146.
- Kasza I, Suh Y, Wollny D, Clark RJ, Roopra A, Colman RJ, MacDougald OA, Shedd TA, Nelson DW, Yen MI, Yen CL & Alexander CM (2014). Syndecan-1 is required to maintain intradermal fat and prevent cold stress. *PLoS Genet* **10**, e1004514.
- Kazak L, Rahbani JF, Samborska B, Lu GZ, Jedrychowski MP, Lajoie M, Zhang S, Ramsay LC, Dou FY, Tenen D, Chouchani ET, Dzeja P, Watson IR, Tsai L, Rosen ED & Spiegelman BM (2019). Ablation of adipocyte creatine transport impairs thermogenesis and causes diet-induced obesity. *Nat Metab* **1**, 360–370.
- Keipert S & Jastroch M (2014). Brite/beige fat and UCP1 - is it thermogenesis?. *Biochim Biophys Acta* **1837**, 1075–1082.
- Keipert S, Lutter D, Schroeder BO, Brandt D, Stahlman M, Schwarzmayr T, Graf E, Fuchs H, de Angelis MH, Tschop MH, Rozman J & Jastroch M (2020). Endogenous FGF21-signaling controls paradoxical obesity resistance of UCP1-deficient mice. *Nat Commun* **11**, 624.
- Kelley DE, Thaete FL, Troost F, Huwe T & Goodpaster BH (2000). Subdivisions of subcutaneous abdominal adipose tissue and insulin resistance. *Am J Physiol Endocrinol Metab* **278**, E941–E948.
- Kern PA, Finlin BS, Zhu B, Rasouli N, McGehee RE, Jr., Westgate PM & Dupont-Versteegden EE (2014). The effects of temperature and seasons on subcutaneous white adipose tissue in humans: evidence for thermogenic gene induction. *J Clin Endocrinol Metab* **99**, E2772–E2779.
- Kershaw EE & Flier JS (2004). Adipose tissue as an endocrine organ. *J Clin Endocrinol Metab* **89**, 2548–2556.
- Kruglikov IL & Scherer PE (2016a). Dermal adipocytes and hair cycling: is spatial heterogeneity a characteristic feature of the dermal adipose tissue depot?. *Exp Dermatol* **25**, 258–262.
- Kruglikov IL & Scherer PE (2016b). Dermal adipocytes: from irrelevance to metabolic targets?. *Trends Endocrinol Metab* **27**, 1–10.
- Kruse V, Neess D & Faergeman NJ (2017). The significance of epidermal lipid metabolism in whole-body physiology. *Trends Endocrinol Metab* **28**, 669–683.
- Kuhn JP, Meffert P, Heske C, Kromrey ML, Schmidt CO, Mensel B, Volzke H, Lerch MM, Hernando D, Mayerle J & Reeder SB (2017). Prevalence of fatty liver disease and hepatic iron overload in a northeastern german population by using quantitative MR imaging. *Radiology* **284**, 706–716.
- Langin D, Dicker A, Tavernier G, Hoffstedt J, Mairal A, Ryden M, Arner E, Sicard A, Jenkins CM, Viguerie N, van Harmelen V, Gross RW, Holm C & Arner P (2005). Adipocyte lipases and defect of lipolysis in human obesity. *Diabetes* **54**, 3190–3197.
- Lee MJ, Wu Y & Fried SK (2013). Adipose tissue heterogeneity: implication of depot differences in adipose tissue for obesity complications. *Mol Aspects Med* **34**, 1–11.
- Liu D, Bordicchia M, Zhang C, Fang H, Wei W, Li JL, Guilherme A, Guntur K, Czech MP & Collins S (2016). Activation of mTORC1 is essential for beta-adrenergic stimulation of adipose browning. *J Clin Invest* **126**, 1704–1716.
- Loh NY, Neville MJ, Marinou K, Hardcastle SA, Fielding BA, Duncan EL, McCarthy MI, Tobias JH, Gregson CL, Karpe F & Christodoulides C (2015). LRP5 regulates human body fat distribution by modulating adipose progenitor biology in a dose- and depot-specific fashion. *Cell Metab* **21**, 262–273.

- Lu Y, Day FR, Gustafsson S, Buchkovich ML, Na J, Bataille V, Cousminer DL, Dastani Z, Drong AW, Esko T, Evans DM, Falchi M, Feitosa MF, Ferreira T, Hedman AK, Haring R, Hysi PG, Iles MM, Justice AE, Kanoni S, Lagou V, Li R, Li X, Locke A, Lu C, Magi R, Perry JR, Pers TH, Qi Q, Sanna M, Schmidt EM, Scott WR, Shungin D, Teumer A, Vinkhuyzen AA, Walker RW, Westra HJ, Zhang M, Zhang W, Zhao JH, Zhu Z, Afzal U, Ahluwalia TS, Bakker SJ, Bellis C, Bonnefond A, Borodulin K, Buchman AS, Cederholm T, Choh AC, Choi HJ, Curran JE, de Groot LC, De Jager PL, Dhonukshe-Rutten RA, Enneman AW, Eury E, Evans DS, Forsen T, Friedrich N, Fumeron F, Garcia ME, Gartner S, Han BG, Havulinna AS, Hayward C, Hernandez D, Hillege H, Ittermann T, Kent JW, Kolcic I, Laatikainen T, Lahti J, Mateo Leach I, Lee CG, Lee JY, Liu T, Liu Y, Lobbens S, Loh M, Lyttikainen LP, Medina-Gomez C, Michaelsson K, Nalls MA, Nielson CM, Oozageer L, Pascoe L, Paternoster L, Polasek O, Ripatti S, Sarzynski MA, Shin CS, Narancic NS, Spira D, Srikanth P, Steinhagen-Thiessen E, Sung YJ, Swart KM, Taittonen L, Tanaka T, Tikkanen E, van der Velde N, van Schoor NM, Verweij N, Wright AF, Yu L, Zmuda JM, Eklund N, Forrester T, Grarup N, Jackson AU, Kristiansson K, Kuulasmaa T, Kuusisto J, Lichtner P, Luan J, Mahajan A, Mannisto S, Palmer CD, Ried JS, Scott RA, Stancakova A, Wagner PJ, Demirkan A, Doring A, Gudnason V, Kiel DP, Kuhnel B, Mangino M, McKnight B, Menni C, O'Connell JR, Oostra BA, Shuldiner AR, Song K, Vandenput L, van Duijn CM, Vollenweider P, White CC, Boehnke M, Boettcher Y, Cooper RS, Forouhi NG, Gieger C, Grallert H, Hingorani A, Jorgensen T, Jousilahti P, Kivimaki M, Kumari M, Laakso M, Langenberg C, Linneberg A, Luke A, McKenzie CA, Palotie A, Pedersen O, Peters A, Strauch K, Tayo BO, Wareham NJ, Bennett DA, Bertram L, Blangero J, Bluher M, Bouchard C, Campbell H, Cho NH, Cummings SR, Czerwinski SA, Demuth I, Eckardt R, Eriksson JG, Ferrucci L, Franco OH, Froguel P, Gansevoort RT, Hansen T, Harris TB, Hastie N, Heliovaara M, Hofman A, Jordan JM, Jula A, Kahonen M, Kajantie E, Knekt PB, Koskinen S, Kovacs P, Lehtimaki T, Lind L, Liu Y, Orwoll ES, Osmond C, Perola M, Perusse L, Raitakari OT, Rankinen T, Rao DC, Rice TK, Rivadeneira F, Rudan I, Salomaa V, Sorensen TI, Stumvoll M, Tonjes A, Towne B, Tranah GJ, Tremblay A, Uitterlinden AG, van der Harst P, Vartiainen E, Viikari JS, Vitart V, Vohl MC, Volzke H, Walker M, Wallaschofski H, Wild S, Wilson JF, Yengo L, Bishop DT, Borecki IB, Chambers JC, Cupples LA, Dehghan A, Deloukas P, Fatemifar G, Fox C, Furey TS, Franke L, Han J, Hunter DJ, Karjalainen J, Karpe F, Kaplan RC, Kooner JS, McCarthy MI, Murabito JM, Morris AP, Bishop JA, North KE, Ohlsson C, Ong KK, Prokopenko I, Richards JB, Schadt EE, Spector TD, Widen E, Willer CJ, Yang J, Ingelsson E, Mohlke KL, Hirschhorn JN, Perspilik JA, Zillikens MC, Lindgren C, Kilpelainen TO & Loos RJ (2016). New loci for body fat percentage reveal link between adiposity and cardiometabolic disease risk. *Nat Commun* **7**, 10495.
- Manolopoulos KN, Karpe F & Frayn KN (2010). Gluteofemoral body fat as a determinant of metabolic health. *Int J Obes* **34**, 949–959.
- Mills EL, Pierce KA, Jedrychowski MP, Garrity R, Winther S, Vidoni S, Yoneshiro T, Spinelli JB, Lu GZ, Kazak L, Banks AS, Haigis MC, Kajimura S, Murphy MP, Gygi SP, Clish CB & Chouchani ET (2018). Accumulation of succinate controls activation of adipose tissue thermogenesis. *Nature* **560**, 102–106.
- Mottillo EP, Balasubramanian P, Lee YH, Weng C, Kershaw EE & Granneman JG (2014). Coupling of lipolysis and de novo lipogenesis in brown, beige, and white adipose tissues during chronic beta3-adrenergic receptor activation. *J Lipid Res* **55**, 2276–2286.
- Neess D, Bek S, Bloksgaard M, Marcher AB, Faergeman NJ & Mandrup S (2013). Delayed hepatic adaptation to weaning in ACBP-/- mice is caused by disruption of the epidermal barrier. *Cell Rep* **5**, 1403–1412.
- Neess D, Kruse V, Marcher AB, Waede MR, Vistisen J, Moller PM, Petersen R, Brewer JR, Ma T, Colleluori G, Severi I, Cinti S, Gerhart-Hines Z, Mandrup S & Faergeman NJ (2020). Epidermal Acyl-CoA-binding protein is indispensable for systemic energy homeostasis. *Mol Metab* **44**, 101144.
- Nicu C, Pople J, Bonsell L, Bhogal R, Ansell DM & Paus R (2018). A guide to studying human dermal adipocytes in situ. *Exp Dermatol* **27**, 589–602.
- Ogasawara J, Izawa T, Sakurai T, Shirato K, Ishibashi Y, Ishida H, Ohno H & Kizaki T (2015). The molecular mechanism underlying continuous exercise training-induced adaptive changes of lipolysis in white adipose cells. *J Obes* **2015**, 473430.
- Perez-Chirinos Buxade C, Sola-Perez T, Castizo-Olier J, Carrasco-Marginet M, Roy A, Marfell-Jones M & Iruiria A (2018). Assessing subcutaneous adipose tissue by simple and portable field instruments: skinfolds versus a-mode ultrasound measurements. *PLoS One* **13**, e0205226.
- Pinnick KE, Nicholson G, Manolopoulos KN, McQuaid SE, Valet P, Frayn KN, Denton N, Min JL, Zondervan KT, Fleckner J, Mol PC, McCarthy MI, Holmes CC & Karpe F (2014). Distinct developmental profile of lower-body adipose tissue defines resistance against obesity-associated metabolic complications. *Diabetes* **63**, 3785–3797.
- Qiao G, Chen M, Bucsek MJ, Repasky EA & Hylander BL (2018). Adrenergic signaling: a targetable checkpoint limiting development of the antitumor immune response. *Front Immunol* **9**, 164.
- Rivera-Gonzalez GC, Shook BA, Andrae J, Holtrup B, Bollag K, Betsholtz C, Rodeheffer MS & Horsley V (2016). Skin adipocyte stem cell self-renewal is regulated by a PDGFA/AKT-signaling axis. *Cell Stem Cell* **19**, 738–751.
- Sampath H, Flowers MT, Liu X, Paton CM, Sullivan R, Chu K, Zhao M & Ntambi JM (2009). Skin-specific deletion of stearoyl-CoA desaturase-1 alters skin lipid composition and protects mice from high fat diet-induced obesity. *J Biol Chem* **284**, 19961–19973.
- Sampath H & Ntambi JM (2014). Role of stearoyl-CoA desaturase-1 in skin integrity and whole body energy balance. *J Biol Chem* **289**, 2482–2488.
- Sbarbati A, Accorsi D, Benati D, Marchetti L, Orsini G, Rigotti G & Panetti P (2010). Subcutaneous adipose tissue classification. *Eur J Histochem* **54**, 48.

- Scheller EL, Khandaker S, Learman BS, Cawthorn WP, Anderson LM, Pham HA, Robles H, Wang Z, Li Z, Parlee SD, Simon BR, Mori H, Bree AJ, Craft CS & MacDougald OA (2019). Bone marrow adipocytes resist lipolysis and remodeling in response to beta-adrenergic stimulation. *Bone* **118**, 32–41.
- Shih MY, Kane MA, Zhou P, Yen CL, Streeper RS, Napoli JL & Farese RV, Jr (2009). Retinol esterification by DGAT1 is essential for retinoid homeostasis in murine skin. *J Biol Chem* **284**, 4292–4299.
- Smith GI, Mittendorfer B & Klein S (2019). Metabolically healthy obesity: facts and fantasies. *J Clin Invest* **129**, 3978–3989.
- Smith SR, Lovejoy JC, Greenway F, Ryan D, deJonge L, de la Bretonne J, Volavafova J & Bray GA (2001). Contributions of total body fat, abdominal subcutaneous adipose tissue compartments, and visceral adipose tissue to the metabolic complications of obesity. *Metabolism* **50**, 425–435.
- Speakman JR (2013). Measuring energy metabolism in the mouse - theoretical, practical, and analytical considerations. *Front Physiol* **4**, 34.
- Storchle P, Muller W, Sengeis M, Lackner S, Holasek S & Furrhapter-Rieger A (2018). Measurement of mean subcutaneous fat thickness: eight standardised ultrasound sites compared to 216 randomly selected sites. *Sci Rep* **8**, 16268.
- Tian XY, Ganeshan K, Hong C, Nguyen KD, Qiu Y, Kim J, Tangirala RK, Tonotonoz P & Chawla A (2016). Thermo-neutral housing accelerates metabolic inflammation to potentiate atherosclerosis but not insulin resistance. *Cell Metab* **23**, 165–178.
- Tschop MH, Speakman JR, Arch JR, Auwerx J, Bruning JC, Chan L, Eckel RH, Farese RV, Jr, Galgani JE, Hambly C, Herman MA, Horvath TL, Kahn BB, Kozma SC, Maratos-Flier E, Muller TD, Munzberg H, Pfluger PT, Plum L, Reitman ML, Rahmouni K, Shulman GI, Thomas G, Kahn CR & Ravussin E (2012). A guide to analysis of mouse energy metabolism. *Nat Methods* **9**, 57–63.
- van der Lans AA, Hoeks J, Brans B, Vijgen GH, Visser MG, Vosselman MJ, Hansen J, Jorgensen JA, Wu J, Mottaghy FM, Schrauwen P & van Marken Lichtenbelt WD (2013). Cold acclimation recruits human brown fat and increases nonshivering thermogenesis. *J Clin Invest* **123**, 3395–3403.
- van Marken Lichtenbelt WD, Vanhommerig JW, Smulders NM, Drossaerts JM, Kemerink GJ, Bouvy ND, Schrauwen P & Teule GJ (2009). Cold-activated brown adipose tissue in healthy men. *N Engl J Med* **360**, 1500–1508.
- Verboven K, Wouters K, Gaens K, Hansen D, Bijnen M, Wetzels S, Stehouwer CD, Goossens GH, Schalkwijk CG, Blaak EE & Jocken JW (2018). Abdominal subcutaneous and visceral adipocyte size, lipolysis and inflammation relate to insulin resistance in male obese humans. *Sci Rep* **8**, 4677.
- Villarroya F & Giral M (2015). The beneficial effects of brown fat transplantation: further evidence of an endocrine role of brown adipose tissue. *Endocrinology* **156**, 2368–2370.
- Volzke H, Alte D, Schmidt CO, Radke D, Lorbeer R, Friedrich N, Aumann N, Lau K, Piontek M, Born G, Havemann C, Ittermann T, Schipf S, Haring R, Baumeister SE, Wallaschofski H, Nauck M, Frick S, Arnold A, Junger M, Mayerle J, Kraft M, Lerch MM, Dorr M, Reffelmann T, Empen K, Felix SB, Obst A, Koch B, Glaser S, Ewert R, Fietze I, Penzel T, Doren M, Rathmann W, Haerting J, Hannemann M, Ropcke J, Schminke U, Jurgens C, Tost F, Rettig R, Kors JA, Ungerer S, Hegenscheid K, Kuhn JP, Kuhn J, Hosten N, Puls R, Henke J, Gloger O, Teumer A, Homuth G, Volker U, Schwahn C, Holtfreter B, Polzer I, Kohlmann T, Grabe HJ, Roszkopf D, Kroemer HK, Kocher T, Biffar R, John U & Hoffmann W (2011). Cohort profile: the study of health in Pomerania. *Int J Epidemiol* **40**, 294–307.
- Wajchenberg BL (2000). Subcutaneous and visceral adipose tissue: their relation to the metabolic syndrome. *Endocr Rev* **21**, 697–738.
- Walker GE, Marzullo P, Ricotti R, Bona G & Prodam F (2014). The pathophysiology of abdominal adipose tissue depots in health and disease. *Horm Mol Biol Clin Invest* **19**, 57–74.
- Wang GX, Zhao XY & Lin JD (2015). The brown fat secretome: metabolic functions beyond thermogenesis. *Trends Endocrinol Metab* **26**, 231–237.
- Wu J, Bostrom P, Sparks LM, Ye L, Choi JH, Giang AH, Khandekar M, Virtanen KA, Nuutila P, Schaart G, Huang K, Tu H, van Marken Lichtenbelt WD, Hoeks J, Enerback S, Schrauwen P & Spiegelman BM (2012). Beige adipocytes are a distinct type of thermogenic fat cell in mouse and human. *Cell* **150**, 366–376.
- Yoneshiro T, Aita S, Matsushita M, Kayahara T, Kameya T, Kawai Y, Iwanaga T & Saito M (2013). Recruited brown adipose tissue as an antiobesity agent in humans. *J Clin Invest* **123**, 3404–3408.
- Yoneshiro T, Matsushita M, Nakae S, Kameya T, Sugie H, Tanaka S & Saito M (2016). Brown adipose tissue is involved in the seasonal variation of cold-induced thermogenesis in humans. *Am J Physiol Regul Integr Comp Physiol* **310**, R999–R1009, ajpregu 00057 02015.
- Zhang Z, Shao M, Hepler C, Zi Z, Zhao S, An YA, Zhu Y, Ghaben AL, Wang MY, Li N, Onodera T, Joffin N, Crewe C, Zhu Q, Vishvanath L, Kumar A, Xing C, Wang QA, Gautron L, Deng Y, Gordillo R, Kruglikov I, Kusminski CM, Gupta RK & Scherer PE (2019). Dermal adipose tissue has high plasticity and undergoes reversible dedifferentiation in mice. *J Clin Invest* **129**, 5327–5342.
- Zwick RK, Guerrero-Juarez CF, Horsley V & Plikus MV (2018). Anatomical, physiological, and functional diversity of adipose tissue. *Cell Metab* **27**, 68–83.

Additional information

Data availability statement

This article is published at BioRxiv, including supplemental data confirming the accuracy of reagents used in this study: <https://www.biorxiv.org/content/10.1101/2020.09.16.300533v2>.

Full datasets are described in the Statistical Summary.

Competing interests

The University of Greifswald is a member of the 'Centre of Knowledge Interchange' programme of Siemens AG. Contrast-enhanced MRI research is part of the entire whole-body MRI study and was supported by Bayer Healthcare. The content of the manuscript is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. This work does not represent the views of the Department of Veterans Affairs or the United States Government.

Author contributions

C.M.A.: Development of strategy, execution of experimental procedures, consultation with analysis, manuscript preparation; I.K.: Design and execution of experimental procedures, data analysis and presentation, manuscript preparation; J.K., H.Z., R.B., D.H.: Design and execution of MRI imaging procedures, data analysis and presentation; A.G., Y.Z., J.S., P.K.: Collection of human skin and fat samples; C.-L.E.Y., D.N., N.R., D.L.: Development of strategy, consultation on mouse metabolic studies, supply of samples from mice on high-fat feeding protocols; O.M.: Development of strategy, manuscript preparation. All authors approved the final version of this manuscript and agree to be accountable for the accuracy and integrity of the work.

Funding

This work was supported by RO1GM113142 (C.M.A., I.K.); a pilot award from the University of Wisconsin Skin Disease Research Center (SDRC) NIAMS P30 AR066524 (I.K.); the University of Wisconsin Carbone Comprehensive Cancer Center (UWCCC) for use of its Shared Services (NIH/NCI P30 CA014520), including the Translational Initiatives in Pathology (TrIP) lab and Translational Science Biocore (TSB); R01 DK112282 and UL1 TR001998 (P.K.); R24DK092759, R01DK121759 and R01DK125513 (O.A.M.), UW Institute on Aging, NIA T32 AG000213 (N.E.R.), NIH/National Institute on Aging, AG056771, AG062328, and AG061635 (D.W.L.), the U.S.

Department of Veterans Affairs, I01-BX004031 (D.W.L.) and the facilities and resources from the William S. Middleton Memorial Veterans Hospital. SHIP is part of the Community Medicine Research Net of the University of Greifswald, Germany, which is funded by the Federal Ministry of Education and Research (01ZZ9603, 01ZZ0103, 01ZZ0403, 01ZZ0701, 03ZIK012), the Ministry of Cultural Affairs as well as the Social Ministry of the Federal State of Mecklenburg-West Pomerania. Whole-body MR imaging was supported by a joint grant from Siemens Healthcare, Erlangen, Germany and the Federal State of Mecklenburg-West Pomerania.

Acknowledgements

We appreciate expert technical assistance from Edgar Ocotl, the University of Wisconsin Translational Research Initiatives in Pathology (TRIP) Laboratory, and the Biobank, supported by the UW Department of Pathology and Laboratory Medicine, UWCCC (P30 CA014520) and the Office of The Director – NIH (S10OD023526). We acknowledge the donation of Sprague–Dawley rat skins from Dr Kumar (Department of Comparative Biosciences, UW).

Keywords

β -adrenergic response, brown adipose tissue, dermal white adipose tissue, dWAT, heat production, lipolysis, obesity, scWAT, skin-associated fat, subcutaneous white adipose tissue, thermogenesis, UCP1

Supporting information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Statistical Summary Document

Data used for Fig. 1. Study of human adiposity and SAF

Data used for Fig. 2B. Study of dWAT accumulation during maturation of C57Bl6 male mice

Data used for Fig. 3. Study of mouse dWAT response to calorie restriction

Translational perspective

This study focuses on the skin-associated adipose depots for mouse and human. We hypothesize that these lipid layers are important to controlling heat loss, which in turn controls whole body energetics and metabolism. We tested whether the adipose depots associated with human and mouse skin had homologous functions. We showed that the body-wide skin-associated adipose depot in human subjects is uniformly responsive to heat production cues, becoming lipolytic in response to adrenergic agents *ex vivo*. Body-wide assay of fat using MRI of >500 subjects revealed this depot to be the single largest adipose depot in lean individuals (estimated at 10–24 kg for a lean woman). Of translational interest is that this depot is likely a direct source of heat production, thus an alternative to the better-studied brown adipose tissue. Since each process of heat production creates a unique endocrine systemic response, it will be important to understand the relative contribution and regulation of brown adipose versus skin-associated adipose during body temperature homeostasis. This body-wide depot is not typically visualized by PET imaging procedures, leaving a knowledge gap. Comparison with mouse skin-associated depots shows that there are functional similarities and differences with the corresponding depot in human, leading us to be able to predict which depot to manipulate to examine the role of skin-associated adipose in whole body energetics.